



**HAL**  
open science

## **CK2 is a key regulator of SLC4A2-mediated Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> – exchange in human airway epithelia**

Salam H. Ibrahim, Mark J. Turner, Vinciane Saint-Criq, James Garnett, Iram J. Haq, Malcolm Brodlie, Chris Ward, Christian Borgo, Mauro Salvi, Andrea Venerando, et al.

### ► To cite this version:

Salam H. Ibrahim, Mark J. Turner, Vinciane Saint-Criq, James Garnett, Iram J. Haq, et al.. CK2 is a key regulator of SLC4A2-mediated Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> – exchange in human airway epithelia. Pflügers Archiv European Journal of Physiology, 2017, 469 (9), pp.1073-1091. 10.1007/s00424-017-1981-3 . hal-02627314

**HAL Id: hal-02627314**

**<https://hal.inrae.fr/hal-02627314>**

Submitted on 26 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# CK2 is a key regulator of SLC4A2-mediated $\text{Cl}^-/\text{HCO}_3^-$ exchange in human airway epithelia

Salam H. Ibrahim<sup>1,2</sup> · Mark J. Turner<sup>3</sup> · Vinciane Saint-Criq<sup>1</sup> · James Garnett<sup>4,5</sup> · Iram J. Haq<sup>4</sup> · Malcolm Brodlie<sup>4</sup> · Chris Ward<sup>4</sup> · Christian Borgo<sup>6</sup> · Mauro Salvi<sup>6</sup> · Andrea Venerando<sup>6</sup> · Michael A. Gray<sup>1</sup>

Received: 19 December 2016 / Revised: 27 March 2017 / Accepted: 10 April 2017  
© The Author(s) 2017. This article is an open access publication

**Abstract** Transepithelial bicarbonate secretion by human airway submucosal glands and surface epithelial cells is crucial to maintain the pH-sensitive innate defence mechanisms of the lung. cAMP agonists stimulate  $\text{HCO}_3^-$  secretion via coordinated increases in basolateral  $\text{HCO}_3^-$  influx and accumulation, as well as CFTR-dependent  $\text{HCO}_3^-$  efflux at the luminal membrane of airway epithelial cells. Here, we investigated the regulation of a basolateral located, DIDS-sensitive,  $\text{Cl}^-/\text{HCO}_3^-$  exchanger, anion exchanger 2 (AE2; SLC4A2) which is postulated to act as an acid loader, and therefore potential regulator of  $\text{HCO}_3^-$  secretion, in human airway epithelial cells. Using intracellular pH measurements performed on Calu-3 cells, we demonstrate that the activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was significantly downregulated by cAMP agonists, via a PKA-independent mechanism and also required  $\text{Ca}^{2+}$  and calmodulin under resting conditions. AE2

contains potential phosphorylation sites by a calmodulin substrate, protein kinase CK2, and we demonstrated that AE2 activity was reduced in the presence of CK2 inhibition. Moreover, CK2 inhibition abolished the activity of AE2 in primary human nasal epithelia. Studies performed on mouse AE2 transfected into HEK-293T cells confirmed almost identical  $\text{Ca}^{2+}$ /calmodulin and CK2 regulation to that observed in Calu-3 and primary human nasal cells. Furthermore, mouse AE2 activity was reduced by genetic knockout of CK2, an effect which was rescued by exogenous CK2 expression. Together, these findings are the first to demonstrate that CK2 is a key regulator of  $\text{Cl}^-$ -dependent  $\text{HCO}_3^-$  export at the serosal membrane of human airway epithelial cells.

**Keywords** CK2 · SLC4A2 · Anion transport ·  $\text{HCO}_3^-$  · Airway epithelia

**Electronic supplementary material** The online version of this article (doi:10.1007/s00424-017-1981-3) contains supplementary material, which is available to authorized users.

✉ Michael A. Gray  
m.a.gray@ncl.ac.uk

- <sup>1</sup> Institute for Cell & Molecular Biosciences, The Medical School, Newcastle University, Framlington Place, Newcastle upon Tyne NE2 4HH, UK
- <sup>2</sup> Department of Basic Science, College of Veterinary Medicine, University of Sulaimani, Sulaimani City, Kurdistan Region, Iraq
- <sup>3</sup> Department of Physiology, McGill University, Montreal, Canada
- <sup>4</sup> Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, UK
- <sup>5</sup> Immunology & Respiratory Diseases Research, Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach an der Riss, Germany
- <sup>6</sup> Department of Biomedical Sciences, CNR Institute of Neuroscience, University of Padova, Padova, Italy

## Abbreviations

AE2 Anion exchanger 2  
ASL Airway surface liquid  
CK2 Casein kinase 2  
PKA Protein kinase A

## Introduction

The secretion of bicarbonate ( $\text{HCO}_3^-$ ) by epithelial cells is essential for maintaining the normal function of many epithelial tissues primarily due to its ability to act as a biological buffer and therefore a key regulator of extracellular pH [22]. In the human airways,  $\text{HCO}_3^-$  plays a major role in the innate defence of the lungs to inhaled pathogens. Transepithelial secretion of  $\text{HCO}_3^-$  and  $\text{Cl}^-$  from serous cells of the submucosal glands, as well as surface epithelial cells, drives isosmotic

water secretion, and this  $\text{HCO}_3^-$ -rich fluid is an important component of the periciliary layer (PCL) of the airway surface liquid (ASL) which lines the conducting airways [4]. The depth of the PCL is important for efficient mucociliary clearance [5, 62] while maintaining the correct ASL pH is vital for efficient bacterial killing by pH-dependent antimicrobials as well as reducing mucus viscosity [17, 44, 53]. In addition,  $\text{HCO}_3^-$  allows for efficient solubilization and transportation of airway mucus [47]. Therefore,  $\text{HCO}_3^-$  secretion enables pH-sensitive components of the innate defence mechanisms of the lung to function efficiently.

Transepithelial  $\text{HCO}_3^-$  secretion is a two-stage process that involves (i) import across the basolateral membrane and accumulation above electrochemical equilibrium and (ii) exit across the apical (serosal) membrane down its electrochemical gradient. It is well established that cAMP-dependent stimulation of the cystic fibrosis transmembrane conductance regulator (CFTR) is crucial for  $\text{HCO}_3^-$  exit across the luminal membrane in human airway epithelia [11, 14, 18, 29, 54, 56]. Loss of functional CFTR causes cystic fibrosis (CF), an autosomal recessive disorder characterized by the accumulation of thick, sticky mucus, bacterial colonization and chronic inflammation of the airways [34]. Impaired cAMP-stimulated  $\text{HCO}_3^-$  secretion has been reported in a wide range of CF tissues [6], and the ASL of CF patients is more acidic than that of healthy patients [10, 13] which is thought to contribute to the pathology of the disease. Although functional CFTR expression is clearly important for airway  $\text{HCO}_3^-$  secretion, members of the SLC26 family of  $\text{Cl}^-/\text{HCO}_3^-$  exchangers, SLC26A3, SLC26A4 and SLC26A6, also transport  $\text{HCO}_3^-$  [57, 61] and previous studies from our laboratory revealed that SLC26A4 (pendrin) participates in  $\text{HCO}_3^-$  secretion by Calu-3 cells [14]. We proposed that a functional interaction existed between the phosphorylated R domain of CFTR, and the STAS domain of pendrin as described by Ko et al. [25] which enabled CFTR to regulate  $\text{HCO}_3^-$  secretion via modulation of pendrin-mediated  $\text{Cl}^-/\text{HCO}_3^-$  exchange [14].

$\text{HCO}_3^-$  transport across the serosal membrane is believed to be governed by members of the SLC4 family of transporters, including importers such as the  $\text{Na}^+/\text{HCO}_3^-$  cotransporters SLC4A4, SLC4A5 and SLC4A7 [11, 26, 54] which act to accumulate  $\text{HCO}_3^-$  inside the cell. In contrast, SLC4A2, an electroneutral  $\text{Cl}^-/\text{HCO}_3^-$  anion exchanger (AE2) that exports  $\text{HCO}_3^-$ , plays an important role in the regulation of intracellular pH during cell alkalization, as well as in the control of cell volume by  $\text{Cl}^-$  uptake [2, 59]. Furthermore, it has been shown that AE2 is expressed in the airways [1, 12, 55] and immunostaining experiments performed on polarized Calu-3 cells have localized the expression of AE2 to the basolateral membrane [32]. Our laboratory [14, 15] and others [24] have recently demonstrated that a functional  $\text{Cl}^-/\text{HCO}_3^-$  exchanger is present at the basolateral surface of Calu-3 cells under resting conditions, with features

consistent with AE2. In addition, Huang et al. [18] showed that intracellular alkalization produced by basolateral  $\text{Cl}^-$  removal was decreased by 80% in SLC4A2 knockdown Calu-3 cells. We found that this exchanger was almost completely inhibited by elevations in intracellular cAMP [14, 15], which we proposed would enhance cAMP-stimulated transepithelial  $\text{HCO}_3^-$  secretion, by reducing  $\text{HCO}_3^-$  export across the basolateral membrane. However, Kim et al. [24] demonstrated that inhibition of CFTR maintained AE2 activity in cAMP-stimulated Calu-3 cells, while Huang et al. [18] suggested that AE2, acting in concert with  $\text{Na}^+/\text{HCO}_3^-$  cotransporters, was involved in basolateral  $\text{Cl}^-$  loading and  $\text{HCO}_3^-$  recycling that helped maintain transepithelial  $\text{Cl}^-$  secretion under cAMP-stimulated conditions in Calu-3 cells. Thus, the role of AE2 in airway  $\text{HCO}_3^-$  and fluid secretion is still controversial and not fully understood. Therefore, the major aim of the present study was to further understand the cellular pathways that regulate basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity in human airway epithelial cells to help provide a better understanding of its role in airways  $\text{HCO}_3^-$  secretion. Our results have uncovered a novel regulation of the  $\text{Cl}^-/\text{HCO}_3^-$  exchanger by  $\text{Ca}^{2+}$ /calmodulin signalling and by the master protein kinase, CK2 (casein kinase 2). Since CK2 has previously been implicated in regulating  $\text{HCO}_3^-$  transport by CFTR in secretory epithelia [65], our findings strongly suggest that CK2 is a key regulatory component of transepithelial  $\text{HCO}_3^-$  transport in the human airways.

## Methods

**Calu-3 cell culture** The human adenocarcinoma-derived cell line, Calu-3 (ATCC HTB-55), was grown in T<sub>75</sub> Costar cell culture flasks (75 cm<sup>2</sup>) in 30 ml of Eagle's Minimum Essential Medium (EMEM) supplemented with 10% foetal bovine serum (FBS), 100 U ml<sup>-1</sup> penicillin and 100 µg ml<sup>-1</sup> streptomycin, 1% non-essential amino acids, 2 mM L-Glutamine (Sigma) and incubated at 37 °C in a humidified air containing 5% (v/v) CO<sub>2</sub>. Cells were initially seeded at 3 × 10<sup>6</sup> cells per flask and passaged every 7 days using 0.05% trypsin and 0.02% ethylenediaminetetraacetic acid (EDTA) in Earle's balanced salt solution. For intracellular pH measurements, cells were seeded at 250,000 cells cm<sup>-2</sup> initial seeding density onto semi-permeable Transwell supports (0.4 µm pore, polyester membrane insert, 1.12 cm<sup>2</sup> surface area, (Corning, UK)). Calu-3 cells generally produced a polarized monolayer after 7 days. All experiments were carried out 8–14 days after seeding.

**Human embryonic kidney-293T (HEK-293T) cell culture** HEK-293T cells were grown in T<sub>75</sub> Costar cell culture flasks (75 cm<sup>2</sup>) with 30 ml of Dulbecco's Modified Eagle's Medium

(DMEM), supplemented as for Calu-3 cells and incubated at 37 °C in a humidified air containing 5% (v/v) CO<sub>2</sub>. Cells were initially seeded at 1 × 10<sup>6</sup> cells per flask and passaged every 7 days using 0.05% trypsin and 0.02% ethylenediaminetetraacetic acid (EDTA) in Earle's balanced salt solution. The knockout CK2 (both the alpha (α) and alpha prime (α prime) catalytic subunits) HEK-293T cells were generated by CRISPR/Cas9 gene editing. Genomic target sequences of guide RNA (gRNA) were 5'-TTACATGTATGAGA TTCTGA-3' (CK2 α) and 5'-GGGTCTACGCCGAG GTGAAC-3' (CK2 α prime). The absence of the catalytic subunits was confirmed by Western blotting (see supplementary Fig. 2). For transient transfection, HEK-293T cells were seeded onto 25 mm glass coverslips (VWR) at 100,000 cells per coverslip and then transiently transfected with cDNA coding for mAE2 or the various CK2 constructs 1 day post-seeding. Intracellular pH measurements were performed 48 h post-transfection.

### Isolation and culture of primary human nasal epithelia

Ethical approval for collection of paediatric human nasal epithelial cells was granted by the NRES Committee North East - Newcastle and North Tyneside 1 (REC reference: 15/NE/0215) and informed written parental consent was obtained. Human nasal epithelial cells were obtained by brushing the inferior nasal turbinate of each nostril from two donors with a cytology brush (Conmed, UK). Nasal epithelial cells were expanded in purified bovine collagen (PureCol; Advanced Biomatrix)-coated T<sub>25</sub> cell culture flasks (25cm<sup>2</sup>) containing bronchial epithelial growth medium supplemented with SingleQuots (Lonza, Basel, Switzerland), penicillin/streptomycin (100 Units ml<sup>-1</sup> penicillin and 0.1 mg ml<sup>-1</sup> streptomycin) and Primocin (InvivoGen, UK). For air liquid interface (ALI) cultures, cells were passaged at 80–90% confluency using 0.05% trypsin and 0.02% EDTA and seeded onto purified collagen coated (PureCol; Advanced Biomatrix) semi-permeable Snapwell supports (0.4 μm pore, polyester membrane insert, 1.12 cm<sup>2</sup> surface area; Corning, UK) and grown until confluent. The apical medium was subsequently removed and cell cultures were fed basolaterally on alternate days. Cultures were incubated at 37 °C in humidified air containing 5% CO<sub>2</sub>. ALI culture differentiation was characterized through the demonstration of mucus production, cilia formation and serial transepithelial resistance (TEER) measurements (STX2 electrodes, EVOM2™ Epithelial Voltohmmeter, World Precision Instruments) to assess tight junction integrity. Experiments were performed on polarized epithelia after 28 days of ALI culture.

**CK2 kinase activity assay** CK2 activity in cell lysates was measured by means of radioactive assays with [γ-33P]ATP towards the specific CK2 substrate peptide CK2-tide (RRRADSDDDDD). Briefly, both Calu-3 and HEK-293T

cells were treated for 5 min with 10 μM of the indicated compound and then lysed in HEPES 50 mM pH 8, NaCl 150 mM, Glycerol 1% (w/v), Triton X-100 1% (v/v), MgCl<sub>2</sub> 45 mM, EGTA 5 mM with a cocktail of anti-proteases and anti-phosphatases. Protein concentration was determined by the Bradford method. CK2-tide peptide (250 μM) were phosphorylated by incubation at 37 °C for 20 min in a 25-μl volume containing 50 mM Tris/HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 100 mM NaCl and 100 μM [γ-33P]ATP (specific radioactivity 1000–1500 c.p.m./pmol). The reaction was started by the addition of 1 or 2 μg of lysate and stopped by ice cooling and absorption on phosphocellulose P81 paper. Papers were washed three times with 75 mM phosphoric acid, dried and counted in a scintillation counter.

**Transfection of HEK-293T cells** Mouse AE2 cDNA was a kind gift from Beth Lee and Ron Kopito, and contained a human haemagglutinin (HA) tag; for more details, see Lee et al. [28]. Empty plasmid (pcDNA 3.1 myc/His), human WT-CK2 (CK2-alpha) and the double CK2 mutant (V66A and I174A; DM-CK2) cDNA were generated as previously described [50, 51]. DNA sequencing analysis confirmed sequence identity of all the constructs. To transfect cDNA constructs into HEK-293T cells, cDNA was pre-complexed with Lipofectamine-2000 (Thermo Fisher) at a ratio of 1:2.28, respectively. Opti-MEM media with GlutaMax (Thermo Fisher) was then added for 15 min at room temperature, and then diluted in culture media to produce a final concentration of 1 μg DNA ml<sup>-1</sup>. This complex media was added to the cells and incubated for 6 h at 37 °C before the complex media was removed and cells were incubated with Opti-MEM plus 10% FBS overnight before being returned back to normal culture media.

**Intracellular pH measurements** For measurements in polarized Calu-3 cells and well-differentiated human nasal epithelia, cells were loaded with 10 or 40 μM, respectively, of the pH sensitive dye, 2'-7'-bis (carboxyethyl)-5(6)-carboxyfluorescein acetoxymethyl ester (BCECF-AM) in Na-HEPES and incubated for 60 min at 37 °C. Cells were mounted on to the stage of a Nikon fluor inverted microscope and viewed at ×60 magnification using a long working distance objective (N.A 0.6). Cells were perfused with Krebs solution at 37 °C gassed with 5% (v/v) CO<sub>2</sub>/95% (v/v) O<sub>2</sub> at a rate of 3 ml min<sup>-1</sup> (apical) and 6 ml min<sup>-1</sup> (basolateral). For measurements in HEK-293T cells, cells were loaded with 10 μM BCECF-AM in Na-HEPES and incubated for 10 min at 37 °C. After dye loading, coverslips were placed in a perfusion chamber and then mounted onto the stage of a Nikon inverted microscope. Cells were viewed at ×40 magnification using an oil immersion objective (N.A 1.2) and perfused with Krebs solution at 37 °C gassed with 5% CO<sub>2</sub>/95% O<sub>2</sub> at a rate of 3 ml min<sup>-1</sup>. Intracellular pH (pH<sub>i</sub>) was measured using a

Life Sciences Microfluorimeter System in which cells were alternatively excited at 490 and 440 nm wavelengths every 1.024 s with emitted light collected at 510 nm. The ratio of 490 nm emission to 440 nm emission was recorded using the PhoCal 1.6b software and calibrated to  $\text{pH}_i$  using the high  $\text{K}^+$ /nigericin technique [16] in which cells were exposed to high  $\text{K}^+$  solutions containing 10  $\mu\text{M}$  nigericin, set to a desired pH, ranging from 6.6 to 8.4. Total buffering capacity ( $\beta_{\text{tot}}$ ) was calculated by addition of the intrinsic buffering capacity ( $\beta_i$ ) to the buffering capacity of the  $\text{CO}_2\text{-HCO}_3^-$  buffer system ( $\beta_{\text{HCO}_3^-}$ ) in which  $\beta_i$  was calculated using the  $\text{NH}_4^+$  technique as described by Roos, Boron [49]. For analysis of  $\text{pH}_i$  measurements, delta  $\text{pH}_i$  ( $\Delta\text{pH}_i$ ) was determined by calculating the mean  $\text{pH}_i$  over 60 s before, during and after treatment. Rate of  $\text{pH}_i$  change ( $\Delta\text{pH}_i/\Delta t$ ) was determined by performing a linear regression over a period of at least 30 s which was converted to a transmembrane  $\text{HCO}_3^-$  flux ( $-J(\text{B})$ ) by multiplying  $\Delta\text{pH}_i/\Delta t$  by  $\beta_{\text{tot}}$ .


### Confocal microscopy to detect expression of HA-tagged mAE2

**mAE2 Control**, untransfected and transfected HEK-293T cells expressing HA-tagged mAE2 were grown on glass coverslips for 2 days and fixed with 4% PFA for 10 min at room temperature. Cells were washed with PBS three times for 5 min, and then with 50 mM  $\text{NH}_4\text{Cl}$  to quench any remaining PFA. After washing, fixed cells were permeabilized using 1% Triton X-100 for 5 min at room temperature and washed in PBS three times for 5 min. To block non-specific binding, cells were incubated with blocking buffer, consisting of 5% goat serum and 1% Na-azide in PBS, at room temperature for 30 min. Blocking buffer was removed and cells were incubated in diluted primary antibody (Anti-HA16B12, 1/1000 in blocking buffer, Abcam) overnight at 4 °C on a shaker. Cells were then rinsed in PBS three times for 15 min to remove any unbound primary antibody, and then incubated with FITC-conjugated goat anti-mouse antibody (1/100 in blocking buffer) for 1 h at room temperature in the dark. Following this, cells were washed with PBS three times for 15 min to remove any unbound secondary antibody. DAPI dye (1  $\mu\text{g ml}^{-1}$ ) was added onto coverslips for 2 min, at room temperature, away from light, to stain the nucleus and gently washed in PBS to remove any remaining DAPI. Coverslips were mounted onto a microscope slide, using mounting medium (VectaShield). Cells were observed under Nikon A1R Confocal microscope at  $\times 60$  magnification (0.1 DIC lens) with a numerical aperture of 1.4. Cells were excited with the DAPI excitation wavelength of 405 nm to visualize DAPI-stained specimens, and imaged at the emission wavelength of 450 nm. To visualize FITC-stained specimens, cells were excited with the FITC-excitation wavelength of

495 nm and imaged at the emission wavelength of 517 nm.

**Solutions and reagents** All reagents and inhibitors were purchased from Sigma-Aldrich (Sigma-Aldrich Company Ltd., UK), apart from forskolin, BAPTA-AM and TBB (R & D Systems); BCECF-AM, DIDS, Lipofectamine 2000, WGA (Thermo Fisher), GlyH-101 and J-8 (Santa Cruz), RpcAMPs (Enzo life science), and Anti-HA antibody (Abcam). CX-4945 was produced by Gliaxx Laboratories (Gliaxx Laboratories, MA, USA). All stock solutions of agonists and inhibitors were made in DMSO, apart from nigericin (made in 100% ethanol), and 8CPT-2Me-cAMP, carbachol and adenosine (dissolved in deionized water). gRNAs designs and reagents for genome editing were supplied by Horizon Discovery ([www.horizondiscovery.com](http://www.horizondiscovery.com)). The  $\text{HCO}_3^-$  free Na-HEPES buffer solution consisted of (in mM) 130 NaCl, 5 KCl, 1  $\text{CaCl}_2$ , 1  $\text{MgCl}_2$ , 10 Na-HEPES and 10 D-Glucose. The  $\text{Cl}^-$ -free HEPES buffer solution consisted of (in mM) 130 Na-gluconate, 2.5  $\text{K}_2\text{SO}_4$ , 6 Ca-gluconate, 1 Mg-gluconate, 10 HEPES (free acid) and 10 D-Glucose. All HEPES-buffered solutions were calibrated to pH 7.4 by addition of 1 M HCl. High  $\text{Cl}^-$  Krebs solution consisted of (in mM) 25  $\text{NaHCO}_3^-$ , 115 NaCl, 5 KCl, 1  $\text{CaCl}_2$ , 1  $\text{MgCl}_2$  and 10 D-Glucose. In the  $\text{Ca}^{2+}$ -free high  $\text{Cl}^-$  Krebs solution, the NaCl concentration was increased to 116 mM, and  $\text{CaCl}_2$  was replaced with  $\text{MgCl}_2$  and 0.5 mM EGTA was added to chelate any remaining  $\text{Ca}^{2+}$ . The  $\text{Cl}^-$ -free Krebs solution consisted of (in mM) 25  $\text{NaHCO}_3^-$ , 115 Na-Gluconate, 2.5  $\text{K}_2\text{SO}_4$ , 6 Ca-gluconate, 1 Mg gluconate and 10 D-Glucose. In the  $\text{Ca}^{2+}$ -free  $\text{Cl}^-$ -free Krebs solution, the Na-gluconate concentration was increased to 124 mM, and 0.5 mM EGTA was added to chelate any remaining  $\text{Ca}^{2+}$ . The intracellular pH calibration solutions consisted of (in mM) 5 NaCl, 130 KCl, 1  $\text{CaCl}_2$ , 1  $\text{MgCl}_2$ , 10 D-Glucose, 10 HEPES (for solutions set at pH 7.6 or below) or 10 TRIS (for solutions set at pH 7.8 or above) as well as 10  $\mu\text{M}$  nigericin. Solutions were set to the desired pH by using either 1 M HCl or 1 M NaOH. The ammonium pulse solutions used to determine intracellular buffering capacity consisted of (in mM) 4.5 KCl, 1  $\text{MgCl}_2$ , 2  $\text{CaCl}_2$ , 5 BaCl, 10 HEPES, 10 D-Glucose as well as varying concentrations of  $\text{NH}_4\text{Cl}/\text{NMDG-Cl}$ , ranging from 0  $\text{NH}_4\text{Cl}/145$   $\text{NMDG-Cl}$  to 30  $\text{NH}_4\text{Cl}/115$   $\text{NMDG-Cl}$ . All solutions were titrated to pH 7.4 at 37 °C using 1 M CsOH.

**Statistical analysis** All results are presented as mean  $\pm$  S.E.M. where  $n$  is the number of experiments. The GraphPad Prism 4 software (GraphPad Software, USA) was used for statistical analysis and either a Student's  $t$  test (paired or unpaired), one-way ANOVA (with Tukey's multiple comparison post-test) or two-way ANOVA (with Bonferroni's post-test), where applicable.  $p$  values of  $<0.05$  were considered statistically significant.




**Fig. 1** Calu-3 cells express a DIDS-sensitive  $\text{Cl}^-$  and  $\text{HCO}_3^-$  dependent anion exchanger on the basolateral membrane: Calu-3 cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 1 h) and  $\text{pH}_i$  was determined by fluorescent microscopy. **a** Shows an example experiment in which basolateral  $\text{Cl}^-$  was removed and replaced by gluconate, basolateral  $\text{HCO}_3^-$  was removed and replaced by HEPES or both  $\text{Cl}^-$  and  $\text{HCO}_3^-$  were removed, as indicated. **b** Summarizes the mean change in  $\text{pH}_i$  resulting from anion replacement. Data represents mean  $\pm$  S.E.M.,  $n = 3-6$ . **c** Shows an example experiment in which the activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$

exchanger was assessed by measuring  $\text{pH}_i$  changes in response to replacement of basolateral  $\text{Cl}^-$  with gluconate in the presence of DIDS (0.01 or 500  $\mu\text{M}$ ). **d** Displays the percent inhibition of the mean change in  $\text{pH}_i$  caused by basolateral  $\text{Cl}^-$  removal and **e** the percent inhibition of the rate of reacidification upon  $\text{Cl}^-$  readdition at different DIDS concentrations. Note that the data points of DIDS concentration of  $10^{-8}$ ,  $10^{-7}$  and  $10^{-6}$  M are plotted but lie directly on the x axis. In each case, a non-linear regression was fit to the data. Data represents mean  $\pm$  S.E.M. ( $n = 3-8$ )

## Results

**Calu-3 cells express a basolateral DIDS-sensitive,  $\text{Cl}^-/\text{HCO}_3^-$  exchanger** Our laboratory [14, 15] and others [24] have previously reported that  $\text{Cl}^-/\text{HCO}_3^-$  exchange occurs across the basolateral membrane in non-stimulated Calu-3 cells. In support of these findings, intracellular pH measurements showed that removal of basolateral  $\text{Cl}^-$  caused an intracellular alkalinization of  $0.36 \pm 0.02$  units ( $n = 8$ ; Fig. 1A) and that readdition of  $\text{Cl}^-$  caused  $\text{pH}_i$  to reacidify at a rate of  $0.57 \pm 0.07$  pH units  $\text{min}^{-1}$  ( $n = 8$ ). Similarly, removal of basolateral  $\text{HCO}_3^-$ , and replacing it with HEPES, induced an intracellular acidification, indicating enhanced basolateral  $\text{HCO}_3^-$  efflux down its concentration gradient. However, there was no measurable pH change in response to removal

of basolateral  $\text{Cl}^-$  in the absence of basolateral  $\text{HCO}_3^-$  (Fig. 1A, B). Overall, these data are consistent with  $\text{Cl}^-/\text{HCO}_3^-$  exchange occurring across the basolateral membrane. The anion exchange inhibitor 4,4'-Diisothiocyano-2,2'-stilbenedisulfonic acid (DIDS) dose-dependently inhibited this basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity, with 500  $\mu\text{M}$  DIDS causing complete inhibition (Fig. 1C). Analysis of the DIDS dose-response curves for inhibition of the change in  $\text{pH}_i$  following basolateral  $\text{Cl}^-$  removal, and rate of reacidification following basolateral  $\text{Cl}^-$  reintroduction, gave a calculated  $\text{IC}_{50}$  value of  $16.5 \pm 1.3$   $\mu\text{M}$  and  $7.5 \pm 1.2$   $\mu\text{M}$ , respectively (Fig. 1D, E). Therefore, these data demonstrate that Calu-3 cells have a DIDS-sensitive, basolateral  $\text{Cl}^-/\text{HCO}_3^-$  anion exchanger with properties consistent with SLC4A2 (AE2), which support previous results from these cells [14, 15, 24].




**Fig. 2** The basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger is negatively regulated by cAMP: Calu-3 cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 1 h) and  $\text{pH}_i$  was determined by fluorescent microscopy. The activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was assessed by measuring  $\text{pH}_i$  changes in response to replacement of basolateral  $\text{Cl}^-$  with gluconate. **a** Shows a representative experiment in which the activity of the  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was assessed in the presence of forskolin (5  $\mu\text{M}$ ). **b** Summarizes the forskolin-stimulated  $\text{HCO}_3^-$  efflux in control cells or in conditions in which  $\text{HCO}_3^-$  was removed and replaced by HEPES, or in cells pretreated with the transmembrane adenylyl cyclase inhibitor SQ 22,536 (500  $\mu\text{M}$ ), the PKA inhibitor H-89 (50  $\mu\text{M}$ ) or the CFTR

inhibitor GlyH-101 (10  $\mu\text{M}$ ) \*Significant difference vs. control ( $p < 0.05$ ; \*\* =  $p < 0.01$ ; \*\*\* =  $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 3$ –22. **c** and **d** summarize the effect of forskolin (Fsk; 5  $\mu\text{M}$ ), adenosine (Ado; 10  $\mu\text{M}$ ), IBMX (1 mM), dibutyl-cAMP (db-cAMP; 800  $\mu\text{M}$ ), MK-571 (10  $\mu\text{M}$ ), Fsk in cells pretreated with RpcAMPs (1 mM; 1 h), Fsk in cells pretreated with H89 (50  $\mu\text{M}$ ; 1 h) and the EPAC agonist 8CPT-2Me-cAMP-AM (10  $\mu\text{M}$ ; 1 h) on the mean change in  $\text{pH}_i$  caused by basolateral  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*Significant effect of treatment vs. control ( $p < 0.05$ ; \*\* =  $p < 0.01$ ; \*\*\* =  $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 3$ –30

**The basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger is inhibited by elevations in cytosolic cAMP** The next series of experiments were designed to investigate the regulation of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger. We first sought to assess the effect of elevated cAMP on  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity, since cAMP is known to stimulate transepithelial  $\text{HCO}_3^-$  secretion from Calu-3 cells in a PKA-dependent manner [14]. As shown in Fig. 2A, addition of forskolin to Calu-3 cells induced an intracellular acidification of  $0.17 \pm 0.01$   $\text{pH}_i$  units ( $n = 22$ ), which gave a calculated  $\text{HCO}_3^-$  efflux of  $9.2 \pm 0.9$   $\text{mM min}^{-1}$  ( $n = 22$ ). This forskolin-stimulated acidification was absent when  $\text{HCO}_3^-$  was replaced by HEPES and was sensitive to the transmembrane adenylyl cyclase inhibitor SQ 22,536, the PKA inhibitor H-89 and the CFTR inhibitor GlyH-101 (Fig. 2B), indicating that this response to forskolin was a result of cAMP-stimulated, CFTR-dependent  $\text{HCO}_3^-$  secretion. Importantly, although forskolin promoted  $\text{HCO}_3^-$  efflux across the apical membrane, it appeared to markedly reduce  $\text{HCO}_3^-$  efflux across the basolateral membrane by the anion exchanger (Fig. 2). Forskolin reduced the mean change in  $\text{pH}_i$  following basolateral  $\text{Cl}^-$  removal by  $85.2 \pm 2.6\%$  compared

to non-stimulated cells ( $p < 0.001$  vs. control;  $n = 10$ ; Fig. 2C) and also reduced the rate of reacidification following basolateral  $\text{Cl}^-$  reintroduction by  $98.4 \pm 1.6\%$  compared to non-stimulated cells ( $p < 0.001$  vs. control;  $n = 10$ ; Fig. 2D). Similar inhibition of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was also observed in cells stimulated with the cAMP-elevating agonist adenosine (Fig. 2C), a key physiological regulator of cAMP-stimulated ion and fluid transport in human airways [48, 63]. In addition, the non-specific phosphodiesterase inhibitor IBMX, and the membrane permeable cAMP analogue dibutyl-cAMP (db-cAMP), both induced almost complete inhibition of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger (Fig. 2C, D). The multidrug resistance protein 4 (MRP4) inhibitor MK-571 also induced an inhibition of the exchanger, but this was less pronounced than for other cAMP agonists (Fig. 2C, D). Therefore, elevations of  $[\text{cAMP}]_i$  via different mechanisms, i.e. (i) activation of adenylyl cyclase (forskolin and adenosine), (ii) inhibition of cAMP breakdown (IBMX) or (iii) inhibition of MRP4-dependent cAMP efflux (MK-571) [67] demonstrated that the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was negatively regulated by cAMP. However,



**Fig. 3** GlyH-101 partially, but not fully, relieves forskolin-induced inhibition of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger: Calu-3 cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 1 h) and  $\text{pH}_i$  was determined by fluorescent microscopy. The activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was assessed by measuring  $\text{pH}_i$  changes in response to replacement of basolateral  $\text{Cl}^-$  with gluconate. **a** Shows a representative experiment in which exchanger activity was assessed under basal conditions and in the presence of GlyH-101 (10  $\mu\text{M}$ ) and forskolin (5  $\mu\text{M}$ ). **b** and **c** summarize the effect of GlyH-101 alone and in the presence of forskolin on the mean change in  $\text{pH}_i$  caused by basolateral  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*Significant effect of treatment vs. control ( $p < 0.05$ ). Data represents mean  $\pm$  S.E.M.,  $n = 4$

inhibition of PKA by two different inhibitors, RpcAMPs and H89, had no effect on the ability of forskolin to reduce AE activity (Fig. 2C, D). Note that this result is in marked contrast to the effect these PKA inhibitors have on the forskolin-activated, CFTR-dependent, apical anion exchanger pendrin, which was reduced by over 80% by H89 [14]. In addition, we also found that the exchange protein activated by cAMP (EPAC) and mTOR, two reported downstream targets of cAMP which are activated independently of PKA [23, 52], were not involved in mediating the effect of cAMP on the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger (Fig. 2C, D). Activation of EPAC by 8CPT-2Me-cAMP-AM failed to mimic the effect of forskolin, with similar  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity compared to control cells ( $n = 3$ ; Fig. 2C, D). Inhibition of mTOR kinase by rapamycin, also had no effect on the forskolin-induced inhibition of exchanger ( $n = 3$ ; data not shown).


In another series of experiments, we tested whether forskolin-stimulated  $\text{HCO}_3^-$  secretion at the apical membrane could potentially mask any basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity, as suggested by Kim et al. [24]. Addition of the CFTR inhibitor, GlyH-101, partially, but not fully, relieved the

forskolin-induced inhibition of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger (Fig. 3). Given that the percent inhibition of the rate of reacidification in response to reintroduction of basolateral  $\text{Cl}^-$  was  $65.2 \pm 14.3\%$  ( $n = 4$ ; Fig. 3C) in the presence of apical GlyH-101 compared to  $97.3 \pm 1.3\%$  in control (+FSK) conditions ( $n = 16$ ; Fig. 2D;  $p < 0.001$ ), suggested that apical  $\text{HCO}_3^-$  secretion under forskolin-stimulated conditions masked some  $\text{pH}_i$  changes resulting from basolateral  $\text{HCO}_3^-$  influx via AE2 in response to removal of basolateral  $\text{Cl}^-$ . However, because there was still a significant decrease of basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity by forskolin, even in the presence of GlyH-101, demonstrates that AE2 is negatively regulated by increases in cAMP. However, this effect appears to be independent of PKA. Thus, the mechanism involved in regulation of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger in Calu-3 cells by cAMP still needs to be resolved.

**The basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger is regulated by resting levels of  $\text{Ca}^{2+}$  and calmodulin signalling** Having shown that the ubiquitous second messenger cAMP played an important role in the regulation of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger in Calu-3 cells, we next investigated its regulation by  $\text{Ca}^{2+}$ , another key second messenger. Not only are  $\text{Ca}^{2+}$  and cAMP signalling generally synergistic in secretory epithelia [9, 27, 38], it has also been demonstrated that the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger in mouse salivary acinar cells is positively regulated by intracellular  $\text{Ca}^{2+}$  [39]. To this end, we first tested whether elevations in intracellular  $\text{Ca}^{2+}$  affected the activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger. Therefore, Calu-3 cells were stimulated with either basolateral carbachol, a muscarinic receptor agonist, (Fig. 4A), or thapsigargin, an inhibitor of sarcoplasmic/endoplasmic reticulum  $\text{Ca}^{2+}$  ATPase (SERCA), (Fig. 4B), which have both been shown to stimulate  $\text{Ca}^{2+}$ -dependent anion transport in Calu-3 cells [37]. However, as shown in Fig. 4C, D, neither carbachol nor thapsigargin had any significant effect on basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity. Therefore, these data suggest that the exchanger is not regulated by  $[\text{Ca}^{2+}]_i$ .

We next tested whether basal levels of  $[\text{Ca}^{2+}]_i$  had any effect on anion exchanger activity. To do this, Calu-3 cells were preloaded with the calcium-chelator, BAPTA-AM, and then basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity assessed (Fig. 5A, B). BAPTA-AM loading did not affect resting  $\text{pH}_i$  (control =  $7.60 \pm 0.06$ ;  $n = 7$  and BAPTA-AM-loaded cells =  $7.70 \pm 0.02$ ;  $n = 5$ ;  $p > 0.05$ ), but did significantly reduce the mean change in  $\text{pH}_i$  following basolateral  $\text{Cl}^-$  removal by  $56.2 \pm 2.9\%$  ( $p < 0.001$   $n = 8$ ; Fig. 5C), as well as the rate of reacidification following basolateral  $\text{Cl}^-$  reintroduction, by  $51.3 \pm 8.9\%$  ( $p < 0.01$ ;  $n = 8$ ; Fig. 5D) compared to control cells. However, BAPTA-AM-loaded cells still showed normal forskolin-induced inhibition of the remaining anion exchange activity. These results suggest that resting levels of intracellular  $\text{Ca}^{2+}$  play an important role in regulating





**Fig. 4** Elevations in intracellular  $Ca^{2+}$  have no effect on the activity of the basolateral  $Cl^-/HCO_3^-$  anion exchanger: Calu-3 cells were loaded with BCECF-AM (10  $\mu$ M; 1 h) and  $pH_i$  was determined by fluorescent microscopy. The activity of the basolateral  $Cl^-/HCO_3^-$  exchanger was assessed by measuring  $pH_i$  changes in response to replacement of basolateral  $Cl^-$  with gluconate. **a** Shows a representative experiment in which basolateral  $Cl^-/HCO_3^-$  anion exchanger activity was assessed in

the presence of basolateral carbachol (Cch; 20  $\mu$ M) and **b** shows a representative experiment in which basolateral  $Cl^-/HCO_3^-$  anion exchanger activity was assessed in the presence of thapsigargin (Thaps; 200 nM). **c** and **d** summarize the effect of carbachol and thapsigargin stimulation on the mean change in  $pH_i$  caused by basolateral  $Cl^-$  removal and the rate of reacidification upon  $Cl^-$  readdition, respectively. Data represents mean  $\pm$  S.E.M.,  $n = 3-13$


basolateral  $Cl^-/HCO_3^-$  exchange activity. Consistent with a  $Ca^{2+}$ -dependent regulation of the basolateral  $Cl^-/HCO_3^-$  exchanger, depletion of intracellular  $Ca^{2+}$  stores (by removal of external  $Ca^{2+}$  and inducing calcium-store depletion with thapsigargin) caused a marked reduction in basolateral  $Cl^-/HCO_3^-$  exchanger activity, with both the change in  $pH_i$  following basolateral  $Cl^-$  removal and the rate of reacidification following  $Cl^-$  readdition reduced by  $57.6 \pm 7.4\%$  and  $65.1 \pm 9.8\%$ , respectively, compared to the control response ( $p < 0.05$ ;  $n = 4$ ; Fig. 5C, D). In contrast, removal of extracellular  $Ca^{2+}$  had no effect on  $Cl^-/HCO_3^-$  exchanger activity (Fig. 5C, D), indicating that intracellular  $Ca^{2+}$  specifically regulated anion exchange activity. Therefore, these data demonstrate that the basal levels of  $[Ca^{2+}]_i$  are sufficient to fully activate the basolateral  $Cl^-/HCO_3^-$  exchanger such that increasing  $[Ca^{2+}]_i$  does not further modulate exchanger activity.

To gain insights into the mechanism underlying the  $Ca^{2+}$ -dependent regulation of the basolateral  $Cl^-/HCO_3^-$  exchanger, we investigated the role of calmodulin (CaM), an important downstream calcium-binding protein. Treatment of cells with the highly specific CaM inhibitor N-(8-aminooctyl)-5-iodonaphthalene-1-sulfonamide (J-8) [64] significantly reduced the mean change in  $pH_i$  following basolateral  $Cl^-$  removal by  $52.1 \pm 7.8\%$  compared to untreated control cells ( $p < 0.01$ ;  $n = 3$ ; Fig. 5C) and also reduced the rate of reacidification following basolateral  $Cl^-$  reintroduction by

$64.0 \pm 13.6\%$  compared to control cells ( $p < 0.05$ ;  $n = 3$ ; Fig. 5D). Thus, these data further support the finding that the resting level of  $Ca^{2+}$  is important in regulating basolateral  $Cl^-/HCO_3^-$  exchanger activity, possibly through  $Ca^{2+}$ -release from thapsigargin-sensitive intracellular stores and subsequent activation of CaM under basal conditions.

#### CK2 plays a major role in the regulation of the basolateral $Cl^-/HCO_3^-$ exchanger in Calu-3 cells

In order to further investigate the mechanism behind  $Ca^{2+}$ /CaM regulation of the basolateral  $Cl^-/HCO_3^-$  exchanger under resting conditions, experiments were performed to assess the role of CK2, a protein kinase that is active under resting conditions in airway cells [65], and which is the main serine/threonine kinase both in vivo and in vitro that can phosphorylate CaM [3]. Furthermore, CK2 has been implicated in the regulation of other ion channels including CFTR [7, 33, 36] and M-type potassium channels, in which it has been reported that CK2-dependent phosphorylation of CaM underlies its binding to the channel [21]. In addition, sequence analysis of SLC4A2 reveals the presence of several potential CK2 phosphorylation sites showing the canonical CK2 consensus sequence pS/pTxxE/D (Supplementary Fig. 1). Calu-3 cells were therefore exposed to the CK2 inhibitor 4,5,6,7-Tetrabromobenzotriazole (TBB) and then basolateral  $Cl^-/HCO_3^-$  exchange activity measured (Fig. 6A). TBB exposure




**Fig. 5** Intracellular  $\text{Ca}^{2+}$  and calmodulin signalling regulate the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger: Calu-3 cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 1 h) and  $\text{pH}_i$  was determined by fluorescent microscopy. The activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was assessed by measuring  $\text{pH}_i$  changes in response to replacement of basolateral  $\text{Cl}^-$  with gluconate. **a** Shows a representative control experiment and **b** shows an experiment in which cells had been preincubated with BAPTA-AM (50  $\mu\text{M}$ ) for 1 h prior to study. **c** and **d**

summarize the effect of BAPTA-AM (50  $\mu\text{M}$ ), depletion of intracellular  $\text{Ca}^{2+}$  stores via stimulation with thapsigargin (200 nM) in the absence of extracellular  $\text{Ca}^{2+}$ , removal of extracellular  $\text{Ca}^{2+}$ , and J-8 (50  $\mu\text{M}$ ), on the mean change in  $\text{pH}_i$  caused by basolateral  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*Significant effect of treatment vs. control ( $p < 0.05$ ; \*\* =  $p < 0.01$ ; \*\*\* =  $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 3\text{--}19$

alone caused a small intracellular acidification ( $7.6 \pm 0.02$  to  $7.5 \pm 0.01$  ( $p < 0.05$ ,  $n = 6$ )), and caused a significant, but fully reversible, inhibition of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger (Fig. 6A–C). Since the mean  $\text{pH}_i$  after TBB exposure was within the normal range for Calu-3 cells, it was unlikely that the fall in  $\text{pH}_i$  caused the decrease in basolateral AE activity. However, to provide further support for a role of CK2 in regulating anion exchange activity, the effect of a more specific CK2 inhibitor, CX4945 [45] was tested. Figure 6D shows that exposure to CX4945 almost completely abolished  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity, an effect which was fully reversible on washout of the inhibitor, and which did not involve a change in  $\text{pH}_i$  (Fig. 6E, F). The use of two different pharmacological inhibitors provided strong evidence that CK2 was essential for basal  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity in Calu-3 cells. The efficacy of a short-term exposure to the two inhibitors on the activity of CK2 was further verified by an in vitro kinase assay of whole cell lysates from Calu-3, as detailed in the ‘Methods’ section. Figure 13A shows that a 5-

min treatment of Calu-3 cells with 10  $\mu\text{M}$  TBB lowered CK2 catalytic activity by more than 40%, whereas the same concentration of CX4945 led to an even more dramatic inhibition. In order to investigate whether CK2 regulation of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger involved CaM, Calu-3 cells were preincubated with the CaM inhibitor J-8 for 60 min, and then cells were acutely exposed to TBB, before basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity was measured. Although TBB + J-8 caused a trend to further reduce AE activity compared to either drug alone, this effect was not statistically significant and, even in the presence of both drugs, a significant amount of AE activity still remained (Fig. 7A, B). Since there was no obvious additive effect of TBB and J-8 suggests that CK2 potentially controls the resting activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger through the downstream target CaM, in Calu-3 cells.

**CK2 inhibition abolishes the activity of basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchange in primary human nasal epithelia**




**Fig. 6** CK2 inhibition by TBB or CX4945 reduces the activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger: Calu-3 cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 1 h) and  $\text{pH}_i$  was determined by fluorescent microscopy. The activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was assessed by measuring  $\text{pH}_i$  changes in response to replacement of basolateral  $\text{Cl}^-$  with gluconate. **a** Shows a representative experiment in which Calu-3 cells were exposed to TBB (10  $\mu\text{M}$ ) for 5 min and  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity measured. **b** and **c** summarize the effect of TBB treatment and TBB reversibility on the mean change in  $\text{pH}_i$  caused by basolateral  $\text{Cl}^-$  removal and the rate of reacidification upon

$\text{Cl}^-$  readdition, respectively. \*Significant effect of TBB vs. untreated controls ( $p < 0.05$ ). Data represents mean  $\pm$  S.E.M.,  $n = 6$ . **d** Shows a representative experiment in which Calu-3 cells were exposed to CX4945 (10  $\mu\text{M}$ ) for 5 min and  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity measured. **e** and **f** summarize the effect of CX4945 treatment and CX4945 reversibility on the mean change in  $\text{pH}_i$  caused by basolateral  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*Significant effect of CX4945 vs. untreated controls ( $p < 0.05$ ). Data represents mean  $\pm$  S.E.M.,  $n = 6$


Having demonstrated that human AE2 was regulated by CK2 in a human airway epithelial cell line, we next assessed whether AE2 activity showed similar CK2-dependent regulation in well-differentiated human nasal epithelial (HNE) cultures. AE2 mRNA expression has previously been identified in the proximal airways and in HNE cells [1, 12, 55], and HNE cells have also been shown to possess a basolateral, DIDS-sensitive  $\text{Cl}^-/\text{HCO}_3^-$  exchanger, indicative of functional expression of AE2 [55]. To this end, intracellular pH measurements were performed on HNE monolayers and the effect of CK2 inhibition on AE2 activity was assessed. In control conditions, removal of basolateral  $\text{Cl}^-$  increased  $\text{pH}_i$  by  $0.08 \pm 0.01$   $\text{pH}_i$  units, and this response was significantly reduced to  $0.01 \pm 0.03$   $\text{pH}_i$  units in the presence of CX4945 ( $n = 5$ ;  $p < 0.05$ ; Fig. 8). The effect of CX4945 was reversible as the response to basolateral  $\text{Cl}^-$  removal could be recovered after wash out of the drug ( $0.09 \pm 0.01$   $\text{pH}_i$  units;  $n = 4$ ;  $p < 0.01$  vs. CX4945; Fig. 8). Therefore, these data indicate that basolateral AE2 activity in primary human nasal epithelia is also positively regulated by CK2 which is consistent with our results from Calu-3 cells.

**Mouse AE2 displays identical regulation to the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger in Calu-3 cells under resting conditions** The current data demonstrated the presence of a DIDS-sensitive  $\text{Cl}^-/\text{HCO}_3^-$  exchanger on the basolateral membrane of Calu-3 cells which was negatively regulated by cAMP, and positively regulated by  $\text{Ca}^{2+}/\text{CaM}$  and CK2. Furthermore, CK2-regulation of a basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was also demonstrated in well differentiated primary cultures of HNE cells. The properties of the exchanger suggested it was SLC4A2 (AE2). Therefore, to study the regulation of AE2 further, we expressed mouse AE2 (mAE2) in HEK-293T cells. Immunocytochemistry revealed successful transfection and expression of the protein at the cell surface of these cells (Fig. 9), although considerable intracellular expression was also observed. Intracellular pH measurements in non-transfected cells showed that HEK cells possessed an endogenous  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity (as previously reported by Sterling et al. [58]). However, this endogenous AE activity was significantly reduced by a low concentration of DIDS (Fig. 10A), which had little effect on exogenously expressed mAE2 activity (Fig. 10B). Therefore, in all further



**Fig. 7** CK2 and CaM regulate the basolateral Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger via similar signalling pathways: Calu-3 cells were loaded with BCECF-AM (10 μM; 1 h) and pH<sub>i</sub> was determined by fluorescent microscopy. The activity of the basolateral Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger was assessed by measuring pH<sub>i</sub> changes in response to replacement of basolateral Cl<sup>-</sup> with gluconate. **a** and **b** display the effect of TBB (10 μM; 5 min exposure), J-8 preincubation (50 μM, 1 h) and J-8 preincubation + acute TBB treatment on the activity of the basolateral Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger, assessed by measuring the mean change in pH<sub>i</sub> caused by basolateral Cl<sup>-</sup> removal and the rate of reacidification upon Cl<sup>-</sup> readdition respectively. \*\*\*Significant effect of treatment ( $p < 0.001$ ); *N.S.* non significant ( $p > 0.05$ ). Data represents mean ± S.E.M.,  $n = 3-6$

experiments, mAE2 activity was assessed in the continuous presence of 25 μM DIDS. Using this approach, after 2 days of transfection, mAE2 expression increased both the magnitude of alkalization in response to removal of basolateral Cl<sup>-</sup>




**Fig. 8** CK2 inhibition abolishes AE2 activity in primary human nasal epithelia: Well-differentiated, primary human nasal epithelia were isolated and cultured as described in the ‘Methods’ section. The activity of the basolateral Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger was assessed by measuring pH<sub>i</sub> changes in response to replacement of basolateral Cl<sup>-</sup> with gluconate. The effect of CX4945 treatment (10 μM; 5 min) and reversibility on the mean change in pH<sub>i</sub> caused by basolateral Cl<sup>-</sup> removal is shown. Note that in these experiments, because the change in pH<sub>i</sub> induced by removal of basolateral Cl<sup>-</sup> was relatively small, it was difficult to obtain accurate rates of reacidification after Cl<sup>-</sup> readdition, particularly for CK2-treated cells, and therefore these data have not been included. \*Significant effect of CX4945 treatment vs. control ( $p < 0.05$ ); \*\*Significant effect of CX4945 wash off (recovery) vs. CX4945 ( $p < 0.01$ ). Data represents mean ± S.E.M.,  $n = 4-5$  from two donors

( $0.13 \pm 0.02$  vs.  $0.69 \pm 0.06$ ;  $p < 0.001$ ;  $n = 10$ ) and the rate of reacidification upon basolateral Cl<sup>-</sup> readdition ( $0.04 \pm 0.01$  pH units min<sup>-1</sup> vs.  $0.46 \pm 0.07$  pH units min<sup>-1</sup>;  $p < 0.001$ ;  $n = 10$ ), compared to non-transfected cells (Fig. 10C, D, respectively). We found mAE2 activity to be sensitive to DIDS, with 100 μM DIDS causing a  $40.0 \pm 4.3\%$  inhibition of the mean pH<sub>i</sub> change in response to extracellular Cl<sup>-</sup> removal ( $p < 0.001$ ;  $n = 3$ ) and a  $80.4 \pm 4.5\%$  inhibition of the rate of reacidification in response to reintroduction of extracellular Cl<sup>-</sup> ( $p < 0.001$ ;  $n = 3$ ).

We next investigated whether mAE2 transiently expressed in HEK-293T cells was regulated in a similar fashion to human AE2 expressed in Calu-3 and HNE cells. Figure 11A–D shows that mAE2 activity in HEK-293T cells was reduced by both BAPTA-AM and J-8, as well as TBB and CX4945, demonstrating that both Ca<sup>2+</sup>/CaM signalling and CK2 regulate mAE2 when exogenously expressed in HEK-293T cells. In addition, short-term exposure to CX4945 also inhibited CK2 catalytic activity in HEK-293T cell lysates, demonstrating that the effect of CX4945 on mAE2 activity was very likely a result of CK2 inhibition (see Fig. 13B).

**Knockout of CK2 catalytic subunits reduces mAE2 activity in HEK-293T cells** To further investigate CK2 regulation of mAE2, we studied the activity of mAE2 in genetically altered HEK-293T cells. CK2 is a tetramer consisting of two catalytically active α subunits and two β subunits. We therefore genetically knocked out either the α catalytic subunit or the α prime catalytic subunit of CK2 (αCK2-KO and αprimeCK2-KO HEK-293T cells; see the ‘Methods’ section and supplementary Fig. 2). Note that α and α prime isoforms are highly homologous proteins with large overlapping functions and we found that CK2 catalytic activity was reduced by ~50% in αCK2-KO cells (Fig. 13B). In both types of CK2 KO cells, mAE2 activity was significantly reduced compared to WT HEK-293T cells, with the mean pH<sub>i</sub> increase in response to removal of extracellular Cl<sup>-</sup> reduced by  $24.6 \pm 6.4\%$  in αCK2-KO cells ( $p < 0.05$ ;  $n = 7$ ; Fig. 12A) and  $24.2 \pm 4.0\%$  in αprimeCK2-KO cells ( $p < 0.05$ ;  $n = 6$ ; Fig. 12A) while the rate of reacidification after extracellular Cl<sup>-</sup> readdition was reduced by  $42.0 \pm 5.9\%$  in αCK2-KO cells ( $p < 0.01$ ;  $n = 7$ ; Fig. 12B) and  $49.4 \pm 5.5\%$  in αprimeCK2-KO cells ( $p < 0.01$ ;  $n = 6$ ; Fig. 12B). Furthermore, transfecting αCK2 KO HEK-293T cells with α-CK2 (to restore α CK2 activity) significantly recovered mAE2 activity in co-transfected cells, compared to ‘control’ co-transfected cells (i.e. mAE2 and empty plasmid) (Fig. 12C, D). Finally, co-transfecting normal HEK-293T cells with a double α catalytic CK2 mutant subunit with reduced sensitivity to TBB (DM CK2; see the ‘Methods’ section) [35, 65] led to a significant decrease in TBB inhibition of



**Fig. 9** Expression and localization of mAE2 transiently transfected into HEK-293T cells: HEK-293T cells were transfected with HA-tagged mAE2 cDNA and studied 2 days post-transfection. **a** and **b** show confocal images in which mAE2 was labelled using a mouse anti-HA primary antibody and a goat anti-mouse secondary antibody conjugated to FITC fluorophore (green fluorescence). The nuclei were stained with DAPI (blue fluorescence). Significant intracellular as well as plasma

membrane staining was observed. In **c**, the plasma membrane was stained with WGA (red fluorescence), and plasma membrane localisation of mAE2 was thus confirmed by yellow staining. **d** Shows non-transfected cells stained with mouse anti-HA primary antibody and goat anti-mouse FITC-conjugated secondary antibody. Data representative of three independent experiments


mAE2 activity, compared to control cells (Fig. 12E, F). Taken together with the pharmacological data, our results clearly indicate that CK2 plays a critical role in the regulation of mAE2 activity under resting conditions. These data are also consistent with the findings from Calu-3 cells and primary HNE cells, which provides further support that the identity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger in airway epithelial cells is hAE2 (SLCA42). Thus, our results have uncovered a key regulatory mechanism for mammalian AE2 by CK2.

## Discussion

SLC4A2 has been identified on the basolateral membrane of human airway epithelia [1, 32] yet its regulation still remains poorly understood. Although the exact role of SLC4A2 in the conducting airways in general, including Calu-3 cells, still needs to be determined, it is likely to be important in regulating intracellular pH via its ability to transport  $\text{HCO}_3^-$  across the basolateral membrane, as shown in this study and by

others [14, 20, 24, 40]. Furthermore, because it transports  $\text{Cl}^-$  under resting conditions, AE2 will also act to accumulate  $\text{Cl}^-$  inside the cell, particularly when working in parallel with the sodium/potassium-dependent chloride (NKCC1) co-transporter that facilitates the influx of  $\text{Na}^+$ ,  $\text{K}^+$  and  $2\text{Cl}^-$  ions across the basolateral membrane of Calu-3 cells [31]. Huang et al. [18] have shown AE2 knockdown in Calu-3 cells reduced transepithelial  $\text{Cl}^-$  secretion by ~60% whereas the NKCC1 inhibitor bumetamide reduced  $\text{Cl}^-$  secretion by ~20%. These results demonstrate the importance of AE2 in transepithelial  $\text{Cl}^-$  secretion. Here, we have explored the cellular mechanisms that regulate the activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger in Calu-3 cells.

We demonstrated that the  $\text{IC}_{50}$  for DIDS inhibition of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was approximately  $17 \mu\text{M}$ , which is in good agreement with the  $\text{IC}_{50}$  of ~ $13 \mu\text{M}$  reported by Humphreys et al. [19] for DIDS inhibition of human AE2 heterologously expressed in *Xenopus oocytes*. However, given that  $500 \mu\text{M}$  DIDS was required to fully inhibit the exchanger suggested that other DIDS-sensitive transporters might also be present in the basolateral membrane of Calu-3




**Fig. 10** mAE2-expressing cells demonstrate enhanced  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity and reduced sensitivity to DIDS compared to non-transfected cells: HEK-293T cells were transfected with HA-tagged mAE2 cDNA and studied 2 days post-transfection, and results compared to non-transfected cells. Cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 10 min) and  $\text{pH}_i$  was determined by fluorescent microscopy. The activity of  $\text{Cl}^-/\text{HCO}_3^-$  exchangers was assessed by measuring  $\text{pH}_i$  changes in response to replacement of extracellular  $\text{Cl}^-$  with gluconate. **a** and **b** show representative experiments in which  $\text{Cl}^-/\text{HCO}_3^-$  exchanger

activity was measured in non-transfected cells and mAE2-transfected cells, respectively. **c** and **d** summarize the effect of both mAE2 transfection and DIDS (25  $\mu\text{M}$ ) on the mean change in  $\text{pH}_i$  caused by extracellular  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*\*Significant effect of DIDS ( $p < 0.01$ ); \*\*\* =  $p < 0.001$ ); †Significant effect of mAE2 transfection ( $p < 0.01$ ). Data represents mean  $\pm$  S.E.M.,  $n = 4$  for non-transfected cells and  $n = 10$  for transfected cells

cells. This could be other SLC4 family members, such as SLC4A9 (AE4) which has recently been demonstrated in mouse submandibular gland acinar cells [41], or even members of the SLC26 family, such as SLC26A7, which play an important role in  $\text{HCO}_3^-$  transport across the basolateral membrane of gastric parietal cells [42] and intercalated cells of the outer medullary collecting duct [43]. We observed that the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger was active in non-stimulated Calu-3 cells yet elevations in cAMP, using a variety of different mechanisms, (i.e. activation of transmembrane adenylyl cyclase, inhibition of phosphodiesterases or inhibition of MRP-dependent cAMP efflux) all significantly inhibited the exchanger, demonstrating that cAMP negatively regulates this anion transporter. This cAMP-dependent inhibition of AE2-dependent  $\text{HCO}_3^-$  efflux at the basolateral membrane would enhance the efficiency of cAMP-stimulated, CFTR-dependent  $\text{HCO}_3^-$  secretion in Calu-3 cells. However, the downstream mechanism underlying AE

inhibition by cAMP appeared not to involve PKA, as it was insensitive to two different PKA inhibitors, nor did it involve a number of other well characterized cAMP-dependent/PKA-independent signalling pathways, including EPAC and mTOR kinase. Although the present study supports our previous work [14], further investigations into the effect of cAMP on the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity demonstrated that some forskolin-induced inhibition of the exchanger was alleviated in the presence of the CFTR inhibitor GlyH-101. These findings are similar, but not identical, to those reported by Huang et al. [18, 54] and Kim et al. [24] who failed to demonstrate any effect of cAMP on basolateral AE activity. These authors suggested that the apparent cAMP-dependent inhibition of the basolateral AE activity we observed was due to the activation of CFTR-dependent  $\text{HCO}_3^-$  efflux across the apical membrane, which ‘swamped’ the basolateral response. Given that we were able to detect basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity in forskolin-stimulated cells when CFTR was




**Fig. 11** mAE2 is regulated by  $\text{Ca}^{2+}$ /CaM signalling and by CK2: HEK-293T cells were transfected with HA-tagged mAE2 cDNA and studied 2 days post-transfection. Cells were loaded with BCECF-AM (10  $\mu\text{M}$ ; 10 min) and  $\text{pH}_i$  was determined by fluorescent microscopy. The activity of mAE2 was assessed by measuring  $\text{pH}_i$  changes in response to replacement of extracellular  $\text{Cl}^-$  with gluconate in the presence of 25  $\mu\text{M}$  DIDS to inhibit endogenous  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity. **a** and **b** display the effect of BAPTA-AM (50  $\mu\text{M}$ ) and J-8 (50  $\mu\text{M}$ ) on the mean change in  $\text{pH}_i$  caused by extracellular  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*Significant effect of treatment vs. control ( $p < 0.05$ ; \*\* =  $p < 0.01$ ; \*\*\* =  $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 4-11$ . **c** and **d** display the effect of TBB (10  $\mu\text{M}$ ) and CX4945 (10  $\mu\text{M}$ ) treatment on the mean change in  $\text{pH}_i$  caused by extracellular  $\text{Cl}^-$  removal and the rate of reacidification upon  $\text{Cl}^-$  readdition, respectively. \*\*Significant effect of treatment vs. control (\*\*\* =  $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 7-16$

inhibited lends support to this explanation. However, it is important to consider that the forskolin-stimulated inhibition of AE2 was not fully relieved, indicating there was still an effect of cAMP on the activity of the exchanger. In addition, our data, and those reported by others [18, 24, 54], do not explain how two different PKA inhibitors failed to affect the cAMP-dependent inhibition of the basolateral AE activity,

particularly as we have previously shown these PKA inhibitors markedly reduced CFTR-dependent  $\text{HCO}_3^-$  efflux in Calu-3 cells [14]. Clearly, further work is still required to unravel the role of cAMP in the regulation of AE2-dependent  $\text{HCO}_3^-$  transport across the basolateral membrane of Calu-3 cells. It was also interesting that GlyH-101 alone affected the rate of reacidification caused by basolateral anion exchanger activity in the absence of cAMP stimulation (Fig. 3B). This effect is likely due to a rise in intracellular  $[\text{Cl}^-]$ , through an inhibition of  $\text{Cl}^-$  efflux by CFTR at the apical membrane, reducing the driving force for  $\text{Cl}^-$  entry at the basolateral membrane (on the reintroduction of external  $\text{Cl}^-$ ), and thereby reducing  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity.

We also demonstrated that the exchanger was  $\text{Ca}^{2+}$ -sensitive, as BAPTA-AM and depletion of thapsigargin-sensitive intracellular  $\text{Ca}^{2+}$  stores all markedly reduced the activity of the exchanger. One possible explanation for the effect seen with  $\text{Ca}^{2+}$  store depletion could involve a rise in cAMP, via the activation of store-operated cAMP signalling (socAMPs) as described by Lefkimmatis et al. [30], in which the endoplasmic reticulum  $\text{Ca}^{2+}$ -sensor STIM couples ER  $\text{Ca}^{2+}$  levels to cAMP production, via activation of transmembrane adenylyl cyclase 8. However, it is worth noting that in these intracellular  $\text{Ca}^{2+}$ -depleted conditions, we observed that cAMP-stimulated  $\text{Cl}^-/\text{HCO}_3^-$  exchange at the apical membrane was also reduced which argues against socAMPs being activated by these conditions in Calu-3 cells (unpublished observations). In addition, inhibition of CaM using J-8 caused a marked reduction in the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity under resting conditions, suggesting  $\text{Ca}^{2+}$ -dependent activation of CaM was important for exchanger activity. To further understand the mechanism of CaM-dependent regulation of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger under resting conditions, the role of CK2 was assessed since it has been shown that CK2 is an important regulator of CaM activity via phosphorylation of three physiological CK2-phosphorylation acceptor sites in this  $\text{Ca}^{2+}$ -binding protein [3, 46]. Treatment of Calu-3 cells with two different selective CK2 inhibitors, TBB and CX4945, caused a significant decrease in basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity under resting conditions. Furthermore, when cells were treated simultaneously with J-8 and TBB, there was very little further decrease in  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity, compared to the presence of each inhibitor alone. These results indicate that CK2 regulates the activity of the basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger under resting conditions, potentially through a CaM-dependent mechanism, although the role of CaM requires further investigation. Therefore, our



**Fig. 12** Genetic knockout of  $\alpha$ CK2 reduces mAE2 activity which was rescued by reintroduction of  $\alpha$ CK2 expression: **a** and **b** summarize the activity of mAE2 transiently expressed in either  $\alpha$ CK2-KO HEK-293T cells or  $\alpha$ primeCK2-KO HEK-293T cells, assessed by measuring  $\text{pH}_i$  changes in response to replacement of extracellular  $\text{Cl}^-$  with gluconate in the presence of 25  $\mu\text{M}$  DIDS to inhibit endogenous  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity. \*Significant effect of  $\alpha$ CK2 KO or  $\alpha$ primeCK2 KO ( $p < 0.05$ ; \*\* =  $p < 0.01$ ). Data represents mean  $\pm$  S.E.M.,  $n = 4-7$ . **c** and **d** summarize the activity of mAE2 transiently expressed in  $\alpha$ CK2 KO

HEK-293T cells that were also co-transfected with WT CK2. Anion exchanger activity was assessed by measuring  $\text{pH}_i$  changes in response to replacement of extracellular  $\text{Cl}^-$  with gluconate in the presence of 25  $\mu\text{M}$  DIDS to inhibit endogenous  $\text{Cl}^-/\text{HCO}_3^-$  exchange activity. \*Significant effect of WT CK2 expression ( $p < 0.05$ ). Data represents mean  $\pm$  S.E.M.,  $n = 4$ . **e** and **f** display the percent inhibition of mAE2 activity by TBB in cells transfected with a CK2 double mutant (DM CK2). \*\*\*Significant difference between DM CK2 and WT CK2 ( $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 7-9$


findings have uncovered an important role for CK2/CaM and calcium in the regulation of  $\text{Cl}^-/\text{HCO}_3^-$  exchange in human airway epithelia.

We also demonstrated that CX4945 significantly reduced basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchanger activity in primary HNE cells that have previously been shown to express AE2 mRNA and a functional, DIDS-sensitive  $\text{Cl}^-/\text{HCO}_3^-$  exchanger [55]. It was interesting to observe that the changes in  $\text{pH}_i$  in response to removal of basolateral  $\text{Cl}^-$  were approximately fourfold lower in HNE cells compared to Calu-3 cells, suggesting that there is more robust expression of AE2 in Calu-3 cells. Although Shin et al. [55] demonstrated  $\text{pH}_i$  changes of  $\sim 0.4$   $\text{pH}_i$  units in response to removal of

basolateral  $\text{Cl}^-$ , their experiments were performed on cells that had not fully differentiated. Indeed, in the same study, they demonstrated that AE2 expression decreased as cells differentiated and this likely underlies the different responses seen in their study and the current study. Nevertheless, these findings demonstrate CK2 regulation of basolateral  $\text{Cl}^-/\text{HCO}_3^-$  exchange also occurs in primary airway epithelial cultures which provide support for CK2 regulation of AE2 in vivo.

To strengthen our hypothesis that the DIDS-sensitive,  $\text{Cl}^-/\text{HCO}_3^-$  exchanger present on the basolateral membrane of human airway epithelia was SLC4A2 (AE2), we overexpressed mouse AE2 in HEK-293T cells and compared its regulation to the findings from Calu-3 cells. Similar to the





**Fig. 13** CK2 catalytic activity is inhibited by short-term exposure to specific inhibitors: Cell lysates were generated from **a** Calu-3 cells treated with TBB (10  $\mu$ M; 5 min) or CX4945 (10  $\mu$ M; 5 min) or **b** HEK-293T cells treated with CX4945 (10  $\mu$ M; 5 min) or the  $\alpha$ CK2-KO HEK-293T and CK2 activity was determined by means of

radioactive assays with [ $\gamma$ -<sup>33</sup>P]ATP towards the specific CK2 substrate peptide CK2-tide (RRRADDSDDDDD).\*\*\*Significant effect of inhibitor vs. untreated control or  $\alpha$ CK2KO vs. control ( $p < 0.001$ ). Data represents mean  $\pm$  S.E.M.,  $n = 4$

data obtained in Calu-3 cells, mAE2 activity in HEK-293T cells was also reduced by chelation of intracellular  $\text{Ca}^{2+}$  and inhibition of CaM. This is consistent with findings from Chernova et al. [8] who showed that both BAPTA-AM and the CaM inhibitor, calmidazolium, significantly reduced  $\text{Cl}^-$  transport by murine AE2, when expressed in *Xenopus* oocytes, findings also reported by Stewart et al. [60]. Collectively, these results indicate that under resting conditions, CaM maintains normal AE2 activity potentially through a  $\text{Ca}^{2+}$ -dependent pathway. Finally, we also demonstrated that pharmacological inhibition, or genetic knockout of CK2, significantly reduced mAE2 activity, consistent with results from Calu-3 and HNE cells. Because CK2 catalytic activity was reduced by  $\sim 50\%$  in HEK-293T KO cells (Fig. 13B) and that mAE2 activity was reduced by a similar amount, (Fig. 12B) provides further evidence that CK2 plays a major role in the regulation of AE2 in airway epithelial cells. It was interesting to observe that the degree of inhibition by CX4945 was significantly less for mAE2 expressed in HEK-293T cells compared to that observed for the exchanger in Calu-3 cells (compare Figs. 6 with 11). This could be explained by the ‘addiction’ phenomenon (reviewed by Venerando et al. [66]) occurring in tumour cell lines, such as Calu-3, in which the level of CK2 is higher than in normal cells such as HEK-293T. Consequently, highly specific inhibition of CK2 by CX4945 in Calu-3 cells causes more pronounced downstream effects than those observed in HEK-293T cells. An interesting result in support of this hypothesis is shown in Fig. 13, where the percent inhibition of CK2 by CX4945 is significantly higher in Calu-3 cells compared to HEK-293T cells ( $69.6 \pm 1.6\%$  vs.  $15.4 \pm 3.0\%$ ;  $p < 0.001$ ;  $n = 4$ ; compare Fig. 13A, B), indicating Calu-3 cells have more active CK2 than HEK-293T cells.

In summary, we have demonstrated for the first time that human airway epithelial cells express a basolateral DIDS-sensitive,  $\text{Cl}^-/\text{HCO}_3^-$  exchanger which is regulated by CK2. Since mouse *slc4A2* was regulated in an identical way when studied in a heterologous expression system suggests that the identity of the transporter in human airway cell was SLC4A2. Therefore, our findings identify CK2 as a new regulator of SLC4A2-dependent anion transport in human airway epithelia and suggest that CK2-dependent phosphorylation of SLC4A2, or an associated regulatory protein such as CaM, is essential for AE2 activity under resting conditions. We suggest that maintenance of basal AE2 activity will ensure efficient  $\text{Cl}^-$  loading into the cell prior to cAMP-stimulated anion secretion, which will help support net transepithelial  $\text{HCO}_3^-$  and fluid secretion in airway epithelial cells [18]. Indeed, since CK2 also positively regulates the activity of CFTR suggests that this protein kinase plays an essential role in coordinating the activity of  $\text{HCO}_3^-$  transporters at both the apical and basolateral membranes of airway epithelial cells.

**Acknowledgements** This work was supported by an overseas studentship to S.H.I. funded by the Higher Committee for Education Development (HCED), Iraq. M.J.T. was funded by a MRC studentship. V.S.-C. was funded by the Cystic Fibrosis Trust (SRC003). J.G. and C.W. were funded by the Medical Research Foundation (Grant Reference: MRF-091-0001-RG-GARNE). I.J.H. was funded by a Wellcome Trust Research Training Fellowship. A.V. was funded by the Italian Cystic Fibrosis Research Foundation (FFC#7/2014 sponsored by Open d’Italia di Golf). The authors would like to thank Dr. Bernard Verdon for technical support and Dr. Anil Mehta for his valuable input into the preparation of this manuscript.

**Author contributions** S.H.I. performed experiments, analysed data and prepared the figures.

M.J.T. performed experiments, analysed data, prepared the figures, drafted the manuscript and edited the manuscript.

V.S.-C. performed experiments, analysed data, prepared the figures and edited the manuscript.

J.G. performed experiments and analysed data.

I.J. H. contributed resources.

M.B. contributed resources.

C.W. contributed resources.

C.B. performed experiments and contributed resources.

M.S. and A.V. contributed resources and edited the manuscript.

M.A.G. conceived and designed the study and edited the manuscript.

**Compliance with ethical standards** Ethical approval for collection of paediatric human nasal epithelial cells was granted by the NRES Committee North East - Newcastle and North Tyneside 1 (REC reference: 15/NE/0215) and informed written parental consent was obtained.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Al-Bazzaz FJ, Hafez N, Tyagi S, Gailey CA, Toofanfard M, Alrefai WA, Nazir TM, Ramaswamy K, Dudeja PK (2001) Detection of Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> and Na<sup>+</sup>/H<sup>+</sup> exchangers in human airways epithelium. *JOP* 2:285–290
- Alper SL, Darman RB, Chernova MN, Dahl NK (2002) The AE gene family of Cl/HCO<sub>3</sub><sup>-</sup> exchangers. *J Nephrol* 15(Suppl 5):S41–S53
- Arrigoni G, Marin O, Pagano MA, Settimo L, Paolin B, Meggio F, Pinna LA (2004) Phosphorylation of calmodulin fragments by protein kinase CK2. Mechanistic aspects and structural consequences. *Biochemistry* 43:12788–12798. doi:10.1021/bi049365c
- Ballard ST, Trout L, Bebek Z, Sorscher EJ, Crews A (1999) CFTR involvement in chloride, bicarbonate, and liquid secretion by airway submucosal glands. *Am J Phys* 277:L694–L699
- Bhaskar KR, Gong DH, Bansil R, Pajevic S, Hamilton JA, Turner BS, LaMont JT (1991) Profound increase in viscosity and aggregation of pig gastric mucin at low pH. *Am J Phys* 261:G827–G832
- Borowitz D (2015) CFTR, bicarbonate, and the pathophysiology of cystic fibrosis. *Pediatr Pulmonol* 50(Suppl 40):S24–S30. doi:10.1002/ppul.23247
- Cesaro L, Marin O, Venerando A, Donella-Deana A, Pinna LA (2013) Phosphorylation of cystic fibrosis transmembrane conductance regulator (CFTR) serine-511 by the combined action of tyrosine kinases and CK2: the implication of tyrosine-512 and phenylalanine-508. *Amino Acids* 45:1423–1429. doi:10.1007/s00726-013-1613-y
- Chernova MN, Stewart AK, Jiang L, Friedman DJ, Kunes YZ, Alper SL (2003) Structure-function relationships of AE2 regulation by Ca<sup>(i)</sup>(2+)-sensitive stimulators NH<sup>(4+)</sup> and hypertonicity. *Am J Physiol Cell Physiol* 284:C1235–C1246. doi:10.1152/ajpcell.00522.2002
- Choi JY, Joo NS, Krouse ME, Wu JV, Robbins RC, Ianowski JP, Hanrahan JW, Wine JJ (2007) Synergistic airway gland mucus secretion in response to vasoactive intestinal peptide and carbachol is lost in cystic fibrosis. *J Clin Invest* 117:3118–3127. doi:10.1172/JCI31992
- Coakley RD, Grubb BR, Paradiso AM, Gatzky JT, Johnson LG, Kreda SM, O'Neal WK, Boucher RC (2003) Abnormal surface liquid pH regulation by cultured cystic fibrosis bronchial epithelium. *Proc Natl Acad Sci U S A* 100:16083–16088. doi:10.1073/pnas.2634339100
- Devor DC, Singh AK, Lambert LC, DeLuca A, Frizzell RA, Bridges RJ (1999) Bicarbonate and chloride secretion in Calu-3 human airway epithelial cells. *J Gen Physiol* 113:743–760
- Dudeja PK, Hafez N, Tyagi S, Gailey CA, Toofanfard M, Alrefai WA, Nazir TM, Ramaswamy K, Al-Bazzaz FJ (1999) Expression of the Na<sup>+</sup>/H<sup>+</sup> and Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger isoforms in proximal and distal human airways. *Am J Phys* 276:L971–L978
- Garland AL, Walton WG, Coakley RD, Tan CD, Gilmore RC, Hobbs CA, Tripathy A, Clunes LA, Bencharit S, Stutts MJ, Betts L, Redinbo MR, Tarran R (2013) Molecular basis for pH-dependent mucosal dehydration in cystic fibrosis airways. *Proc Natl Acad Sci U S A* 110:15973–15978. doi:10.1073/pnas.1311999110
- Garnett JP, Hickman E, Burrows R, Hegyi P, Tiszlavicz L, Cuthbert AW, Fong P, Gray MA (2011) Novel role for pendrin in orchestrating bicarbonate secretion in cystic fibrosis transmembrane conductance regulator (CFTR)-expressing airway serous cells. *J Biol Chem* 286:41069–41082. doi:10.1074/jbc.M111.266734
- Garnett JP, Hickman E, Tunkammerdthai O, Cuthbert AW, Gray MA (2013) Protein phosphatase 1 coordinates CFTR-dependent airway epithelial HCO<sub>3</sub><sup>-</sup> secretion by reciprocal regulation of apical and basolateral membrane Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchangers. *Br J Pharmacol* 168:1946–1960. doi:10.1111/bph.12085
- Hegyi P, Gray MA, Argent BE (2003) Substance P inhibits bicarbonate secretion from guinea pig pancreatic ducts by modulating an anion exchanger. *Am J Physiol Cell Physiol* 285:C268–C276. doi:10.1152/ajpcell.00574.2002
- Hoegger MJ, Fischer AJ, McMenimen JD, Ostedgaard LS, Tucker AJ, Awadalla MA, Moninger TO, Michalski AS, Hoffman EA, Zabner J, Stoltz DA, Welsh MJ (2014) Impaired mucus detachment disrupts mucociliary transport in a piglet model of cystic fibrosis. *Science* 345:818–822. doi:10.1126/science.1255825
- Huang J, Shan J, Kim D, Liao J, Evagelidis A, Alper SL, Hanrahan JW (2012) Basolateral chloride loading by the anion exchanger type 2: role in fluid secretion by the human airway epithelial cell line Calu-3. *J Physiol* 590:5299–5316. doi:10.1113/jphysiol.2012.236919
- Humphreys BD, Jiang L, Chernova MN, Alper SL (1994) Functional characterization and regulation by pH of murine AE2 anion exchanger expressed in *Xenopus* oocytes. *Am J Phys* 267:C1295–C1307
- Inglis SK, Finlay L, Ramminger SJ, Richard K, Ward MR, Wilson SM, Olver RE (2002) Regulation of intracellular pH in Calu-3 human airway cells. *J Physiol* 538:527–539
- Kang S, Xu M, Cooper EC, Hoshi N (2014) Channel-anchored protein kinase CK2 and protein phosphatase 1 reciprocally regulate KCNQ2-containing M-channels via phosphorylation of calmodulin. *J Biol Chem* 289:11536–11544. doi:10.1074/jbc.M113.528497
- Kellum JA (2000) Determinants of blood pH in health and disease. *Crit Care* 4:6–14. doi:10.1186/cc644
- Kim HW, Ha SH, Lee MN, Huston E, Kim DH, Jang SK, Suh PG, Houslay MD, Ryu SH (2010) Cyclic AMP controls mTOR through regulation of the dynamic interaction between Rheb and phosphodiesterase 4D. *Mol Cell Biol* 30:5406–5420. doi:10.1128/MCB.00217-10
- Kim D, Kim J, Burghardt B, Best L, Steward MC (2014) Role of anion exchangers in Cl<sup>-</sup> and HCO<sub>3</sub><sup>-</sup> secretion by the human airway

- epithelial cell line Calu-3. *Am J Physiol Cell Physiol* 307:C208–C219. doi:10.1152/ajpcell.00083.2014
25. Ko SB, Zeng W, Dorwart MR, Luo X, Kim KH, Millen L, Goto H, Naruse S, Soyombo A, Thomas PJ, Muallem S (2004) Gating of CFTR by the STAS domain of SLC26 transporters. *Nat Cell Biol* 6: 343–350. doi:10.1038/ncb1115
  26. Kreindler JL, Peters KW, Frizzell RA, Bridges RJ (2006) Identification and membrane localization of electrogenic sodium bicarbonate cotransporters in Calu-3 cells. *Biochim Biophys Acta* 1762:704–710. doi:10.1016/j.bbadis.2006.06.005
  27. Lee RJ, Foskett JK (2010) cAMP-activated  $Ca^{2+}$  signaling is required for CFTR-mediated serous cell fluid secretion in porcine and human airways. *J Clin Invest* 120:3137–3148
  28. Lee BS, Gunn RB, Kopito RR (1991) Functional differences among nonerythroid anion exchangers expressed in a transfected human cell line. *J Biol Chem* 266:11448–11454
  29. Lee MC, Penland CM, Widdicombe JH, Wine JJ (1998) Evidence that Calu-3 human airway cells secrete bicarbonate. *Am J Phys* 274: L450–L453
  30. Lefkimmiatis K, Srikanthan M, Maiellaro I, Moyer MP, Curci S, Hofer AM (2009) Store-operated cyclic AMP signalling mediated by STIM1. *Nat Cell Biol* 11:433–442. doi:10.1038/ncb1850
  31. Liedtke CM, Cody D, Cole TS (2001) Differential regulation of  $Cl^{-}$  transport proteins by PKC in Calu-3 cells. *Am J Physiol Lung Cell Mol Physiol* 280:L739–L747
  32. Loffing J, Moyer BD, Reynolds D, Shmukler BE, Alper SL, Stanton BA (2000) Functional and molecular characterization of an anion exchanger in airway serous epithelial cells. *Am J Physiol Cell Physiol* 279:C1016–C1023
  33. Luz S, Kongsuphol P, Mendes AI, Romeiras F, Sousa M, Schreiber R, Matos P, Jordan P, Mehta A, Amaral MD, Kunzelmann K, Farinha CM (2011) Contribution of casein kinase 2 and spleen tyrosine kinase to CFTR trafficking and protein kinase A-induced activity. *Mol Cell Biol* 31:4392–4404. doi:10.1128/MCB.05517-11
  34. Matsui H, Randell SH, Peretti SW, Davis CW, Boucher RC (1998) Coordinated clearance of periciliary liquid and mucus from airway surfaces. *J Clin Invest* 102:1125–1131. doi:10.1172/JCI2687
  35. Meggio F, Pagano MA, Moro S, Zagotto G, Ruzzene M, Sarno S, Cozza G, Bain J, Elliott M, Deana AD, Brunati AM, Pinna LA (2004) Inhibition of protein kinase CK2 by condensed polyphenolic derivatives. An in vitro and in vivo study *Biochemistry* 43:12931–12936. doi:10.1021/bi048999g
  36. Mehta A (2008) Cystic fibrosis as a bowel cancer syndrome and the potential role of CK2. *Mol Cell Biochem* 316:169–175. doi:10.1007/s11010-008-9815-4
  37. Moon S, Singh M, Krouse ME, Wine JJ (1997) Calcium-stimulated  $Cl^{-}$  secretion in Calu-3 human airway cells requires CFTR. *Am J Phys* 273:L1208–L1219
  38. Namkung W, Finkbeiner WE, Verkman AS (2010) CFTR-adenylyl cyclase I association responsible for UTP activation of CFTR in well-differentiated primary human bronchial cell cultures. *Mol Biol Cell* 21:2639–2648. doi:10.1091/mbc.E09-12-1004
  39. Nguyen HV, Stuart-Tilley A, Alper SL, Melvin JE (2004)  $Cl^{-}/HCO_3^{-}$  exchange is acetazolamide sensitive and activated by a muscarinic receptor-induced  $[Ca^{2+}]_i$  increase in salivary acinar cells. *Am J Physiol Gastrointest Liver Physiol* 286:G312–G320. doi:10.1152/ajpgi.00158.2003
  40. Paradiso AM, Coakley RD, Boucher RC (2003) Polarized distribution of  $HCO_3^{-}$  transport in human normal and cystic fibrosis nasal epithelia. *J Physiol* 548:203–218. doi:10.1113/jphysiol.2002.034447
  41. Pena-Munzenmayer G, Catalan MA, Kondo Y, Jaramillo Y, Liu F, Shull GE, Melvin JE (2015) Ae4 (Slc4a9) anion exchanger drives  $Cl^{-}$  uptake-dependent fluid secretion by mouse submandibular gland acinar cells. *J Biol Chem* 290:10677–10688. doi:10.1074/jbc.M114.612895
  42. Petrovic S, Ju X, Barone S, Seidler U, Alper SL, Lohi H, Kere J, Soleimani M (2003) Identification of a basolateral  $Cl^{-}/HCO_3^{-}$  exchanger specific to gastric parietal cells. *Am J Physiol Gastrointest Liver Physiol* 284:G1093–G1103. doi:10.1152/ajpgi.00454.2002
  43. Petrovic S, Barone S, Xu J, Conforti L, Ma L, Kujala M, Kere J, Soleimani M (2004) SLC26A7: a basolateral  $Cl^{-}/HCO_3^{-}$  exchanger specific to intercalated cells of the outer medullary collecting duct. *Am J Physiol Renal Physiol* 286:F161–F169. doi:10.1152/ajprenal.00219.2003
  44. Pezzulo AA, Tang XX, Hoegger MJ, Alaiwa MH, Ramachandran S, Moninger TO, Karp PH, Wohlford-Lenane CL, Haagsman HP, van Eijk M, Banfi B, Horswill AR, Stoltz DA, McCray PB Jr, Welsh MJ, Zabner J (2012) Reduced airway surface pH impairs bacterial killing in the porcine cystic fibrosis lung. *Nature* 487: 109–113. doi:10.1038/nature11130
  45. Pierre F, Chua PC, O'Brien SE, Siddiqui-Jain A, Bourbon P, Haddach M, Michaux J, Nagasawa J, Schwaebe MK, Stefan E, Vialesse A, Whitten JP, Chen TK, Darjania L, Stansfield R, Andres K, Bliesath J, Drygin D, Ho C, Omori M, Proffitt C, Streiner N, Trent K, Rice WG, Ryckman DM (2011) Discovery and SAR of 5-(3-chlorophenylamino)benzo[c][2,6]naphthyridine-8-carboxylic acid (CX-4945), the first clinical stage inhibitor of protein kinase CK2 for the treatment of cancer. *J Med Chem* 54: 635–654. doi:10.1021/jm101251q
  46. Quadroni M, James P, Carafoli E (1994) Isolation of phosphorylated calmodulin from rat liver and identification of the in vivo phosphorylation sites. *J Biol Chem* 269:16116–16122
  47. Quinton PM (2008) Cystic fibrosis: impaired bicarbonate secretion and mucoviscidosis. *Lancet* 372:415–417. doi:10.1016/S0140-6736(08)61162-9
  48. Rollins BM, Burn M, Coakley RD, Chambers LA, Hirsh AJ, Clunes MT, Lethem MI, Donaldson SH, Tarran R (2008)  $A_{2B}$  adenosine receptors regulate the mucus clearance component of the lung's innate defense system. *Am J Respir Cell Mol Biol* 39: 190–197. doi:10.1165/rcmb.2007-0450OC
  49. Roos A, Boron WF (1981) Intracellular pH. *Physiol Rev* 61:296–434
  50. Salvi M, Sarno S, Marin O, Meggio F, Itarte E, Pinna LA (2006) Discrimination between the activity of protein kinase CK2 holoenzyme and its catalytic subunits. *FEBS Lett* 580:3948–3952. doi:10.1016/j.febslet.2006.06.031
  51. Sarno S, Salvi M, Battistutta R, Zanotti G, Pinna LA (2005) Features and potentials of ATP-site directed CK2 inhibitors. *Biochim Biophys Acta* 1754:263–270. doi:10.1016/j.bbapap.2005.07.043
  52. Schmidt M, Dekker FJ, Maarsingh H (2013) Exchange protein directly activated by cAMP (epac): a multidomain cAMP mediator in the regulation of diverse biological functions. *Pharmacol Rev* 65: 670–709. doi:10.1124/pr.110.003707
  53. Shah VS, Meyerholz DK, Tang XX, Reznikov L, Abou Alaiwa M, Ernst SE, Karp PH, Wohlford-Lenane CL, Heilmann KP, Leidinger MR, Allen PD, Zabner J, McCray PB Jr, Ostedgaard LS, Stoltz DA, Randak CO, Welsh MJ (2016) Airway acidification initiates host defense abnormalities in cystic fibrosis mice. *Science* 351:503–507. doi:10.1126/science.aad5589
  54. Shan J, Liao J, Huang J, Robert R, Palmer ML, Fahrenkrug SC, O'Grady SM, Hanrahan JW (2012) Bicarbonate-dependent chloride transport drives fluid secretion by the human airway epithelial cell line Calu-3. *J Physiol* 590:5273–5297. doi:10.1113/jphysiol.2012.236893

55. Shin JH, Son EJ, Lee HS, Kim SJ, Kim K, Choi JY, Lee MG, Yoon JH (2007) Molecular and functional expression of anion exchangers in cultured normal human nasal epithelial cells. *Acta Physiol (Oxf)* 191:99–110. doi:[10.1111/j.1748-1716.2007.01731.x](https://doi.org/10.1111/j.1748-1716.2007.01731.x)
56. Smith JJ, Welsh MJ (1992) cAMP stimulates bicarbonate secretion across normal, but not cystic fibrosis airway epithelia. *J Clin Invest* 89:1148–1153. doi:[10.1172/JCI115696](https://doi.org/10.1172/JCI115696)
57. Soleimani M, Greeley T, Petrovic S, Wang Z, Amlal H, Kopp P, Burnham CE (2001) Pendrin: an apical  $\text{Cl}^-/\text{OH}^-/\text{HCO}_3^-$  exchanger in the kidney cortex. *Am J Physiol Renal Physiol* 280:F356–F364
58. Sterling D, Brown NJ, Supuran CT, Casey JR (2002) The functional and physical relationship between the DRA bicarbonate transporter and carbonic anhydrase II. *Am J Physiol Cell Physiol* 283:C1522–C1529. doi:[10.1152/ajpcell.00115.2002](https://doi.org/10.1152/ajpcell.00115.2002)
59. Stewart AK, Chernova MN, Shmukler BE, Wilhelm S, Alper SL (2002) Regulation of AE2-mediated  $\text{Cl}^-$  transport by intracellular or by extracellular pH requires highly conserved amino acid residues of the AE2 NH2-terminal cytoplasmic domain. *J Gen Physiol* 120:707–722
60. Stewart AK, Kurschat CE, Vaughan-Jones RD, Shmukler BE, Alper SL (2007) Acute regulation of mouse AE2 anion exchanger requires isoform-specific amino acid residues from most of the transmembrane domain. *J Physiol* 584:59–73. doi:[10.1113/jphysiol.2007.136119](https://doi.org/10.1113/jphysiol.2007.136119)
61. Stewart AK, Shmukler BE, Vandorpe DH, Reimold F, Heneghan JF, Nakakuki M, Akhavan A, Ko S, Ishiguro H, Alper SL (2011) SLC26 anion exchangers of guinea pig pancreatic duct: molecular cloning and functional characterization. *Am J Physiol Cell Physiol* 301:C289–C303. doi:[10.1152/ajpcell.00089.2011](https://doi.org/10.1152/ajpcell.00089.2011)
62. Tarran R, Trout L, Donaldson SH, Boucher RC (2006) Soluble mediators, not cilia, determine airway surface liquid volume in normal and cystic fibrosis superficial airway epithelia. *J Gen Physiol* 127:591–604. doi:[10.1085/jgp.200509468](https://doi.org/10.1085/jgp.200509468)
63. Tarran R, Button B, Boucher RC (2006) Regulation of normal and cystic fibrosis airway surface liquid volume by phasic shear stress. *Annu Rev Physiol* 68:543–561. doi:[10.1146/annurev.physiol.68.072304.112754](https://doi.org/10.1146/annurev.physiol.68.072304.112754)
64. Tian Y, Kongsuphol P, Hug M, Ousingasawat J, Witzgall R, Schreiber R, Kunzelmann K (2011) Calmodulin-dependent activation of the epithelial calcium-dependent chloride channel TMEM16A. *FASEB J* 25:1058–1068. doi:[10.1096/fj.10-166884](https://doi.org/10.1096/fj.10-166884)
65. Treharne KJ, Xu Z, Chen JH, Best OG, Cassidy DM, Gruenert DC, Hegyi P, Gray MA, Sheppard DN, Kunzelmann K, Mehta A (2009) Inhibition of protein kinase CK2 closes the CFTR  $\text{Cl}^-$  channel, but has no effect on the cystic fibrosis mutant  $\Delta\text{F508}$ -CFTR. *Cell Physiol Biochem* 24:347–360. doi:[10.1159/000257427](https://doi.org/10.1159/000257427)
66. Venerando A, Ruzzene M, Pinna LA (2014) Casein kinase: the triple meaning of a misnomer. *Biochem J* 460:141–156. doi:[10.1042/BJ20140178](https://doi.org/10.1042/BJ20140178)
67. Xie M, Rich TC, Scheitrum C, Conti M, Richter W (2011) Inactivation of multidrug resistance proteins disrupts both cellular extrusion and intracellular degradation of cAMP. *Mol Pharmacol* 80:281–293. doi:[10.1124/mol.111.071134](https://doi.org/10.1124/mol.111.071134)