

## Acute stress and an electrolyte-imbalanced diet, but not chronic hypoxia, increase oxidative stress and hamper innate immune status in a Rainbow trout (Oncorhynchus mykiss) isogenic line

Leonardo J. Magnoni, Sara C. Novais, Ep Eding, Isabelle Leguen, Marco F. L. Lemos, Rodrigo O. A. Ozório, Inge Geurden, Patrick Prunet, Johan W. Schrama

### ▶ To cite this version:

Leonardo J. Magnoni, Sara C. Novais, Ep Eding, Isabelle Leguen, Marco F. L. Lemos, et al.. Acute stress and an electrolyte-imbalanced diet, but not chronic hypoxia, increase oxidative stress and hamper innate immune status in a Rainbow trout (Oncorhynchus mykiss) isogenic line. Frontiers in Physiology, 2019, 10, pp.1-14. 10.3389/fphys.2019.00453 . hal-02627473

## HAL Id: hal-02627473 https://hal.inrae.fr/hal-02627473

Submitted on 26 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License





# Acute Stress and an Electrolyte-Imbalanced Diet, but Not Chronic Hypoxia, Increase Oxidative Stress and Hamper Innate Immune Status in a Rainbow Trout (*Oncorhynchus mykiss*) Isogenic Line

Leonardo J. Magnoni<sup>1\*</sup>, Sara C. Novais<sup>2</sup>, Ep Eding<sup>3</sup>, Isabelle Leguen<sup>4</sup>, Marco F. L. Lemos<sup>2</sup>, Rodrigo O. A. Ozório<sup>1</sup>, Inge Geurden<sup>5</sup>, Patrick Prunet<sup>4</sup> and Johan W. Schrama<sup>3</sup>

<sup>1</sup> CIIMAR – Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Matosinhos, Portugal,
<sup>2</sup> MARE – Marine and Environmental Sciences Centre, ESTM, Instituto Politécnico de Leiria, Peniche, Portugal,
<sup>3</sup> Aquaculture and Fisheries Group, Wageningen Institute of Animal Sciences, Wageningen University, Wageningen, Netherlands,
<sup>4</sup> Laboratoire de Physiologie et Génomique des Poissons, Institut National de la Recherche Agronomique, Rennes, France,
<sup>5</sup> Nutrition Metabolisme Aquaculture (NuMeA)- Institut National de la Recherche Agronomique (INRA), Saint-Pée-sur-Nivelle, France

In aquaculture, fish may be exposed to sub-optimal rearing conditions, which generate a stress response if full adaptation is not displayed. However, our current knowledge of several coexisting factors that may give rise to a stress response is limited, in particular when both chronic and acute stressors are involved. This study investigated changes in metabolic parameters, oxidative stress and innate immune markers in a rainbow trout (Oncorhynchus mykiss) isogenic line exposed to a combination of dietary (electrolyteimbalanced diet, DEB 700 mEq Kg<sup>-1</sup>) and environmental (hypoxia, 4.5 mg O<sub>2</sub> L<sup>-1</sup>) challenges and their respective controls (electrolyte-balanced diet, DEB 200 mEq Kg<sup>-1</sup> and normoxia, 7.9 or mg O<sub>2</sub> L<sup>-1</sup>) for 49 days. At the end of this period, fish were sampled or subjected to an acute stressor (2 min of handling/confinement) and then sampled. Feeding trout an electrolyte-imbalanced diet produced a reduction in blood pH, as well as increases in cortisol levels, hepato-somatic index (HSI) and total energy content in the liver. The ratio between the lactate dehydrogenase (LDH) and isocitrate dehydrogenase (IDH) activities decreased in the liver of trout fed the DEB 700 diet, but increased in the heart, suggesting a different modulation of metabolic capacity by the dietary challenge. Several markers of oxidative stress in the liver of trout, mainly related to the glutathione antioxidant system, were altered when fed the electrolyteimbalanced diet. The dietary challenge was also associated with a decrease in the alternative complement pathway activity (ACH<sub>50</sub>) in plasma, suggesting an impaired innate immune status in that group. Trout subjected to the acute stressor displayed reduced blood pH values, higher plasma cortisol levels as well as increased levels of metabolic markers associated with oxidative stress in the liver. An interaction between

#### **OPEN ACCESS**

#### Edited by:

José Luis Soengas, University of Vigo, Spain

#### Reviewed by:

Lluis Tort, Autonomous University of Barcelona, Spain Bernardo Baldisserotto, Universidade Federal de Santa Maria, Brazil Saichiro Yokoyama, Kagoshima University, Japan

\*Correspondence:

Leonardo J. Magnoni Imagnoni@ciimar.up.pt

#### Specialty section:

This article was submitted to Aquatic Physiology, a section of the journal Frontiers in Physiology

Received: 15 November 2018 Accepted: 01 April 2019 Published: 24 April 2019

#### Citation:

Magnoni LJ, Novais SC, Eding E, Leguen I, Lemos MFL, Ozório ROA, Geurden I, Prunet P and Schrama JW (2019) Acute Stress and an Electrolyte- Imbalanced Diet, but Not Chronic Hypoxia, Increase Oxidative Stress and Hamper Innate Immune Status in a Rainbow Trout (Oncorhynchus mykiss) Isogenic Line. Front. Physiol. 10:453. doi: 10.3389/fphys.2019.00453

1

diet and acute stressor was detected for oxidative stress markers in the liver of trout, showing that the chronic electrolyte-imbalance impairs the response of rainbow trout to handling/confinement. However, trout reared under chronic hypoxia only displayed changes in parameters related to energy use in both liver and heart. Taken together, these results suggest that trout displays an adaptative response to chronic hypoxia. Conversely, the dietary challenge profoundly affected fish homeostasis, resulting in an impaired physiological response leading to stress, which then placed constraints on a subsequent acute challenge.

Keywords: rainbow trout, dietary imbalance, metabolic capacity, fish homeostasis, chronic hypoxia, stressors

#### INTRODUCTION

Farming fish involves following certain practices and procedures which may include handling, low water levels, confinement, and crowding (Conte, 2004). These procedures can act as stressors when habituation is not present (Pickering, 1993; Wendelaar Bonga, 1997; Bratland et al., 2010; Nilsson et al., 2012). Additionally, chronic sub-optimal rearing conditions may also give rise to stress responses if full adaptation is not displayed. However, the information available on existing conditions that may act as stressors during rearing, along with their interactions, is very limited. This is an area of increasing interest, as stressors have been linked to a reduction in both growth performance and immune condition; hampering health and welfare in fish (Portz et al., 2006).

One of the known existing factors of stress in cultured fish species is the persistent low oxygen concentration (chronic hypoxia). Nevertheless, its consequences are still poorly documented. In addition, long-term dietary imbalances affect fish homeostasis and energy balance. The induction of stress by dietary imbalances have been described in fish, including responses linked to partial or total replacement of fish meal and fish oil by alternative sources (Montero et al., 2003, 2008, 2010; Gómez-Requeni et al., 2004). Previous studies have shown that growth performance, feed intake and nutrient digestibility are altered when fish are fed electrolyte imbalanced diets (Dersjant-Li et al., 1999, 2000; Saravanan et al., 2013b; Magnoni et al., 2016). However, the induction of stress by this type of dietary imbalance is yet to be investigated in fish.

The dietary electrolyte balance (DEB) is defined as the sum of the mineral cations minus the sum of mineral anions present in the diet. Differences in DEB may occur when feed ingredients containing different quantities of cations (Na, K, Ca, and Mg) and anions (Cl and P) are included in the diet formulation (Patience and Wolynetz, 1990). It was previously shown that rainbow trout (*Oncorhynchus mykiss*) fed an electrolyte imbalanced diet (DEB 700) were energetically less efficient than those fed an electrolyte balanced diet (DEB 200). In spite of this change in energy balance, feed intake and growth performance were not affected (Magnoni et al., 2018).

The objective of this study was to determine whether electrolyte imbalanced diet may affect the ability of trout to cope with chronic hypoxia conditions. The assessment of physiological impacts of combined stress factors (diet and hypoxia) was analyzed in fish before and after applying an acute stressor (handling/confinement). This acute stress condition is commonly present in aquaculture, and was used to determine the physiological coping ability of the fish.

A set of plasma parameters related to stress and innate immune response were determined, as they are commonly associated with fish welfare. Additionally, several parameters were analyzed in liver and heart, since their energy use are altered when subjected to stress conditions (Wendelaar Bonga, 1997; Hermes-Lima et al., 2001), including changes in oxidative stress and metabolic response. An isogenic heterozygous family of rainbow trout was used as fish model, due to its genetic uniformity providing low intra-specific variability and high reproducibility.

#### MATERIALS AND METHODS

#### **Fish and Housing**

The isogenic heterozygous family of rainbow trout (R23) obtained by crossing two homozygote isogenetic lines was produced by INRA/PEIMA (France) experimental fish facilities (Sadoul et al., 2015). Fish were housed in the Aquatic Metabolic Unit (AMU) tanks of Aquaculture and Fisheries group, Wageningen University, The Netherlands. Thirty rainbow trout (115.2  $\pm$  2.0 g) were randomly assigned to each of the twelve experimental tanks (200 L). The tanks were connected to a water recirculation system consisting of a trickling filter, an oxygenation unit, a sump, a drum filter (Hydrotech 500<sup>(6)</sup>), and a cooling/heating system for maintaining uniform water quality throughout the study. The oxygenation unit maintained the DO levels by injecting oxygen into the water and was facilitated with separate automatic probes for the detection of water flow and oxygen consumption. Water temperature was set at  $14 \pm 1^{\circ}$ C. Photoperiod was maintained at 12:12 (Light: Dark) with daybreak set at 07:00 h.

#### **Experimental Diets and Feeding**

Two isoproteic (45% DM) and isoenergetic (22 kJ gDM<sup>-1</sup>) diets, floating pellets of 4 mm, were extruded by Research Diet Services (Wijk bij Duurstede, The Netherlands). Diets, upon arrival to the AMU at University of Wageningen, were stored in a room under controlled conditions during the trial. The two diets were formulated to provide a contrast in electrolyte content (DEB); 200 or 700 mEq Kg<sup>-1</sup>. This difference was created by adding different amounts of  $Na_2CO_3$  and diamol (inert filler) in the diets. Fish were fed the experimental diets to apparent satiation, twice a day for 49 days. Ingredients and proximate composition of the experimental diets are included in **Supplementary Table S1**.

#### **Experimental Conditions**

The difference in DO levels was induced by adjusting the rate of water flow into the tanks as described by Saravanan et al. (2013a). The water volume was kept constant at 200 L in all tanks. For the normoxia groups, the rate of water inflow into each tank was kept at 7.2  $\pm$  0.0 L min<sup>-1</sup> (mean  $\pm$  SEM) with a mean water DO level of  $10.2 \pm 0.1$  mg O<sub>2</sub> L<sup>-1</sup>. The DO level in the outflowing water remained at 7.9  $\pm$  0.1 mg O<sub>2</sub> L<sup>-1</sup>. Hypoxia conditions were created by gradually reducing the rate of water inflow into the tank (2.2  $\pm$  0.0 L min<sup>-1</sup>) with a DO level of 10.2  $\pm$  0.1 mg O<sub>2</sub> L<sup>-1</sup> for the first 3 days after the start of the experiment. DO level in the outflowing water remained at  $4.5 \pm 0.1 \text{ mg O}_2 \text{ L}^{-1}$  and was kept equal in all the hypoxic tanks. The DO level in the water outlet was considered to be equivalent to ambient DO level of the tanks, as differences between both DO levels (outlet versus inside the tank) shown in several previous experiments were negligible ( $<0.2 \text{ mg O}_2 \text{ L}^{-1}$ , e.g., Tran-Duy et al., 2008). The force of the evenly distributed water supplied by the inlet in each tank together with the swimming activity prevented the occurrence of water stratification within the tanks. The DO level applied in the hypoxia treatment is recognized as an environmental challenge, with the value decided based on the reported incipient DO level of 6.0 mg  $O_2 L^{-1}$  for rainbow trout (Pedersen, 1987).

#### **Experimental Design**

Employing a 2  $\times$  2 design (DEB diets and DO levels), the experimental tanks (12) were divided into three experimental blocks, with four tanks randomly assigned within each of three blocks (n = 3 tanks per treatment). Fish were not fed on the day prior to sampling.

Fish from each experimental group were divided into 2 sub-groups for the application of a standardized handling/ confinement protocol (acute stress) and posterior sampling. A control sub-group in which three fish per tank (9 per treatment) were sampled by reducing the exposure to potential stressors. Subsequently, a group of 3 fish per tank were netted and exposed to a confinement stress (2 min at a density of 200 kg/m<sup>3</sup>) and transferred back to their original tank (empty) for 1 h. Then fish from both experimental sub-groups were netted and euthanized for sampling. A sampling effect was prevented by sampling the tanks following the order tanks were installed in the room, starting with the control sub-group and with water from sampled tanks on flow through to prevent water re-entering the RAS circulation. A lethal dose of 2-phenoxy ethanol solution (1 mg/L) was used in all the sampling procedures.

Blood was drawn from the caudal region with a heparinized syringe. Blood pH was immediately measured (pH meter, WTW pH 320; pH electrode, WTW SenTix Sp). The duration of procedure (blood withdrawal and pH measurement) was strictly standardized to 1 min for all fish to minimize blood pH variation (Saravanan et al., 2013b). Blood was centrifuged at 3000 g for 20 min at 4°C, and plasma samples were frozen and stored for later analyses. Fish, liver, and heart were weighed and sampled. Samples were frozen in liquid nitrogen and stored at  $-80^{\circ}$ C for further analyses.

#### **Plasma Metabolite Content**

Lactate concentration in plasma was quantified using a commercial kit (LO-POD, Spinreatc, Sant Esteve de Bas, Spain). Glucose concentration in plasma was quantified using a commercial kit (GOD-POD, Spinreatc, Sant Esteve de Bas, Spain). An ELISA kit based on the competitive link between cortisol and related monoclonal antibodies was used to quantify cortisol levels in plasma of rainbow trout (RE 52061, IBL International, Hamburg, Germany). Validation of the kit for determination of cortisol in plasma of trout was performed. The intra- and inter-assay coefficient of variation in plasma samples were <9% and <10%, respectively. The linearity showed an  $r^2$ -value of 0.96. Results from this validation indicated the suitability for the use of this kit to quantify changes in cortisol levels in rainbow trout. All measurements were performed in triplicates, following the recommendations provided by the manufacturers.

#### **Energy Balance in Tissues**

Total protein, glycogen and lipids contents were measured spectrophotometrically in the tissue homogenates according to De Coen and Janssen (1997, 2003). Liver and heart tissues were homogenized in phosphate buffer (1/10 vol, 0.1 M, pH = 7.4). Proteins in the homogenates were precipitated with the addition of cold 15% trichloroacetic acid and incubated at -20°C for 10 min. After a centrifugation at 1000 g for 10 min, the resulting supernatant was used for glycogen quantification by adding phenol (5%) and H<sub>2</sub>SO<sub>4</sub> (conc.). After 30 min incubation at 20°C, glucose was quantified by measuring the absorbance at 490nm. The protein pellet obtained after trichloroacetic acid precipitation was resuspended in NaOH (1N), incubated at 60°C for 30 min and then neutralized with HCl (1.67 M). The resuspended pellet was used to quantify total protein content by the Bradford (1976) method measuring absorbance at 600 nm, using bovine serum albumin as standard.

As for lipids extraction, chloroform, methanol, and Mili-Q water were added to homogenates in a 2:2:1 proportion, respectively. The organic phase was separated after centrifugation, treated with  $H_2SO_4$  at 200°C, and used to quantify total lipid content at 400 nm, using tripalmitin as standard. Total protein, glycogen, and lipid content of each sample was expressed as mg per g of wet weight tissue. Total energy content was calculated as the sum of the protein, glycogen and lipids contents in each tissue, transformed into energetic equivalents using values for the enthalpy of combustion (24 kJ/g proteins, 17.5 kJ/g carbohydrates, and 39.5 kJ/g lipids), as described by De Coen and Janssen (1997), and expressed as kJ g<sup>-1</sup> wet weight wet tissue.

An aliquot of homogenate was centrifuged for 5 min, at 3000 g ( $4^{\circ}$ C) and the supernatant was used for lactate dehydrogenase (LDH) and isocitrate dehydrogenase (IDH) activity measurements. LDH was measured following the method described by Vassault (1983) and adapted to microplate. IDH

was measured following the method described by Ellis and Goldberg (1971) with the adaptations of Lima et al. (2007). For the normalization by protein content, Bradford (1976) method was applied to quantify protein in the supernatant fraction, using bovine *G*-globulin as standard and reading absorbance at 600 nm. Energy consumption rate (Ec: measured through the electron transport system -ETS- activity) was performed according to the procedure described by De Coen and Janssen (1997, 2003). Measurements were performed at 25°C in triplicates using proper reaction blanks in a synergy H1 Hybrid Multi-Mode microplate reader (Biotek<sup>®</sup> Instrument, Vermont, United States).

#### **Oxidative Stress Markers in Liver**

Liver and heart samples were homogenized in phosphate buffer (1/10 vol., 0.1 M pH 7.4). Enzymatic analyses were all carried out with the reaction mixtures and homogenate dilution established in preliminary tests. Protein concentration was assayed in homogenates using bovine serum albumin as standard (Bradford, 1976). Lipid peroxidation (LPO) was determined by quantifying the presence of formed thiobarbituric acid reactive substances (TBARS) (Ohkawa et al., 1979). Glutathione reductase (GR) (EC1.8.1.7) and glutathione peroxidase (GPx) (EC 1.11.1.9.) were evaluated based on NADPH (Sigma, Portugal) oxidation at 340 nm (Mohandas et al., 1984; Cribb et al., 1989). Total glutathione (TG) and oxidized glutathione (GSSG) were evaluated by the formation of 5-thio-2-nitrobenzoic acid at 412 nm (Baker et al., 1990). Reduced glutathione (GSH) was calculated as the difference between TG and GSSG. Changes in absorbance were measured at 22°C in a Power-Wave<sup>TM</sup> microplate spectrophotometer (BioTek Instruments), and reactions were performed in triplicates. Substrate was omitted in reaction blanks and background activity was subtracted from that measured in the presence of substrate.

#### **Innate Immune Parameters in Plasma**

Immune status was assessed through the determination of key parameters, namely lysozyme, peroxidase and the alternative complement pathway (ACH<sub>50</sub>) activities. Lysozyme activity was evaluated according to Lie et al. (1986), employing hen egg white lysozyme (Sigma, Germany) as standard and *Micrococcus lysodeikticus* (0.5 mg mL<sup>-1</sup>; 0.05 M sodium phosphate buffer; pH 6.2) as bacterial suspension. Peroxidase activity (U mL<sup>-1</sup> plasma) was determined based on the methodology described by Quade and Roth (1997). ACH<sub>50</sub> activity was analyzed according to Sunyer and Tort (1995) using a concentration of  $2.8 \times 10^8$  cells mL<sup>-1</sup> rabbit erythrocytes. The reciprocal of the plasma dilution giving 50% haemolysis of erythrocytes equals to one ACH<sub>50</sub> unit.

#### **Measurements and Calculations**

Weight gain (WG) was calculated as:

 $WG(g) = BW_f - BW_i$ 

where  $BW_i$  and  $BW_f$  are the initial and the final body weight of fish in the trial, respectively.

Feed intake (FI) was calculated as: FI (g DM) = FI<sub>TOT</sub>/ n  $\times$  t,

where  $FI_{TOT}$  is the total FI per tank over the experimental period corrected for dead fish and uneaten feed, n is the number of fish per tank and t is the experimental period (days).

Uneaten feed (pellets) were collected on the surface at the end of each feeding period (1 h). Pellets remaining on the bottom of the tank were collected by a decantation unit. All the uneaten pellets were counted and the amount calculated by taking and average weight of pellets in each diet production lot. The amount of feed was registered daily and accounted for in the feed intake calculation. No mortalities were recorded during the trial, except one fish fed the DEB 200 diet in hypoxia (98.9% survival).

Feed conversion ratio (FCR) was calculated as:

$$FCR = FI/(BW_{\rm f} - BW_{\rm i})$$

Specific growth rate (SGR) was calculated as:

 $SGR(\%BWday^{-1}) = (lnBW_{\rm f} - lnBW_{\rm i})/time(days) \times 100$ 

The hepato-somatic index was calculated as:

$$HSI(\%) = 100 \times [liverweight(g)/BW(g)]$$

The cardio-somatic-index was calculated as:

$$CSI(\%) = 100 \times [heartweight(g)/BW(g)]$$

The effects of DEB 200 and DEB 700 on growth performance and feed intake of this trout isogenic line subjected chronic hypoxia or normoxia has been published in Magnoni et al. (2018). However, to assist with the interpretation of the results presented in the current study, growth performance and feed intake parameters were included as **Supplementary Table S2**.

#### **Statistical Analysis**

Final body weight (BW<sub>f</sub>), weight gain (WG), and specific growth rate (SGR) were analyzed for the effects of the diet and DO levels along with their interactions by a two-way ANOVA analysis (triplicate tanks). All the other parameters in fish were analyzed for the effects of diet, DO levels and acute stress, along with their interactions by a three-way ANOVA analysis (n = 9). When differences in mean values were detected by the ANOVA analysis, *post hoc* tests were applied (Holm-Sidak method). In all these analyses, data sets were tested for normality (Shapiro–Wilk) and for equal variance (Levene's test), and if these assumptions were not met, data transformation was applied (Natural logarithm). Statistical analysis was made using SigmaPlot 12.0 (Systat Software, Inc. 2011) and significant differences were considered for P < 0.05.

#### RESULTS

#### Metabolic Markers in Blood and Plasma

Blood pH remained similar in fish reared at different DO levels (**Table 1**). Feeding trout an electrolyte-imbalanced diet (DEB

TABLE 1 | Effect of dietary electrolyte balance (DEB), dissolved oxygen levels, and acute stress on blood parameters of rainbow trout.

		DEE	3 200			DEB	700		I	Facto	ors	In	terac	ctions	6
	Nor	moxia	Нур	ooxia	Norn	noxia	Нур	ooxia							
	с	S	с	S	С	S	с	S	D	0	s	DOS	DO	DS	os
pН	$7.06 \pm 0.2$	$6.99\pm0.02$	$7.08 \pm 0.02$	$7.02 \pm 0.02$	$7.0 \pm 0.01$	$6.95\pm0.03$	$7.07 \pm 0.02$	$6.92 \pm 0.03$	**	ns	**	ns	ns	ns	ns
Cortisol	$6.63 \pm 1.95$	$28.00\pm4.88$	$6.58 \pm 1.84$	$34.88 \pm 8.50$	$11.66\pm1.80$	$41.62\pm4.52$	$8.55\pm1.68$	$31.84\pm6.35$	*	ns	**	ns	ns	ns	ns
Lactate	$0.26\pm0.07$	$0.45\pm0.11$	$0.31\pm0.03$	$0.65\pm0.13$	$0.27\pm0.06$	$0.60\pm0.10$	$0.29\pm0.07$	$0.63\pm0.05$	ns	ns	**	ns	ns	ns	ns
Glucose	$5.40\pm0.31$	$7.22\pm0.60$	$5.39\pm0.45$	$6.83\pm0.47$	$5.34\pm0.30$	$7.29\pm0.16$	$5.31\pm0.44$	$6.32\pm0.47$	ns	ns	**	ns	ns	ns	ns

Plasma cortisol levels in ng mL<sup>-1</sup>. Plasma lactate and glucose concentrations in mM. Values are mean of fish (n = 9)  $\pm$  SE. C, Control; S, Acute stress. Three-way ANOVA analysis results. D, Diet; O, dissolved oxygen levels; DOS, DO, DS, and OS, Interactions; ns, not significant P > 0.1; \*P < 0.05; \*\*P < 0.01.

700) or subjecting the fish to acute stress induced a reduction on blood pH (P < 0.001).

Plasma cortisol levels were increased in trout subjected to acute stress or when fed an electrolyte-imbalanced diet (P < 0.001 and P < 0.005, respectively). However, cortisol levels remained unaltered when fish were subjected to chronic hypoxia. Plasma lactate and glucose concentrations were increased in fish subjected to acute stress (P < 0.001) but remained similar when fed diets with varied DEB or when maintained at different DO levels.

#### **Energy Use in Tissues**

The hepato-somatic index (HSI) of trout was 16% higher in fish fed the DEB 700 diet (P < 0.05) (**Table 2**). All the parameters related to energy content measured in liver of trout remained similar at different DO levels. However, total energy content in the liver was increased by 6% in fish fed the electrolyte-imbalanced diet (P < 0.05).

The cardiac-somatic index (CSI) of trout was reduced by 6% in fish subjected to hypoxia (P < 0.05) (**Table 3**). The energy stored as glycogen in the heart of trout was increased by 27% in fish reared in hypoxia (P < 0.01) and decreased by 26% when subjected to acute stress (P < 0.01).

A decrease in IDH activity (P < 0.01), with an increase in LDH to IDH activity ratio (P < 0.05) was detected in the liver of trout reared in hypoxia (**Table 4**). On this tissue a decrease in LDH activity (P < 0.05), as well as in the LDH to IDH activity ratio (P < 0.01) were detected in trout fed the DEB 700 diet. However, all these metabolic parameters remained similar when acute stress was applied.

On the other hand, enzyme activities and parameters related to energy-use in the heart of trout remained similar in fish reared at different DO levels (**Table 5**). Both electrolyte-imbalanced diet and acute stress increased LDH to IDH activity ratio in the heart of trout (P < 0.05). In addition, acute stress decreased the IDH activity measured in the trout's heart (P < 0.05).

#### **Oxidative Stress Markers in Liver**

All the oxidative stress parameters analyzed in this tissue remained similar in fish reared at different DO levels (**Table 6**). However, in trout fed with the DEB 700 diet, a decrease in the GPX activity (P < 0.05), as well as an increase in the TG and GSSG levels (P < 0.05) were observed. In addition, acute stress

induced increases in GR activity (P < 0.05) and GSH content (P < 0.05) in trout's liver.

#### Innate Immune Status

All the parameters analyzed in plasma remained similar in fish reared at different DO levels (**Table 7**). However, feeding trout an electrolyte-imbalanced diet produced a decrease in ACH<sub>50</sub> levels (P < 0.05). Acute stress increased lysozyme activity (P < 0.05).

#### Interactions Between Experimental Factors

No interaction between dietary treatment and DO levels was observed in parameters related to fish performance (**Supplementary Table S2**). In addition, no interactions were observed between the factors investigated when analyzing markers related to stress, metabolism or innate immune status in plasma (**Tables 1**, 7, respectively). Furthermore, the energy content of the heart of trout did not display any interaction between the experimental factors investigated (**Table 3**).

By contrast, the total energy content of the liver was affected by the diet and the DO levels (P < 0.05), as well as by the DO levels and the acute stressor (P < 0.01) (**Table 2**). The diet also influenced the energy consumption (Ec) value in the liver when the acute stressor was considered (P < 0.05). Additionally, the dietary treatment influenced DO levels (P < 0.05), acute stressor (P < 0.01), and all the three factors together (P < 0.05) when the marker of oxidative stress, GPx activity, was analyzed in liver (**Table 6**). The three factors together displayed significant interaction when several markers of energy use in heart of trout were investigated (LDH, IDH, LDH/IDH, Ec) (**Table 5**). DO levels and acute stress revealed significant interactions when glycogen content was analyzed in liver (P < 0.05) (**Table 2**).

#### DISCUSSION

This study analyzed the prolonged effects of dietary electrolyteimbalance and environmentally low DO levels, along with an acute stressor (handling/confinement), on various functions including: (i) acute stress responses at the level of plasma or cellular stress, (ii) tissue metabolism (liver and heart), and (iii) innate immunity. By implementing such approach, the aim

		DE	B 200			DEB	200			actors		5	teract	suo	
	Norm	bixia	Hy	oxia	Norr	noxia	Hy	poxia							
	U	S	υ	S	U	S	U	S	۵	0	s	DOS	DO	DS	SO
HSI	1.42 ± 0.22	1.31 ± 0.15	1.25 ± 0.17	1.39 ± 0.23	1.56 ± 0.18	1.73 ± 0.19	1.29 ± 0.13	1.81 ± 0.32	*	SU	SU	SU	SU	su	ns
Protein	$61.87 \pm 5.65$	$61.50 \pm 2.61$	$61.82 \pm 7.63$	$54.51 \pm 4.56$	$59.35 \pm 3.12$	$58.09 \pm 3.34$	$61.54 \pm 3.92$	$53.68 \pm 6.41$	SU	ns	ns	SU	SU	SU	ns
Lipids	$24.47 \pm 1.95$	$22.10 \pm 1.17$	$25.33 \pm 1.86$	$28.06 \pm 2.89$	$26.37 \pm 2.80$	$26.86 \pm 3.31$	$23.43 \pm 0.73$	28.83 ± 3.46	SU	ns	ns	SU	SU	ns	ns
Glycogen	$95.18 \pm 14.62$	$74.50 \pm 8.59$	88.15 ± 14.31	$103.74 \pm 10.19$	$109.81 \pm 6.32$	$105.52 \pm 10.08$	$83.47 \pm 7.69$	$112.72 \pm 10.72$	SU	ns	ns	SU	SU	SU	*
Total energy	4.12 土 0.13	$3.65 \pm 0.14$	$4.03 \pm 0.16$	4.23 ± 0.17	4.39 ± 0.13	4.30 ± 0.21	$3.86 \pm 0.82$	$4.40 \pm 0.16$	×	ns	ns	ns	×	ns	* *

was to provide a refined tool to evaluate stress response in fish. The prolonged conditions implemented, DEB 700 diet and hypoxia, were designed to produce deep physiological challenges in trout. These conditions were chosen over other conditions, such as repeated physical handling, as this may increase the likelihood of skin, mucus or fin damages, and could make interpretation of chronic stress responses difficult, as this may increase the likelihood of skin, mucus or fin damages, and could make interpretation of chronic stress responses difficult due to co-factors that could be present (e.g., opportunistic pathogens), leading to erroneous conclusions.

# The Electrolyte-Imbalanced Diet Acts as a Chronic Stressor in Rainbow Trout

Growth performance and FI remains unchanged when trout was fed an electrolyte-imbalanced diet at satiety for 49 days (Supplementary Table S2). In spite of the lack of changes in these parameters, markers of energy use, oxidative stress response, and innate immune status were all altered in trout. In particular, the DEB 700 diet induced increases in HSI and total energy content in the liver of the trout. These changes could be related to an energetic constrain imposed by electrolyte-imbalanced diet, as mechanisms managing acid-base homeostasis may involve an increased energy expenditure. In fact, feeding a DEB 700 diet on this isogenic trout line has shown to produce a 65% increase in the metabolizable energy requirements for maintenance, suggesting higher energy demand to maintain acid-base balance in that group (Magnoni et al., 2018). Such scenario is supported by a postprandial decrease in blood pH of trout fed the DEB 700 diet (Table 1). An alteration on the energy balance is shown by the increase in the total energy content stored in liver of trout fed the DEB 700 diet, possibly related as well to changes in the relative importance of aerobic and anaerobic metabolism to provide energy to this organ.

As any stress condition carries additional energetic requirements, biomarkers on anaerobic and aerobic pathways of energy production may provide important indications in terms of global energetic demands toward a stress condition. In particular, LDH activity has a key role in anaerobic pathways and changes on this enzyme activity are usually associated to increased energy demand, e.g., due to stressful conditions (Dahlhoff, 2004). On the other hand, IDH, an enzyme involved in energy production through the aerobic pathways, catalyzes a key step of the Krebs cycle, and is also involved in antioxidant defenses, where it is crucial for the regeneration of NADPH required for glutathione conjugation pathways (Jo et al., 2001; Lee et al., 2002). For example, in juvenile European seabass (Dicentrarchus labrax) an increased IDH activity was detected in liver when exposed to high temperature, which has been linked to a stress response (Almeida et al., 2015). An increased reliance in aerobic metabolism (low LDH/IDH activities ratio) usually enables a more efficient energy metabolism than the anaerobic pathways. In our study, LDH/IDH activities ratio was decreased in liver of trout by the electrolyte-imbalanced diet. In contrast, the heart of trout displayed an increase in LDH/IDH activities ratio when fed TABLE 3 | Effect of dietary electrolyte balance (DEB), dissolved oxygen levels and acute stress on cardiac-somatic index (CSI), protein, lipids, glycogen, and energy content in heart of rainbow trout.

		DEE	3 200			DEB	3 700		F	Facto	ors	In	terac	ction	s
	Norn	noxia	Нур	oxia	Norr	noxia	Нур	oxia							
	С	S	С	S	С	S	С	S	D	ο	s	DOS	DO	DS	os
CSI	$0.15\pm0.01$	$0.15\pm0.01$	$0.13\pm0.01$	$0.14 \pm 0.01$	0.14 ± 0.01	0.15 0.01	$0.13\pm0.01$	0.14 ± 0.01	ns	*	ns	ns	ns	ns	ns
Protein	$57.23 \pm 4.17$	$51.06\pm2.61$	$51.47 \pm 2.24$	$53.57\pm3.98$	$55.29\pm3.61$	$52.16\pm4.72$	$46.58\pm3.76$	$51.51 \pm 4.27$	ns	ns	ns	ns	ns	ns	ns
Lipids	$23.93\pm3.31$	$17.83 \pm 1.28$	$18.64\pm2.02$	$21.04 \pm 1.94$	$17.92 \pm 1.24$	$19.63\pm1.49$	$16.90\pm1.66$	$18.59 \pm 1.55$	ns	ns	ns	ns	ns	ns	ns
Glycogen	$4.35\pm0.51$	$2.79\pm0.27$	$4.80\pm0.22$	$4.34\pm0.40$	$4.12\pm0.32$	$2.61\pm0.34$	$4.57\pm0.32$	$4.20\pm0.56$	ns	**	**	ns	ns	ns	ns
Total energy	$2.40\pm0.19$	$1.98\pm0.92$	$2.25\pm0.11$	$2.38\pm0.14$	$2.11\pm0.10$	$2.07\pm0.16$	$1.87\pm0.13$	$2.04\pm0.16$	ns	ns	ns	ns	ns	ns	ns

Protein, lipids, and glycogen expressed as mg  $g^{-1}$  weight wet tissue. Total energy content expressed as kJ  $g^{-1}$  weight wet tissue. Values are mean of fish (n = 9)  $\pm$  SE. C, Control; S, Acute stress. Three-way ANOVA analysis results. D, Diet; O, dissolved oxygen levels; DOS, DO, DS, and OS, Interactions; ns, not significant P > 0.1; \*P < 0.05; \*\*P < 0.01.

the electrolyte-imbalanced diet, indicating that the metabolic capacity of liver and heart are modulated differently when subjected to the dietary challenge.

Key components of the glutathione system keeping redox balance, including GPX activity, TG, and GSSG levels, were all altered by diet in the liver of trout. In particular, GPx activity has been suggested as a reliable marker of stress in fish subjected to prolonged dietary or environmental challenges (Winston, 1991; Martínez-Álvarez et al., 2005; Sitjà-Bobadilla et al., 2005; Fontagné-Dicharry et al., 2018). Therefore, this study was able to confirm importance of this stress marker in the liver of trout subjected to a prolonged dietary challenge.

Haemolytic activity due to alternative complement pathway, measured as ACH<sub>50</sub>, is interpreted by numerous authors as a sign of a prompter innate immune system, improving the resistance to pathogens (Chiu et al., 2008; Biller-Takahashi et al., 2012). It is known that ACH<sub>50</sub> activity in fish changes with stress conditions (Boshra et al., 2006). Furthermore, decreased ACH<sub>50</sub> activity has been linked to chronic stress produced by crowding in gilthead seabream (Sparus aurata) juveniles (Montero et al., 1999b). Trout fed the DEB 700 diet displayed a decreased ACH<sub>50</sub> activity in plasma, which reinforces the conclusion that the disruption on fish homeostasis generated by the electrolyte-imbalanced diet could be evoking a stress response. Fish are able to mount successful and robust innate responses (Tort et al., 2003), such as the measured in this study. Previous studies have shown that the innate immune response in fish is altered by chronic crowding stress (Rotllant et al., 1997; Montero et al., 1999a) and by a combination of dietary supplementations and chronic crowding stress (Montero et al., 1999b). However, future studies should investigate the response of adaptive immune parameters to combined stressors such as those implemented in this study.

#### Rainbow Trout Shows a Full Adaptive Response to Chronic Hypoxia

Hypoxia tolerance appears to be variable in close phylogenetic related species of fish (Mandic et al., 2009), also displaying marked changes during development (Ishibashi et al., 2005). In this sense, the heterozygous isogenic line of rainbow trout used in this study, reared for many generations, provides a unique model with low intraspecific variability, adequate to test the effect of chronic hypoxia. This study is a follow-up of a previous one (Magnoni et al., 2018), in which it was shown that both feed intake (FI) and growth performance were decreased when this trout line was exposed to chronic hypoxia. In particular, rainbow trout maintained under hypoxia (4.5  $\pm$  0.1 mg O<sub>2</sub> L<sup>-1</sup>) reduced their feed intake and specific growth rate by 33 and 22%, respectively (Supplementary Table S2). In a previous study, Glencross (2009) also showed a decrease in FI in rainbow trout exposed to hypoxia (28 days at 4.29 mg  $O_2$  L<sup>-1</sup>). Therefore, the decrease in FI during chronic hypoxia detected in the present study is probably not linked to chronic stress, but more likely to a reduction in appetite triggered by the inhibition of food-anticipatory behavior, as has been suggested by previous studies (Glencross, 2009; Folkedal et al., 2012a,b). Corticotropinreleasing factor in the forebrain appears to mediate sustained anorexia during chronic hypoxia (Bernier and Craig, 2005), suggesting that this mechanism may be an essential energysaving strategy due to the reduction in fish maintenance energy. Also suppression of appetite may save energy by reducing the postprandial increase in O2 consumption associated with digestion, absorption, transformation, and storage of nutrients (Jobling, 1981, 1993). In this study, we cannot completely rule out the possibility that chronic hypoxia could act as a chronic stressor in trout, inhibiting feed intake by altering feeding behavior. However, as will be discussed in the next section, we propose that this may not be the case for this study, since the cortisol response was not altered after chronic low DO levels, suggesting that trout is adapted to these particular conditions.

There is increasing evidence that rearing fish under intensive aquaculture conditions may alter their cardiovascular physiology (Gamperl and Farrell, 2004). The DO level applied in the hypoxia treatment during this study (4.5 mg O<sub>2</sub> L<sup>-1</sup>) is recognized as an environmental challenge, with the value decided based on the critical DO level for feed consumption and for both growth rate and feed conversion efficiency (6 and 7 mg O<sub>2</sub> L<sup>-1</sup>, respectively) for rainbow trout (Pedersen, 1987). However, a previous study showed that rainbow trout reared at hatcheries, with sub-optimal water quality conditions, appear to have a remarkable degree of tolerance to short-term severe hypoxia, which has been

	DEB	3 200			DEE	3 700		Facto	s	Inte	eracti	ons	Interactions
-	Normoxia	Hype	oxia	Norn	noxia	Hyp	oxia						
U	S	O	S	υ	S	O	S	0	s	l soc	0	SC	) SO DO SO
LDH 2699.12 ± 27. IDH 84.64 ± 4.0	4.64 3349.83 ± 437.29 5 85 80 ± 8 50	2990.73 ± 298.11 70.13 ± 4.08	3092.40 ± 315.63 74.58 ± 4.00	2287.73 ± 281.52 04.00 ± 3.82	2579.46 ± 204.78 03.10 ± 4.61	2754.66 ± 114.60 80.05 ± 3.12	2606.28 ± 297.18 76 34 ± 4 20	* US	SL 3	SU 30	SU SU	SU SU	SU SU SU
LDH/IDH 32.43 ± 3.4	8 39.30 ± 3.26	43.01 ± 3.18	42.69 ± 4.69	25.08±3.64	28.20 ± 2.58	$34.83 \pm 1.96$	33.56 ± 2.89	* 20 *	su su	e su	21	s su	I SU SU SU
Ec 211.37 ± 23.	23 175.51 ± 12.78	268.48 ± 28.93	182.61 ± 22.62	183.38 ± 13.03	$205.08 \pm 15.22$	199.70 土 18.66	177.93 ± 22.04	ns ns	SU	ns	ns	×	ns ns *
9 fish ≟ SEM. C, Control; Pfish ≟ SEM. C, Control; TABLE 5   Effect of dietary	DH) and iso-citrate dehyo S, Acute stress. Three-we electrolyte balance (DEB)	trogenase (IDH) activi ay ANOVA analysis re: ay ANOVA analysis re: dissolved oxygen le	ties are expressed as sults; D, Diet; O, diss sults, and acute stress	nmol min <sup>-1</sup> trg <sup>-1</sup> t olved oxygen levels; son energy use in he	issue. Energy consu. DOS, DO, DS, and i and i rainbow trout.	mption value (Ec) is e OS, Interactions: ns, ı	xpressed as mJ mg tot significant P > (	7 <sup>−1</sup> weight v 0.1; *P < 0.1	vet tissu 35, ** P	e. Valu < 0.01	• •	e me	e. Values are mex < 0.01.
	DEB	200			DEB 700		Ĕ	actors		Inter	actior	sr	Interactions

									'					2	
	Norn	noxia	Hyp	oxia	Norm	loxia	Hyp	oxia							I
	U	S	υ	S	U	S	U	S	۵	0	s DO	S	ă	٥ س	s
LDH	$596.40 \pm 55.20$	506.88 ± 41.59	736.26 ± 79.99	580.91 ± 53.58	596.66 ± 37.47	627.02 ± 42.02	556.57 ± 42.57	676.77 ± 45.19	ns	SU	su su	su	*	Ċ	S
HQI	$452.34 \pm 28.98$	380.32 ± 22.44	$443.31 \pm 23.85$	$427.57 \pm 20.33$	$440.96 \pm 16.12$	$364.53 \pm 30.59$	$390.43 \pm 19.87$	$407.13 \pm 17.12$	SU	ns	*	su	SU SU	*	
LDH/IDH	$1.32 \pm 0.08$	$1.34 \pm 0.08$	$1.52 \pm 0.09$	$1.35 \pm 0.08$	$1.35 \pm 0.07$	$1.78 \pm 0.12$	$1.42 \pm 0.06$	$1.67 \pm 0.10$	*	ns	*	su	*	Ċ	S
Ec	192.29 ± 11.73	182.52 ± 10.64	$190.61 \pm 22.37$	160.44 ± 11.72	$159.85 \pm 18.85$	141.11 ± 9.69	$148.62 \pm 9.36$	$208.08 \pm 21.60$	NS	ns	* *	×	ы		S
Lactate de	hydrogenase (LDH) .	and iso-citrate dehy	idrogenase (IDH) aci	tivities are expressed	d as nmol min <sup>-1</sup> mg	γ <sup>-1</sup> tissue. Energy c	consumption value	(Ec) is expressed as	m Lm	g−1 we	ght wet tis:	sue. Valı	ues are	e mear	of

fish (n = 9) ± SE. C, Control; S, Acute stress. Three-way ANOVA analysis results. D, Diet; O, dissolved oxygen levels; DOS, DO, DS, and OS, Interactions; ns, not significant P > 0.1; \*P < 0.05; \*\*P < 0.01.

		DEB	200			DEB	1 700			Factor	s		Intera	ctions		
	Norn	рхіа	Hyp	oxia	Norr	moxia	Нур	oxia								1
	U	S	v	S	U	S	U	S	۵	0	S	DOS	Q	DS	so	-
GPx	0.64 ± 0.09	0.41 ± 0.03	$0.55 \pm 0.04$	0.46 ± 0.03	0.34 ± 0.04	0.48 ± 0.07	$0.50 \pm 0.03$	0.47 ± 0.08	*	su	SU	*	*	* *	SU	
GR	3.27 ± 0.44	$4.09 \pm 0.33$	$2.56 \pm 0.25$	$3.13 \pm 0.25$	$3.34 \pm 0.32$	$4.59 \pm 0.57$	$4.11 \pm 0.40$	$3.72 \pm 0.53$	SU	SU	×	SU	NS	ns	ns	
TG	$1.14 \pm 0.18$	$1.24 \pm 0.09$	$1.06 \pm 0.07$	$1.20 \pm 0.07$	$1.28 \pm 0.07$	1.45 土 0.11	$1.28 \pm 0.12$	1.34 ± 0.13	*	SU	SU	SU	ns	SU	ns	
GSSG	$0.53 \pm 0.08$	$0.45 \pm 0.05$	$0.57 \pm 0.06$	$0.41 \pm 0.07$	$0.55 \pm 0.06$	$0.61 \pm 0.04$	$0.64 \pm 0.05$	$0.54 \pm 0.04$	*	SU	SU	SU	ns	SU	ns	
GSH	$0.61 \pm 0.11$	$0.79 \pm 0.08$	$0.56 \pm 0.06$	$0.79 \pm 0.10$	$0.73 \pm 0.07$	$0.84 \pm 0.08$	$0.64 \pm 0.10$	$0.86 \pm 0.13$	ns	su	×	SU	SU	SU	ns	
LPO	$5.62 \pm 0.43$	$5.81 \pm 0.52$	$6.38 \pm 0.93$	$5.70 \pm 0.41$	$5.48 \pm 0.28$	$6.07 \pm 0.79$	$5.28\pm0.71$	$5.11 \pm 0.46$	ns	SU	NS	SU	SU	NS	SU	
Glutathic mg prot *P < 0.C TABLE 7	ne peroxidase (GF -1. Values are mea 5; **P < 0.01.	$x_{0}$ and Glutathions in of fish ( $n = 9$ ) $\pm$ alectrolyte balance	s reductase (GA) ¿ SE. C, Control; S (DEB), dissolved d	), Acute stress. Tr.	min-1 mg prot-1. nree-way ANOVA . acute stress on ir	. Total glutathione analysis results. D, nate immune statt	(TG), oxidized glu Diet; O, dissolvei Las in plasma of rai	d oxygen levels; D inbow trout.	educed OS, DO,	glutathio DS, ano	ne (GSH), I OS, Inte	, and lipid p ractions; ns	, not sig	e (LPO) nificant F	n nmol > 0.1;	*
			DEB 200				DEB 700				Facto	ors	Inte	ractions		
		Normoxia		Hypoxia		Normoxia		Hypoxi								

	Norn	noxia	Hyp	oxia	Norr	noxia	Нур	oxia							
	U	S	υ	S	U	S	U	S	۵	0	s	DOS	DO	DS	so
ACH50	13.64 ± 2.12	15.80 ± 2.28	16.25 ± 3.25	20.43 ± 2.16	10.90 ± 2.09	10.19 ± 2.20	8.59 ± 0.96	10.72 土 1.59	*	su	SU	SU	SU	SU	SU
Lysozyme	$522.22 \pm 24.26$	$645.37 \pm 44.03$	$633.33 \pm 64.67$	$596.30 \pm 50.29$	$570.37 \pm 60.26$	$815.74 \pm 138.37$	$544.79 \pm 63.27$	$576.85 \pm 35.05$	ns	ns	*	NS	ns	ns	ns
Peroxidase	116.53 土 4.46	$123.04 \pm 4.87$	$120.26 \pm 5.80$	119.74 土 4.90	126.42 土 3.91	$106.08 \pm 14.32$	128.59 土 4.58	$125.60 \pm 2.25$	NS	ns	ns	ns	ns	ns	ns
Alternative cc dissolved oxy	omplement pathway /gen levels; DOS, D	/ ACH50, Lysozyme 0, DS, and OS, Inte	e and Peroxidase a	stivities expressed in splificant P > 0.1; *F	n U mL <sup>-1</sup> . Values é > < 0.05; **P < 0.0	are mean of fish (n = )1.	= 9) $\pm$ SE. C, Contro	ol; S, Acute stress.	Three	-way,	ANOVA	analysi	is result	's. D, I	Diet; O,

linked to advantageous cardiovascular adaptations to enhance the animal's function (Faust et al., 2004). Previous studies have shown that chronic hypoxia in both zebrafish (Danio rerio) and the cichlid Haplochromis piceatus lead to decreases in the ventricular space, suggesting a profound restructuration of this organ (Marques et al., 2008). Our results also suggest that the trout population used in the trial displayed a good capacity to adapt to the chronic hypoxic conditions. In particular, the rainbow trout subjected to 49 days of hypoxia displayed a reduction in heart to body mass index (CSI), but increased the glycogen content compared to the normoxic group. Therefore, the rainbow trout isogenic line appears to have the capacity to fully adapt to the applied hypoxic levels, which was confirmed by the lack of significant change in all the markers of oxidative stress (liver) and innate immune status (plasma) markers investigated on this study.

# Reliability of Markers for Acute and Chronic Stress

Changes in plasma cortisol levels are commonly used to assess acute stress conditions in fish, as a peak in the release of this hormone is associated with stress response. As expected, in this study, the cortisol levels in plasma of trout subjected to acute stressor were significantly increased when compared to their controls. However, the utility of changes in the levels of this cortisol to evaluate chronic stress in fish is not always possible due to several mechanisms reducing cortisol levels, including negative feedback (Pickering and Stewart, 1984; Mommsen et al., 1999) and the increased metabolic clearance rate of cortisol, all mechanisms which overall result in suppression of the cortisol response (see reviews Wendelaar Bonga, 1997; Gorissen and Flik, 2016). Thus, in rainbow and brown trout, chronic confinement stress for 6 weeks resulted in an elevation of plasma cortisol levels up to 4 weeks before returning to basal level after 42 days (Pickering and Pottinger, 1989; Person-Le Ruyet et al., 2008). A similar response was observed in Atlantic salmon parr or smolt exposed to repeated chasing stress over 3 weeks, when a decline in plasma cortisol levels was observed after 1 week, suggesting a rapid habituation/desensibilization of the HPI axis (Madaro et al., 2016). However, cortisol response to chronic stressors may vary according to species and/or the type of stressor. For example, seabass exposed for 57 days to water quality deterioration did not show any significant changes in the basal cortisol levels (Santos et al., 2010), whereas in other study in seabream and seabass, the same parameter increased after 3 weeks when exposed to three levels of chronic stress (confinement/chasing/air exposure) (Samaras et al., 2018). Another approach to characterize chronic stress status is the analysis of cortisol response to acute confinement stress, which affects the HPI axis responsiveness. This approach has been implemented in rainbow trout and salmon by applying chronic crowding stress or repeated acute stress over several weeks, resulting in a less pronounced cortisol response to acute stress (Madaro et al., 2015, 2016; Moltesen et al., 2016). In contrast, seabass exposed to water quality deterioration showed an increased cortisol response to acute stress (Santos et al., 2010)

and a similar response was also observed in trout exposed to chronic hypoxia (Leguen and Prunet, unpublished). Overall, this data indicates that chronic stress generally leads to dysregulation of cortisol production (basal and/or acute stress levels) which varies according to species and stressor. In the present study, exposure of rainbow trout to chronic low DO levels did not modify cortisol levels nor cortisol response, which fits with our previous conclusions that indicate fish were adapted to these hypoxic conditions.

In this study, the prolonged dietary challenge increased the cortisol levels in plasma of trout 1.23 times. However, such increase was modest compared to the spike of 4.08 times in plasma cortisol levels of trout when subjected to the acute stressor. In spite of these observed changes, cortisol levels were not significantly different in trout fed the diets with different DEB and then subjected to acute stress (P < 0.057), also indicated by the lack of interactions between both factors (P = 0.284). These results may suggest that the capacity of trout to mount a proper cortisol response when subjected to an acute stressor remained unaltered by the dietary treatment. These results contrast with studies presented in the previous paragraph where chronic stress leads to a clear dysregulation of cortisol responsiveness in fish. Therefore, additional studies are required to further appraise how the cortisol response and the allostatic load of trout may be altered under chronic stressors, for example when subjected to dietary challenges.

No detectable changes were observed in LPO, a marker of oxidative damage, in the liver of trout exposed to the experimental conditions, even when facing the acute stressor. However, components of the glutathione system (GR and GSH) were increased in liver of trout subjected to the acute stressor, indicating an alteration on the balance between prooxidants/antioxidants in this organ. In addition, another component of the glutathione system, the GPx activity in liver, displayed an interaction between dietary treatment and acute stress, as well as for all three factors together (P < 0.01), suggesting that the prolonged dietary challenge places a constraint in trout exposed to a subsequent acute stressor.

The innate immune system in fish is constitutive, reacting at short time scale to non-specific cues and induced by several external molecules (Tort et al., 2003; Tort, 2011). In particular, lysozyme activity is used as an indicator of non-specific immune response (Tort et al., 1996, 1998), rapidly induced in fish exposed to acute stressors (Demers and Bayne, 1997; Rotllant and Tort, 1997). Although lysozyme activity can vary considerably between fish species, this enzymatic activity in plasma appears to be modulated in response to both physical and nutritional cues (Möck and Peters, 1990; Montero et al., 1999b). The increased lysozyme activity in plasma of trout subjected to acute stress observed in this study suggest a link between stress and suppression of the innate immune response, which has important implications for aquaculture. As it was suggested earlier by Fevolden et al. (1999), it is possible to hypothesize that the enhanced lysozyme activity after exposure to the acute stressor may not be directly related to pathogen resistance, and may rather reflect disease-susceptible fish. In this study ACH<sub>50</sub>



activity was decreased by the dietary imbalance, where lysozyme activity was decreased by the acute stressor in plasma of rainbow trout. Surprisingly, no interaction between the two factors was detected for any of the innate immune status parameters analyzed. Therefore, it will be necessary to further investigate how chronic stressors may be affecting not only the innate but also the adaptive immune response of rainbow trout exposed to acute challenges.

A summarized description of the results obtained from this study is presented in Figure 1. This study suggest that the trout isogenic line displays a complete adaptive response to chronic hypoxia, whereas the challenge imposed by feeding an electrolyte-imbalanced diet profoundly affected fish homeostasis after 49 days. Such physiological condition in fish appear to lead to stress response as indicated by several markers of oxidative stress and decreased innate immune response. Although no interactions in the markers for innate immune response were detected between the different factors analyzed, the dietary challenge, by feeding trout the DEB 700 diet, suggests an impairment of the physiological response in this fish to a subsequent acute stressor. This is shown in this study by an increase in the levels of several oxidative stress markers detected in liver of trout exposed to the acute stressor in trout fed the electrolyte-imbalance diet. Nevertheless, the results presented here were obtained in an isogenic line and may not necessarily reflect the response from other rainbow trout used in aquaculture.

### CONCLUSION

In conclusion, this study suggests that dietary imbalances present during rearing may act as chronic stressors, interacting with other well-established stressors such as the acute stressors applied on this study. These results are of great interest to the aquaculture industry as both of these factors may be present in cultured fish and could be linked to negative effects. Future studies should further investigate the physiological and behavioral aspects associated to such stress responses, and how to reduce their occurrence.

### ETHICS STATEMENT

The fish trials were approved and carried out according to the Wageningen University Ethics Board for experimentation with animals (DEC, Registration protocol 2014056.a), under Dutch and EU legislation on the handling of experimental animals.

### **AUTHOR CONTRIBUTIONS**

LM, IG, PP, and JS conceptualized the study. LM, SN, EE, ML, RO, IL, PP, IG, and JS were involved in the methodology and formal analysis. LM, SN, PP, and JS were involved in writing the original draft. All authors reviewed the manuscript.

### FUNDING

The aquatic metabolic unit used in this study was cofounded by The Netherlands Organization for Scientific Research (code 805-34.025). Currently it is part of a large-scale EU research facility program. The study was funded by the EU-FP7 project AQUAEXCEL (262336). LM is currently supported by a Fundação para a Ciência e a Tecnologia (FCT) research position (IF/01314/2014/CP1231/CT0001) from Portugal. This study had also the support of FCT through strategic projects UID/Multi/04423/2019 and UID/MAR/04292/2013 granted to CIIMAR and MARE, respectively. SN has support from FCT grant (SFRH/BPD/94500/2013). This work was also supported by the Integrated Programme of SR&TD "Smart Valorization of Endogenous Marine Biological Resources Under a Changing Climate" (reference Centro-01-0145-FEDER-000018), co-funded by Centro 2020 program, Portugal 2020, European Union, through the European Regional Development Fund.

#### ACKNOWLEDGMENTS

We thank Menno ter Veld and the staff at the aquatic research facility "De Haar Vissen" of Wageningen University for their technical support in conducting the experiments. We also would like to thank Ronald Booms and Tino Leffering (WIAS, WUR,

#### REFERENCES

- Almeida, J. R., Gravato, C., and Guilhermino, L. (2015). Effects of temperature in juvenile seabass (*Dicentrarchus labrax* L.) biomarker responses and behaviour: implications for environmental monitoring. *Estuaries Coasts* 38, 45–55. doi: 10.1007/s12237-014-9792-7
- Baker, M. A., Cerniglia, G. J., and Zaman, A. (1990). Microtiter plate assay for the measurement of glutathione and glutathione disulfide in large numbers of biological samples. *Anal. Biochem.* 190, 360–365. doi: 10.1016/0003-2697(90) 90208-Q
- Bernier, N. J., and Craig, P. M. (2005). CRF-related peptides contribute to stress response and regulation of appetite in hypoxic rainbow trout. Am. J. Physiol. Regul. Integr. Comp. Physiol. 289, R982–R990. doi: 10.1152/ajpregu.00668.2004
- Biller-Takahashi, J. D., Takahashi, L. S., Marzocchi-Machado, C. M., Zanuzzo, F. S., Sabioni, R. E., and Urbinati, E. C. (2012). Hemolytic activity of alternative complement pathway as an indicator of innate immunity in pacu (*Piaractus* mesopotamicus). Revista Brasileira de Zootecnia 41, 237–241. doi: 10.1590/ S1516-35982012000200001
- Boshra, H., Li, J., and Sunyer, J. O. (2006). Recent advances on the complement system of teleost fish. *Fish Shellfish Immunol.* 20, 239–262. doi: 10.1016/j.fsi. 2005.04.004
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254. doi: 10.1016/0003-2697(76)90527-3
- Bratland, S., Stien, L. H., Braithwaite, V. A., Juell, J.-E., Folkedal, O., Nilsson, J., et al. (2010). From fright to anticipation: using aversive light stimuli to investigate reward conditioning in large groups of Atlantic salmon (*Salmo salar*). Aquac. Int. 18, 991–1001. doi: 10.1007/s10499-009-9317-8
- Chiu, S.-T., Tsai, R.-T., Hsu, J.-P., Liu, C.-H., and Cheng, W. (2008). Dietary sodium alginate administration to enhance the non-specific immune responses, and disease resistance of the juvenile grouper *Epinephelus fuscoguttatus*. *Aquaculture* 277, 66–72. doi: 10.1016/j.aquaculture.2008.01.032
- Conte, F. S. (2004). Stress and the welfare of cultured fish. Appl. Anim. Behav. Sci. 86, 205-223. doi: 10.1016/j.applanim.2004.02.003
- Cribb, A. E., Leeder, J. S., and Spielberg, S. P. (1989). Use of a microplate reader in an assay of glutathione reductase using 5,5'-dithiobis(2-nitrobenzoic acid). *Anal. Biochem.* 183, 195–196. doi: 10.1016/0003-2697(89)90188-7
- Dahlhoff, E. P. (2004). Biochemical indicators of stress and metabolism: applications for marine ecological studies. *Annu. Rev. Physiol.* 66, 183–207. doi: 10.1146/annurev.physiol.66.032102.114509
- De Coen, W., and Janssen, C. R. (1997). The use of biomarkers in *Daphnia magna* toxicity testing. IV: cellular energy allocation: a new methodology to assess the energy budget of toxicant-stressed *Daphnia* populations. *J. Aquat. Ecosyst. Stress Recover.* 6, 43–55. doi: 10.1023/A:1008228517955
- De Coen, W. M., and Janssen, C. R. (2003). The missing biomarker link: relationships between effects on the cellular energy allocation biomarker of toxicant-stressed *Daphnia magna* and corresponding population characteristics. *Environ. Toxicol. Chem.* 22, 1632–1641. doi: 10.1002/etc.5620220727
- Demers, N. E., and Bayne, C. J. (1997). The immediate effects of stress on hormones and plasma lysozyme in rainbow trout. *Dev. Comp. Immunol.* 21, 63–73. doi: 10.1016/S0145-305X(97)00009-8
- Dersjant-Li, Y., Verreth, J. A. J., Evers, F., Tijssen, P. A. T., Booms, R., and Verstegen, M. W. A. (1999). The influence of dietary cation-anion differences on acid-base balance, food intake, growth and nutrient utilisation of juvenile

Netherlands) for their assistance during sampling, as well as to Carla Silva (MARE, Portugal), Francisca Brito and Luis Pereira (CIIMAR, Portugal) for laboratory work.

#### SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fphys. 2019.00453/full#supplementary-material

African catfish *Clarias gariepinus* (Burchell). *Fish Physiol. Biochem.* 20, 305–311. doi: 10.1023/A:1007707507228

- Dersjant-Li, Y., Verreth, J. A. J., Tijssen, P. A. T., Booms, R., Verstegen, M. W. A., and Huisman, E. A. (2000). Metabolic costs of changing the cation-anion difference in the diet of juvenile African catfish *Clarias gariepinus* (Burchell). *Aquac. Nutr.* 6, 39–45. doi: 10.1046/j.1365-2095.2000.00126.x
- Ellis, G., and Goldberg, D. M. (1971). An improved manual and semi-automatic assay for NADP dependent isocitrate dehydrogenase activity, with a description of some kinetic properties of human liver and serum enzyme. *Clin. Biochem.* 2, 175–185. doi: 10.1016/S0009-9120(71)91363-4
- Faust, H. A., Gamperl, A. K., and Rodnick, K. J. (2004). All rainbow trout (Oncorhynchus mykiss) are not created equal: intra-specific variation in cardiac hypoxia tolerance. J. Exp. Biol. 207, 1005–1015. doi: 10.1242/jeb.00824
- Fevolden, S.-E., Røed, K. H., Fjalestad, K. T., and Stien, J. (1999). Poststress levels of lysozyme and cortisol in adult rainbow trout: heritabilities and genetic correlations. J. Fish Biol. 54, 900–910. doi: 10.1111/j.1095-8649.1999.tb02040.x
- Folkedal, O., Stien, L. H., Torgersen, T., Oppedal, F., Olsen, R. E., Fosseidengen, J. E., et al. (2012a). Food anticipatory behaviour as an indicator of stress response and recovery in Atlantic salmon post-smolt after exposure to acute temperature fluctuation. *Physiol. Behav.* 105, 350–356. doi: 10.1016/j.physbeh. 2011.08.008
- Folkedal, O., Torgersen, T., Olsen, R. E., Fernö, A., Nilsson, J., Oppedal, F., et al. (2012b). Duration of effects of acute environmental changes on food anticipatory behaviour, feed intake, oxygen consumption, and cortisol release in Atlantic salmon parr. Physiol. Behav. 105, 283–291. doi: 10.1016/j.physbeh. 2011.07.015
- Fontagné-Dicharry, S., Larroquet, L., Dias, K., Cluzeaud, M., Heraud, C., and Corlay, D. (2018). Effects of dietary oxidized fish oil supplementation on oxidative stress and antioxidant defense system in juvenile rainbow trout (*Oncorhynchus mykiss*). Fish Shellfish Immunol. 74, 43–51. doi: 10.1016/j.fsi. 2017.12.039
- Gamperl, A. K., and Farrell, A. P. (2004). Cardiac plasticity in fishes: environmental influences and intraspecific differences. J. Exp. Biol. 207, 2539–2550. doi: 10. 1242/jeb.01057
- Glencross, B. D. (2009). Reduced water oxygen levels affect maximal feed intake, but not protein or energy utilization efficiency of rainbow trout (*Oncorhynchus mykiss*). Aquac. Nutr. 15, 1–8. doi: 10.1111/j.1365-2095.2007.00562.x
- Gómez-Requeni, P., Mingarro, M., Calduch-Giner, J. A., Médale, F., Martin, S. A. M., Houlihan, D. F., et al. (2004). Protein growth performance, amino acid utilisation and somatotropic axis responsiveness to fish meal replacement by plant protein sources in gilthead sea bream (*Sparus aurata*). Aquaculture 232, 493–510. doi: 10.1016/S0044-8486(03)00532-5
- Gorissen, M., and Flik, G. (2016). "Endocrinology of the stress response in fish," in *Biology of Stress in Fish*, eds C. B. Schreck, L. Tort, A. P. Farrell, and C. J. Brauner (San Diego, CA: Academic press), 75–111. doi: 10.1016/B978-0-12-802728-8.00003-5
- Hermes-Lima, M., Storey, J. M., and Storey, K. B. (2001). "Antioxidant defenses and animal adaptation to oxygen availability during environmental stress," in *Cell* and Molecular Responses to Stress, eds K. B. Storey and J. M. Storey (Amsterdam: Elsevier), 263–287.
- Ishibashi, Y., Inoue, K., Nakatsukasa, H., Ishitani, Y., Miyashita, S., and Murata, O. (2005). Ontogeny of tolerance to hypoxia and oxygen consumption of larval and juvenile red sea bream. *Pagrus major. Aquaculture* 244, 331–340. doi: 10.1016/j.aquaculture.2004.11.019

- Jo, S.-H., Son, M.-K., Koh, H.-J., Lee, S.-M., Song, I.-H., Kim, Y.-O., et al. (2001). Control of mitochondrial redox balance and cellular defense against oxidative damage by mitochondrial NADP+-dependent isocitrate dehydrogenase. J. Biol. Chem. 276, 16168–16176. doi: 10.1074/jbc.M010120200
- Jobling, M. (1981). The influences of feeding on the metabolic rate of fishes: a short review. J. Fish Biol. 18, 385–400. doi: 10.1111/j.1095-8649.1981.tb03780.x
- Jobling, M. (1993). "Bioenergetics: feed intake and energy partitioning," in *Fish Ecophysiology*, eds J. C. Rankin and F. B. Jensen (Dordrecht: Springer).
- Lee, S. M., Koh, H.-J., Park, D.-C., Song, B. J., Huh, T.-L., and Park, J.-W. (2002). Cytosolic NADP+-dependent isocitrate dehydrogenase status modulates oxidative damage to cells. *Free Radic. Biol. Med.* 32, 1185–1196. doi: 10.1016/S0891-5849(02)00815-8
- Lie, O., Syed, M., and Solbu, H. (1986). Improved agar plate assays of bovine lysozyme and haemolytic complement activity. Acta Vet. Scand. 27, 23–32.
- Lima, I., Moreira, S. M., Osten, J. R., Soares, A. M. V. M., and Guilhermino, L. (2007). Biochemical responses of the marine mussel *Mytilus galloprovincialis* to petrochemical environmental contamination along the northwestern coast of Portugal. *Chemosphere* 66, 1230–1242. doi: 10.1016/j.chemosphere.2006.07.057
- Madaro, A., Fernö, A., Kristiansen, T. S., Olsen, R. E., Gorissen, M., Flik, G., et al. (2016). Effect of predictability on the stress response to chasing in Atlantic salmon (*Salmo salar* L.) parr. *Physiol. Behav.* 153, 1–6. doi: 10.1016/j.physbeh. 2015.10.002
- Madaro, A., Olsen, R. E., Kristiansen, T. S., Ebbesson, L. O. E., Nilsen, T. O., Flik, G., et al. (2015). Stress in Atlantic *salmon*: response to unpredictable chronic stress. *J. Exp. Biol.* 218, 2538–2550. doi: 10.1242/jeb.120535
- Magnoni, L. J., Eding, E., Leguen, I., Prunet, P., Geurden, I., Ozório, R. O. A., et al. (2018). Hypoxia, but not an electrolyte-imbalanced diet, reduces feed intake, growth and oxygen consumption in rainbow trout (*Oncorhynchus mykiss*). Sci. Rep. 8:4965. doi: 10.1038/s41598-018-23352-z
- Magnoni, L. J., Salas-Leiton, E., Peixoto, M. J., Pereira, L., Silva-Brito, F., Fontinha, F., et al. (2016). Dietary electrolyte balance affects growth performance, amylase activity and metabolic response in the meagre (*Argyrosomus regius*). Comp. Biochem. Physiol. B Biochem. Mol. Biol. 211, 8–15. doi: 10.1016/j.cbpb.2017.03.001
- Mandic, M., Todgham, A. E., and Richards, J. G. (2009). Mechanisms and evolution of hypoxia tolerance in fish. Proc. R. Soc. B Biol. Sci. 276, 735–744. doi: 10.1098/ rspb.2008.1235
- Marques, I. J., Leito, J. T. D., Spaink, H. P., Testerink, J., Jaspers, R. T., Witte, F., et al. (2008). Transcriptome analysis of the response to chronic constant hypoxia in zebrafish hearts. *J. Comp. Physiol. B* 178, 77–92. doi: 10.1007/s00360-007-0201-4
- Martínez-Álvarez, R. M., Morales, A. E., and Sanz, A. (2005). Antioxidant defenses in fish: biotic and abiotic factors. *Rev. Fish Biol. Fish.* 15, 75–88. doi: 10.1007/ s11160-005-7846-4
- Möck, A., and Peters, G. (1990). Lysozyme activity in rainbow trout, Oncorhynchus mykiss (Walbaum), stressed by handling, transport and water pollution. J. Fish Biol. 37, 873–885. doi: 10.1111/j.1095-8649.1990.tb03591.x
- Mohandas, J., Marshall, J. J., Duggin, G. G., Horvath, J. S., and Tiller, D. J. (1984). Differential distribution of glutathione and glutathione-related enzymes in rabbit kidney. Possible implications in analgesic nephropathy. *Biochem. Pharmacol.* 33, 1801–1807. doi: 10.1016/0006-2952(84)90353-8
- Moltesen, M., Laursen, D. C., Thörnqvist, P.-O., Andersson, M. Å, Winberg, S., and Höglund, E. (2016). Effects of acute and chronic stress on telencephalic neurochemistry and gene expression in rainbow trout (*Oncorhynchus mykiss*). *J. Exp. Biol.* 219, 3907–3914. doi: 10.1242/jeb.139857
- Mommsen, T. P., Vijayan, M. M., and Moon, T. W. (1999). Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. *Rev. Fish Biol. Fish.* 9, 211–268. doi: 10.1023/A:1008924418720
- Montero, D., Grasso, V., Izquierdo, M. S., Ganga, R., Real, F., Tort, L., et al. (2008). Total substitution of fish oil by vegetable oils in gilthead sea bream (*Sparus aurata*) diets: effects on hepatic Mx expression and some immune parameters. *Fish Shellfish Immunol.* 24, 147–155. doi: 10.1016/j.fsi.2007.08.002
- Montero, D., Izquierdo, M. S., Tort, L., Robaina, L., and Vergara, J. M. (1999a). High stocking density produces crowding stress altering some physiological and biochemical parameters in gilthead seabream, *Sparus aurata*, juveniles. *Fish Physiol. Biochem.* 20, 53–60. doi: 10.1023/A:1007719928905
- Montero, D., Marrero, M., Izquierdo, M. S., Robaina, L., Vergara, J. M., and Tort, L. (1999b). Effect of vitamin E and C dietary supplementation on some

immune parameters of gilthead seabream (*Sparus aurata*) juveniles subjected to crowding stress. *Aquaculture* 171, 269–278. doi: 10.1016/S0044-8486(98) 00387-1

- Montero, D., Kalinowski, T., Obach, A., Robaina, L., Tort, L., Caballero, M. J., et al. (2003). Vegetable lipid sources for gilthead seabream (*Sparus aurata*): effects on fish health. *Aquaculture* 225, 353–370. doi: 10.1007/s11745-016-4191-4
- Montero, D., Mathlouthi, F., Tort, L., Afonso, J. M., Torrecillas, S., Fernández-Vaquero, A., et al. (2010). Replacement of dietary fish oil by vegetable oils affects humoral immunity and expression of pro-inflammatory cytokines genes in gilthead sea bream Sparus aurata. *Fish Shellfish Immunol.* 29, 1073–1081. doi: 10.1016/j.fsi.2010.08.024
- Nilsson, J., Stien, L. H., Fosseidengen, J. E., Olsen, R. E., and Kristiansen, T. S. (2012). From fright to anticipation: reward conditioning versus habituation to a moving dip net in farmed Atlantic cod (*Gadus morhua*). Appl. Anim. Behav. Sci. 138, 118–124. doi: 10.1016/j.applanim.2012.02.014
- Ohkawa, H., Ohishi, N., and Yagi, K. (1979). Assay for lipid peroxides in animal tissues bythiobarbituric acid reaction. Anal. Biochem. 95, 351–358. doi: 10.1016/ 0003-2697(79)90738-3
- Patience, J. F., and Wolynetz, M. S. (1990). Influence of dietary undetermined anion on acid-base status and performance in pigs. J. Nutr. 120, 579–587. doi: 10.1093/jn/120.6.579
- Pedersen, C. L. (1987). Energy budgets for juvenile rainbow trout at various oxygen concentrations. Aquaculture 62, 289–298. doi: 10.1016/0044-8486(87) 90171-2
- Person-Le Ruyet, J., Labbé, L., Le Bayon, N., Sévère, A., Le Roux, A., Le Delliou, H., et al. (2008). Combined effects of water quality and stocking density on welfare and growth of rainbow trout (*Oncorhynchus mykiss*). Aquat. Living Res. 21, 185–195. doi: 10.1051/alr:2008024
- Pickering, A. D. (1993). Growth and stress in fish production. Aquaculture 111, 51-63. doi: 10.1016/0044-8486(93)90024-S
- Pickering, A. D., and Pottinger, T. G. (1989). Stress responses and disease resistance in salmonid fish: effects of chronic elevation of plasma cortisol. *Fish Physiol. Biochem.* 7, 253–258. doi: 10.1007/BF00004714
- Pickering, A. D., and Stewart, A. (1984). Acclimation of the interrenal tissue of the brown trout, *Salmo trutta* L., to chronic crowding stress. *J. Fish Biol.* 24, 731–740. doi: 10.1111/j.1095-8649.1984.tb04844.x
- Portz, D. E., Woodley, C. M., and Cech, J. J. (2006). Stress-associated impacts of short-term holding on fishes. *Rev. Fish Biol. Fish.* 16, 125–170. doi: 10.1007/ s11160-006-9012-z
- Quade, M. J., and Roth, J. A. (1997). A rapid, direct assay to measure degranulation of bovine neutrophil primary granules. *Vet. Immunol. Immunopathol.* 58, 239–248. doi: 10.1016/S0165-2427(97)00048-2
- Rotllant, J., Pavlidis, M., Kentouri, M., Abad, M. E., and Tort, L. (1997). Nonspecific immune responses in the red porgy *Pagrus pagrus* after crowding stress. *Aquaculture* 156, 279–290. doi: 10.1016/S0044-8486(97)00075-6
- Rotllant, J., and Tort, L. (1997). Cortisol and glucose response after acute stress in the sparid red porgy (*Pagrus pagrus*) previously subjected to crowding stress. *J. Fish Biol.* 51, 21–28. doi: 10.1111/j.1095-8649.1997.tb02510.x
- Sadoul, B., Leguen, I., Colson, V., Friggens, N. C., and Prunet, P. (2015). A multivariate analysis using physiology and behavior to characterize robustness in two isogenic lines of rainbow trout exposed to a confinement stress. *Physiol. Behav.* 140, 139–147. doi: 10.1016/j.physbeh.2014. 12.006
- Samaras, A., Espírito Santo, C., Papandroulakis, N., Mitrizakis, N., Pavlidis, M., Höglund, E., et al. (2018). Allostatic load and stress physiology in European seabass (*Dicentrarchus labrax* L.) and *Gilthead Seabream (Sparus aurata* L.). *Front. Endocrinol.* 9:451. doi: 10.3389/fendo.2018.00451
- Santos, G. A., Schrama, J. W., Mamauag, R. E. P., Rombout, J. H. W. M., and Verreth, J. A. J. (2010). Chronic stress impairs performance, energy metabolism and welfare indicators in European seabass (*Dicentrarchus labrax*): the combined effects of fish crowding and water quality deterioration. *Aquaculture* 299, 73–80. doi: 10.1016/j.aquaculture.2009.11.018
- Saravanan, S., Geurden, I., Figueiredo-Silva, A. C., Nusantoro, S., Kaushik, S., Verreth, J. A. J., et al. (2013a). Oxygen consumption constrains food intake in fish fed diets varying in essential amino acid composition. *PLoS One* 8:e72757. doi: 10.1371/journal.pone.0072757
- Saravanan, S., Geurden, I., Orozco, Z. G. A., Kaushik, S. J., Verreth, J. A. J., and Schrama, J. W. (2013b). Dietary electrolyte balance affects the nutrient

digestibility and maintenance energy expenditure of *Nile tilapia. Br. J. Nutr.* 110, 1948–1957. doi: 10.1017/S0007114513001323

- Sitjà-Bobadilla, A., Peña-Llopis, S., Gómez-Requeni, P., Médale, F., Kaushik, S., and Pérez-Sánchez, J. (2005). Effect of fish meal replacement by plant protein sources on non-specific defence mechanisms and oxidative stress in gilthead sea bream (*Sparus aurata*). Aquaculture 249, 387–400. doi: 10.1016/j.aquaculture. 2005.03.031
- Sunyer, J. O., and Tort, L. (1995). Natural hemolytic and bactericidal activities of sea bream Sparus aurata serum are effected by the alternative complement pathway. Vet. Immunol. Immunopathol. 45, 333–345. doi: 10.1016/0165-2427(94) 05430-Z
- Tort, L. (2011). Stress and immune modulation in fish. Dev. Comp. Immunol. 35, 1366–1375. doi: 10.1016/j.dci.2011.07.002
- Tort, L., Balasch, J. C., and Mackenzie, S. (2003). Fish immune system. A crossroads between innate and adaptive responses. *Inmunologia* 22, 277–286.
- Tort, L., Rotllant, J., and Rovira, L. (1998). Immunological suppression in gilthead sea bream Sparus aurata of north-west Mediterranean at low temperatures. Comp. Biochem. Physiol. 120A, 175–179. doi: 10.1016/S1095-6433(98) 10027-2
- Tort, L., Sunyer, J. O., Gómez, E., and Molinero, A. (1996). Crowding stress induces changes in serum haemolytic and agglutinating activity in the gilthead sea bream Sparus aurata. Vet. Immunol. Immunopathol. 51, 179–188. doi: 10.1016/ 0165-2427(95)05502-9

- Tran-Duy, A., Schrama, J. W., Van Dam, A. A., and Verreth, J. A. J. (2008). Effects of oxygen concentration and body weight on maximum feed intake, growth and hematological parameters of Nile tilapia. *Oreochromis niloticus. Aquaculture* 275, 152–162. doi: 10.1016/j.aquaculture.2007.12.024
- Vassault, A. (1983). "Lactate dehydrogenase," in *Methods of Enzymatic Analysis. Enzymes: Oxireductases, Transferases*, 3rd Edn, ed. H. O. Bergmeyer (New York, NY: Academic Press).
- Wendelaar Bonga, S. E. (1997). The stress response in fish. *Physiol. Rev.* 77, 591–625. doi: 10.1152/physrev.1997.77.3.591
- Winston, G. W. (1991). Oxidants and antioxidants in aquatic animals. Comp. Biochem. Physiol. C Comp. Pharmacol. 100, 173–176. doi: 10.1016/0742-8413(91)90148-M

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Magnoni, Novais, Eding, Leguen, Lemos, Ozório, Geurden, Prunet and Schrama. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.