



HAL
open science

The 4p1000 initiative: Opportunities, limitations and challenges for implementing soil organic carbon sequestration as a sustainable development strategy

Cornelia Rumpel, Farshad Amiraslani, Claire Chenu, Magaly Garcia Cardenas, Martin Kaonga, Lydie-Stella Koutika, Jagdish Ladha, Beata Madari, Yasuhito Shirato, Pete Smith, et al.

► To cite this version:

Cornelia Rumpel, Farshad Amiraslani, Claire Chenu, Magaly Garcia Cardenas, Martin Kaonga, et al.. The 4p1000 initiative: Opportunities, limitations and challenges for implementing soil organic carbon sequestration as a sustainable development strategy. *AMBIO: A Journal of Environment and Society*, 2020, 49, pp.:350-360. 10.1007/s13280-019-01165-2. hal-02627512

HAL Id: hal-02627512

<https://hal.inrae.fr/hal-02627512>

Submitted on 11 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

UC Davis

UC Davis Previously Published Works

Title

The 4p1000 initiative: Opportunities, limitations and challenges for implementing soil organic carbon sequestration as a sustainable development strategy.

Permalink

<https://escholarship.org/uc/item/7071t744>

Journal

Ambio, 49(1)

ISSN

0044-7447

Authors

Rumpel, Cornelia
Amiraslani, Farshad
Chenu, Claire
[et al.](#)

Publication Date

2020

DOI

10.1007/s13280-019-01165-2



Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



The 4p1000 initiative: Opportunities, limitations and challenges for implementing soil organic carbon sequestration as a sustainable development strategy

Cornelia Rumpel , Farshad Amiraslani, Claire Chenu, Magaly Garcia Cardenas, Martin Kaonga, Lydie-Stella Koutika , Jagdish Ladha, Beata Madari, Yasuhito Shirato, Pete Smith, Brahim Souidi, Jean-François Soussana, David Whitehead, Eva Wollenberg

Received: 8 October 2018 / Revised: 20 October 2018 / Accepted: 28 February 2019 / Published online: 23 March 2019

Abstract Climate change adaptation, mitigation and food security may be addressed at the same time by enhancing soil organic carbon (SOC) sequestration through environmentally sound land management practices. This is promoted by the “4 per 1000” Initiative, a multi-stakeholder platform aiming at increasing SOC storage through sustainable practices. The scientific and technical committee of the Initiative is working to identify indicators, research priorities and region-specific practices needed for their implementation. The Initiative received its name due to the global importance of soils for climate change, which can be illustrated by a thought experiment showing that an annual growth rate of only 0.4% of the standing global SOC stocks would have the potential to counterbalance the current increase in atmospheric CO₂. However, there are numerous barriers to the rise in SOC stocks and while SOC sequestration can contribute to partly offsetting greenhouse gas emissions, its main benefits are related to increased soil quality and climate change adaptation. The Initiative provides a collaborative platform for policy makers, practitioners, scientists and stakeholders to engage in finding solutions. Criticism of the Initiative has been related to the poor definition of its numerical target, which was not understood as an aspirational goal. The objective of this paper is to present the aims of the initiative, to discuss critical issues and to present challenges for its implementation. We identify barriers, risks and trade-offs and advocate for collaboration between multiple parties in order to stimulate innovation and to initiate the transition of agricultural systems toward sustainability.

Keywords Carbon sequestration · Climate change · Food security · Soil

INTRODUCTION

In recent years, with rising atmospheric CO₂ concentrations, the role of soils in the global carbon cycle has been increasingly acknowledged. As a result and as a supplement to immediate and aggressive emissions reduction, an increase of soil organic carbon (SOC) sequestration has been promoted by scientists and policy makers as a prospective additional opportunity to partly counterbalance increasing atmospheric CO₂ concentrations (e.g. Lal 2004; <https://www.4p1000.org/>). The SOC pool of the terrestrial biosphere is estimated to be around 1500 Gt C to a depth of 1 m. Changes of this large pool may affect atmospheric CO₂ concentrations. Consequently, increasing SOC sequestration through environmentally sound agricultural practices has been advocated as an option to remove CO₂ from the atmosphere (Smith et al. 2016).

In 2015, the French government launched the “4 per 1000” (4p1000) Initiative at the 21st Conference of Parties of the United Nations Framework Convention on Climate Change (UNFCCC) as part of the Lima-Paris Action Plan. The Initiative promotes an innovative model for helping to mitigate climate change, through increase in SOC and contributing to climate change adaptation and food security. It is believed that increasing SOC enhances certain soil functions, thereby benefitting agricultural production (Lal 2004).

As agricultural activities and land use change account for about 25% of the CO₂, 50% of the CH₄ and 70% of the N₂O anthropogenic emissions (Hutchinson et al. 2007), enhanced SOC sequestration could help offset these emissions (Paustian et al. 2016). SOC sequestration could also help to fill the gap between the intended national contributions and the reality to achieve the Paris climate goal (Rumpel et al. 2018).

Moreover, increased SOC sequestration is likely to generate co-benefits helping to achieve several sustainable development goals, in particular those related to reducing hunger (SDG 2), extreme poverty (SDG1, 3), and improving the protection of the environment (SDGs 6, 11, 12, 14, 15) and the global climate (SDG 13) (Soussana et al. 2019). Particularly, the Initiative may have the possibility to contribute to SDG 15.3, by combatting desertification and restoring degraded lands through increasing SOC storage.

The 4p1000 Initiative mainly focuses on agricultural soils with low levels of SOC due to continuous cultivation and often unsustainable crop intensification practices (Pingali 2012). The Initiative encourages farm management practices that preserve and build SOC stocks while limiting carbon trade-offs. Adoption of these practices may lead to a transition towards sustainable agricultural production (Tilman et al. 2011; <https://futurepolicy.org>).

The objectives of this paper are to (1) discuss the aims of the 4p1000 Initiative and controversial issues concerning the Initiative, (2) highlight the potential of the 4p1000 Initiative to provide collaborative platform for policy-science-practice interaction and (3) proposes an implementation pathway from policy to action.

CRITIQUES OF THE 4P1000 INITIATIVE

The 4p1000 initiative was launched based on a thought experiment suggesting that a small increase of the SOC stocks of global soils (4 per 1000 or 0.4% of the standing SOC stock) would remove a significant proportion of CO₂ from the atmosphere, while simultaneously augmenting the capability of agricultural systems to adapt to climate change and to provide food security. The achievability of the Initiative's target of an annual increase in agricultural SOC stocks of 0.4% to a depth of 0.3–0.4 m globally has been intensively discussed and criticised (de Vries et al. 2018; VandenBygaert 2018).

As a policy goal, a single number, i.e. a quantity of carbon to be stored in soils that appeared to be easily attainable was clear and thus easier to communicate than multiple numbers for different regions or conditions. Articulation of a clear target by prominent promoters of the Initiative including well-respected scientists and policy makers was necessary to ensure inclusion of SOC on the global political agenda (Kon Kam King et al., 2018). The selection of this simplified 4p1000 target for increasing SOC sequestration may be interpreted as analogous to the selection of targets to limit global temperature increase to 2 or 1.5 °C above pre-industrial levels set by the UNFCCC and to targets for Sustainable Development Goals established by the United Nations in 2015. These are broad

aspirational goals with much uncertainty about what is achievable, especially in relation to specific geographical locations. The climate science community was faced with similar criticisms when global warming targets were announced. We suggest that some of the controversy regarding the 4p1000 Initiative is attributable to the initial setting of an aspirational target of an annual SOC increase of 0.4% of the standing stock. The initial criticism was related to the suggestion that this could offset *all* fossil fuel emission and that it could therefore be used as an excuse not to drastically reduce CO₂ and other greenhouse gas emissions. This was seen as a complete exaggeration and dangerous. Moreover, the target was interpreted as a strong commitment rather than an aspirational goal. Criticism has also focused on the number, its calculation, significance and achievability. Further, there was ambiguity related to the presentation of the calculation of the quantity of SOC needed to partly offset anthropogenic CO₂ emissions without considering other greenhouse gas emissions (de Vries et al. 2018; Minasny et al. 2018). The initial statements were thus not framed precisely in scientific terms, which made the nature and the role of the target difficult to interpret.

More specific criticisms of the Initiative in relation to biophysical, agronomic and socio-economic issues are presented in Table 1 and discussed below. These include (1) biophysical limits (demands in terms of water, nutrients and energy), and other barriers such as (2) the trade-off effects, (3) climate change effects and (4) the socio-economic implications for the agricultural sector, including cultural issues and governance (van Groeningen et al. 2017; Baveye et al. 2018; de Vries et al. 2018; van den Bygaert 2018; White et al. 2018; Poulton et al. 2018).

Biophysical limits and barriers

Under given constant conditions, SOC stocks will approach an equilibrium level depending on carbon inputs and outputs determined by pedoclimatic conditions, land use and management practices (Fig. 1). Regulation of SOC storage under equilibrium conditions is increasingly ascribed to SOC input (Fujisaki et al. 2018), soil-inherent pedologic characteristics (Barré et al. 2017) and the state of soil development (Schiefer et al. 2018). When land management changes, the equilibrium may be disturbed leading to SOC gain or loss. Following land use change (e.g. agriculture), SOC losses generally occur through increased microbial decomposition rates and through soil erosion (Sanderman et al. 2017). Agricultural practices also often decrease organic matter inputs. For example, in many regions of the world, biomass input into soil is reduced through burning of crop residues (<http://www.fao.org/faostat/en/#data/GB>), when these could otherwise be used

Table 1 Classification of the criticisms of the 4 per 1000 Initiative's target and explanation and proposed actions to respond to the criticisms

Criticism	Articles	Proposed explanation and action	Associated research needs
Poor calculation of target			
Inconsistent inputs for calculation	(de Vries et al. 2018)	Consistent communication and clear explanation of calculations	na
Global emissions number only reflect CO ₂ , not CH ₄ and N ₂ O, so the calculation of the offset is too low	(de Vries et al. 2018), (Baveye et al. 2018)	Explanation of calculations: only anthropogenic CO ₂ emissions are targeted in the calculation of the Initiative, not all anthropogenic greenhouse gas emissions, Actions: non CO ₂ GHG emissions should not be increased	na
Biophysical			
C storage is limited. Storage reaches an equilibrium value and the rate of storage starts to decrease once storage is initiated, so the potential for sequestering carbon sequestered will decrease rapidly over time.	White et al. (2018), Baveye et al. (2018), Schiefer et al. (2018)	Even additional storage over a few decades would help mitigate CO ₂ emissions. Predictions must account for these dynamics	Assessments of the local/ regional/national C stocks and C storage potential considering time limits
Non-permanence of SOC storage	Baveye et al. (2018), Poulton et al. (2018)	Encourage the maintenance of best management practices	Vulnerability of SOC stocks
4p1000 per year (rate of sequestration over time) is not feasible quantitatively: estimates are too high globally but also locally	de Vries et al. (2018), White et al. (2018)	Even an additional storage, less than 4% would contribute to mitigate CO ₂ emissions. Large variability of SOC storage rates depending on pedoclimatic conditions and management options implemented	Assessments of the local/ regional/national C stocks and C storage potential, using long-term observations and experimental farm plots
Insufficient biomass available	Poulton et al. (2018)	Implementation has to be spatially differentiated. Promote recycling and valuation of waste (circular economy)	SOC storage potential of organic wastes
Insufficient nitrogen and phosphorus available	van Groenigen (2017), White et al. (2018), Baveye et al. (2018)	Where possible, N-use efficiency needs to be improved. Implementation has to be spatially differentiated. Avoid use of synthetic or mined fertilisers by alternative practices (e.g. mycorrhizae, legumes, Plant Growth Promoting Rhizobacteria, rotations, waste management and circular economy)	Effects of nitrogen fertiliser on SOC storage in grasslands (has been better studied in cropland). Global estimation of the nitrogen fixing potential of agro-ecosystems. Development of new fertilisation strategies
Need for comprehensive greenhouse gas accounting (i.e. include non-CO ₂ emissions such as N ₂ O, CH ₄)	White et al. (2018), Baveye et al. (2018)	A net greenhouse gas balance must be provided for all projects. Avoid or adapt SOC storage strategies in situations with high risk (e.g. inhibitors, liming, timing nitrogen additions, slow release fertilisers, paddy water management)	Conditions conducive to N ₂ O emissions (nature of organic matter, pH, soil structure)
Not accounting for climate change (temperature increase)	Baveye et al. (2018)	Reinforces the need for the Initiative	Temperature sensitivity estimates have been based mostly on disturbed soil and laboratory incubations. Perform more in situ measurements
Enhanced mineralisation on addition of easily decomposable carbon (priming effect) could release more CO ₂	Baveye et al. (2018)	Measure changes in SOC storage rates under field conditions, integrate enhanced priming effect if any	Modelling and experiments to quantify and reduce priming effects
Not all carbon is organic; inorganic carbon could release large amounts of CO ₂ with temperature rise or microbial activity	Baveye et al. (2018)	Inorganic C dynamics must be accounted for in climate change modelling	Model temperature and microbial activity to assess climate impacts of inorganic carbon in soils

Table 1 continued

Criticism	Articles	Proposed explanation and action	Associated research needs
Better measurement and monitoring are needed to implement the initiative	White et al. (2018)	Use best available methods for measurement and activity. Improve and disseminate measurement guidelines.	Developing high through-put and low cost methods to monitor changes in SOC stocks
Many soils are already well managed therefore presenting limited opportunities to increase SOC storage	White et al. (2018)	Concerns only certain regions; the majority of agricultural soils is not managed sustainably	Maintain best management practices. Identify most promising sites
Socio-economic			
Farmers will not be able to adopt practices due to social and institutional and economic constraints (costs, need for continuous financial incentives)	White et al. (2018), Poulton et al. (2018), Baveye et al. (2018)	Address first farm sustainability (SOC storage is likely to also lead to success in sustainable production). Demonstrate the benefits of soil carbon and related incentives. Identify whether benefits outweigh costs. Capacity building. Develop policies.	Quantify the benefits of SOC increase on productivity and resilience, so that a monetary value can be attributed to SOC increases. Show levels of sequestration possible based on different carbon costs.
Political			
The 4p1000 is proposed to avoid making any changes in community lifestyle	White et al. (2018)	A strategy reducing the fossil fuel consumption of communities is out of scope for the Initiative but the Initiative contributes to the much broader Paris agreement of the UNFCCC	na
Overall credibility of the soil science community is weakened	Baveye et al. (2018)	Even additional storage of less than 4% would help mitigate CO ₂ emissions. The 4p1000 Initiative is an aspirational target to contribute to climate change mitigation	Improve estimates of SOC sequestration potential at the local to the global scale

to increase organic carbon inputs. We suggest that improved management practices of agricultural systems are required in order to recycle carbon back to soil. These can be achieved through permanent soil cover, reduced carbon exports (e.g. recycling rather than burning crop residues) or following input of exogenous organic amendments (Chabbi et al. 2017; Chenu et al. 2019).

When management practices leading to increasing SOC stocks are applied, the sequestration rate will decrease as the SOC stock approaches a new equilibrium, beyond which further sequestration will be negligible (Fig. 1; Sommer and Bossio 2014; Chenu et al. 2019). Modelling has shown that increases in SOC sequestration can continue for 20 years globally (Sommer and Bossio 2014) and even up to 120 years for specific agricultural practices and pedoclimatic conditions (Poeplau and Don 2015). However, it is likely that SOC sequestration will not continue indefinitely and that its contribution to mitigating climate warming is time-limited. Permanence of SOC storage will not only depend on the continuity of best management practices but also on the forms of carbon that comprise SOC stocks and stability of pedoclimatic conditions, which may be compromised by climate change. SOC sequestration is only part of the solution to mitigate climate change

and must be complemented with other mitigation initiatives that will lead to aggressive and urgent reductions in all greenhouse gas emissions.

Several authors have raised concerns about the nutrients needed for increasing SOC sequestration (de Vries et al. 2018; van Groenigen et al. 2017). In mineral soils, nutrients are needed to achieve increases in SOC sequestration because they (1) increase plant production and therefore carbon input into soil (Ladha et al. 2011) and (2) build up stable (mineral associated) SOC (Kirkby et al. 2014). In particular, estimates of the amounts of nitrogen and phosphorus required to increase SOC stocks on agricultural land globally were deemed unrealistic (van Groenigen et al. 2017; de Vries et al. 2018). The nutrient cost of SOC sequestration may be addressed by (1) optimising nutrient management through improved farm management practices (Ditzler et al. 2018), (2) incorporating spatially-differentiated SOC sequestration strategies into precision agriculture and (3) using green manure legumes instead of mineral fertilisers (Soussana et al. 2017). Use of exogenous amendments in the form of farm manure and compost may be part of improved nutrient management practices while additionally contributing to increasing SOC stocks (Diacono and Montemurro 2010).




Fig. 1 SOC trajectories after adoption of improved management practices. Adapted from Lal (2004)

However, their local application could result in major carbon and nutrient transfers from other locations with no net increase in SOC sequestration, and possible increases in other greenhouse gas emissions (Powelson et al. 2011; Poulton et al. 2018). Exceptions are where the biomass would otherwise be burned or deposited into landfills. In this context, the recycling of organic wastes from domestic activities and urban areas as organic fertilisers is an opportunity to transfer organic carbon in ways that enhance SOC storage, ameliorate the nutrient content of soils and close nitrogen and phosphorus cycles at regional scales (Chabbi et al. 2017; Minasny et al. 2018; Nath et al. 2018). Use of amendments containing organic carbon in thermally stable forms, (biochar), while being a practical way of recycling organic wastes, may avoid inputs of nitrogen and phosphorus to form SOC because of their low concentrations of both elements. Peatland restoration is another option for sequestering SOC with minimal nitrogen inputs due to the high carbon to nitrogen ratios of peatland plants (Leifeld and Menichietti 2018).

Important biophysical issues that possibly limit SOC storage potential are related to the (1) inherent capacity of soil to store carbon in a stable form, (2) longevity of the additional stored carbon, (3) reversibility if C retaining practices are not maintained and (4) scarcity of crop residues or other biomass and nutrient inputs for soil amendment. We acknowledge these limitations, but suggest that there are many possibilities for improving nutrient and organic residue management at farm, region and national scales, which could be exploited to maintain and if possible increase SOC stocks and improve soil quality. As concluded by van Groeningen et al. (2017), a spatially diversified strategy is needed for climate change mitigation from agricultural soils. Research to develop new innovative technologies is also required.

Socio-economic barriers

The feasibility of SOC increases will depend on the abilities of farmers to implement changes to management practices as driven by their equipment, skills, operational and economic constraints. Farmers are likely to implement management changes only if there are clear co-benefits, in terms of yields and long-term economic profitability. Some authors have suggested that the achievement of 0.4% SOC increase will not be feasible since farmers are unlikely to adopt new management practices given the low trading price of carbon and more profitable alternative uses of carbon-rich materials (White et al. 2018; Poulton et al. 2018). However, the trading price of carbon is likely to increase with increasing focus on climate change mitigation and adaptation policies providing strong incentives for farmers (Frank et al. 2017). Adoption of novel practices or systems may also require cultural adaptation, as new practices present risks for farmers, when there is insufficient support from farm advisors or where there are vested interests. Smallholder farmers in developing countries may be less interested in change because they are more vulnerable to impacts on food security and community well-being (Lal 2019). In some developing countries, gender inequality, social exclusion, lack of land rights and/or tenure security, and lack of education impede the adoption of new practices, compounded by the lack of financial resources (Nath et al. 2018; Corbeels et al. 2019). However, there are documented ways to overcome these constraints in at least some locations (Pan et al. 2017). Support for information exchange, finance and capacity building can also enable farmers to adopt more innovative practices. One example is the adoption of biochar technology which, despite being a promising option to improve soil quality and increase SOC stocks (Marousek et al. 2017), remains unknown to many farmers and uneconomic to implement due to high demand for organic residues from other sectors and high transportation costs.

RISKS AND TRADE-OFFS

Emissions of greenhouse gases and water use

Non-CO₂ greenhouse gas emissions with a much higher global warming potential may limit the climate change mitigation potential of SOC sequestration. These include N₂O emissions following mineral fertilisation, CH₄ and N₂O emissions from ruminant livestock and CH₄ and N₂O emissions from rice production systems. Practices promoted by the 4p1000 Initiative need to take them into account to ensure that net greenhouse emissions do not exceed the offset benefit from increased SOC

sequestration. The trade-off effects between greenhouse gas emissions and SOC sequestration may be dynamic. For example, if fertiliser applications are not reduced, increases in SOC sequestration may no longer offset N₂O emissions when the system is approaching a new equilibrium for SOC storage (Lugato et al. 2018). These dynamic processes need to be evaluated carefully, and should be considered when actions to increase SOC stocks are undertaken.

One critical issue, not yet addressed, is the effect of SOC sequestration on the water balance of (agro-) ecosystems. For example, Jackson et al. (2005) showed that C sequestration in woody biomass reduced water availability for consumption because of increased water loss from the evaporation of intercepted rainfall. In many agricultural systems, irrigation is used to enhance productivity with variable impacts on SOC sequestration (Trost et al. 2013). Especially under arid conditions, water is needed for (1) additional biomass production and thus carbon release into soils, (2) microbial activity to transform plant litter compounds into refractory SOC and (3) compensation of water loss in plants, due to high evapotranspiration, as water is needed for photosynthesis. On the other hand, improvements in soil structure when increasing soil organic matter content have positive effects on soil water retention and infiltration (Pittelkow et al. 2015). These interrelationships need to be considered as well as the fact that water shortage following climate change may put at risk SOC in systems with permanent water-logging (exp. Paddy rice).

Avoiding emissions from SOC-rich soils

SOC-rich soils and organic soils are among the most fertile sites but some are heavily exploited for agricultural production, often at the expense of maintaining SOC stocks, leading to large releases of CO₂ to the atmosphere (Leifeld and Menichetti 2018). Globally, peatlands occupy only 3% of land area but are estimated to store about 600 Gt of SOC. This corresponds to around 20% of SOC stored in the first 30 centimetres of soils globally (Scharlemann et al. 2014). Natural peatlands are characterised by continuous water-logging, limiting organic matter decomposition because of low oxygen supply. For this reason, avoiding further drainage of intact peatland soils should be a priority. Many of these soils are under agricultural management and major contributors to greenhouse gas emissions. A recent analysis showed that degraded peatlands globally store ~ 80.8 Gt of soil C with emissions dominantly from tropical regions of ~ 1.91 (range 0.31–3.38) Gt CO₂-eq. year⁻¹ (Leifeld and Menichetti, 2018). The authors also showed that the global greenhouse gas emissions estimated from cultivated peatlands may completely offset the SOC sequestration potential of mineral soils. Therefore, in humid regions, careful management of water-logging may

be required to ensure that losses from the large amounts of SOC stored in peatland soils are minimised.

THE 4P1000 INITIATIVE AS A COLLABORATIVE PLATFORM FOR POLICY-SCIENCE-PRACTICE INTERACTIONS

Increasing terrestrial biosphere carbon sinks could contribute to achieving the ambitious climate change mitigation target of limiting the increase in global average temperature to well below 2 °C above pre-industrial levels by offsetting emissions. The use of bioenergy with carbon capture and storage (BECCS), biochar and SOC sequestration have been presented as possibilities (IPCC 2006). It is apparent that SOC sequestration is the most viable option because it (1) has been tested, (2) is feasible at large spatial scales, (3) does not constrain the use of land and (4) provides potential co-benefits to meet other SDGs (Smith 2016). The 4p1000 Initiative attracted attention because it addresses many social issues related to agriculture that impact widely on communities and integrates engagement from many disciplines and sectors. The Initiative addresses global issues to mitigate greenhouse gas emissions and food security and, at the same time, local issues to improve soil quality and agricultural production. However, this broad application also leads to difficulties in engaging adoption to implement the necessary actions. While there are already other initiatives to promote SOC sequestration and improve soil quality, such as the Global Soil Partnership, the 4p1000 Initiative provides a platform to encourage interactions among scientists, policy makers and practitioners (farmers, NGOs, funders...). This tripartite collaboration is important to ensure that policy decisions are based on credible research and that scientific findings are implemented to meet local needs. The biggest challenge to the success of the 4p1000 Initiative is to stimulate collaboration across the breadth of collaborators to agree on actions and their implementation to achieve the target of the Initiative. It should serve as a catalyst to enhance information exchange and collaboration, leading to joint actions by a wide range of stakeholders.

THE WAY FORWARD

The controversy resulting from the initial articulation of the goal of the Initiative has been helpful to promote scientific rigour and policy debate to formulate action. After successful engagement with stakeholders, and elaboration of criteria to assess management actions by the Scientific and Technical Committee of the Initiative (Fig. 2), the next challenge is to build on tripartite engagement between

policy makers, scientists and practitioners to promote implementation of best practices. To support the implementation, the 4p1000 Initiative must provide linkages with action plans, contributions and agricultural development projects at national scales. Progress was made at COP of the UNFCCC in Bonn in 2017, where discussion of agriculture and the role of soil carbon stocks were included for the first time in the Koronivia Decision on joint work of the subsidiary body for scientific and technological advice (SBSTA) and the subsidiary body for implementation (SBI) (UNFCCC 2018). Eight steps for achieving increased SOC sequestration were recently presented. These include protection of existing SOC stocks, e.g. in organic soils, promotion of C uptake through new practices and regulations, monitoring, reporting and verifying impact through advanced analytical techniques and data harmonisation. New strategies need to be tested and communities must be involved. Further, education, identification and coordination of policies as well as provision of financial support to help farmers, who use sustainable SOC improving practices is required (Rumpel et al. 2018). To increase public awareness about the necessity to increase SOC stocks, the Initiative promotes SOC sequestration to a wide audience, including farmers and land managers, agricultural suppliers of resources, other contributors to the supply chain, central and local governments, urban waste managers and consumers, etc. The 4p1000 Initiative will take advantage of existing online tools and create an interactive platform to support exchange between multiple partners with different roles and from different geographical regions and cultures. It is essential to communicate success stories of increasing SOC sequestration in different pedoclimatic conditions and different agricultural management systems. Moreover, further investment in research and the development of innovative technologies will be needed to provide stronger support for the 4p1000 Initiative. In addition, the Scientific

and Technical Committee of the Initiative established a research programme (STC 2017). This programme comprises four pillars: (1) Estimation of the SOC storage potential, (2) Development of management practices, (3) Definition of the enabling environment and (4) Monitoring, reporting and verification. Within each of these pillars, key knowledge gaps have been identified and these need to be promoted to engage activities by research organisations and promote investment in these areas. To initiate implementation of C sequestering options that are relevant to local conditions and embrace farmer knowledge along with research findings, innovative learning networks linking farmers, technical assistance organisations, scientists and policy makers are also required. This can be achieved by establishing living labs and networks of demonstration farms to better communicate successful management practices based on rigorous research findings. The 4p1000 Initiative, as an international multi-participant programme, will facilitate adoption of the best management practices and innovative technologies by providing information and promoting international collaboration at all levels (Rumpel et al. 2018; Lal 2019).

CONCLUSIONS

The ‘4 per 1000’ Initiative aims to increase carbon storage in agricultural soils and therefore contributes to mitigating climate change, adapting to climate change and increasing food security (<http://www.4p1000.org>). The Initiative has potential as an international multi-disciplinary platform combining a recommended research programme with a multi-stakeholder action plan to link scientific research and action. It aims to communicate and promote management actions to increase SOC sequestration through implementation of sustainable development practices. The main




Fig. 2 Criteria that need to be met by management actions implemented under the 4p1000 Initiative (STC, 2017)

strength of the Initiative is that it provides a collaborative space for engagement and discussion between contributors (scientists, practitioners, NGOs, private sector and policy makers) from different educational and cultural backgrounds. With its simple message, the Initiative encourages widespread participation and adoption by many partners. Recent clarification of the initial message has strengthened the rationale for the Initiative. It is clear that SOC sequestration has the potential to offset greenhouse gas emissions to contribute to aggressive, large-scale, urgent reductions in greenhouse gas emissions, as well as to improving food security and climate change adaptation. However, the potential of soils to sequester SOC is limited by biophysical, socio-economic and political barriers. These need to be overcome by region-specific actions and the development and implementation of innovative technologies. While SOC sequestration can make a significant contribution to climate change mitigation, the more certain and principal benefits, especially those on degraded land, will be improvements in soil quality, contributing to food security and agricultural systems that are more resilient to climate change. To achieve this, priorities will need to be decided to ensure that actions are focused on sites and conditions where opportunities to increase soil carbon stocks are most likely to be successful. We conclude that the 4p1000 Initiative is likely to facilitate findings from site-specific studies, practical experiences and model predictions to be incorporated into future policy actions to encourage long-term adoption and implementation of sustainable development strategies.

Acknowledgements Authors would like to acknowledge the executive secretariat of the 4p1000 initiative, Charlotte Verger and Claire Weill for their valuable contributions during the preparation of this manuscript. The input of PS contributes to the UK NERC-funded Soils-R-GGREAT project (NE/P019455/1).

REFERENCES

- Barre, P., H. Durand, C. Chenu, P. Meunier, D. Montagne, G. Castel, D. Billiou, L. Soucemarianadin, et al. 2017. Geological control of soil organic carbon and nitrogen stocks at the landscape scale. *Geoderma* 285: 50–56.
- Baveye, P.C., J. Berthelin, D. Tessier, and G. Lemaire. 2018. The “4 per 1000” initiative: A credibility issue for the soil science community? *Geoderma* 309: 118–123.
- Chabbi, A., J. Lehmann, P. Ciais, H.W. Loescher, M.F. Cotrufo, A. Don, M. SanClements, L. Schipper, et al. 2017. Aligning agriculture and climate policy. *Nature Climate Change* 7: 307–309.
- Chenu, C., D.A. Angers, P. Barré, D. Derrien, D. Arrouays, and J. Balesdent. 2019. Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations. *Soil and Tillage Research* 118: 42–51.
- Corbeels, M., K. Naudin, H. Guibert, E. Torquebiau, and R. Cardinael. 2019. Is 4 per 1000 soil carbon storage attainable with agroforestry and conservation agriculture in sub-Saharan Africa? *Soil & Tillage Research* 188: 16–26.
- de Vries, W. 2018. Soil carbon 4 per mille: A good initiative but let's manage not only the soil but also the expectations: Comment on Minasny et al. (2017). *Geoderma* 292: 59–86.
- Diacono, M., and F. Montemurro. 2010. Long-term effects of organic amendments on soil fertility. A review. *Agriculture for Sustainable Development* 30: 401–422.
- Ditzler, L., T.A. Breland, C. Francis, M. Chakraborty, D.K. Singh, A. Srivastava, F. Eyhorn, J.C.J. Groot, et al. 2018. Identifying viable nutrient management interventions at the farm level: The case of smallholder organic Basmati rice production in Uttarakhand, India. *Agricultural Systems* 161: 61–71.
- Frank, S., P. Havlík, J.F. Soussana, A. Levesque, H. Valin, L. Wollenberg, U. Kleinwechter, O. Fricko, et al. 2017. Reducing greenhouse gas emissions in agriculture without compromising food security? *Environmental Research Letters* 12: 105004.
- Fujisaki, K., T. Chevallier, L. Chapuis-Lardy, A. Albrecht, T. Razafimbelo, D. Masse, and J.-L. Chotte. 2018. Soil carbon stock changes in tropical croplands are mainly driven by carbon inputs: A synthesis. *Agriculture, Ecosystems & Environment* 259: 147–158.
- Hutchinson, J.J., C.A. Campbell, and R. Desjardins. 2007. Some perspectives on carbon sequestration in agriculture. *Agricultural and Forest Meteorology* 142: 288–302.
- IPCC. 2006. 2006 IPCC guidelines for national greenhouse gas inventories. In *Prepared by the national greenhouse gas inventories programme*, eds. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe. Japan: IGES.
- Jackson, R.B., E.G. Jobbágy, R. Avissar, S.R. Baidya, D.F. Barrett, C.W. Cook, K.A. Farley, D.C. le Maitre, et al. 2005. Trading water for carbon with biological carbon sequestration. *Science* 5756: 1944–1947.
- Kirkby, C.A., A.E. Richardson, L.J. Wade, J.B. Passioura, G.D. Batten, C. Blanchard, and J.A. Kirkegaard. 2014. Nutrient availability limits carbon storage in agricultural soils. *Soil Biology and Biochemistry* 68: 204–209.
- Kon Kam King, J., C. Granjou, J. Fournil, and L. Cecillon. 2018. Soil sciences and the French 4 per 1000 Initiative—The promises of underground carbon. *Energy Research & Social Science* 45: 144–152.
- Ladha, J.K., C.K. Reddy, A.T. Padre, and C.V. Kessel. 2011. Role of nitrogen fertilization in sustaining organic matter in cultivated soils. *Journal of Environmental Quality* 40: 1756–1766.
- Lal, R. 2004. Soil carbon sequestration impacts on global climate change and food security. *Science* 304: 1623–1627.
- Lal, R. 2019. Promoting “4 Per Thousand” and “Adapting African Agriculture” by south-south cooperation: Conservation agriculture and sustainable intensification. *Soil and Tillage Research* 118: 27–34.
- Leifeld, J., and L. Menichetti. 2018. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nature Communications* 9: 1071.
- Lugato, E., A. Leip, and A. Jones. 2018. Mitigation potential of soil carbon management overestimated by neglecting N₂O emissions. *Nature Climate Change* 8: 219–223.
- Maroušek, J., M. Vochozka, J. Plachý, and J. Žák. 2017. Glory and misery of biochar. *Clean Technologies and Environmental Policy* 19: 311–317.
- Minasny, B., D. Arrouays, A.B. McBratney, D.A. Angers, A. Chambers, V. Chaplot, and L. Winowiecki. 2018. Rejoinder to Comments on Minasny et al., 2017 Soil carbon 4 per mille. *Geoderma* 292: 59–86.
- Nath, J.A., R. Lal, G.W. Siles, K. Dasa, and A.K. Das. 2018. Managing India's small landholder farms for food security and

- achieving the “4 per Thousand” target. *The Science of the Total Environment* 634: 1024–1033.
- Pan, W.L., W.S. Schillinger, F.L. Young, E. Kirby, and G.G. Yorgey, et al. 2017. Integrating old principles and new technologies into win-win scenarios for farm and climate. *Frontiers in Environmental Science*. <https://doi.org/10.3389/fenvs.2017.00076>.
- Paustian, K., J. Lehmann, S. Ogle, D. Reay, G.P. Robertson, and P. Smith. 2016. Climate-smart soils. *Nature* 532: 49–57.
- Pingali, P.L. 2012. Green revolution: impacts, limits and the path ahead. *PNAS* 109: 12302–12308.
- Pittelkow, C.M., X. Liang, B.A. Linquist, K.J. van Groenigen, J. Lee, M.E. Lundy, N. van Gestel, J. Six, et al. 2015. Productivity limits and potentials of the principles of conservation agriculture. *Nature* 517: 365–368. <https://doi.org/10.1038/nature13809>.
- Poepflau, C., and A. Don. 2015. Carbon sequestration in agricultural soils via cultivation of cover crops—A meta-analysis. *Agriculture, Ecosystems & Environment* 200: 33–41.
- Poulton, P., J. Johnston, A. MacDonald, R. White, and D. Powlson. 2018. Major limitations to achieving “4 per 1000” increases in soil organic carbon stock in temperate regions: Evidence from long-term experiments at Rothamsted Research. *UK. Global Change Biology* 12: 3218–3221.
- Powlson, D.S., A.P. Whitmore, and A.W.T. Goulding. 2011. Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science* 62: 42–55.
- Rumpel, C., F. Amiraslani, L.-S. Koutika, P. Smith, D. Whitehead, and E. Wollenberg. 2018. Put more carbon in soils to meet Paris climate pledges. *Nature* 564: 32–34.
- Sanderman, J., C. Creamer, W.T. Baisden, M. Farrell, and S. Fallon. 2017. Greater soil carbon stocks and faster turnover rates with increasing agricultural productivity. *Soil* 3: 1–16.
- Scharlemann, J.P.W., E.V.J. Tanner, R. Hiederer, and V. Kapos. 2014. Global soil carbon: understanding and managing the largest terrestrial carbon pool. *Carbon Manag.* 5: 81–91.
- Schiefer, J., G.J. Lair, C. Lütgens, E.M. Wild, P. Steiner, and W.H. Blum. 2018. The increase of soil organic carbon as proposed by the “4/1000 initiative” is strongly limited by the status of soil development—A case study along a substrate age gradient in Central Europe. *The Science of the Total Environment* 628–629: 840–847.
- Smith, P. 2016. Soil carbon sequestration and biochar as negative emission technologies. *Global Change Biology* 22: 1315–1324.
- Smith, P., S.J. Davis, F. Creutzig, S. Fuss, J. Minx, B. Gabrielle, et al. 2016. Biophysical and economic limits to negative CO₂ emissions. *Nature Climate Change* 6: 42–50.
- Sommer, R., and D. Bossio. 2014. Dynamics and climate change mitigation potential of soil organic carbon sequestration. *Journal Environmental Management* 144: 83–87.
- Soussana, J.F., S. Lutfalla, R. Lal, C. Chenu, and P. Ciais. 2017. Letter to the editor: answer to the viewpoint “sequestering soil organic carbon: a nitrogen dilemma” by van Groenigen et al. (2017). *Environmental Science and Technology* 51: 11502.
- Soussana, J.F., S. Lutfalla, F. Ehrhardt, T. Rosenstock, C. Lamanna, P. Havlík, and R. Lal. 2019. Matching policy and science: Rationale for the “4 per 1000 - soils for food security and climate” initiative. *Soil and Tillage Research* 188: 3–15.
- STC. 2017. The ‘4 per 1000’ Research Priorities. https://www.4p1000.org/sites/default/files/content/gov_cst_en_consortium_3-4-4p1000_research_priorities.pdf.
- Tilman, D., C. Balzer, J. Hill, and B.L. Befort. 2011. Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Science USA* 108: 20260–20264.
- Trost, B., A. Prochnow, K. Drastig, A. Meyer-Aurich, F. Ellmer, and M. Baumecker. 2013. Irrigation, soil organic carbon and N₂O emissions. *A review. Agronomy for Sustainable Development* 33: 733–749.
- UNFCCC. 2018. Koronivia Joint Work on Agriculture (decision 4/CP.23).
- Van Groenigen, J.W., C. Van Kessel, B.A. Hungate, O. Oenema, D.S. Powlson, and K.J. Van Groenigen. 2017. Sequestering soil organic carbon: A nitrogen dilemma. *Environmental Science and Technology* 51: 4738–4739.
- VandenBygaart, A.J. 2018. Comments on soil carbon 4 per mille by Minasny et al. 2017. *Geoderma* 309: 113–114.
- White, R.E., B. Davidson, S.K. Lam, and D. Chen. 2018. A critique of the paper “Soil carbon 4 per mille” by Minasny et al. (2017). *Geoderma* 309: 115–117.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

AUTHOR BIOGRAPHIES

Cornelia Rumpel (✉) is a research director working for the French National Research Center (CNRS) at the Institute of Ecology and Environmental Sciences Paris. She studied forestry in Germany and Scotland and received a master diploma from the Ludwig-Maximilians University at Munich, Germany in 1994. Afterwards, she worked as a research scientist at the Brandenburg University of Technology in Cottbus, Germany, where she obtained her PhD degree in natural sciences in 1999. She studied the origin and fate of terrestrial organic matter, aiming to understand the mechanisms controlling carbon sequestration in soils. Her studies concerned various spatial and temporal scales in different environments ranging from soils and sediments in mining areas to natural as well as managed ecosystems, including those affected by fire. Her work was carried out in temperate and tropical climates and the results of her research changed of a number of paradigms. She published >160 papers, which were cited more than 7500 times. In 2016 and 2017 she was listed as a highly cited researcher. She was nominated ambassador of the Technical University of Munich and has been the chair of the STC of the 4p1000 initiative since 2018.

Address: CNRS, Institute for Ecology and Environmental Sciences, Thiverval-Grignon, France.

e-mail: cornelia.rumpel@inra.fr

Farshad Amiraslani has been involved in dryland management and research over the last 19 years. He received his PhD from the University of Sydney and is currently serving as Associate Professor (University of Tehran). He has been a Cheney Fellow (University of Leeds) and Research Fellow at Chinese Academy of Sciences (UNEP-IEMP). He has been elected or invited for various consultancies and publication assignments for the UN and international organisations over the past. He has won several international awards and fellowships and published over 60 publications.

Address: Department of RS/GIS, Faculty of Geography, University of Tehran, Tehran, Iran.

Claire Chenu is professor of soil science at AgroParisTech, the leading French technical University in the field of biology, agronomy, food and environmental sciences. Her research deals with soil organic carbon: dynamics, stabilisation processes and the effect of cropping practices on soil C stocks. She is very involved in the science-policy-practice interface and in awareness raising activities on soils. She has been nominated Special Ambassador of soils in 2015 by the FAO. She is member of several international committees, including being a vice

chair of the Scientific and Technical Committee of the 4 per 1000 initiative.

Address: AgroParisTech, UMR Ecosys INRA, AgroParisTech, Université Paris-Saclay, Thiverval-Grignon, France.

Magaly Garcia Cardenas is a Bolivian researcher specialized on Climate change and trends analysis of impacts on local communities. She obtained her PhD and previous post-graduate diplomas at Dutch and Belgian Universities. She works especially on farmers' capacity for response and adaptation to climate change and strategies to enhance farmers reactions and better decisions systems through the availability of adequate climatic information. Lately, she is strongly involved in evaluation of Soil Organic Carbon contents in high altitude (>3800 m) agricultural watersheds to suggest strategies for management and identification of Carbon sinks potentialities in tropical high altitude environments.

Address: Faculty of Engineering, Universidad Mayor de San Andrés, La Paz, Bolivia.

Martin Kaonga is Director of Cambridge Centre for Environment in the United Kingdom. He obtained a PhD in agroforestry carbon biogeochemistry from Cambridge University (UK). Martin has more than 15 years' experience in terrestrial ecosystem carbon science and is the author of more than 15 peer-reviewed papers, and the editor of a book entitled 'Agroforestry for Biodiversity and Ecosystem Services: Science and Practice.' Following a PhD in plant and soil carbon storage and fluxes in agroforestry systems, he took up a position as Director of Conservation Projects at A Rocha International (ARI) in the United Kingdom. Managing a climate change mitigation projects in northern Ghana, Peru and South Africa, he validated baseline plant and soil carbon stocks and simulated changes over a 50-year project cycle using the CO2Fix model. He also developed an aggregator model for recruitment of new project sites and for carbon accounting. After two years in post at ARI, Martin was appointed as the Director of Science and Conservation, responsible for conservation research and community-based conservation projects in 19 countries. He designed and directed four programmes: Tropical Forest Programme; Mediterranean Conservation Science Programme; Marine and Coastal Research Programme and European Conservation Programme involving nine countries in western Europe. He directed over 50 projects in 19 countries, including terrestrial and aquatic biodiversity and carbon research projects. These projects had strong components of sustainable agriculture and food security, and climate change adaptation and mitigation. Martin designed and led three carbon research projects, which assessed altitudinal and climate variability effects on biodiversity and carbon dynamics (France); vegetation carbon dynamics and floristic diversity in dry forests of in Eastern Ghats (India) and SOC storage in forest/agroforest ecosystems in Ghana.

Address: Cambridge Center for Environment, Cambridge, UK.

Lydie-Stella Koutika is a soil scientist and Deputy Director at the Forestry Research Centre (CRDPI), Pointe-Noire, Congo. She works on soil organic matter (both C and N) and phosphorus dynamics in the mixed-species of eucalypts and Acacia mangium plantations established in the Congolese coastal plains. She obtained an engineer degree in agronomy (Timiriachev Institute, Moscow) and a PhD in soil science (Université Nancy I, merged to Université de Lorraine, France). She is a former fellow "Research in Brussels" (2003–2004), Belgium and Rothamsted International (2009–2010), UK. She is currently a TWAS-ENEA International Fellow. She was a laureate of the AU 'Kwame Nkrumah Regional Scientific Award for Women' (2014) and TWAS-AI-Kharafi Prize (2018).

Address: CRDPI, Av. Ma Loango Moe Poaty, BP 1291 Pointe-Noire, Republic of the Congo.

e-mail: ls_koutika@yahoo.com

Jagdish Ladha has devoted more than 35 years to aspects of sustainable management of agriculture and natural resources for increasing food security and environmental quality in developing countries. He is an expert of soil fertility and plant nutrition; serving at different positions since 1980. He served at the International Rice Research Institute (IRRI) from 1980 to 2017 was an adjunct senior scientist at the Columbia University until 2016. Currently, he is an Adjunct Professor in the Department of Plant Sciences at the University of California-Davis. He is a fellow of the American Association for the Advancement of Science (AAAS), American Society of Agronomy (ASA), the Soil Science Society of America (SSA), the Crop Science Society of America (CSSA), the Indian Academy of Agricultural Sciences (NAAS), and an associate member of the Philippine Council of Agricultural Research (PARC).

Address: Department of Plant Sciences, University of California, Davis, USA.

Beata Madari is agronomist (1994) with PhD (1999) in Soil Science from the Szent István University, Gödöllő, Hungary. She worked as researcher at the National Soil Research Center of the Brazilian Agricultural Research Corporation (Embrapa) between 2002 and 2005. Since 2005 she is a scientist at the Embrapa National Research Center for Rice and Bean and is Professor in post-graduate training at the School of Agronomy of Federal University of Goiás, Brazil. She was leader of the Embrapa Research Network on Greenhouse Gas Emissions from Grain Crop Production Systems (Embrapa Fluxus Network) and is presently member of the Executive Committee of the Climate Change Portfolio of Projects of Embrapa. Accordingly, she has experience in carbon and nitrogen cycling in terrestrial ecosystems, particularly regarding tropical acid soils under annual crops, but also in integrated crop-livestock-forestry systems. She has knowledge on soil carbon dynamics and physical and chemical fractionation of soil organic matter. She has worked with methods of soil carbon determination using wet and dry combustion and infrared spectroscopy. She is Fellow Scientist of the Brazilian Council on Science and Technology Development (CNPq) and reviewer of several international journals. She has also contributed to the IPCC on HWP, Wetlands and Soil N₂O and to the UN Global Compact Initiative (unglobalcompact.org). She is member of the Scientific and Technical Committee of the 4 per 1000 Initiative and of FAO's Livestock Environmental Assessment Partnership's technical advisory group on soil organic carbon stock change.

Address: Brazilian Agricultural Research Corporation, National Rice and Bean Research Center (Embrapa Arroz e Feijão), Santo Antônio de Goiás, Brazil.

Yasuhito Shirato is a soil scientist and currently Research Manager for Climate Change, Institute for Agro-Environmental Sciences, NARO (National Agriculture and Food Science Organization), Japan. Dr. Shirato has been working on soil carbon dynamics mainly on modelling. Dr. Shirato developed country-scale calculation system of soil carbon in agricultural land, which is now used for Japanese National Inventory Report of Greenhouse Gases for United Nations Framework Convention on Climate Change (UNFCCC). Dr. Shirato is a member of the Scientific and technical Committee of "the 4 per 1000 Initiative".

Address: National Agriculture and Food Research Organization, Tsukuba, Japan.

Pete Smith is the Professor of Soils and Global Change at the Institute of Biological and Environmental Sciences at the University of Aberdeen (Scotland, UK) and Science Director of the Scottish Climate Change Centre of Expertise (ClimateXChange). Since 1996, he has served as Convening Lead Author, Lead Author and Author for the Intergovernmental Panel on Climate Change (IPCC), which was awarded the Nobel Peace Prize in 2007. He was the Convening Lead

Author of the Agricultural Mitigation chapter of the IPCC Fourth Assessment Report and for the Agriculture and Forestry Mitigation chapter of the IPCC Fifth Assessment, and is currently Convening Lead Author for the IPCC Special report on Climate Change and Land. He has coordinated and participated in many national and international projects on soils, agriculture, bioenergy, food security, greenhouse gases, climate change, mitigation and impacts, and ecosystem modelling. In addition to his role in ClimateXChange, he is a former member of Defra's Science Advisory Council, and a current member of DfID's Research Advisory Group and the Global Food Security Science Advisory Board, and has been an advisor to the Committee on Climate Change. He is a Fellow of the Royal Society of Biology, a Fellow of the Institute of Soil Scientists, a Fellow of the Royal Society of Edinburgh, a Foreign Fellow of the Indian National Science Academy, a Fellow of the European Science Academy, and a Fellow of the Royal Society (London). He has published >440 peer-reviewed journal papers with total citations of >23000 with an H-index of 82. He has been a Highly Cited Researcher each year since 2015.

Address: Institute of Biological & Environmental Sciences, University of Aberdeen, Aberdeen, UK.

Brahim Soudi born in the Province of Taroudant, in 1955, is Agronomist from Hassan II Institute of Agronomy and Veterinary Medicine (IAV), in 1982 and he obtained his PhD degree at the Catholic University of Leuven, Belgium in soil Sciences and biological chemistry Department. He stayed at the USA (University of Minnesota and University of Cornell) in 1991 as part of his Post-Doctorate on modelling of Nitrogen and Organic Matter mineralization. He is a professor at the IAV since 1982 where he is responsible for several courses including: basic soil science, management of soil organic matter, biogeochemical cycling, and recycling of organic waste, composting and recycling compost, soil and water monitoring systems under intensive agriculture, environmental assessments, etc. He also coordinated several training modules for the benefit of technical staff within Ministry of Agriculture and Ministry of Environment. In the area of scientific research, he participated and coordinated several research projects for national institutions and in the framework of international cooperation, particularly in the following fields: impact of intensive agriculture on soil quality and water, the status of organic matter in agricultural systems in arid and semi-arid zones, modelling of the transfer and transformation of nitrogen, the dynamics of organic matter in salt-affected lands, establishment and optimising soil and water quality monitoring in irrigated areas, soil degradation under irrigation, etc. In connection with this research activity, he supervised many works graduation and PhD theses. He is author and co-author of over 60 scientific papers and author of books and guidance manuals. It is also reviewer for some scientific journals.

Address: Institut Agronomique et Vétérinaire Hassan II, Rabat, Morocco.

Jean-François Soussana Since March 2017, Jean-François Soussana is Vice-Chair for international affairs at INRA, Paris, France. He obtained his PhD in plant physiology at USTL Montpellier in 1986 after an engineer degree in agronomy. After becoming a senior

scientist, he led a research lab. on grassland ecosystems and global change and was Scientific Director for Environment at INRA (2010–2017). Dr. Soussana is member of the Working Group II of IPCC since 1998 and is currently Lead Author for the Special Report on Climate Change and Land. He contributes to scientific expertise for FAO (State of Food and Agriculture, 2016), co-chairs the Integrative Research Group of the Global Research Alliance on agricultural greenhouse gases (56 countries) and the Steering Council of AgMIP, an international modeling program on climate change impacts on agriculture. Dr. Soussana is a Governing Board member of the joint programming of research on agriculture, food security and climate change (FACCE JPI) and of the Climate KIC of the European Institute of Technology. He is also vice-chair of the Scientific and Technical Committee of the “4 per 1000. Soils for Food Security and Climate” initiative and of the TempAg Collaborative Research Network on temperate agriculture and a member of the French high-level committee on climate. He coordinates European (EC FP5, FP7 and H2020) research projects on climate change and agriculture, has published close to 150 research papers in international journals and is a highly cited researcher (Clarivate analytics). Honours: Shared Nobel Prize for Peace in 2007 with all IPCC co-authors; shared Gerbier-Mumm prize of WMO; gold medal of the French academy for agriculture; commander of the French order of agricultural merit.

Address: INRA, Institut National de la Recherche Agronomique, Paris, France.

David Whitehead works as a scientist at Manaaki Whenua – Landcare Research and the New Zealand Agricultural Greenhouse Gas Research Centre to promote ‘climate-smart agriculture’. David leads a collaborative research programme to investigate ways for management practices to manipulate soil carbon inputs to grasslands that will lead to maintaining soil carbon stocks but reduce leaching losses of nitrogen and nitrous oxide emissions. David received a certificate acknowledging his contribution to the Nobel Peace Prize for 2007 awarded to the Intergovernmental Panel on Climate Change and was elected as a Fellow of the Royal Society of New Zealand in 2012.

Address: Manaaki Whenua – Landcare Research, Lincoln, New Zealand.

Eva Wollenberg leads the low emissions development research programme for CCAFS, the CGIAR Research Program on Climate Change, Agriculture and Food Security, and is a Research Professor at the Gund Institute for Ecological Economics, University of Vermont. She previously worked with the Center for International Forestry Research (CIFOR) in Indonesia for 11 years, and the Ford Foundation. Lini received her BS, MSc and PhD degrees in natural resource management from the University of California, Berkeley, USA. Her expertise includes climate change mitigation, land use and natural resource governance. She has worked primarily in Asia, especially Indonesian Borneo

Address: Gund Institute for Environment, University of Vermont and CGIAR Research Program on Climate Change, Agriculture and Food Security, Burlington, USA.