



HAL
open science

Long-term phytomanagement with compost and a sunflower - Tobacco rotation influences the structural microbial diversity of a Cu-contaminated soil

Aritz Burges, Virgil Fievet, Nadège Oustrière, Lur Epelde, Carlos Garbisu, Jose María Becerril, Michel Mench

► To cite this version:

Aritz Burges, Virgil Fievet, Nadège Oustrière, Lur Epelde, Carlos Garbisu, et al.. Long-term phytomanagement with compost and a sunflower - Tobacco rotation influences the structural microbial diversity of a Cu-contaminated soil. *Science of the Total Environment*, 2020, 700, pp.1-10. 10.1016/j.scitotenv.2019.134529 . hal-02627846

HAL Id: hal-02627846

<https://hal.inrae.fr/hal-02627846v1>

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Long-term phytomanagement with compost and a sunflower – tobacco rotation**
2 **influences the structural microbial diversity of a Cu-contaminated soil**

3

4 Aritz Burges^{a,d,*}, Virgil Fievet^a, Nadège Oustriere^b, Lur Epelde^c, Carlos Garbisu^c, Jose
5 María Becerril^d, Michel Mench^a

6

7 ^aUMR BIOGECO INRA 1202, University of Bordeaux, Bât. B2, allée Geoffroy St-
8 Hilaire, F-33615 Pessac cedex, France

9 ^bLaboratoire Génie Civil et Géoenvironnement (LGCGE), Yncréa Hauts-de-France,
10 Institut Supérieur d'Agriculture, 48 Bld Vauban, 59046 Lille cedex, France

11 ^cNEIKER-Tecnalia, Department of Ecology and Natural Resources, Soil Microbial
12 Ecology Group, c/ Berreaga 1, E-48160 Derio, Spain

13 ^dUniversity of the Basque Country (UPV/EHU), Department of Plant Biology and
14 Ecology, P.O. Box 644, E-48080 Bilbao, Spain

15 *Corresponding author: Dr. Aritz Burges

16 UMR BIOGECO INRA 1202

17 Biodiversity, Genes & Communities

18 University of Bordeaux, Bât. 2

19 Allée Geoffroy St-Hilaire

20 F-33615 Pessac cedex

21 FRANCE

22 E-mail: aritzburges@gmail.com

23

24 **Abstract**

25 At a former wood preservation site contaminated with Cu, various phytomanagement
26 options have been assessed in the last decade through physicochemical,
27 ecotoxicological and biological assays. In a field trial at this site, phytomanagement
28 with a crop rotation based on tobacco and sunflower, combined with the incorporation
29 of compost and dolomitic limestone, has proved to be efficient in Cu-associated risk
30 mitigation, ecological soil functions recovery and net gain of economic and social
31 benefits. To demonstrate the long-term effectiveness and sustainability of
32 phytomanagement, we assessed here the influence of this remediation option on the
33 diversity, composition and structure of microbial communities over time, through a
34 metabarcoding approach. After 9 years of phytomanagement, no overall effect was
35 identified on microbial diversity; the soil amendments, notably the repeated compost
36 application, led to shifts in soil microbial populations. This phytomanagement option
37 induced changes in the composition of soil microbial communities, promoting the
38 growth of microbial groups belonging to *Alphaproteobacteria*, many being involved in
39 N cycling. Populations of *Nitrososphaeria*, which are crucial in nitrification, as well as
40 taxa from phyla *Planctomycetacia*, *Chloroflexi* and *Gemmatimonadetes*, which are
41 tolerant to metal contamination and adapted to oligotrophic soil conditions, decreased in
42 amended phytomanaged plots. Our study provides an insight into population dynamics
43 within soil microbial communities under long-term phytomanagement, in line with the
44 assessment of soil ecological functions and their recovery.

45

46 **Keywords:** phytoremediation; metal pollution; soil functions; organic amendments;
47 metabarcoding.

48

49 **1. Introduction**

50 Long-term phytomanagement of metal(loid)-contaminated soils can (1) reduce soil
51 phytotoxicity, promoting ecological soil functions while preserving the soil resource
52 (Quintela-Sabaris et al., 2017; Mench et al., 2018), and (2) produce raw materials for
53 local biomass processing technologies and incomes for the local communities (Mench et
54 al., 2010; Evangelou et al., 2012; Witters et al., 2012; Van Slycken et al., 2013; Cundy
55 et al., 2016). This was notably shown for Cu-contaminated soils due to mining and
56 wood preservation activities (Kidd et al., 2015; Touceda-González et al., 2017a;
57 Touceda-González et al., 2017b; Mench et al., 2018), which are often characterized by
58 unfavourable soil properties, e.g. lack of structure with low OM content, low nutrient
59 availability and acidic pH (Mench and Bes, 2009; Bes et al., 2010; Hattab-Hambli et al.,
60 2016; Oustriere et al., 2016), .

61 The Biogeco site (a former wood preservation site, St-Médard d'Eyrans, France;
62 Cu-contaminated soils) has received a lot of attention with the aim of demonstrating the
63 benefits of long-term phytomanagement (Kolbas et al., 2011; Kumpiene et al., 2011;
64 Marchand et al., 2011; Hattab-Hambli et al., 2016; Oustriere et al., 2016; Mench et al.,
65 2018). Here, soil amendments, i.e. a single incorporation of compost combined with
66 dolomitic limestone in year 1 (OMDL) and this OMDL treatment followed by compost
67 application renewed in year 6 (OM2DL), and high yielding crops (sunflower, tobacco)
68 improved soil organic matter (OM) and nutrient contents, soil pH and CEC and
69 sustainably decreased Cu availability in year 9 (Mench et al., 2018). Both OMDL and
70 OM2DL treatments led to higher shoot DW yields and Cu removals than the untreated
71 treatment (UNT). Similarly, at the Touro site, NW Spain, compost incorporation into
72 Cu-rich mine tailings, in combination with planting of *Salix* spp., *Populus nigra* L. and
73 *Agrostis capillaris* L, improved soil properties, i.e. pH, CEC and fertility, and decreased

74 soil Cu availability, which notably promoted the growth of *Salix viminalis* L. and *A.*
75 *capillaris* (Touceda-González et al., 2017a).

76 Beside crop biomass and soil physico-chemical properties, there are increasing
77 evidences that phytomanagement, combined with soil amendments, can influence soil
78 microbial communities in the long term. Soil microorganisms are pivotal in the delivery
79 of numerous soil functions and underlying ecosystem services and the success of any
80 gentle remediation option (GRO), such as phytomanagement, should be evaluated in
81 terms of soil function recovery, through the assessment of soil microbial properties
82 (Epelde et al., 2009; Kumpiene et al., 2009; Burges et al., 2018). Phytomanagement
83 increased soil microbial biomass and activity at three out of six field trials with
84 metal(loid)-contaminated soils across Europe, obtaining the most pronounced effects at
85 the Biogeco site (Touceda-González et al., 2017b). At this one, enzyme activities
86 involved in the biogeochemical cycles of C, N, P, and S were 2 to 11-fold higher in
87 amended soils as compared to untreated soils. Furthermore, according to Touceda-
88 González et al. (2017b), changes in specific phylogenetic microbial groups could be
89 more responsive and informative about the effect of phytomanagement on microbial
90 communities than those observed in the community as a whole. In this respect,
91 phytomanagement also induced shifts in the microbial community structure and
92 increased the abundance of genes involved in the N cycle (*nirK*, *nirS*, *nosZ*, and *amoA*).
93 Similarly, at the Touro site, both compost-amendment and plant root activity induced
94 shifts in the bacterial community structure in year 3, along with enzyme activities
95 stimulation (Touceda-González et al., 2017a).

96 Other datasets were gained at the Biogeco site, over time and in field trials with
97 various plant covers. Biochemical activity and functional gene diversity were studied in
98 field plots revegetated with a mixed stand of willows, black poplar, and false indigo

99 bush, and amended or not with OMDL (Lagomarsino et al., 2011; Xue et al., 2018). In
100 year 6, the OMDL treatment reduced Cu availability and soil toxicity, and increased
101 microbial biomass and activity, as well as microbial functional diversity, including
102 genes encoding for metal resistance, as compared to the unamended soil (UNT) (Xue et
103 al., 2018).

104 Considering the key role of microbes in soil ecological processes, larger,
105 biochemically more active and genetically more diverse microbial communities, as
106 observed in above-mentioned studies, could suggest the recovery in soil functioning
107 with phytomanagement. At the Biogeco site, one remaining question is the influence of
108 the vegetation cover and soil amendments on soil microbial communities over time.
109 Gathering long-term data based on evidence from the field is essential to demonstrate
110 the sustainability and efficiency of phytotechnologies (Mench et al., 2010; Kidd et al.,
111 2015; Cundy et al., 2016). Here, we decided to contribute to the monitoring of microbial
112 communities in long-term phytomanaged sites with a metabarcoding approach. We
113 assessed, at year 9, the effect of an application of compost and dolomitic limestone,
114 with and without a renewed compost application , and annual rotation crop on the
115 diversity, composition (the presence or the absence of microbial taxa) and structure (the
116 relative abundance of the microbial taxa) of soil microbial communities.

117

118 **2. Materials and methods**

119 *2.1 Site and experimental design*

120 The wood preservation site (about 10 ha, only 2 ha remaining in activity) is located at
121 Saint-Médard d'Eyrans, Gironde, SW France (N 44°43.353, W 000°30.938) with a
122 temperate Atlantic climate (variable mean rainfall and temperature; in 2017: 736 mm,
123 14.4 °C). Site history, soil characterization and zoning of soil ecotoxicity were

124 previously reported (Mench and Bes, 2009; Bes et al., 2010; Kolbas et al., 2011). Plant
125 communities were characterized in Bes et al. (2010). Copper is the major inorganic
126 contaminant in topsoil at the P1-3 sub-site with a high spatial variation (163-1170 mg
127 Cu kg⁻¹), while As, Zn, Cr and other metal(loid)s were at their background levels, and
128 some polycyclic aromatic hydrocarbons (PAH) reached high concentrations (in mg kg⁻¹
129 soil DW): fluoranthene (1.9), indeno[1,2,3-cd]pyrene (0.95), benzo[g,h,i]perylene (0.8),
130 and benzo[b]fluoranthene (0.8) (Mench and Bes, 2009; Kolbas et al., 2011). Soil texture
131 is sandy, i.e. 85.8% sand, 5.9% clay, and 8.3% silt, with 1.6% OM, C/N 17, soil pH 7,
132 and a low CEC (3.5 cmol kg⁻¹) (Kolbas et al., 2011).

133 The field trial, located at the P1-3 sub-site, consists of 4 blocks (2m x 10m): B1,
134 B2, B3 and B4 (Fig. 1). In March 2008 (year 1), plots in B1, B2 and B3 were amended
135 with compost (5% w/w, made from poultry manure and pine bark chips) and dolomitic
136 limestone (0.2% w/w). The amendment was mixed in the topsoil (0-0.25 m) with a
137 stainless steel spade after soil loosening (Marchand et al., 2011). In 2013, a second
138 compost dressing (5% w/w, made with green waste) was incorporated to half of the
139 plots in B1-B3 (hereafter referred to as OM2DL), while the remaining plots were not
140 additionally amended (OMDL), making 4 replicated plots per treatment and per block.
141 The composition of OMDL and OM2DL soil amendments are detailed in Mench et al.
142 (2018). The plot from B4 remained unamended throughout the whole study (UNT). All
143 plots were cultivated with an annual tobacco -sunflower rotation since 2008.

144

145 *2.2 Soil sampling and physicochemical characterization*

146 In March 2017 (year 9) four soil samples were collected from the topsoil (0–10 cm) of
147 each plot with a stainless sampling cylinder (ø 3.6 cm X 11.5 cm). Once fresh weight
148 was determined for soil bulk density, the four replicated samples were combined to

149 produce a composite soil sample (1 kg fresh weight, FW) for each plot of B1-B3, while
150 the four samples from the UNT plot in B4 were treated as separate samples, making a
151 total of 28 samples for analysis. Then, samples were air-dried, sieved at 2 mm and
152 manually homogenized. Subsamples (150 g FW) for soil microbial analysis were stored
153 fresh at -20°C until analysis. Element concentrations in soil samples, determined using
154 ICP-AES, and other physicochemical parameters were analysed following standard
155 methods and standard quality assurance employed by INRA LAS, Arras, France (2014)
156 (Mench et al., 2018).

157

158 *2.3 DNA extraction and sequencing, and bioinformatic analysis*

159 DNA was extracted from soil samples (0.25 g dry weight, DW) using Power Soil™ kits
160 (Mo Bio). Prior to DNA extraction, soil samples were washed twice in 120 mM
161 K₂HPO₄ (pH 8.0) to remove extracellular DNA (Kowalchuk et al., 1997).

162 Metabarcoding (amplicon) library preparations were carried out using a dual
163 indexing approach (Lanzen et al. (2016). Briefly, the V3-V4 hypervariable region of the
164 small subunit ribosomal RNA (SSU rRNA) was amplified from prokaryotes using the
165 primers 519F (CAGCMGCCGCGGTAA) adapted from Øvreås et al. (1997), and 806R
166 (GGACTACHVGGGTWTCTAAT) from Caporaso et al. (2012). Adapter-linked primer
167 pairs were used during a first PCR, followed by cleaning and a second PCR with
168 adapters linked to sample-specific barcodes. Pair-ended sequencing was carried out
169 using an Illumina MiSeq with the V2 kit at Tecnia Corporation (Miñano, Spain).

170 Sequence read-pairs were quality-filtered and overlapped using vsearch (default
171 parameters; Rognes et al., 2016) and resulting sequences trimmed to remove N5 and
172 primer sequences, using cutadapt (Martin, 2011). Sequences were then truncated to 253
173 nt using vsearch, removing shorter sequences or those low quality based on expected

174 incorrect read calls (fastq_maxee=0.5). All quality-filtered overlapped sequences were
175 then clustered into OTUs using Swarm v2 (Mahé et al., 2015). Singleton Swarm OTUs
176 were removed and the remaining subjected to both *de novo* and reference based chimera
177 filtering (with the rdp_gold reference database), using vsearch (UCHIME algorithm).
178 Remaining Swarm OTUs were further clustered into fixed similarity OTUs, taking into
179 account total read abundances, and using a maximum sequence divergence threshold of
180 3%, again using vsearch (Rognes et al., 2016). OTU abundances were obtained by
181 mapping reads back to the representative OTU sequences.

182 Taxonomic classification was carried out by aligning representative OTU
183 sequences to the SilvaMod database (v128) using blastn (v.2.2.25 + task megablast) and
184 the LCAClassifier of CREST (default parameters; Lanzén et al., 2012;
185 <https://github.com/lanzen/CREST>). Unclassified OTUs below the alignment threshold
186 and those classified as belonging to eukaryotic organellar rRNA were excluded from
187 further analysis.

188

189 *2.4 Bioinformatic and statistical analysis*

190 The effect of soil treatments on soil physicochemical properties, diversity indices, and
191 relative abundance of classified bacterial taxa (per phylogenetic level) was assessed by
192 means of one-way analysis of variance (ANOVA) using R. When significant differences
193 occurred between soil treatments, multiple comparisons of mean values were made
194 using post-hoc Tukey HSD test. A high variance in sequencing depth (number of reads)
195 occurred between samples, indicating that total OTU richness may not be an unbiased
196 measure of α -diversity in our study. Therefore, rarefied richness was evaluated, using
197 the read number of the smallest included dataset (7850 reads), along with Shannon (H')
198 and Simpson diversity indices. The influence of soil treatments on α -diversity indices

199 was further assessed accounting for the considerable variability in soil Cu across the
200 replicated plots (Fig. 1): (i) taking total soil Cu as co-variable, by means of analysis of
201 covariance (ANCOVA) and (ii) considering only plots with similarly higher values of
202 total soil Cu in the analysis, i.e. B3 and B4, by means of ANOVA. Even though total
203 soil Cu does not directly reflect the bioavailable fraction subject to interact with
204 organisms, values of 1 M NH₄NO₃-extractable Cu in the UNT plot were far beyond the
205 range of variability of values from OMDL and OM2DL plots (Fig. 1.b), making this
206 parameter unreliable, both as co-variable and categorizing factor, for the comparison of
207 α -diversity metrics among soil treatments.

208 Multivariate statistics, calculation of diversity indices and visualization of the
209 amplicon sequencing data was performed using the R package *vegan* (Oksanen et al.,
210 2019). Function *decostand* was used to transform OTU distributions into relative
211 abundances. Bray-Curtis dissimilarity matrices comparing community composition
212 between samples were calculated, as described by Lanzén et al. (2016), and were
213 subsequently used to perform non-metric multidimensional scaling (NMDS) with
214 function *metaMDS*. These matrices were also used to assess the significance of the
215 effect of soil treatments on the composition of microbial communities by means of
216 permutational multivariate analysis of variance (PERMANOVA).

217 We also explored the relationship between soil microbial communities and
218 changes in soil physicochemical properties induced by soil treatments. Firstly, a
219 selection of soil physicochemical parameters was fitted to the resulting NMDS space
220 using the function *envfit*. We used analysis of similarities (ANOSIM) based on Bray-
221 Curtis dissimilarities to evaluate the significance of the influence of soil
222 physicochemical properties on soil microbial composition. These correlation analyses
223 were subjected to Bonferroni correction and not reported unless $p < 0.05$ after correction.

224 Only significantly correlated physicochemical properties were used in further analyses.
225 In parallel, function *bioenv* was used to find the subset of physicochemical parameters
226 that together showed maximum correlation with community dissimilarity. Secondly,
227 multivariate analyses were performed by means of redundancy analysis (RDA) and
228 variation partitioning analysis, using Canoco 5 (ter Braak and Šmilauer, 2012), with
229 physicochemical properties and (i) α -diversity metrics, and (ii) abundance of the main
230 bacterial taxa, at class level, that showed significant differences due to soil treatments.

231

232 **3. Results**

233 *3.1 Analysis of soil microbial α -diversity*

234 In total, amplicon sequencing resulted in 1,042,643 16S rRNA reads, which clustered
235 into 10,164 OTUs, after quality filtering and removal of singletons. No direct
236 correlation between soil treatments and α -diversity estimates could be identified (Table
237 1). Accounting for the background influence of soil Cu variability among the plots
238 showed that soil treatments did not significantly affect α -diversity metrics either (Fig.
239 2). An interaction between soil treatment and total soil Cu was found for Shannon
240 diversity index, corroborating that its values increased in the OM2DL plots with
241 increasing levels of total soil Cu, while they decreased in the OMDL and UNT plots.
242 When considering only plots with similarly higher soil Cu levels, higher values of
243 Shannon diversity were also found in the OM2DL plots (Supplementary Table 2),
244 reflecting the influence of this soil treatment on microbial diversity.

245

246 *3.2 Analysis of soil microbial composition*

247 PERMANOVA analysis of Bray-Curtis dissimilarities revealed a significant effect of
248 the soil treatments on the composition of soil microbial communities (Table 2), the

249 post-hoc analysis corroborating significant differences in microbial composition
250 between the soil treatments. This is illustrated by the NMDS ordination space (Fig. 3),
251 which shows the clustering of soil microbial communities according to soil treatments.
252 Consistently, the ANOSIM analysis also confirmed the composition discrimination
253 driven by soil treatments ($R = 0.68$; $P = 0.001$). Based on the NMDS plot, the bacterial
254 communities of OMDL and OM2DL plots clustered more closely, clearly separated
255 from the UNT ones. This is corroborated by Bray-Curtis dissimilarities average values
256 between soil treatments (mean \pm SD; 0.62 ± 0.16 for OMDL-OM2DL, 0.71 ± 0.11 for
257 OMDL-UNT, 0.70 ± 0.11 for OM2DL-UNT), indicating that the microbial composition
258 of UNT soils is most different as compared to that from amended soils.

259

260 3.3 Analysis of soil microbial structure

261 Bacteria dominated the soil microbial community, representing 98.9-99.6% of the
262 quality-filtered reads, whereas archaea accounted for only 0.4-1.0% (Supplementary
263 Table 3). About 99, 98 and 83% of the 16S rRNA amplicons could be taxonomically
264 classified to phylum, class and order level, respectively, resulting in 23 phyla, 61 classes
265 and 76 orders. Fig. 4 illustrates the relative abundance of the most abundant taxa at class
266 level. Among them, *Alphaproteobacteria*, *Betaproteobacteria*, *Gammaproteobacteria*
267 and *Deltaproteobacteria*, which belong to phylum *Proteobacteria*, accounted for 39-
268 47% of the total amplified amplicons dominating microbial communities in all plots,
269 followed by classes belonging to phyla *Acidobacteria* (14-16%), *Bacteroidetes* (9-11%)
270 and *Actinobacteria* (7-10%). Remaining predominant classes fell within phyla
271 *Verrucomicrobia* (4-5%), *Planctomycetes* (4-5%), *Chloroflexi* (3-6%),
272 *Gemmatimonadetes* (3-4%), *Firmicutes* (1-2%) and *Epsilonbacteraeota* (1%).

273 Out of the resulting classified microbial taxa, 13 phyla, 32 classes and 42 orders
274 showed significant differences in their relative abundance due to soil treatments
275 (Supplementary Table 3). Overall, the abundance of bacteria showed a trend to increase
276 in amended pots, while archaeal population, represented mainly by the class
277 *Nitrososphaeria*, significantly decreased to less than half in the OM2DL plots. Taking a
278 closer look at class level (Fig. 5), soil amendments, particularly in the OM2DL soils,
279 increased the population of *Aphaproteobacteria*, while *Thermoleophilia* (phylum
280 *Actinobacteria*), *Blastocatellia* (*Acidobacteria*), *Spartobacteria* (*Verrucomicrobia*) and
281 *Desulfurellia* (*Epsilonbacteraeota*) increased mainly in the OMDL soils. Soil
282 amendments resulted in a reduction of the less predominant bacterial groups mostly
283 belonging to phyla *Planctomycetes*, *Chloroflexi* and *Gemmatimonadetes*, as well as
284 classes *Cytophagia* (phylum *Bacteroidetes*), *OPB35 soil group* (*Verrucomicrobia*) and
285 *Acidobacteriia* (*Acidobacteria*).

286

287 *3.4 Relationship between soil physicochemical and microbial properties*

288 The influence of soil treatments on soil microbial communities was further explored
289 through changes in physicochemical soil properties. Soil CEC, total and organic soil C,
290 total N and total P influenced microbial diversity, being correlated to richness and
291 Shannon diversity index (Supplementary Fig. 1; $F = 2.6$, $P < 0.05$;). Based on microbial
292 composition, most physicochemical soil parameters strongly correlated to the NMDS
293 ordination space (Table 3): soil CEC, WHC, total N and organic C, followed by total C,
294 pH, Olsen's extractable and total P, correlated with microbial communities from
295 amended soils, mainly the OM2DL ones; while total and extractable soil Cu correlated
296 with those from the UNT soils (Fig. 3). BIOENV analysis indicated that microbial
297 community dissimilarity is best explained by a combination of total and Olsen's

298 extractable P, total and extractable soil Cu, and soil WHC ($R = 0.78$). Regarding the
299 influence of soil physicochemical properties on soil microbial structure, most soil
300 physicochemical parameters positively correlated with *Alphaproteobacteria*, whereas
301 taxa belonging to phyla *Gemmatimonadetes*, *Planctomycetes* and *Chloroflexi*, and
302 archaea correlated with parameters related to soil Cu contamination, i.e., total and
303 NH_4NO_3 -extractable soil Cu (Fig. 6).

304

305 **4. Discussion**

306 Phytomanagement of metal-contaminated soils combines phytoremediation options with
307 a sustainable site management, aiming at recovering soil functions and ecosystem
308 services, as well as achieving effective risk management (Kidd et al., 2015; Cundy et
309 al., 2016). After 9 years of phytomanagement in this field trial, Mench et al. (2018)
310 reported a long lasting positive effect of soil amendments and annual plant cultivation
311 on the improvement of soil physicochemical properties and the reduction of Cu
312 availability (Supplementary Table 1). In addition to the nutrient incorporation, this
313 allowed a higher production for sunflower and tobacco crops in amended plots, when
314 the annual climatic conditions (i.e. spring and summer drought, heat waves) were not
315 challenging it (Kolbas et al., 2011; Mench et al., 2018). Shoot Cu concentrations of
316 OMDL and OM2DL plants fitted into their common range and can be used by biomass
317 processing technologies and oilseeds as well. Several other studies performed at this
318 site, assessed through chemical and ecotoxicological assays, also corroborated the
319 effectiveness and sustainability of phytomanagement options for the remediation of Cu-
320 contaminated soils (Kolbas et al., 2011; Kumpiene et al., 2011; Marchand et al., 2011;
321 Hattab-Hambli et al., 2016; Oustriere et al., 2016; Mench et al., 2018). Likewise, other
322 studies at this site have evaluated soil microbial properties of activity, biomass and

323 diversity to provide further knowledge on the influence of these remediation options on
324 the recovery of soil functions (Touceda-González et al., 2017b; Xue et al., 2018). Here,
325 we contribute to assess the long-term effect of phytomanagement on soil microbial
326 communities with a metabarcoding approach.

327

328 *4.1 Effect of phytomanagement on microbial α -diversity*

329 As previously reported for this field trial (Kolbas et al., 2011; Hattab-Hambli et al.,
330 2016; Mench et al., 2018), both total and extractable soil Cu displayed an increasing
331 gradient across the plots, from B1 to B4. This spatial variability in soil Cu
332 contamination, mainly attributed to variability in cumulative wood washings resulting
333 from long-term storage of preserved wood (Oustriere et al., 2016; Mench et al., 2018),
334 may account for a large part of the variance in α -diversity metrics among soil
335 treatments, leading to no significant differences. Nevertheless, when we took into
336 account the background influence of total soil Cu variability across the plots (Fig. 2), no
337 significant effect of the soil treatments on microbial diversity was observed either.

338 Focusing on plots with higher soil Cu contamination, the OM2DL treatment had
339 a stimulating effect on microbial diversity, only reflected in higher Shannon values. In
340 fact, Shannon and Simpson indices are diversity metrics that account for both richness
341 and evenness (Bent and Forney, 2008), being more sensitive to changes in microbial
342 populations than only richness-based metrics. Interestingly, the response of microbial
343 diversity to the renewed compost application in the most contaminated plots followed
344 the same pattern as crop yield: plant biomass in the OM2DL plots remained steady as
345 levels of total soil Cu increased, whereas it decreased in the OMDL plots (Mench et al.,
346 2018). Plant biomass can strongly affect and promote soil microbial communities
347 (Epelde et al., 2010) and enhance microbial diversity (Burgess et al., 2017). Root

348 exudates are an excellent source of energy and nutrients for soil microorganisms, and
349 differences in their amount and quality, due to changes in plant biomass, induce changes
350 in soil microbial diversity (Lucisine et al., 2014; Lopez et al., 2019). Accordingly, the
351 renewed compost application may have counterbalanced the detrimental effects of the
352 increasing levels of total soil Cu on microbial diversity, partly by promoting plant
353 growth. This also demonstrates the benefits to renew compost supply not just to produce
354 higher yield of annual crops at high total soil Cu but also to stimulate microbial
355 diversity through the improvement of soil revegetation.

356 In any case, no overall effect of soil treatments on microbial diversity across all
357 plots could be identified. However, considering the evidence of season variability in soil
358 microbial communities (Bouskill et al., 2010; Pereira et al., 2014), microbial diversity at
359 the sampling time (right before the growing season) may not be corresponding to its full
360 potential. Importantly, we must bear in mind that diversity is relative and constrained by
361 method of measurement, and without sufficient context it can turn out to be
362 uninformative or liable to misinterpretation (Shade, 2016). In addition, α -diversity
363 metrics are simplified estimations of microbial diversity calculated from the vast
364 amount of data provided by next generation sequencing techniques. They are still useful
365 tools that facilitate interpretation of metabarcoding results and provide valuable
366 information that could serve as a first step to provide key insights into underlying
367 ecological processes that drive microbial community patterns (Shade, 2016). Therefore,
368 it is fundamental to assess soil microbial composition and structure, along with
369 microbial diversity, for determining the effect of long-term phytomanagement on soil
370 microbial communities.

371

372 4.2 *Effect of phytomanagement on the composition and structure of soil microbial*
373 *communities*

374 As opposed to α -diversity, compositional analysis showed higher differences among
375 soil treatments (Fig. 3, Table 2), revealing information on soil microbial communities
376 not implied by diversity metrics alone. These changes in microbial composition could
377 reflect the influence of the soil amendments on microbe-mediated soil ecological
378 processes. For instance, in the same field trial, 6 years after the application of the
379 OMDL amendment, Touceda-Gonzalez et al. (2017b) observed an increase in microbial
380 biomass and activity, as well as in the abundance of genes involved in N cycling.
381 Furthermore, organic amendments, like the ones used here, can increase genes encoding
382 for metal and antibiotic resistance (Caban et al., 2018; Garbisu et al., 2018). At this site,
383 in a field trial phytomanaged with the OMDL treatment combined with a mixed stand of
384 willows, poplar and false indigo bush, Xue et al. (2018) reported higher abundance of
385 functional genes encoding for resistance to metals and antibiotics. They suggested that
386 this could be due to proliferation of certain microbial groups caused by OM
387 mineralization, as well as introduction of exogenous metal-resistant microbes with the
388 soil amendment.

389 Compositional analysis also allowed a more comprehensive interpretation of the
390 effect of soil treatments based on differences in the chemical composition and
391 application rate of soil amendments. The compost incorporated in both OMDL and
392 OM2DL plots in year 1 was made with poultry manure and pine bark chips, the latter
393 known to contain, along with cellulose and xylans, a high lignin content (Vane et al.,
394 2006; Xue et al., 2018). The second compost dressing, incorporated in the OM2DL
395 plots in year 6, was made from green waste and sandy soils (Jones et al., 2016),
396 containing more easily biodegradable OM. Since the biochemical nature of the plant

397 residue can determine the mineralization rate of green manures (Tejada et al., 2008),
398 this would certainly shape soil microbial communities. For instance, 6 years after the
399 incorporation of the OMDL soil amendment in the field trial phytomanaged with trees,
400 Xue et al. (2018) reported a stimulation of microbial functional genes involved in both
401 labile (e.g. cellulose) and recalcitrant (e.g. lignin and aromatic components) C
402 decomposition. Here, we hypothesize that by year 9 much of the relatively labile C in
403 the OMDL soils may be mineralized, with the bulk of soil OM accounting for the more
404 recalcitrant forms of C; while the second compost dressing, in contrast, contributed with
405 a more recent input of nutrients and easily degradable OM (Oustriere et al., 2016) in the
406 OM2DL soils. Based on this, we could suggest that the active fraction of the microbes
407 in the OMDL soils may be mainly represented by decomposers of lignin-rich OM;
408 whereas microbial populations in the OM2DL soils may have shifted to more easily
409 biodegradable OM-decomposers.

410 After 9 years of phytomanagement, plots under different soil treatments
411 sheltered genetically diverse microbial communities, reflected in changes in microbial
412 composition (Fig. 3), i.e. the presence or the absence of microbial taxa. Likewise,
413 organic amendments will inexorably promote the growth of certain microbial groups in
414 detriment of others, resulting in changes in microbial structure, i.e. the proportion of the
415 different microbial taxa.

416 The incorporation of soil amendments, particularly the second compost dressing,
417 stimulated the growth of several microbial groups of the class *Alphaproteobacteria*.
418 Among them, the order *Rhizobiales*, typically N₂-fixing bacteria living in root nodules
419 of legumes (Hartwig, 1998), was notably abundant in OM2DL plots where the green
420 waste compost promoted the development of clover vegetation in winter time (Mench et
421 al., 2018). The OM2DL soil treatment also increased the populations of

422 *Rhodospirillales*, another N₂-fixer that can live both free in soil or associated with the
423 rhizosphere of host plants, and *Sphingomonadales*, whose several members can play
424 various roles in microbe-assisted metal phytoremediation (Waigi et al., 2017). Another
425 taxon from *Alphaproteobacteria* whose abundance was higher in the OM2DL plots was
426 *Caulobacterales*, which can be positively influenced by root exudates, along with
427 *Rhizobiales* and *Sphingomonadales* (Shi et al., 2011). The fact that these taxa are
428 generally associated with plants may justify their predominance in plots that received
429 the second compost dressing, where plant biomass was the highest and contributed to
430 their growth, corroborating that above and belowground communities are tightly
431 interlinked in many soil processes. In addition, *Alphaproteobacteria* belong to
432 *Proteobacteria*, a phylum with great importance to global C, N and S cycling (Spain et
433 al., 2009; Li et al., 2019), which explains that they are the main microbial group
434 profiting from the amendment-derived benefits on soil properties (Fig. 6).

435 Archaeal populations, on the contrary, decreased with the incorporation of soil
436 amendments, reflected in the lower abundance of *Nitrososphaeria*, a class that
437 encompasses several ammonia oxidizing archaea (AOA). At year 6, Touceda-González
438 et al. (2017b) reported an increase in denitrification genes and urease activity with the
439 OMDL treatment, whereas no differences in AOA nitrification genes were found.
440 However, in a phytoextraction experiment with *Sedum alfredii* growing on a Cd/Zn
441 contaminated soil, Luo et al. (2019) observed a decrease in AOA groups belonging to
442 *Nitrososphaeria* with successive crops, resulting in a reduction in the potential
443 nitrification rate and nitrogen loss. They also indicated a negative correlation between
444 the nitrification rate and root exudates. Accordingly, the lower abundance of
445 *Nitrososphaeria* observed here in year 9 in amended plots, along with the higher plant
446 biomass, could indicate a decrease in the nitrification rate with time. This, paired with

447 the increase in N₂-fixing bacteria, suggests a beneficial effect of the incorporation of
448 soil amendments on the ecological processes involved in N cycling partly through the
449 physiological improvement of the annual rotation crop.

450 The abundance of bacterial taxa belonging to *Planctomycetes*, *Chloroflexi* and
451 *Gemmatimonadetes*, was also negatively affected by soil amendments. These phyla are
452 generally abundant in metal-contaminated soils owing to their tolerance to metal excess,
453 including Cu (DeBruyn et al., 2011; Chodak et al., 2013; Singh et al., 2014; Azarbad et
454 al., 2015; Jiang et al., 2019), and contain slow-growing bacteria adapted to oligotrophic
455 habitats, like *Chloroflexi* (Davis et al., 2011; Barton et al., 2014), which enables them to
456 successfully occupy extreme environments (Durand et al., 2018). This may explain that
457 the most predominant groups of these phyla were favoured by the rather oligotrophic
458 and highly contaminated conditions of the unamended plots (Fig. 6). The ecological
459 traits that make these phyla highly adaptive and efficient in oligotrophic, contaminated
460 soils may put them, however, in disadvantage against fast-growing bacterial groups,
461 more efficient in competing under the more favourable conditions of the amended plots.

462 These results indicate there was a consistency in the direction and magnitude of
463 the response to soil treatments, both at class and phylum level. However, the phyla
464 *Acidobacteria*, *Actinobacteria* and *Verrucomicrobia* include several taxa, e.g.
465 *Blastocatellia*, *Thermoleophilia* and *Spartobacteria*, whose abundance increased mainly
466 in the OMDL plots, demonstrating the long lasting effect of the first soil amendment;
467 whereas they also include groups, e.g. *OP35 soil group* and *Acidobacteriia*, that
468 decreased. These microbial groups present a wide range of lifestyle, metal-resistance
469 and metabolic properties. For instance, *Acidobacteria* has a diversity of members with
470 varying tolerance to Cu contamination (Pereira et al., 2014; Singh et al., 2014) and
471 metabolic activity, with many groups reported to be degraders of plant-derived-OM,

472 while others, like *Blastocatellia*, prefer oligotrophic environments (Navarrete et al.,
473 2015; Li et al., 2019). *Actinobacteria* are also a heterogeneous group among the
474 metabolically active bacteria in metal(loid)-contaminated soils (Gremion et al., 2004).
475 *Verrucomicrobia* are relatively abundant in subsurface horizons due to their preference
476 for rather oligotrophic soils (Bergmann et al., 2011). The ecological diversity of these
477 microbial groups may account for the divergent response among their members to soil
478 treatments, which results in no overall changes at phylum level. As suggested by Spain
479 et al. (2009), this highlights the detailed resolution and the importance of subphylum
480 phylogenetic analysis of metabarcoding datasets.

481 Finally, the relationship between patterns in physicochemical and microbial
482 properties demonstrated the influence of soil treatments on soil microbial communities
483 through changes in soil physicochemical properties. Soil organic amendments can
484 incorporate OM and available nutrients, reduce metal bioavailability, and modify pH
485 and other physicochemical properties (Alvarenga et al., 2009a, 2009b; Epelde et al.,
486 2009), that strongly affect microbial communities. pH has often been highlighted as a
487 most important factor in metal-contaminated soils as it highly affects metal mobility, as
488 well as availability of nutrients, soil OC and OM, N, etc. (Kumpiene et al., 2011; Jiang
489 et al., 2016; Mench et al., 2018; Jiang et al., 2019). However, the pH effect here was
490 rather limited and nutrients level, OM, water holding capacity or CEC would influence
491 soil microbial communities in a greater way (Fig. 3; Table 2; Fig. 6; Supplementary Fig.
492 1). Indeed, Xue et al. (2018) reported the amelioration of soil properties such as nutrient
493 content and aggregate formation as major drivers of soil biochemical activity.

494 In any case, the reported influence of the soil treatments was mainly reflected in
495 microbial composition and structure, in the same way that other studies have reported
496 that soil metal contamination had more impact on compositional and structural diversity

497 than on diversity metrics *per se* (Pereira et al., 2014; Azarbad et al., 2015; Epelde et al.,
498 2015; Jiang et al., 2016). This sensitive response of microbial composition and structure
499 may be reflecting an eventual positive influence on soil ecological processes, such as N
500 cycling, demonstrating the potential benefits of phytomanagement on Cu-contaminated
501 soils even in the long-term. Based on this, the monitoring of soil microbial functional
502 genes, along with compositional and structural diversity, should be considered for
503 evaluating the effectiveness of long-term phytomanagement in Cu-contaminated soils,
504 like this site. Lastly, and considering that Cu is a well-known fungicide (Singh et al.,
505 2014), the composition and structure of fungal communities should also be assessed.

506

507 **5. Conclusions**

508 The long-term phytomanagement of Cu-contaminated soils, based on a crop rotation
509 with tobacco and sunflower and the incorporation of compost with dolomitic limestone
510 (compost dressing being renewed or not in year 6), contributed to shift the composition
511 and structure of soil microbial communities in year 9, even though it had no effect on
512 microbial diversity. This phytomanagement option induced changes in soil
513 physicochemical properties in the long term that led to genetically diverse microbial
514 communities even 9 years after the compost application. The input of organic matter
515 and nutrients, the reduction in Cu availability and the improvement of soil properties,
516 particularly with the renewed compost application in year 6, enhanced the growth of
517 specific soil microbial groups, which are involved in different soil ecological processes.
518 The stimulation of plant biomass in the amended plots, induced by the direct and
519 indirect effects of amendments on soil quality, also contributed to shape soil microbial
520 communities. Further research is needed to determine the long-term sustainability and
521 effectiveness of phytomanagement in Cu-contaminated soils by (i) monitoring microbial

522 functional diversity, (ii) assessing also the genetic structure of soil fungal communities,
523 (iii) evaluating the influence of temporal variability through the sampling of plots
524 during and after the growth season, and (iv) exploring the relationship between above
525 and belowground communities through the assessment of plant physiological status.

526

527 **Acknowledgements**

528 The authors thank Dr. Anders Lanzén for bioinformatics and statistic support. This
529 work has been financially supported by the European Regional Development Fund
530 through the PhytoSUDOE Interreg. Project (SOE1/P5/E0189). Aritz Burges thanks the
531 Basque Government for funding his research.

532

533 **References**

534 Alvarenga P, Gonçalves AP, Fernandes RM, de Varennes A, Vallini G, Duarte E, et al.
535 Organic residues as immobilizing agents in aided phytostabilization: (I) effects on
536 soil chemical characteristics. *Chemosphere* 2009a; 74: 1292-1300.

537 Alvarenga P, Palma P, Gonçalves AP, Fernandes RM, de Varennes A, Vallini G, et al.
538 Organic residues as immobilizing agents in aided phytostabilization: (II) effects
539 on soil biochemical and ecotoxicological characteristics. *Chemosphere* 2009b; 74:
540 1301-1308.

541 Azarbad H, Niklinska M, Laskowski R, van Straalen NM, van Gestel CA, Zhou J, et al.
542 Microbial community composition and functions are resilient to metal pollution
543 along two forest soil gradients. *FEMS Microbiol. Ecol.* 2015; 91: 1-11.

544 Barton HA, Giarrizzo JG, Suarez P, Robertson CE, Broering MJ, Banks ED, et al.
545 Microbial diversity in a Venezuelan orthoquartzite cave is dominated by the

546 *Chloroflexi* (Class *Ktedonobacterales*) and *Thaumarchaeota* Group I.1c. Front.
547 Microbiol. 2014; 5: 615.

548 Bent SJ, Forney LJ. The tragedy of the uncommon: understanding limitations in the
549 analysis of microbial diversity. ISME J. 2008; 2: 689-95.

550 Bergmann GT, Bates ST, Eilers KG, Lauber CL, Caporaso JG, Walters WA, et al. The
551 under-recognized dominance of *Verrucomicrobia* in soil bacterial communities.
552 Soil Biol. Biochem. 2011; 43: 1450-1455.

553 Bes CM, Mench M, Aulen M, Gaste H, Taberly J. Spatial variation of plant
554 communities and shoot Cu concentrations of plant species at a timber treatment
555 site. Plant Soil 2010; 330: 267-280.

556 Bouskill NJ, Barker-Finkel J, Galloway TS, Handy RD, Ford TE. Temporal bacterial
557 diversity associated with metal-contaminated river sediments. Ecotoxicology
558 2010; 19: 317-28.

559 Burges A, Alkorta I, Epelde L, Garbisu C. From phytoremediation of soil contaminants
560 to phytomanagement of ecosystem services in metal contaminated sites. Int. J.
561 Phytoremediat. 2018; 20: 384-397.

562 Burges A, Epelde L, Blanco F, Becerril JM, Garbisu C. Ecosystem services and plant
563 physiological status during endophyte-assisted phytoremediation of metal
564 contaminated soil. Sci. Total Environ. 2017; 584-585: 329-338.

565 Caban JR, Kuppusamy S, Kim JH, Yoon Y-E, Kim SY, Lee YB. Green manure
566 amendment enhances microbial activity and diversity in antibiotic-contaminated
567 soil. Appl. Soil Ecol. 2018; 129: 72-76.

568 Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-
569 high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
570 platforms. ISME J. 2012; 6: 1621-4.

571 Cundy AB, Bardos RP, Puschenreiter M, Mench M, Bert V, Friesl-Hanl W, et al.
572 Brownfields to green fields: realising wider benefits from practical contaminant
573 phytomanagement strategies. *J. Environ. Manag.* 2016; 184: 67-77.

574 Chodak M, Gołębiowski M, Morawska-Płoskonka J, Kuduk K, Niklińska M. Diversity
575 of microorganisms from forest soils differently polluted with heavy metals. *Appl.*
576 *Soil Ecol.* 2013; 64: 7-14.

577 Davis KE, Sangwan P, Janssen PH. *Acidobacteria*, *Rubrobacteridae* and *Chloroflexi* are
578 abundant among very slow-growing and mini-colony-forming soil bacteria.
579 *Environ. Microbiol.* 2011; 13: 798-805.

580 DeBruyn JM, Nixon LT, Fawaz MN, Johnson AM, Radosevich M. Global
581 biogeography and quantitative seasonal dynamics of *Gemmatimonadetes* in soil.
582 *Appl. Environ. Microbiol.* 2011; 77: 6295-300.

583 Durand A, Maillard F, Alvarez-Lopez V, Guinchard S, Bertheau C, Valot B, et al.
584 Bacterial diversity associated with poplar trees grown on a Hg-contaminated site:
585 community characterization and isolation of Hg-resistant plant growth-promoting
586 bacteria. *Sci. Total Environ.* 2018; 622-623: 1165-1177.

587 Epelde L, Becerril JM, Barrutia O, Gonzalez-Oreja JA, Garbisu C. Interactions between
588 plant and rhizosphere microbial communities in a metalliferous soil. *Environ.*
589 *Pollut.* 2010; 158: 1576-1583.

590 Epelde L, Becerril JM, Mijangos I, Garbisu C. Evaluation of the efficiency of a
591 phytostabilization process with biological indicators of soil health. *J. Environ.*
592 *Qual.* 2009; 38: 2041-2049.

593 Epelde L, Lanzen A, Blanco F, Urich T, Garbisu C. Adaptation of soil microbial
594 community structure and function to chronic metal contamination at an abandoned
595 Pb-Zn mine. *FEMS Microbiol. Ecol.* 2015; 91: 1-11.

596 Evangelou MWH, Conesa HM, Robinson BH, Schulin R. Biomass Production on trace
597 element-contaminated land: a review. *Environ. Eng. Sci.* 2012; 29: 823-839.

598 Garbisu C, Garaiurrebaso O, Lanzen A, Alvarez-Rodriguez I, Arana L, Blanco F, et al.
599 Mobile genetic elements and antibiotic resistance in mine soil amended with
600 organic wastes. *Sci. Total Environ.* 2018; 621: 725-733.

601 Gremion F, Chatzinotas A, Kaufmann K, Von Sigler W, Harms H. Impacts of heavy
602 metal contamination and phytoremediation on a microbial community during a
603 twelve-month microcosm experiment. *FEMS Microbiol. Ecol.* 2004; 48: 273-283.

604 Hartwig UA. The regulation of symbiotic N₂ fixation: a conceptual model of N
605 feedback from the ecosystem to the gene expression level. *Perspect. Plant Ecol.*
606 *Evol. Syst.* 1998; 1: 92-120.

607 Hattab-Hambli N, Motelica-Heino M, Mench M. Aided phytoextraction of Cu, Pb, Zn,
608 and As in copper-contaminated soils with tobacco and sunflower in crop rotation:
609 mobility and phytoavailability assessment. *Chemosphere* 2016; 145: 543-50.

610 Jiang B, Adebayo A, Jia J, Xing Y, Deng S, Guo L, et al. Impacts of heavy metals and
611 soil properties at a Nigerian e-waste site on soil microbial community. *J. Hazard.*
612 *Mater.* 2019; 362: 187-195.

613 Jiang L, Song M, Yang L, Zhang D, Sun Y, Shen Z, et al. Exploring the influence of
614 environmental factors on bacterial communities within the rhizosphere of the Cu-
615 tolerant plant, *Elsholtzia splendens*. *Sci. Rep.* 2016; 6: 36302.

616 Jones S, Bardos RP, Kidd PS, Mench M, de Leij F, Hutchings T, et al. Biochar and
617 compost amendments enhance copper immobilisation and support plant growth in
618 contaminated soils. *J. Environ. Manag.* 2016; 171: 101-112.

619 Kidd P, Mench M, Alvarez-Lopez V, Bert V, Dimitriou I, Friesl-Hanl W, et al.
620 Agronomic practices for improving gentle remediation of trace element-
621 contaminated soils. *Int. J. Phytoremediat.* 2015; 17: 1005-37.

622 Kolbas A, Mench M, Herzig R, Nehnevajova E, Bes CM. Copper phytoextraction in
623 tandem with oilseed production using commercial cultivars and mutant lines of
624 sunflower. *Int. J. Phytoremediat.* 2011; 13 Suppl 1: 55-76.

625 Kowalchuk GA, Stephen JR, DeBoer W, Prosser JI, Embley TM, Woldendorp JW.
626 Analysis of ammonia-oxidizing bacteria of the beta subdivision of the class
627 *Proteobacteria* in coastal sand dunes by denaturing gradient gel electrophoresis
628 and sequencing of PCR-amplified 16S ribosomal DNA fragments. *Appl. Environ.*
629 *Microbiol.* 1997; 63: 1489-1497.

630 Kumpiene J, Guerri G, Landi L, Pietramellara G, Nannipieri P, Renella G. Microbial
631 biomass, respiration and enzyme activities after in situ aided phytostabilization of
632 a Pb- and Cu-contaminated soil. *Ecotoxicol. Environ. Saf.* 2009; 72: 115-119.

633 Kumpiene J, Mench M, Bes CM, Fitts JP. Assessment of aided phytostabilization of
634 copper-contaminated soil by X-ray absorption spectroscopy and chemical
635 extractions. *Environ. Pollut.* 2011; 159: 1536-1542.

636 Lagomarsino A, Mench M, Marabottini R, Pignataro A, Grego S, Renella G, et al.
637 Copper distribution and hydrolase activities in a contaminated soil amended with
638 dolomitic limestone and compost. *Ecotoxicol. Environ. Saf.* 2011; 74: 2013-2019.

639 Lanzén A, Epelde L, Blanco F, Martin I, Artetxe U, Garbisu C. Multi-targeted
640 metagenetic analysis of the influence of climate and environmental parameters on
641 soil microbial communities along an elevational gradient. *Sci. Rep.* 2016; 6:
642 28257.

643 Lanzén A, Jørgensen SL, Huson DH, Gorfer M, Grindhaug SH, Jonassen I, et al.
644 CREST – classification resources for environmental sequence tags. PLOS ONE
645 2012; 7: e49334.

646 Li B, Cao Y, Guan X, Li Y, Hao Z, Hu W, et al. Microbial assessments of soil with a
647 40-year history of reclaimed wastewater irrigation. Sci. Total Environ. 2019; 651:
648 696-705.

649 Lopez S, Goux X, van der Ent A, Erskine PD, Echevarria G, Calusinska M, et al.
650 Bacterial community diversity in the rhizosphere of nickel hyperaccumulator
651 species of Halmahera Island (Indonesia). Appl. Soil Ecol. 2019; 133: 70-80.

652 Lucisine P, Echevarria G, Sterckeman T, Vallance J, Rey P, Benizri E. Effect of
653 hyperaccumulating plant cover composition and rhizosphere-associated bacteria
654 on the efficiency of nickel extraction from soil. Appl. Soil Ecol. 2014; 81: 30-36.

655 Luo J, Liu Y, Tao Q, Hou Q, Wu K, Song Y, et al. Successive phytoextraction alters
656 ammonia oxidation and associated microbial communities in heavy metal
657 contaminated agricultural soils. Sci. Total Environ. 2019; 664: 616-625.

658 Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M. Swarm v2: highly-scalable
659 and high-resolution amplicon clustering. PeerJ 2015; 3: e1420.

660 Marchand L, Mench M, Marchand C, Le Coustumer P, Kolbas A, Maalouf JP.
661 Phytotoxicity testing of lysimeter leachates from aided phytostabilized Cu-
662 contaminated soils using duckweed (*Lemna minor* L.). Sci. Total Environ. 2011;
663 410-411: 146-53.

664 Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
665 EMBnet J. 2011; 17: 3.

666 Mench M, Bes C. Assessment of ecotoxicity of topsoils from a wood treatment site.
667 Pedosphere 2009; 19: 143-155.

668 Mench M, Lepp N, Bert V, Schwitzguébel J-P, Gawronski SW, Schröder P, et al.
669 Successes and limitations of phytotechnologies at field scale: outcomes,
670 assessment and outlook from COST Action 859. *J. Soil Sediment* 2010; 10: 1039-
671 1070.

672 Mench MJ, Dellise M, Bes CM, Marchand L, Kolbas A, Le Coustumer P, et al.
673 Phytomanagement and remediation of Cu-contaminated soils by high yielding
674 crops at a former wood preservation site: sunflower biomass and ionome. *Front.*
675 *Ecol. Evol.* 2018; 6.

676 Navarrete AA, Venturini AM, Meyer KM, Klein AM, Tiedje JM, Bohannan BJM, et al.
677 Differential response of *Acidobacteria* subgroups to forest-to-pasture conversion
678 and their biogeographic patterns in the western Brazilian Amazon. *Front.*
679 *Microbiol.* 2015; 6: 1443-1443.

680 Oksanen J, Blanchet G, Kindt R, Legendre P, O'Hara RB, Simpson G, et al. *Vegan:*
681 *community ecology package. R Package Version 2.5-4.* 2019.

682 Oustriere N, Marchand L, Galland W, Gabbon L, Lottier N, Motelica M, et al. Influence
683 of biochars, compost and iron grit, alone and in combination, on copper solubility
684 and phytotoxicity in a Cu-contaminated soil from a wood preservation site. *Sci.*
685 *Total Environ.* 2016; 566-567: 816-825.

686 Ovreås L, Forney L, Daae FL, Torsvik V. Distribution of bacterioplankton in
687 meromictic Lake Saelenvannet, as determined by denaturing gradient gel
688 electrophoresis of PCR-amplified gene fragments coding for 16S rRNA. *Appl.*
689 *Environ. Microbiol.* 1997; 63: 3367-73.

690 Pereira LB, Vicentini R, Ottoboni LM. Changes in the bacterial community of soil from
691 a neutral mine drainage channel. *PLOS One* 2014; 9: e96605.

692 Quintela-Sabaris C, Marchand L, Kidd PS, Friesl-Hanl W, Puschenreiter M, Kumpiene
693 J, et al. Assessing phytotoxicity of trace element-contaminated soils
694 phytomanaged with gentle remediation options at ten European field trials. *Sci.*
695 *Total Environ.* 2017; 599-600: 1388-1398.

696 Rognes T, Flouri T, Nichols B, Quince C, Mahe F. VSEARCH: a versatile open source
697 tool for metagenomics. *PeerJ* 2016; 4: e2584.

698 Shade A. Diversity is the question, not the answer. *ISME J.* 2016; 11: 1.

699 Shi S, Richardson AE, O'Callaghan M, DeAngelis KM, Jones EE, Stewart A, et al.
700 Effects of selected root exudate components on soil bacterial communities. *FEMS*
701 *Microbiol. Ecol.* 2011; 77: 600-610.

702 Singh BK, Quince C, Macdonald CA, Khachane A, Thomas N, Al-Soud WA, et al.
703 Loss of microbial diversity in soils is coincident with reductions in some
704 specialized functions. *Environ. Microbiol.* 2014; 16: 2408-20.

705 Spain AM, Krumholz LR, Elshahed MS. Abundance, composition, diversity and
706 novelty of soil *Proteobacteria*. *ISME J.* 2009; 3: 992-1000.

707 Tejada M, Gonzalez JL, García-Martínez AM, Parrado J. Effects of different green
708 manures on soil biological properties and maize yield. *Bioresour. Technol.* 2008;
709 99: 1758-1767.

710 ter Braak CJF, Šmilauer P. Canoco reference manual and user's guide: software for
711 ordination (version 5.0). Microcomputer power, Itaca, USA., 2012.

712 Touceda-González M, Álvarez-López V, Prieto-Fernández Á, Rodríguez-Garrido B,
713 Trasar-Cepeda C, Mench M, et al. Aided phytostabilisation reduces metal toxicity,
714 improves soil fertility and enhances microbial activity in Cu-rich mine tailings. *J.*
715 *Environ. Manag.* 2017a; 186, Part 2: 301-313.

716 Touceda-González M, Prieto-Fernandez A, Renella G, Giagnoni L, Sessitsch A, Brader
717 G, et al. Microbial community structure and activity in trace element-
718 contaminated soils phytomanaged by Gentle Remediation Options (GRO).
719 Environ. Pollut. 2017b; 231: 237-251.

720 Van Slycken S, Witters N, Meiresonne L, Meers E, Ruttens A, Van Peteghem P, et al.
721 Field evaluation of willow under short rotation coppice for phytomanagement of
722 metal-polluted agricultural soils. Int. J. Phytoremediat. 2013; 15: 677-89.

723 Vane CH, Drage TC, Snape CE. Bark decay by the white-rot fungus *Lentinula edodes*:
724 polysaccharide loss, lignin resistance and the unmasking of suberin. Int. Biodeter.
725 Biodeg. 2006; 57: 14-23.

726 Waigi MG, Sun K, Gao Y. *Sphingomonads* in microbe-assisted phytoremediation:
727 tackling soil pollution. Trends Biotechnol. 2017; 35: 883-899.

728 Witters N, Mendelsohn RO, Van Slycken S, Weyens N, Schreurs E, Meers E, et al.
729 Phytoremediation, a sustainable remediation technology? Conclusions from a case
730 study. I: energy production and carbon dioxide abatement. Biomass Bioenerg.
731 2012; 39: 454-469.

732 Xue K, Zhou J, Van Nostrand J, Mench M, Bes C, Giagnoni L, et al. Functional activity
733 and functional gene diversity of a Cu-contaminated soil remediated by aided
734 phytostabilization using compost, dolomitic limestone and a mixed tree stand.
735 Environ. Pollut. 2018; 242: 229-238.

736
737

Figure captions

Fig. 1: Spatial variability of (a) total soil Cu (mg Cu kg⁻¹ DW), and (b) 1M NH₄NO₃-extractable Cu (µg Cu kg⁻¹ DW) within the field trial.

Fig. 2. Influence of soil treatments on α -diversity indices, assessed using ANCOVA, considering total soil Cu concentration as co-variable. Probability values from ANCOVA (ns: non-significant; * represents significance of $P < 0.05$) for rarefied richness: amendment, ns; soil Cu, ns; amendment x soil Cu, ns; Shannon diversity index: amendment, ns; soil Cu, ns; amendment x soil Cu, *; and Simpson diversity index: amendment, ns; soil Cu, ns; amendment x soil Cu, ns.

Fig. 3. Non-metric multidimensional scaling (NMDS) based on Bray-Curtis dissimilarities of the composition of soil microbial communities. Fitted physicochemical soil parameters with significant correlation to the multidimensional NMDS space are included. Tot_N: total N; Tot_P: total P; P_Olsen: Olsen's extractable P; Tot_Cu: total Cu; Ext_Cu: 1 M NH₄NO₃-extractable Cu; Tot_C: total C; OC: organic C; WHC: water holding capacity; CEC: cationic exchange capacity.


Fig. 4. Average relative abundance of the main microbial taxa, at class level, per soil treatment.

Fig. 5. Heat map based on relative abundance of the top 15 microbial taxa, at class level, that showed significance differences among soil treatments.


Fig. 6. Biplot of redundancy analysis (RDA) performed between soil physicochemical properties and relative abundance of the main microbial taxa, at class level, considering only taxa showing significant differences in their abundance due to soil treatment ($F = 3.0$, $P < 0.01$). RDA1 and RDA2 account for 29 and 15% of the total variance, respectively. Tot_N: total N; Tot_P: total P; P_Olsen: Olsen's extractable P; Tot_Cu: total Cu; Ext_Cu: NH_4NO_3 -extractable Cu; Tot_C: total C; OC: total organic C; WHC: water holding capacity; CEC: cationic exchange capacity




Supplementary Fig. 1. Biplot of the redundancy analysis (RDA) from α -diversity metrics explained by physicochemical topsoil parameters ($F = 2.6$, $P < 0.05$); RDA1 and RDA2 account for 61 and 1% of the variance, respectively. Richness: rarefied total richness; Shannon: Shannon diversity index; Simpson: Simpson diversity index; Tot_N: total N; P_Olsen: Olsen's extractable P; Tot_P: total P; Tot_Cu: total Cu; Ext_Cu: extractable Cu; Org_C: organic C; WHC: water holding capacity; CEC: cationic exchange capacity.


(a)





(b)














Table 1. Values of amplified reads and α -diversity metrics per soil treatment (Mean values \pm standard deviation). Values followed by different letters are significantly different ($P < 0.05$) according to Tukey's test.

	UNT	OMDL	OM2DL
Reads	10956 \pm 1092 ^a	46058 \pm 32597 ^a	40557 \pm 33994 ^a
Rarefied richness	1736 \pm 148 ^a	1862 \pm 87 ^a	1891 \pm 113 ^a
Shannon index (H')	6.57 \pm 0.15 ^a	6.70 \pm 0.11 ^a	6.75 \pm 0.12 ^a
Simpson index	0.996 \pm 0.001 ^a	0.996 \pm 0.000 ^a	0.997 \pm 0.001 ^a

Table 2. (a) Effect of soil treatments, total soil Cu, and their interaction, and (b) pair-wise comparisons between soil treatments, on the composition of soil microbial communities, assessed using PERMANOVA (asterisks represent strength of significance).

	Df	F	R²	Pr(>F)	
a Amendment	2	7.0002	0.3327	0.001	**
Soil Cu	1	4.7321	0.1124	0.001	**
Amendment x Soil Cu	2	1.1723	0.0557	0.252	
Residuals	21				
b OMDL - OM2DL	22	7.886	0.273	0.006	**
OMDL - UNT	15	18.0655	0.5633	0.006	**
OM2DL - UNT	14	27.5414	0.6793	0.006	**

Df: degree of freedom; F: Fisher value; R²: coefficient of determination;

Pr: probability value.

Table 3. Correlation of physicochemical soil properties to NMDS coordinates (asterisks represent strength of significance)

	R²	Pr(>r)	
CEC	0.630	0.001	***
Water Holding Capacity	0.620	0.001	***
Total N	0.593	0.001	***
1M NH ₄ NO ₃ -extractable Cu	0.544	0.001	***
Total Cu	0.541	0.001	***
Organic C	0.519	0.001	***
Total C	0.496	0.001	***
pH	0.459	0.002	**
Olsen's extractable P	0.367	0.003	**
Total P	0.312	0.013	*

R²: coefficient of determination; Pr: probability value.

Supplementary Table. 1. Soil physicochemical parameters per block and per soil treatments (Mean values \pm standard deviation). Values followed by different letters are significantly different ($P < 0.05$) according to Tukey's test. No letter in a row indicates no significant difference.

	BLOCK 4		BLOCK 3		BLOCK 2		BLOCK 1	
	UNT	OMDL	OM2DL	OMDL	OM2DL	OMDL	OM2DL	
pH water	6.8 \pm 0.3 ^b	7.0 \pm 0.2 ^{ab}	7.0 \pm 0.2 ^{ab}	7.0 \pm 0.1 ^{ab}	7.1 \pm 0.1 ^{ab}	7.0 \pm 0 ^{ab}	7.2 \pm 0.1 ^a	
CEC (cmol kg ⁻¹)	3.3 \pm 0.2 ^d	7.0 \pm 3.3 ^{bcd}	10.4 \pm 2.6 ^{abc}	6.4 \pm 0.2 ^{cd}	10.8 \pm 0.5 ^a	6.0 \pm 1.1 ^d	10.4 \pm 1.1 ^{ab}	
Organic C (g kg ⁻¹)	8.3 \pm 2.1 ^c	19.0 \pm 8.2 ^{ab}	23.7 \pm 8.5 ^{ab}	14.5 \pm 0.6 ^{abc}	24.3 \pm 1.7 ^a	14.0 \pm 1.4 ^{bc}	22.8 \pm 1.0 ^{ab}	
OM (g kg ⁻¹)	17 \pm 1 ^c	33 \pm 14 ^{abc}	41 \pm 15 ^{ab}	25 \pm 1 ^{abc}	42 \pm 3 ^a	24 \pm 3 ^{bc}	40 \pm 2 ^{ab}	
Total C (g kg ⁻¹)	9.8 \pm 0.5 ^c	19.0 \pm 8.2 ^{abc}	23.7 \pm 8.5 ^{ab}	14.8 \pm 0.5 ^{abc}	24.3 \pm 1.7 ^a	14.0 \pm 1.4 ^{bc}	23.3 \pm 1.3 ^{ab}	
Total N (g kg ⁻¹)	0.7 \pm 0.0 ^c	1.2 \pm 0.5 ^{bc}	1.7 \pm 0.6 ^{ab}	1.0 \pm 0.0 ^{bc}	2.0 \pm 0.0 ^a	1.0 \pm 0.0 ^c	2.0 \pm 0.0 ^a	
C/N	13.8 \pm 1.3	14.0 \pm 0.0	13.3 \pm 0.6	13.8 \pm 0.5	13.8 \pm 0.5	13.5 \pm 0.6	14.0 \pm 0.0	
Olson extractable P (mg kg ⁻¹)	72 \pm 2 ^c	78 \pm 9 ^{bc}	84 \pm 5 ^{abc}	72 \pm 6 ^c	96 \pm 10 ^a	75 \pm 4 ^c	92 \pm 7 ^{ab}	
Total CaCO ₃ (g kg ⁻¹) ¹	<1	1.0 \pm 0.7 ^b	1.7 \pm 0.6 ^{ab}	1.0 \pm 0.0 ^b	1.5 \pm 0.6 ^{ab}	1.0 \pm 0.0 ^b	2.5 \pm 0.6 ^a	
Water holding capacity (%)	6.8 \pm 0.6 ^d	7.8 \pm 1.7 ^{cd}	8.0 \pm 0.0 ^{cd}	10.0 \pm 0.3 ^{bc}	12.8 \pm 0.9 ^a	9.3 \pm 1.2 ^{cd}	11.9 \pm 1.4 ^{ab}	
Elements (mg kg⁻¹)								
Total Cu	780 \pm 112 ^a	842 \pm 69 ^a	836 \pm 70 ^a	514 \pm 48 ^b	316 \pm 52 ^c	307 \pm 60 ^c	237 \pm 41 ^c	
Extractable Cu (μ g kg ⁻¹) ²	6274 \pm 2639 ^a	3398 \pm 879 ^b	3667 \pm 770 ^{ab}	1543 \pm 215 ^b	2025 \pm 270 ^b	889 \pm 161 ^b	1903 \pm 320 ^b	
Nutrients (mg kg⁻¹)								
Ca	1391 \pm 50 ^{ab}	2292 \pm 1728 ^{ab}	3072 \pm 1563 ^{ab}	1611 \pm 92 ^{ab}	3284 \pm 235 ^a	1285 \pm 250 ^b	3304 \pm 388 ^a	
Fe	6199 \pm 316	6397 \pm 301	6209 \pm 238	6516 \pm 253	6399 \pm 116	6271 \pm 19	6403 \pm 322	
K ¹	7893	7645 \pm 269 ^{abc}	7322 \pm 219 ^c	7760 \pm 118 ^{ab}	7501 \pm 158 ^{bc}	7992 \pm 28 ^a	7372 \pm 101 ^c	
Mg	771 \pm 26 ^b	919 \pm 83 ^a	909 \pm 36 ^a	887 \pm 26 ^a	885 \pm 34 ^a	828 \pm 17 ^{ab}	840 \pm 51 ^{ab}	
Na ¹	2015	1925 \pm 50 ^b	1942 \pm 42 ^b	2098 \pm 31 ^a	1907 \pm 93 ^b	2115 \pm 31 ^a	2005 \pm 47 ^{ab}	
P	259 \pm 14 ^b	940 \pm 128 ^a	910 \pm 68 ^a	956 \pm 63 ^a	739 \pm 435 ^a	878 \pm 37 ^a	874 \pm 139 ^a	
Texture (g kg⁻¹)								
Clay ¹	54	60 \pm 6	65 \pm 9	63 \pm 1	70 \pm 3	64 \pm 4	70 \pm 2	
Silt ¹	109	96 \pm 8	97 \pm 9	99 \pm 3	97 \pm 3	101 \pm 3	101 \pm 5	
Sand ¹	837	844 \pm 13	839 \pm 17	838 \pm 3	833 \pm 5	836 \pm 5	829 \pm 5	

¹n = 1 for UNT; ²1M NH₄NO₃-extractable soil Cu.

Supplementary Table 2. Values of α -diversity metrics per soil treatment in B3 and B4 (Mean values \pm standard deviation). Values followed by different letters are significantly different ($P < 0.05$) according to Tukey's test. No letter in a row indicates no significant difference.

	UNT	OMDL	OM2DL
Rarefied richness	1877 \pm 159	1980 \pm 65	2052 \pm 48
Shannon's index (H')	6.56 \pm 0.15 ^b	6.60 \pm 0.16 ^b	6.84 \pm 0.05 ^a
Simpson index	0.996 \pm 0.000 ^a	0.994 \pm 0.001 ^a	0.997 \pm 0.000 ^a

Supplementary Table 3. Effect of the soil treatments on the relative abundance of classified soil microbial taxa.

TAXA	UNT		OMDL		OM2DL		P val
<i>Domain</i>							
Bacteria	9.89E-01	b	9.95E-01	a	9.96E-01	a	0.000
Archaea	1.06E-02	a	4.65E-03	b	4.25E-03	b	0.000
<i>Phylum</i>							
Proteobacteria	3.83E-01	b	4.31E-01	ab	4.70E-01	a	0.003
Verrucomicrobia	4.57E-02	ab	4.97E-02	a	4.11E-02	b	0.002
Planctomycetes	5.49E-02	a	3.74E-02	b	3.63E-02	b	0.000
Gemmatimonadetes	4.42E-02	a	3.48E-02	ab	2.57E-02	b	0.012
Chloroflexi	5.97E-02	a	3.15E-02	b	2.81E-02	b	0.000
Epsilonbacteraeota	9.21E-03	b	1.23E-02	a	9.27E-03	b	0.004
Armatimonadetes	4.12E-03	b	7.21E-03	a	5.47E-03	b	0.004
Thaumarchaeota	9.92E-03	a	4.51E-03	b	3.91E-03	b	0.001
Ca. Latescibacteria WS3	9.39E-03	a	2.77E-03	b	2.57E-03	b	0.000
Chlorobi	1.31E-03	b	2.39E-03	a	1.97E-03	ab	0.032
Ca. Tectomicrobia	2.39E-03	a	1.14E-03	b	8.33E-04	b	0.000
Ca. Dependuntiae TM6	1.25E-03	a	7.25E-04	b	6.81E-04	b	0.002
Ca. Parcubacteria	1.56E-03	a	5.83E-04	b	7.82E-04	b	0.000
<i>Class</i>							
Alphaproteobacteria	1.57E-01	b	1.85E-01	b	2.21E-01	a	0.000
Blastocatellia	2.27E-02	b	3.86E-02	a	3.23E-02	ab	0.006
Thermoleophilia	1.94E-02	ab	2.77E-02	a	1.76E-02	b	0.008
Gemmatimonadetes	3.60E-02	a	2.76E-02	ab	2.07E-02	b	0.009
Cytophagia	3.67E-02	a	2.32E-02	b	3.09E-02	a	0.001
Spartobacteria	1.02E-02	b	2.10E-02	a	1.38E-02	b	0.004
Planctomycetacia	3.00E-02	a	2.05E-02	b	2.17E-02	ab	0.003
OPB35 soil group	2.24E-02	a	1.75E-02	b	1.52E-02	c	0.000
Desulfurellia	9.21E-03	b	1.23E-02	a	9.27E-03	b	0.004
Phycisphaerae	1.46E-02	a	1.15E-02	ab	9.96E-03	b	0.009
Acidobacteriia	2.12E-02	a	6.45E-03	b	3.33E-03	c	0.000
Chloroflexi Subdivision 2	1.40E-02	a	5.95E-03	b	6.81E-03	b	0.000
Nitrososphaeria	9.92E-03	a	4.51E-03	b	3.91E-03	b	0.001
Chloroflexi Subdivision 8	7.12E-03	a	3.75E-03	b	2.90E-03	b	0.001
Chloroflexia	4.06E-03	a	3.16E-03	ab	1.88E-03	b	0.027
OM190	6.93E-03	a	2.76E-03	b	2.89E-03	b	0.000
Armatimonadetes Group 4	1.61E-03	b	2.47E-03	a	1.64E-03	b	0.023
Chlorobia	1.31E-03	b	2.39E-03	a	1.97E-03	ab	0.032
Clostridia	2.20E-03	ab	2.35E-03	b	3.10E-03	a	0.023
S085	4.06E-03	a	2.23E-03	b	2.57E-03	b	0.002
Anaerolineae	4.59E-03	a	1.93E-03	b	2.21E-03	b	0.000
Chthonomonadetes	7.50E-04	b	1.85E-03	a	9.68E-04	b	0.000
Verrucomicrobiae	1.94E-03	ab	1.72E-03	b	2.33E-03	a	0.040

JG30 KF CM66	3.72E-03	a	1.69E-03	b	1.26E-03	b	0.000
Saprospira	2.34E-03	ab	1.50E-03	b	2.85E-03	a	0.018
Thermomicrobia	2.69E-03	a	1.22E-03	b	6.08E-04	b	0.003
vadinHA49	1.21E-03	a	1.15E-03	a	5.93E-04	b	0.002
Acidobacteria clade RB25	2.42E-03	a	1.04E-03	b	1.01E-03	b	0.000
Ktedonobacteria	5.95E-03	a	9.86E-04	b	5.30E-04	b	0.000
BD2 11 terrestrial group	1.30E-03	a	4.85E-04	b	9.83E-04	a	0.000
Acidobacteria group 2	3.33E-03	a	4.27E-04	b	3.21E-04	b	0.000
Caldilineae	7.68E-04	a	4.15E-04	b	6.37E-04	a	0.001
Order							
Rhizobiales	6.80E-02	b	8.43E-02	ab	9.38E-02	a	0.032
Sphingomonadales	3.96E-02	b	6.06E-02	a	6.85E-02	a	0.013
Xanthomonadales	6.71E-02	a	5.46E-02	b	6.67E-02	a	0.001
Blastocatellales	2.26E-02	b	3.86E-02	a	3.22E-02	ab	0.006
Gemmatimonadales	3.60E-02	a	2.76E-02	ab	2.07E-02	b	0.009
Rhodospirillales	2.93E-02	ab	2.43E-02	b	3.57E-02	a	0.002
Cytophagales	3.66E-02	a	2.31E-02	b	3.08E-02	a	0.001
Chthoniobacterales	9.91E-03	b	2.08E-02	a	1.31E-02	b	0.003
Planctomycetales	3.00E-02	a	2.05E-02	b	2.17E-02	ab	0.003
Solirubrobacterales	8.73E-03	b	1.66E-02	a	1.18E-02	b	0.010
Desulfurellales	9.21E-03	b	1.23E-02	a	9.27E-03	b	0.004
Caulobacterales	1.29E-02	ab	1.11E-02	b	1.78E-02	a	0.002
Frankiales	6.43E-03	ab	8.80E-03	a	5.52E-03	b	0.009
TRA3 20	8.17E-03	ab	7.41E-03	b	9.66E-03	a	0.002
Propionibacterales	5.64E-03	b	7.35E-03	b	1.06E-02	a	0.000
Gaiellales	9.63E-03	a	7.24E-03	a	4.19E-03	b	0.000
Acidobacteriales	2.12E-02	a	6.45E-03	b	3.33E-03	c	0.000
Nitrososphaerales	8.02E-03	a	4.46E-03	b	3.88E-03	b	0.013
SC I 84	4.11E-03	a	3.57E-03	a	2.19E-03	b	0.000
Kineosporiales	1.15E-03	b	3.31E-03	a	1.34E-03	b	0.000
Chloroflexales	2.74E-03	a	2.67E-03	a	1.42E-03	b	0.037
Chlorobiales	1.31E-03	b	2.39E-03	a	1.97E-03	ab	0.032
Clostridiales	2.09E-03	ab	2.12E-03	b	2.86E-03	a	0.021
Anaerolineales	4.59E-03	a	1.93E-03	b	2.21E-03	b	0.000
Chthonomonadales	7.50E-04	b	1.85E-03	a	9.68E-04	b	0.000
Verrucomicrobiales	1.89E-03	ab	1.69E-03	b	2.27E-03	a	0.033
Saprosirales	2.34E-03	ab	1.50E-03	b	2.85E-03	a	0.018
Streptosporangiales	9.94E-04	b	1.44E-03	b	2.79E-03	a	0.012
Phycisphaerales	3.20E-03	a	1.25E-03	b	1.62E-03	b	0.000
Rhodobacterales	2.85E-03	a	1.11E-03	b	9.63E-04	b	0.000
Acidobacteria group 22	2.42E-03	a	1.04E-03	b	1.01E-03	b	0.000
Acidobacteria group 7a	6.09E-04	ab	9.19E-04	a	3.71E-04	b	0.006
JG30 KF CM45	2.17E-03	a	8.72E-04	b	4.45E-04	b	0.003
Lineage IIa	3.49E-04	a	7.50E-04	a	5.07E-04	a	0.025
C0119	1.55E-03	a	7.36E-04	b	3.86E-04	c	0.000
NB1 j	1.86E-03	a	7.05E-04	b	7.56E-04	b	0.000

Methylophilales	2.30E-03	a	6.77E-04	b	4.09E-04	b	0.000
Gitt GS 136	1.21E-03	a	6.41E-04	ab	9.26E-04	a	0.024
Vicinamibacter order incertae sedis	2.39E-04	b	6.14E-04	b	1.29E-03	a	0.000
HTA4	1.65E-03	a	6.12E-04	b	3.64E-04	b	0.000
Oceanospirillales	6.28E-04	b	5.68E-04	b	1.35E-03	a	0.002
Caldilineales	7.68E-04	a	4.15E-04	b	6.37E-04	a	0.001

Ca., candidate

OM2DL


OMDL

UNT

Microbial taxa

Associated with

Relative abundance



Rhizobiales
Rhodospirillales

N₂ fixation



Sphingomonadales
Caulobacterales

Plant exudates production



Nitrososphaeria

Nitrification



Planctomycetacia
Chloroflexi
Gemmatimonadetes

Cu contamination
Oligotrophic soils

Soil properties



pH / CEC / OM and nutrient contents



Available Cu