



Role of CFTR in epithelial physiology

Vinciane Saint-Criq, Michael A. Gray

► To cite this version:

Vinciane Saint-Criq, Michael A. Gray. Role of CFTR in epithelial physiology. Cellular and Molecular Life Sciences, 2017, 74 (1), pp.93-115. 10.1007/s00018-016-2391-y . hal-02628029

HAL Id: hal-02628029

<https://hal.inrae.fr/hal-02628029>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Role of CFTR in epithelial physiology

Vinciane Saint-Criq¹ · Michael A. Gray¹

Received: 27 September 2016 / Accepted: 28 September 2016 / Published online: 6 October 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Salt and fluid absorption and secretion are two processes that are fundamental to epithelial function and whole body fluid homeostasis, and as such are tightly regulated in epithelial tissues. The CFTR anion channel plays a major role in regulating both secretion and absorption in a diverse range of epithelial tissues, including the airways, the GI and reproductive tracts, sweat and salivary glands. It is not surprising then that defects in CFTR function are linked to disease, including life-threatening secretory diarrhoeas, such as cholera, as well as the inherited disease, cystic fibrosis (CF), one of the most common life-limiting genetic diseases in Caucasian populations. More recently, CFTR dysfunction has also been implicated in the pathogenesis of acute pancreatitis, chronic obstructive pulmonary disease (COPD), and the hyper-responsiveness in asthma, underscoring its fundamental role in whole body health and disease. CFTR regulates many mechanisms in epithelial physiology, such as maintaining epithelial surface hydration and regulating luminal pH. Indeed, recent studies have identified luminal pH as an important arbiter of epithelial barrier function and innate defence, particularly in the airways and GI tract. In this chapter, we will illustrate the different operational roles of CFTR in epithelial function by describing its

characteristics in three different tissues: the airways, the pancreas, and the sweat gland.

Keywords CFTR · Physiology · Epithelial transport · Chloride · Bicarbonate

Introduction

The basic toolkit for epithelial electrolyte and fluid absorption and secretion

Epithelial tissues are composed of one or more layers of closely assembled cells that cover a surface or that line a cavity. The main characteristic of epithelial cells is that they are polarised; the plasma membrane in contact with the external environment is called the mucosal or apical membrane, whereas the basolateral (serosal) membrane faces toward the interstitium. Both membranes have distinct roles due to their localization and differential expression of proteins (Fig. 1a). The basolateral membrane uptakes nutrients, ions and oxygen from the blood, and disposes of the cellular waste products. Because the apical membrane is in contact with the external environment, it serves both as a physical and chemical barrier to prevent potential pathogens or toxic matter reaching the bloodstream. The epithelial cells are joined together by points of contacts separating the apical and basolateral membrane: the interstitium is separated from the apical external milieu by junctional proteins. These proteins were originally described by histologists and grouped into four different structures: the zona occludens (tight junctions), zona adherens, macula adherens, and gap junctions. These are fundamental to maintain the polarity of the epithelia and, therefore, for the directional movement of ions and fluid

✉ Michael A. Gray
m.a.gray@newcastle.ac.uk
Vinciane Saint-Criq
vinciane.saint-criq@newcastle.ac.uk

¹ Epithelial Research Group, Institute for Cell and Molecular Biosciences, University Medical School, Newcastle University, Framlington Place, Newcastle upon Tyne NE2 4HH, UK




Fig. 1 Basic characteristics of an epithelial layer. **a** Epithelial cells are joined together by junctions [tight junctions (TJ), gap junctions (GJ)]. Uptake of nutrients and oxygen, and removal of cellular waste products occur on the basolateral surface. Water and ion transport can occur through the transcellular or paracellular pathways. **b** Absorption is mainly driven by active Na^+ absorption through ENaC in the apical membrane and the Na^+/K^+ -ATPase in the basolateral membrane creating an electrochemical driving force for paracellular passive Cl^- transport. Water then follows either through aquaporins or the

paracellular pathway. **c** Secretion is mainly driven by Cl^- secretion through CFTR and other Cl^- channels in the apical membrane. NKCC1 and the coupled action of an anion exchanger and NBC in the basolateral membrane accumulate Cl^- in the cell. Active Cl^- secretion creates the driving force for Na^+ movement across the epithelium through the paracellular pathway and water transport occurs paracellularly and/or transcellularly. Red and green arrows show active transport, and yellow arrows show passive transport

that underscore the processes of absorption and secretion [1]. Gap junctions also allow electrical communication between cells as well as the diffusion of low molecular weight solutes.

The overall mechanism for electrolyte and fluid transport depends on the tissue and mainly on the differential expression, and distribution, of ion channels, exchangers, cotransporters, and pumps. Salt and fluid move either through the paracellular (between the cells) or transcellular (through the cells) pathways (Fig. 1a). The transcellular route of electrolyte transport requires active (ATP-dependent) or passive (following electrochemical gradients) transport of ions. The paracellular route is a passive process that is ultimately controlled by the prevailing transepithelial electrochemical gradients, as well as the cation-to-anion selectivity of the tight junctions. Indeed, the differential expression of claudins, which have been extensively studied in kidney epithelia [2–4], have been shown to regulate paracellular permeability, through their ability to modulate the charge, water, and size selectivity of the tight junctions [5]. Furthermore, recent patch clamp studies from polarised cultures of kidney epithelial cells have shown that claudin-4 appears to act as a chloride-selective pore, with properties very similar to conventional membrane

spanning ion channels [6] Transepithelial electrical resistance (TEER) measurement is used to assess the barrier function of epithelial tissues, or cells grown on semi-permeable supports, and reflects the paracellular as well as transcellular conductivity. It is known that specific tight junction proteins largely influence epithelial resistance [7], and in the early 1970s, Frömter and Diamond used this measurement to classify epithelia into leaky (with a low TEER) and tight ($\text{TEER} > 500 \Omega \text{ cm}^{-2}$). Leaky epithelia also generate a small transepithelial voltage (V_t) that is due to a low junctional resistance as well as to electroneutral transport (i.e., cotransport of a cation with an anion or exchange of an anion for another anion). In contrast, tight epithelia can produce high V_t , and are able to maintain large ion concentration and osmotic gradients [8]. Therefore, leaky epithelia are mainly involved in isotonic electrolyte and fluid transport, whereas tight epithelia are mainly involved in hypo- or hypertonic salt and fluid absorption or secretion.

Generally, salt absorption depends on the active transepithelial absorption of sodium (Na^+) ions, which creates an electrochemical driving force for passive chloride (Cl^-) transport in the luminal to basolateral direction (Fig. 1b). This, in term, creates a salt concentration gradient

across the epithelium provoking water to be absorbed passively by osmosis. Na^+ absorption is generally governed by the activity of the apically located epithelial Na^+ channel (ENaC), a heterotrimer composed of alpha, beta, and gamma subunits. Na^+ efflux across the basolateral membrane is then driven by the Na^+/K^+ -ATPase, which ultimately maintains an inwardly directed Na^+ gradient necessary for absorption. In addition, both apical and basolateral K^+ channels are essential to maintain a suitable negative membrane potential for Na^+ influx to occur (Fig. 1b).

In contrast, transcellular salt secretion is mainly controlled by Cl^- exit across the apical membrane and this occurs predominantly via CFTR, although a number of distinct Cl^- channels, such as calcium-activated Cl^- channels (CaCC), and solute carrier (SLC)26A9, are also involved, depending on tissue. Secretion depends on active Cl^- transport, which creates the driving force for Na^+ movement across the epithelium through the paracellular pathway (Fig. 1c). The increased salt concentration on the luminal surface generates an osmotic driving force for water to be secreted, producing an isotonic secretion. For both processes, water moves either passively through the paracellular pathway or transcellularly via aquaporins, or by both routes [9–11] (Fig. 1b, c). It is important to note that in addition to Cl^- , CFTR and ANO1 also conduct HCO_3^- and that transport of this anion will also contribute to transepithelial fluid secretion (not shown on Fig. 1). Under normal physiological electrochemical gradients, and because CFTR is about 5 times more conductive for Cl^- than HCO_3^- , transport of Cl^- by CFTR accounts for the majority of fluid secretion in most secretory tissues. However, the secretion of HCO_3^- does play a key role in modulating the pH of the secreted fluid, which is discussed in more detail below.

In CFTR expressing tissues, fluid secretion is primarily controlled by the extent of transcellular Cl^- transport, with the rate of Cl^- exit across the apical membrane being the rate-limiting step. Chloride secretion is essentially a two-stage process, which begins with the active accumulation of Cl^- across the basolateral membrane, through the $\text{Na}^+ \text{K}^+ 2\text{Cl}^-$ cotransporter (NKCC1), a secondary-active transporter that uses the inwardly directed Na^+ gradient, established by the Na^+/K^+ -ATPase, to accumulate Cl^- above electrochemical equilibrium. In addition, recent evidence suggests that a basolateral $\text{Cl}^-/\text{HCO}_3^-$ anion exchanger (most likely SLC4A2) working in parallel with an Na^+ -bicarbonate cotransporter (NBC) can also accumulate Cl^- within some epithelial cells [12]. In many epithelial tissues, Cl^- exit across the luminal membrane occurs via CFTR. The total CFTR Cl^- conductance of the apical membrane is dependent on three parameters: the activity, or open state probability (P_o) of CFTR, which is

controlled predominantly via cAMP/PKA phosphorylation as described in chapter “Biochemistry and physiology of CFTR”; the number or density of CFTR channels (N), and finally the single channel conductance (G_s), which itself is governed by the electrochemical gradient across the apical membrane (membrane potential and Cl^- concentration). The net transport of Cl^- across the epithelium then drives passive Na^+ transport via a cation-selective paracellular route, and water follows osmotically. Importantly, as Cl^- exits the cells the apical membrane will depolarise which will ultimately limit Cl^- secretion, as the apical membrane potential moves towards the Cl^- equilibrium potential. In some epithelial cells, intracellular Cl^- also falls due to CFTR activity, which can lead to activation of several protein kinases that alter CFTR anion permeability (see section on pancreas).

The role of CFTR in epithelia has been extensively studied in relation to the affected organs in cystic fibrosis (CF). As stated in Chapter “Cystic Fibrosis: a clinical view”, CF is the most common genetically inherited disease in Caucasian populations (1 in 3500 newborns in Europe) [13, 14] and 70–90 % of CF individuals harbour the F508del mutation on at least one allele [15], which results in misfolding and incorrect processing of CFTR to the apical membrane. One of the first symptoms associated with CF, and occurring in around 17 % of CF patients, is meconium ileus; an obstruction of the bowel due to thick meconium and 98 % of babies with meconium ileus has CF. Distal intestinal obstruction syndrome also has a lifetime prevalence of 8 % in children with CF and 16 % in adult CF patients, and constipation occurs in around 50 % of CF patients [16] (median of 16 % in the general population [17]). Moreover, historically, some of the first suspected descriptions of CF come from the XVI century and associated a salty skin and pancreatic damage with early childhood death [18]. Finally, the main cause of mortality and morbidity in CF is the lung pathology in which thick and sticky mucus blocks the airways promoting the development and persistence of virulent pathogens. To develop further the fundamental, but contrasting, roles of CFTR in the physiology of epithelia, we will describe its role in three different CF-affected organs; the lungs, the pancreas, and the sweat gland. However, it is important to note that CFTR plays a fundamental role in electrolyte transport throughout most of the intestinal tract, as well as in the female and male reproductive tracts, but due to space limitations, these topics will not be discussed further. For the interested reader, several recent reviews describe in detail the role of CFTR in anion (Cl^- and HCO_3^-) and fluid transport in these tissues, together with diseases associated with defective CFTR, such as CF secretory diarrhoeas and infertility [19–24].

Role of CFTR in the lungs

The main function of the lung is to provide oxygen to the bloodstream and, therefore, to all organs in the body, and to remove carbon dioxide from the bloodstream out to the atmosphere. The lung is composed of the conducting airways and the respiratory airways (Fig. 2). Both entities serve a particular function. The conducting airways composed of the nose, mouth, pharynx, larynx, trachea, bronchi and conducting bronchioles, conduct and warm up the air from the upper airways to the respiratory airways (respiratory bronchioles, alveolar ducts, alveoli) where gas exchange occurs. The epithelia lining the surfaces of the conducting and the respiratory airways have different cell composition: the bronchial epithelium is mainly composed of ciliated cells, goblet cells and basal cells (Fig. 2a), whereas the alveolar epithelium consists of alveolar type I and alveolar type II cells (Fig. 2b). It is also important to note that submucosal glands are found in the cartilaginous conducting airways and are composed of distinct regions with different types of cells, including ciliated and non-ciliated serous and mucous cells [25], with different functions (not shown in the figure but described below).

The conducting airways also have a role of preventing any noxious matter from reaching the alveolar gas exchange site. To accomplish this role, the epithelial cells lining the airways are protected from the inspired air by a thin ($\sim 10\ \mu\text{m}$) liquid layer called the airway surface liquid (ASL) that is composed of a lower periciliary layer (PCL) ($\sim 7\ \mu\text{m}$), in which cilia beat to remove the inhaled

particles and pathogens trapped in the upper mucus layer (Fig. 2a). To maintain an efficient mucociliary clearance, the process that allows the removal of trapped particulate matter by the coordination of fluid secretion and ciliary beating, ASL hydration is tightly regulated by the transport of ions and water across the surface epithelia as well as by fluid secretion arising from submucosal glands. Indeed, Ballard et al. provided evidence that the ASL is mainly secreted by the glands lying below the mucosal layer. They showed in pig airways that the rate of fluid secretion induced by acetylcholine was unaffected by the removal of the surface epithelium [26]. In submucosal glands, mucous cells are mainly localised in the tubules distal from the collecting duct, whereas the CFTR-expressing serous cells are predominantly present in the distal ducts and acini. Mucous cells secrete gel-forming mucins (mainly MUC5B) as well as fluid, but the serous cells are the primary origin of the fluid component of the gland secretions, secreting solutes, fluid and innate defence molecules [27]. Although, in the upper airways, much of the ASL originates from the submucosal glands, surface epithelia have a critical role in regulating final ASL volume and composition. This is because surface airway epithelial cells express both ENaC and CFTR and, therefore, are able to absorb Na^+ as well as secrete Cl^- . Studies using thin film cultured human bronchial epithelial cells (HBEs) have shown that both Na^+ absorption and Cl^- secretion regulate ASL height and composition [28, 29]. In CF airways, the ASL is depleted which strongly affects mucociliary clearance and, therefore, the eradication of bacterial infections. Moreover, it




Fig. 2 Cellular components of the airways. The airways are composed of the conducting airways (a) and the respiratory airways (b). **a** The conducting airways are covered by an aqueous film called the airway surface liquid (ASL) which is composed of the periciliary layer (PCL) and the mucus layer (M). Three types of cells constitute

the conducting airways: the ciliated cells (CC), goblet cells (GC), and basal cells (BC). **b** The respiratory airways are composed of the alveolar type I (ATI) and type II (ATII) cells and are covered by the alveolar lining fluid (ALF) that prevent alveoli from collapsing

was shown by Joo et al. that cAMP agonists failed to induce secretion by submucosal glands from CF patients [30]. As stated above, because of the importance of the submucosal glands in the formation of the ASL, it is likely that any defect in CFTR in these glands will strongly affect ASL homeostasis. Although CFTR is apparently not expressed in mucous cells [31], the defect in fluid secretion from serous cells affects mucin secretion (by mucous cells) as well as its hydration [32] and leads to severe mucus plugging within the glands, probably because of an acidic environment within the glands (see below).

In surface epithelia, ASL depletion was thought to be due mainly to a lack of Cl^- secretion that leads to a lack of an osmotic gradient, which, in turn, prevents fluid secretion, but it may also involve enhanced absorption through ENaC. Indeed, it has been shown that CFTR regulates other ion channels and transporters, and critically is involved in ASL pH regulation and innate immunity through regulation of secretion of antioxidant and antimicrobial molecules. All these functions of CFTR act together to maintain a close to sterile environment in the lungs that prevents pathogens and noxious agents from entering the blood circulation. Indeed, proper hydration of the ASL is crucial for an efficient mucociliary clearance, and recent studies have shown that ASL pH homeostasis is essential for bacterial killing [33, 34], mucus rheology [35] and fluid homeostasis [36].

Regulation of airway surface fluid composition and hydration

Surface airway epithelial cells, as well as serous cells of the submucosal glands, secrete Cl^- and HCO_3^- in response to agents increasing intracellular cAMP (VIP, adenosine and noradrenaline) and/or Ca^{2+} (acetylcholine, histamine or ATP) and this controls ASL volume and composition. As described in the introduction, for many epithelial cells, the Na^+/K^+ -ATPase actively transports Na^+ out of the cells producing a transmembrane electrochemical gradient allowing for the cotransport of Na^+ , K^+ , and Cl^- through NKCC1 at the basolateral membrane, thereby increasing intracellular Cl^- concentration above equilibrium. In serous cells, it has been shown that Cl^- can also be accumulated at the basolateral membrane by two other mechanisms: (1) through the coupled action of a Na^+/H^+ exchanger (NHE) and a $\text{Cl}^-/\text{HCO}_3^-$ exchanger and (2) parallel operation of a $\text{Cl}^-/\text{HCO}_3^-$ exchanger with NBC [12, 37]. At the apical membrane, Cl^- is secreted through CFTR, CaCCs or SLC transporters and channels.

In CF, an enhanced Na^+ absorption has also been reported and one hypothesis to explain this finding is that CFTR normally downregulates ENaC activity in non-CF cells. This was first suggested from studies in which

measurements of nasal potentials in CF patients showed a large response to amiloride (a highly selective ENaC inhibitor) that was not seen in non-CF patients [38, 39]. Later work from a β -ENaC overexpressing mouse [40] provided further support, since these transgenic mice exhibited a lung phenotype characteristic of human CF airways. Indeed, airway epithelia isolated from this model showed CF ASL depletion, dehydrated mucus, neutrophilic inflammation, and poor bacterial clearance. However, the regulation of ENaC by CFTR is an ongoing debate as some studies do not report the same results [41]. In an attempt to reverse the β -ENaC mouse phenotype, Grubb et al. [42] developed another mouse strain overexpressing human CFTR that were bred with the β -ENaC mice. The hypothesis was that increasing the amount of CFTR would restore a “normal” CFTR/ENaC ratio and reverse the CF-like phenotype of β -ENaC mice airways. In this study, the double transgenic mice (hCFTR/ β -ENaC) showed the same phenotype as β -ENaC mice, characterised by a reduced survival rate, airways obstruction and depleted ASL showing that in this model, CFTR was not able to rescue the lung phenotype observed in the β -ENaC mice [42]. There are a large number of studies supporting the regulatory effect of CFTR on ENaC in human airways (Fig. 3), but the mechanisms involved are still unclear due to contradictory results in different studies. A 2011 patch-clamp study from Lazrak et al. reported that CFTR regulates ENaC activity in isolated type II alveolar cells, even when CFTR protein levels were minimal [43]. On the other hand, the recent generation of CF pigs (both CFTR $^{-/-}$ and F508del) do not show any increased Na^+ or water absorption compared to wild-type animals, in spite of an increased amiloride-sensitive voltage, and short-circuit current, observed in airway epithelial cultures from these animals [44]. Although the direct or indirect regulatory effect of CFTR on ENaC still needs to be confirmed, several groups are pursuing ENaC inhibitors as a therapeutic approach to increase ASL depth, and the available data show that interfering with ENaC activity could potentially be beneficial in CF to rehydrate the ASL [45].

To maintain an optimally hydrated ASL, CFTR also regulates other Cl^- channels and transporters, some of which belong to the SLC26 family [46]. The mammalian SLC26 family of anion exchangers and channels is composed of 10 genes that encode proteins with a C- and N-termini that frame a transmembrane domain. In the C-terminus, there is a regulatory region called the STAS (sulphate transporter and anti-sigma factor antagonist) domain. Mutations in human SLC26 genes lead to tissue specific diseases such as autosomal recessive non-syndromic deafness, DFNB4, and Pendred Syndrome in the SLC26A4 gene [47], and studies in mice models have shed light on the organ distribution of other SLC26 transporters.




Fig. 3 Ion transport in the airways. On the apical surface, CFTR drives Cl^- and HCO_3^- secretion and regulates Na^+ absorption by inhibiting ENaC. CFTR also positively regulates the Cl^- channels ANO1 and SLC26A9 as well as the anion exchanger SLC26A4, increasing Cl^- and HCO_3^- secretion and therefore increasing ASL hydration and pH. Na^+ and water follow the electrochemical gradient through the paracellular pathway. On the basolateral membrane, NKCC1 accumulates Cl^- intracellularly supported by the Na^+/K^+ -ATPase. K^+ recycling across the basolateral membrane occurs for proper function of the Na^+/K^+ -pump

Among the 10 identified members of the SLC26 family, two have been shown to be expressed in airway cells, SLC26A4 and SLC26A9. Importantly, genome-wide association studies have recently demonstrated a significant association between single nucleotide polymorphisms (SNPs) in SLC genes and CF disease severity. Correlations were found with susceptibility to meconium ileus, *Pseudomonas* infections and decline in pulmonary function in CF children (SLC9A3; Na^+/H^+ exchanger) and CF-related diabetes (SLC26A9; $\text{Cl}^-/\text{HCO}_3^-$ transporter) [48, 49]. Among the latest discovered SLC26 are the SLC26A7, A8, and A9 [50]. SLC26A9 is predominantly expressed in the brain and lungs, and recent work suggests that it plays an important role in airways hydration and mucus homeostasis as a polymorphism in the 3'UTR region has been shown to be associated with asthma [51]. To study the function of this transporter, Lohi et al. injected *Xenopus laevis* oocytes with SLC26A9 cRNA and measured the uptake of radio-labelled sulphate, Cl^- and oxalate. They reported the transport of all three anions by SLC26A9 and an inhibition of sulphate transport by DIDS and thiosulphate, implying that the protein acted as an anion exchanger, similar to several other SLC26 members (A3, A4 and A6). Further

characterisation of this transporter reported a small effect of the specific CFTR inhibitor, CFTR-Inh172, in patch-clamp experiments from oocytes [52] but a more significant effect of GlyH-101 (another CFTR inhibitor, but with lower selectivity) in SLC26A9-transfected HEK 293 [53]. However, it was shown in 2009 that SLC26A9 was regulated by CFTR in HBEs and that the forskolin response was enhanced in cells coexpressing CFTR and SLC26A9, when compared to HEK-293 expressing CFTR alone. Since the increase in current was much greater than the addition of each response alone, this suggested that both channels interacted synergistically in response to forskolin. Moreover, combined data from patch-clamp experiments in HEK 293 cells and short-circuit current experiments in HBE cells showed that the constitutive basal Cl^- current observed in HBE cells was likely to be due to SLC26A9. This current was absent from HBE cells cultured at air liquid interface (ALI) from CF patients, and furthermore, these cells lacked the signature currents, typical of SLC26A9 alone or coexpressed with WT-CFTR in HEK 293 coexpressing F508del-CFTR and SLC26A9 [53], suggesting that the mutant CFTR prevented SLC26A9 activity. However, in the same year, Chang et al. showed that this anion transporter was inhibited by the regulatory domain of CFTR and that this effect occurred through an interaction with the STAS domain of SLC26A9 [54]. Clearly, the effect of CFTR on SLC26A9 activity requires further investigation, but in the airways at least, it seems both channels work together to maintain ASL hydration (Fig. 3).

CFTR has also been shown to regulate Ca^{2+} activated Cl^- secretion, potentially via changes in CaCCs (Fig. 3). The histamine or UTP peak secretory response was enhanced in cultured CF airway epithelial cells and freshly excised nasal tissues from CF patients, respectively [55, 56]. In CF mice, enhanced Ca^{2+} -dependent Cl^- secretion was also detected [57]. In a study from 2011, soon after the identification of the CaCC as being encoded by TMEM16A (also called Anoctamin 1, ANO1) [58–60], Ousingawatt et al. reported that the activation of CFTR inhibited ANO1 in a bronchial epithelial cell line [61] and also showed a molecular interaction between the two channels in an overexpressing model using HEK293 cells. These are a few examples of how CFTR regulates other channels to maintain a fully hydrated ASL. However, CFTR is also involved in regulating the movement of water across the epithelium through the transcellular pathway by regulating aquaporins such as aquaporin 3 in non-CF cells [62]. Studies performed in CF and non-CF cell lines, by two different groups, also showed that CFTR modulates the paracellular pathway. When CFTR was activated in non-CF cells, the transepithelial electrical resistance decreased and paracellular conductance

increased. CFTR inhibition led to a disorganisation of F-actin and α -tubulin that was also observed in CF epithelia [63] and involved a myosin II dependent mechanism [64].

Another hallmark of CF airway disease that is also found in COPD is the accumulation of mucus that forms mucus plugs on the surface of airway epithelial cells. The composition of mucus includes water, ions, and macromolecules with protective functions such as antimicrobial, anti-protease, and anti-oxidant activity. Mucins are glycoproteins that are responsible for the viscoelastic property of mucus, which is crucial for an effective mucociliary clearance. Because organs affected in CF are mainly mucus producing epithelia, it has been proposed that a defect in CFTR expression and/or activity would affect mucus production and properties [65]. However, the fact that some mucus producing organs do not show complete obstruction in CF, such as the salivary and lacrimal glands, suggest an indirect role for CFTR in the regulation of mucus production. Moreover, the phenotype of the overexpressing β -ENaC mice (described earlier in this chapter) suggested that CFTR affects mucus by activating and/or inhibiting other factors, such as pH and HCO_3^- .

In the lower airways, alveolar cells must remove water from their surface to permit gas exchange. This is a major function after birth when the lungs are filled with amniotic fluid, but it is also involved in the clearance of fluid in the resolution of pulmonary oedema in adulthood. Cardiogenic lung oedema is due to an elevated hydrostatic pressure secondary to a high pulmonary venous pressure. It was thought that fluid movement into the alveolar lumen was due to a passive movement of water through the paracellular pathway due to the elevated pressure. However, a recent study has shown that active signalling could be involved in this process and that CFTR could mediate alveolar Cl^- and fluid secretion [66]. However, other studies have shown that this anion channel is also involved in fluid resorption in alveolar cells. Na^+ absorption plays a crucial role in this mechanism [67–70], but recent evidence points towards a control of fluid absorption by CFTR. Using in situ perfused lungs from wild-type mice, Fang et al. showed that inhibition of Cl^- channels, as well as Cl^- substitution by gluconate, inhibited fluid clearance by 50 % [71]. Increasing intracellular cAMP led to an increase in fluid clearance, and this could be inhibited by addition of glibenclamide, an inhibitor of CFTR, confirming the role of CFTR in alveolar fluid resorption. The same group validated their findings in primary human alveolar type II cells cultured at air–liquid interface. Inhibition of CFTR with CFTR-Inh172 did not have any effect on basal fluid absorption but inhibited ~ 35 % of cAMP-induced fluid absorption [72]. A more recent study by

Korbmacher et al., in which they used a novel Deuterium oxide (D_2O) dilution method in conjunction with short-circuit measurements in Ussing chamber experiments, showed that inhibition of ENaC abolished more than 70 % of water resorption. Inhibition of CFTR (using CFTR-Inh172 or glibenclamide) reduced basal water resorption by ~ 15 –20 %, whereas a more general inhibition of Cl^- channels by NPPB blocked almost 60 % of this process, demonstrating the involvement of CFTR and other apical Cl^- channels in water resorption across the alveolar epithelium [73].

Regulation of the ASL pH by CFTR

Many recent studies have highlighted the importance of ASL pH regulation in airways homeostasis. Indeed, it was first demonstrated in the 1970s that the pancreatic secretions in patients with CF were acidic (see section on pancreas), and some years later, it was hypothesized that the defect in HCO_3^- secretion caused by defective CFTR could also occur in the lungs, and therefore decrease the ASL pH [74]. An acidic ASL pH has also been demonstrated in other lung pathologies such as asthma [75], COPD [76] and acute respiratory distress syndrome [77]. Although the link between CFTR and asthma is still controversial [78–81], it is now well established that COPD shares some common features with CF, including bronchiectasis, mucus plugging, and inflammation. The primary cause of COPD is a chronic exposure to oxidative insults such as cigarette smoking or passive exposure to cigarette smoke. Cigarette smoke exposure leads to an increase in inflammation and a decrease in CFTR activity. It was first reported in the early 1980s, before the identification of the CFTR gene, that cigarette smoke decreased Cl^- secretion which could not be prevented by administration of antioxidants [82]. In 2006, Cantin et al. showed that this effect was due to an inhibition of CFTR [83], and it was later demonstrated that cigarette smoke actually induced CFTR internalisation, and this led to ASL dehydration [84], potentially contributing to the development of COPD pathology.

There is now more evidence to show that the lack of HCO_3^- secretion in CF airways leads to a reduced ASL pH, and this further deteriorates the lung pathophysiology. An acidic ASL pH was demonstrated in the early 2000s in primary HBE cell cultures from CF and non-CF patients [85]. This study also showed that activating CFTR by increasing intracellular cAMP, increased ASL pH in non-CF epithelia but had the opposite effect in CF monolayers. Because normal submucosal glands are capable of secreting HCO_3^- and CFTR conducts this anion, it has been hypothesized that submucosal gland fluid, and therefore ASL pH, would be affected in CF. Indeed, Song et al.

showed that freshly collected fluid secreted by submucosal glands from CF patients was more acidic than glands from non-CF individuals [86]. Also in a 2012 study on newborn CF pigs, Pezzulo et al. showed that the ASL was more acidic when compared to wild-type piglets. Although the antimicrobial composition of the ASL was similar in CF and non-CF piglets, the airways from the CF pigs showed a reduced efficiency in bacterial killing, and they suggested that the acidic pH was the cause for the lack of activity of antimicrobial molecules [34]. Moreover, in primary HBEs, the decreased ASL pH also increased Na^+ and fluid absorption, because the secreted protein short palate lung and nasal epithelial clone 1 (SPLUNC1) effectively was not able to inhibit ENaC-dependent Na^+ absorption and preserve ASL volume, when ASL pH was acidic [36]. Finally, mucus properties are also regulated by pH, and different groups have shown the negative impact of an acidic pH on mucus rheology [65, 87–89]. In a review published in 2008, Quinton hypothesized that the original cause of the different organ pathologies in CF would be the loss of HCO_3^- , which would affect mucin expansion and mucus gel formation. Indeed, before they are released, mucins are in a very condensed form in granules. This compact form is maintained within the granules by a high concentration of cations, specifically Ca^{2+} and H^+ [90, 91]. Upon release of mucins, HCO_3^- would complex with Ca^{2+} and H^+ , allowing mucin chains to fully expand and form a gel [65].

As stated above, CFTR regulates extracellular (luminal) pH. It appears to do this by its ability to conduct HCO_3^- as well as Cl^- . However, as discussed in more detail in the section on the pancreas, CFTR may also be able to regulate ASL pH via an indirect route through regulating the activity of SLC26 $\text{Cl}^-/\text{HCO}_3^-$ exchangers, such as pendrin and potentially other members of the SLC26 family (Fig. 3). The reciprocal regulatory interaction between the SLC26 and CFTR involves binding of the STAS domain to the regulatory domain of CFTR (see section on pancreas). The SLC26 transporter SLC26A4, pendrin, has been shown to be upregulated in inflammatory conditions in airway epithelia such as chronic rhinosinusitis [92, 93], upon stimulation with IL-17 [94] and in response to bacterial infections [95]. The other mechanism that involves CFTR in ASL pH regulation is its capacity to regulate $\text{Cl}^-/\text{HCO}_3^-$ anion exchange, and more particularly, SLC26A4 in airway epithelial cells. In serous cells, an increase in cAMP triggered HCO_3^- secretion through the apical membrane. Using genetically modified cell lines (Calu-3 WT, CFTR knockdown and pendrin knockdown), the authors showed that this effect was CFTR-dependent but that although fluid secretion was mainly mediated through CFTR, the cAMP-induced rise in HCO_3^- secretion was mediated via pendrin [96].

Regulation of pulmonary innate immunity

ASL pH is also a crucial factor for an efficient immune response to eradicate trapped pathogens and prevent their propagation. Indeed, it has been shown that the activity of molecules such as the protease inhibitor SPLUNC1, which regulates antimicrobial peptides (AMPs) and shows itself antimicrobial activity [97], depends on optimal ASL pH [36, 98], as does the activity of AMPs themselves [34]. On a more general level, when pathogens enter the airways, they are recognised by epithelial cells that activate inflammatory pathways to recruit immune cells, specifically neutrophils, into the airway lumen to eradicate the infection. It has been shown that CFTR regulates this process and when CFTR is absent, there is an upregulation of proinflammatory molecules. The current debate is whether the exacerbated inflammation in CF is due to the sustained infection or to the defective CFTR channel. In 2007, Verhaeghe et al. showed the overexpression of proinflammatory proteins in the airways of 24 weeks old CF fetuses [99]. Moreover, Tirouvanziam showed in 2000 and 2002 that a graft of non-infected tracheas or distal airways from CF fetuses on severe combined immunodeficiency (SCID) mice induced an increase in IL-8 and the recruitment of neutrophils [100, 101]. However, since the development of the CF pig model, it has been shown that inflammatory markers in bronchoalveolar lavages (BALs) of the lung did not differ from non-CF piglets, although the CF pigs showed a marked increased susceptibility to infection [102]. However, a recent transcriptomic study showed that although levels of inflammatory markers did not change, CF pig airways responded with a diminished host defence response after infection with *S. aureus*, when compared to non-CF pigs [103].

In addition, CFTR transports glutathione (GSH) [104] and thiocyanate (SCN^-) [105], and both molecules have been shown to have a crucial role in the regulation of the immune response in the lungs. GSH is a sulfhydryl containing tripeptide that can bind and inactivate oxidants. Reactive oxygen species (ROS) can be produced endogenously (from inflammatory cells) or come from exogenous sources (pollution, cigarette smoke). Their endogenous production is important for the antimicrobial defence of the lungs, but a sustained increase in ROS can lead to increased inflammation and tissue damage. It is, therefore, important for the epithelium to be able to inactivate ROS once the infection has been eradicated. As a GSH transporter, it has been shown that a defective CFTR leads to a reduced concentration of GSH in the ASL [106] which will impair the redox balance in the airways. SCN^- is an anion that plays an important role in the antimicrobial defence of the lungs by reducing tissue-damaging species such as hydrogen peroxide (H_2O_2) and hypochlorite (OCl^-) by

subjecting itself to oxidation by lactoperoxidase, producing hypothiocyanite (OSCN^-). OSCN^- itself has been shown to have antimicrobial activity [107] and SCN^- protected a lung cell line from injury caused by H_2O_2 and from OCl^- [108]. The absence of CFTR in CF and other inflammatory lung pathologies leads to an abnormal antioxidant composition of the ASL, worsening airway inflammation and damaging the epithelium. Epithelial tissue damage triggers signalling cascades that lead to wound repair to prevent potential pathogens to enter the bloodstream. In a process called epithelial restitution, the injured epithelial cells go through different de-differentiation and re-differentiation stages to proliferate, fill the “gap”, and recover their full functionality [109]. It has been shown that CFTR plays a critical role in this mechanism. Although it is not clear whether a defective CFTR decreases or enhances wound healing, many studies show that bronchial epithelial cells lacking CFTR are not able to reconstitute a fully differentiated epithelium [110–112].

In conclusion, CFTR has a fundamental role in the physiology of the airways. It is involved directly and indirectly in anion secretion that is crucial for airways hydration, ASL pH and defence homeostasis (Fig. 3). Potential therapies for CF have proven difficult to implement, possibly because CFTR is engaged in so many different functions in the airways. However, as correctors and potentiators of CFTR are being developed, another possible way to compensate for defective CFTR would be to target alternative ion channels or transporters such as the SLC26 family members (A4 or A9), or the anoctamins, short-circuiting their regulation by CFTR and rebalancing ASL homeostasis. One important advantage of this ‘alternate non-CFTR approach’ is that it would benefit all CF patients regardless of genotype, and would potentially be useful for other pathologies in which CFTR has been shown to be downregulated such as COPD.

Role of CFTR in the exocrine pancreas

The pancreas is composed of both exocrine and endocrine glands. The exocrine pancreas secretes ~ 2 L/day of an isotonic, HCO_3^- rich fluid, containing a complex mixture of digestive enzymes (zymogens), classically known as pancreatic juice. The exocrine pancreas is composed mainly of two types of epithelial cells; acinar cells which make up ~ 90 % by volume of the gland, and ductal cells which make up the remaining 10 %, with a small proportion of mucus secreting and endocrine cells. The ductal cells form a complex, tubular, network which ramifies throughout the gland [113] (Fig. 4). CFTR is highly expressed in the apical membrane of ductal cells but is not present in acinar cells [114, 115], and is essential for

electrolyte and fluid secretion from the gland. The importance of this fluid secretion to pancreatic function is well demonstrated in CF, where the lack of CFTR-dependent fluid secretion leads to near complete destruction of the gland at birth in ~ 85 % of people with CF [116, 117].

In terms of functional roles, the acinar cells secrete a small volume of an NaCl-rich secretion plus many types of digestive enzymes (8–20 g per day) in an inactive form, together with various factors (e.g., ATP) that contribute to cell signalling within the ductal system [118]. Acinar cells secrete in response to stimulation by acetylcholine, cholecystokinin (CCK), and several other agonists. These agonists cause a rise in intracellular calcium which stimulates NaCl and fluid secretion, as well the regulated exocytosis of enzyme-containing secretory granules [119]. Recent studies have also shown that following regulated exocytosis the local pH falls to approximately 6.8, due to proton secretion that accompanies exocytosis [120] (Fig. 5a).

The ductal cells forming the intercalated, and/or intralobular and interlobular ducts express CFTR (varies with species) and are responsible for secreting an isotonic, highly alkaline, fluid containing up to 160 mM NaHCO_3^- . However, the composition of the secretion is very much flow dependent in all species studied [113, 118, 121]. There is a reciprocal relationship between HCO_3^- and Cl^- concentrations: as fluid secretory rate increases, so does HCO_3^- concentration with a corresponding reduction in Cl^- concentration. At maximal flow rates, HCO_3^- concentration peaks around 140–160 mM in most species except the rat and mouse, where the maximum is around 70 mM [113, 118]. The major physiological regulator of pancreatic HCO_3^- secretion is the peptide hormone, secretin, which is released from intestinal cells mainly in response to acid chyme entering the intestine. Binding of secretin to ductal cells stimulates a rise in intracellular cAMP and activation of PKA, which leads to an increase in CFTR activity through phosphorylation of the channel, as described in detail in the chapter “Biochemistry and Physiology of CFTR”. Other transporters required for HCO_3^- secretion are also regulated by cAMP/PKA (see below).

While the role of the enzymes in digestion is clear, the functions of pancreatic electrolyte secretion are less precise, but include acting as a vehicle for transporting the inactive digestive enzymes to the small intestine where the HCO_3^- helps to neutralise gastric acid and so elevate duodenal pH to the optimal value required by the digestive enzymes. Since HCO_3^- is a chaotropic anion, it will also aid disaggregation of secreted enzymes following exocytosis, as well as help in mucin secretion and gel formation as described in the section on the airways. In addition, HCO_3^- acts to neutralise the protons that are cosecreted

Fig. 4 Simplified structure of the pancreas. The exocrine pancreas is composed of acini that surround a central lumen open to the duct system. Acinar cells (AC) secrete digestive enzymes into small intercalated ducts (ICD) where the pancreatic duct cells (PDC) raise the pH of the pancreatic juice (PJ). These ducts are directly connected to increasingly larger intralobular (intraLD) and interlobular (interLD) ducts that join the main pancreatic duct





Fig. 5 Ion transport in acinar (a) and pancreatic duct cells (b). **a** Upon stimulation with acetylcholine, cholecystokinin or other agonists, intracellular $[Ca^{2+}]$ increases and stimulates $NaCl$ and fluid secretion, as well as the exocytosis of enzyme-containing secretory granules. As these granules also contain H^+ , the local pH falls to approximately 6.8. Cl^- secretion occurs through a $CaCC$ on the apical membrane. **b** $CFTR$ conducts Cl^- and HCO_3^- and works in concert with a Cl^-/HCO_3^- apical exchanger, to mediate net transepithelial HCO_3^- secretion, with Cl^- recycling across the apical membrane. Na^+ moves paracellularly in response to transepithelial HCO_3^-

secretion, and water follows osmotically, to produce a HCO_3^- -rich isotonic fluid. Cl^- accumulates across the basolateral membrane via $NKCC1$ and accumulation of HCO_3^- inside the cells occurs through the hydration of CO_2 to HCO_3^- and H^+ by carbonic anhydrase (CA), together with backward transport of H^+ via the basolateral Na^+/H exchanger (NHE). This is driven by the Na^+ gradient established by the Na^+/K^+ -ATPase. The Na^+ -Bicarbonate cotransporter (NBC) helps accumulate HCO_3^- within the cell and maintain an electrical driving force for efflux of HCO_3^- across the apical membrane. It also works with K^+ channels to maintain a negative membrane potential

with enzymes from the acinar cells. Maintaining a neutral to alkaline luminal pH is important for the membrane dynamics of acinar cells and, consequently, the exocytosis of zymogen granules [122]. Furthermore, lower extracellular pH has been shown to enhance secretagogue-induced

zymogen activation within acinar cells [123]. Finally, recent studies have provided strong evidence that adequate HCO_3^- secretion is essential for preventing premature autoactivation of trypsin by protons within the ductal tree, and therefore inhibiting autodigestion of the gland. Indeed,

trypsin itself has been shown to reduce CFTR-dependent HCO_3^- secretion from duct cells [124], through activation of apical proteinase-activated receptor-2, which leads to further trypsin autoactivation. Furthermore, a fall in ductal HCO_3^- secretion appears particularly important in protecting the pancreas from developing acute pancreatitis induced by stresses such as bile and alcohol [125], the two most common causes of pancreatitis in man (see below).

Role of CFTR in pancreatic HCO_3^- secretion

As described in the section on the airways, CFTR is known to conduct a variety of anions, including Cl^- and HCO_3^- . Early patch clamp studies from rat and human pancreatic duct cells demonstrated that under near physiological gradients, CFTR was approximately 5 times more permeable to Cl^- than to HCO_3^- [126–129]. This finding, together with microelectrode studies from intact ducts that demonstrated the presence of a SITS-sensitive $\text{Cl}^-/\text{HCO}_3^-$ exchange activity on the apical membrane together with an NPPB-sensitive Cl^- conductance [130, 131], combined with intracellular pH measurements [132], led to the first model of HCO_3^- secretion that is shown in Fig. 5b [126, 130, 131]. In this model, CFTR was seen to act purely as Cl^- secretory channel that worked in concert with an apical $\text{Cl}^-/\text{HCO}_3^-$ exchanger, to mediate net transepithelial HCO_3^- secretion from the duct cells, with Cl^- recycling across the apical membrane. Na^+ then moves paracellularly in response to transepithelial HCO_3^- secretion, and water follows osmotically, to produce a HCO_3^- -rich isotonic fluid. In essence, CFTR's role is to help maintain the activity of the exchanger by providing a source of luminal Cl^- , as well as limiting intracellular Cl^- accumulation, through the operation of the exchanger, which would ultimately put a break on HCO_3^- efflux by the exchanger. In this early model, Cl^- was shown to be accumulated across the basolateral membrane via NKCC1, and active accumulation of HCO_3^- inside the duct cells was achieved through the hydration of CO_2 to HCO_3^- and H^+ by carbonic anhydrase, together with backward transport of H^+ , via the basolateral NHE. The latter step being driven by the inwardly directed Na^+ gradient established by the basolateral Na^+/K^+ -ATPase. In contrast to the rat, the pig was shown to express a basolateral H^+ -ATPase that served a similar purpose to the NHE [133]. While this CFTR/anion-exchanger model for HCO_3^- secretion satisfied the early electrophysiological results from the rat gland (which only secretes about 70 mM NaHCO_3), it was soon apparent that this model had serious deficiencies and could not explain the secretion of near isotonic (140–160 mM) NaHCO_3 juice secreted by the human pancreas, as well as many other species, such as the cat, pig, and guinea-pig [113, 118]. Indeed,

computer modelling of pancreatic HCO_3^- secretion clearly showed that this arrangement of transporters would only be capable of achieving HCO_3^- levels of ~ 80 mM [134, 135]. Partial resolution to this apparent problem came when the molecular identity of the putative $\text{Cl}^-/\text{HCO}_3^-$ exchanger was identified and found to be due to expression of several members of the SLC26 family of anion transporters, SLC26A3 and SLC26A6 [136–138], which were introduced in the section on the airways. Both these exchangers are electrogenic with A3 transporting 2Cl^- for 1HCO_3^- (therefore causing net Cl^- accumulation per transport cycle), while A6 transported 2HCO_3^- out of the cell for one Cl^- into the cell [139]. Therefore, the direction and magnitude of anion transport is not only dependent on the prevailing chemical gradients for Cl^- and HCO_3^- across the apical membrane, but is also affected by membrane potential. However, it is generally believed that SLC26A6 mediates the majority of the $\text{Cl}^-/\text{HCO}_3^-$ exchange at the apical membrane based on inhibitor profile (DIDS-sensitivity) and studies from transgenic *slc26a6* knock-out mice [140]. Furthermore, pioneering studies from the Muallem lab showed that both SLC26A exchangers were regulated by CFTR [141–144] through a physical interaction between the phosphorylated R domain of CFTR and the highly conserved STAS domain of the SLC26 transporter. This physical coupling between CFTR and the exchanger not only functionally activated anion exchange activity, but also had a positive effect on CFTR, and led to an increase in P_o [136, 143], although the mechanism for both these effects is currently not understood. In addition, the physical coupling of these two anion transporters was shown to be facilitated by their binding to several scaffold proteins such as NHERF1 and CAP50, via their respective C-terminal PDZ binding motifs, which brings CFTR and SLC26A exchangers in close proximity, essentially forming a multimeric, anion transporting, complex [136, 141, 145]. In relation to HCO_3^- secretion, and particularly in the case of SLC26A6, computer modelling showed that having a 2:1 electrogenic $\text{Cl}^-/\text{HCO}_3^-$ transporter on the apical membrane, working with CFTR, could theoretically increase HCO_3^- to ~ 120 mM [135]. Although this value was significantly higher than could be achieved by a 1:1 $\text{Cl}^-/\text{HCO}_3^-$ exchanger, it was still less than the measured values in human pancreatic juice (140 mM). The reason for this was that the capacity of the exchanger to increase HCO_3^- was found to be limited once luminal HCO_3^- levels increased above 120 mM, since it would be operating close to equilibrium (i.e., very slowly), during the secretion of 140 mM HCO_3^- . Taken together with the observation that guinea-pig ducts can still secrete HCO_3^- into a luminal fluid nominally free of Cl^- [146], this suggested that neither of the SLC26 transporters provided

the main route for HCO_3^- efflux across the apical membrane during *maximal* secretion. The final answer to the puzzle came from several observations. First, it was found that stimulation of HCO_3^- secretion by cAMP agonists in intact guinea-pig ducts led to a very marked drop in intracellular Cl^- [147], and this reduction in intracellular $[\text{Cl}^-]$ was subsequently shown to activate a kinase cascade involving the WNK1-OSR1/SPAK pathway, that ultimately led to phosphorylation of CFTR (or a regulatory protein) and a marked increase in the relative permeability of CFTR to HCO_3^- [148]. In addition, the low Cl^- concentration in the lumen also caused CFTR to shift its selectivity in favour of HCO_3^- ions [149]. All of these factors, therefore, ensure that a HCO_3^- -rich secretion is produced with much of the secreted HCO_3^- entering the lumen via CFTR.

This increase in the relative HCO_3^- conductance of CFTR, together with the observed membrane potential of ~ 60 mV of stimulated duct cells [146, 150], was found to be capable of further increasing ductal HCO_3^- levels to ~ 140 – 160 mM. However, when comparing ductal secretion from different species, it was also apparent that those species that produced a relatively low HCO_3^- -containing secretion (rat, mouse, and rabbit), had significant levels of NKCC1 activity, while high HCO_3^- secretors did not, and instead expressed mainly the Na^+ -dependent HCO_3^- transporter, NBCe1 (SLC4A4) [151], an electrogenic HCO_3^- importer with a 2HCO_3^- to 1Na^+ stoichiometry [113, 152]. In the latter case, the NBC not only helped accumulate HCO_3^- within the cell, by utilising the inward-directed sodium gradient across the basolateral membrane, but it also importantly helped maintain an electrical driving force for efflux of HCO_3^- across the apical membrane, by offsetting the depolarisation of the cell due to electrogenic anion secretion [151, 153, 154]. The NBC, therefore, works in concert with potassium channels that are found in ductal cells, to maintain a negative membrane potential [155]. Furthermore, like CFTR, NBC activity is positively regulated by cAMP/PKA phosphorylation [156], and both proteins are synergistically regulated in a coordinated fashion by intracellular IRBIT, in a cAMP and calcium-dependent manner [157, 158]. Together, these data suggest that the flow-dependent changes in luminal $\text{Cl}^-/\text{HCO}_3^-$ concentrations observed both in vivo and in vitro studies [113, 118] can be attributed to dynamic alterations in the mechanism of anion secretion through regulatory changes in CFTR and SLC26A6 activities. However, because HCO_3^- secretion is controlled by the activity of CFTR, the production of a high luminal HCO_3^- fluid must occur relatively high up in the ductal tree (intercalated/intralobular ducts), where CFTR is expressed [114, 115].

It should also be noted that there are inhibitory pathways that limit the extent of pancreatic HCO_3^- and fluid secretion, probably to prevent excessive hydrostatic pressure building up within the ductal tree at high secretory flow rates [113]. Several studies have shown that these inhibitory pathways work through the release of neurotransmitters such as Substance P, that lead to a reduction in HCO_3^- secretion by selectively reducing apical $\text{Cl}^-/\text{HCO}_3^-$ exchange activity through a PKC-dependent pathway [159–161].

Defects in CFTR-dependent HCO_3^- secretion predispose to pancreatic disease

Several studies have indicated that insufficient ductal HCO_3^- and fluid secretion leads to the destruction of the gland, as observed in the inherited disease CF [116, 162, 163]. An important consequence of impaired HCO_3^- secretion is an acidic pancreatic juice (less than 6.5) that increases mucus viscosity, and decreases the solubility of secreted digestive enzymes. Both these factors predispose to the formation of mucin/protein plugs and eventually cysts within the ductal tree, as well as premature activation of digestive enzymes. This ultimately leads to the destruction of the gland which is one of the characteristic pathological features of CF of the pancreas [65]. As discussed in Chapter “Cystic Fibrosis: a clinical view”, mutations in the CFTR gene cause CF. There are over 2000 disease causing mutations which have been grouped into 5–6 classes based on the functional consequences of the mutation. Classes 1–3 are severe mutations, while Classes 4–5 are mild mutations. In relation to pancreatic pathology, $\sim 85\%$ of people with CF are born pancreatic insufficient (PI), which equates to a reduction in pancreatic function of more than 95% . In these people, there is a very good correlation between disease severity and the class of mutation [164–166] with ‘severe’ CF mutations, such as the most common CF mutation, F508del and the class 3 gating mutant, G511D, strongly correlating with PI. For those with ‘milder’ mutations (some residual channel activity such as Class 4, R117H), pancreatic function is preserved (pancreatic sufficient, PS), albeit to differing extents. However, in general, these PS individuals require less enzyme supplements, but can become PI with age. As described above, the increase in activity of the apical SLC26A6 anion exchanger during secretion appears to be strongly dependent on its interaction with CFTR, and interestingly, the exchanger is activated by a number of CFTR mutants that lack Cl^- channel activity [167]. This correlates with a good retention of pancreatic function in patients carrying those mutations [168]. Furthermore, anion transport studies from polarised cultures of the human CF pancreatic ductal cell line CFPAC, which is homozygous

for F508del, showed that in addition to a lack of CFTR, apical SLC26A $\text{Cl}^-/\text{HCO}_3^-$ exchange activity was also absent, despite evidence for mRNA expression. Importantly, anion exchange activity was restored upon viral-mediated CFTR transduction of the CFPAC cells [169]. Taken together, these results strongly suggest that a functional CFTR at the apical plasma membrane is a prerequisite for SLC26A-mediated anion exchange, and that mild CFTR mutations are likely to preserve $\text{Cl}^-/\text{HCO}_3^-$ exchange activity, although this needs more formal demonstration.

Several recent investigations have also shown that two agents that classically induce stress/inflammation of the exocrine pancreas (pancreatitis), bile and alcohol, caused marked changes in HCO_3^- and fluid secretion based on in vitro intracellular pH and fluid transport studies from isolated microdissected ducts. At low concentrations, both agents increased HCO_3^- secretion, a response that required CFTR and $\text{Cl}^-/\text{HCO}_3^-$ exchange activity [170–172]. However, higher levels of these agents led to a severe inhibition of CFTR-dependent HCO_3^- secretion, which was due to profound mitochondrial damage and a consequent reduction in intracellular ATP levels [173, 174]. These studies were the first to suggest that ductal HCO_3^- secretion could play a protective role against these noxious agents not hitherto thought of. Further support for this hypothesis came from studying the extent of pancreatic inflammation and necrosis caused by these agents in vivo, using either CFTR knock-out mice or a mouse model with reduced CFTR expression (NHERF1 KO mouse) [182]. In both cases, the extent of pancreatic pathology induced by administration of these noxious agents was significantly increased, highlighting a key role of CFTR in pancreatic protection. Furthermore, patients with autoimmune [175] or acute and chronic alcohol-induced pancreatitis, showed marked abnormalities in membrane localisation and expression levels of CFTR [176]. Finally, there is a significant correlation between the development of pancreatitis and variants in the CFTR gene that do not cause a typical CF phenotype, but appear to have impaired HCO_3^- secretion [177–179].

In summary, CFTR plays an essential role in HCO_3^- and fluid secretion in the exocrine pancreas. It regulates HCO_3^- secretion in two fundamentally different ways; first, as a regulator of SLC26A-mediated $\text{Cl}^-/\text{HCO}_3^-$ exchange, and second, as a direct exit pathway for HCO_3^- secretion. Defects in CFTR-mediated HCO_3^- transport lead to severe pancreatic dysfunction. Strategies for improving HCO_3^- secretion in the CF pancreas are limited because of the marked tissue destruction at birth in the majority of people with CF. However, preliminary results from measurements of pancreatic function in young children with CF taking the Class 3 CFTR potentiator,

Ivacaftor (see Chapter “Cystic Fibrosis: a clinical view”) over 24 weeks, have shown a significant restoration of enzyme-secreting capacity (increased faecal elastase-1 levels), and by inference, pancreatic tissue regeneration, which is an extremely exciting finding [180] that warrants further research. Furthermore, it is known that variants (SNPs) in the SLC26A9 anion transporter can influence disease severity in the CF lungs and gut (meconium ileus) and, therefore, act as gene modifiers. Importantly, a recent study has suggested that SNPs in SLC26A9 can also influence the degree of pancreatic insufficiency [181]. This opens up the possibility of targeting this anion transporter as a potential therapeutic target to slow the progression of exocrine dysfunction in CF (in addition to the lungs and ileum).

In terms of pancreatitis, recent animal studies have suggested that strategies that help maintain levels of HCO_3^- secretion would limit the extent of pathology induced by bile and alcohol [174, 176, 182]. Furthermore, the effects of ethanol and ethanol metabolites on CFTR are consistent with reduced biogenesis, accelerated plasma membrane turnover, as well as channel inhibition [176]. Thus, restoring cell surface expression and activity of CFTR may partly alleviate the ethanol-induced damage. This potentially could be through the use of the FDA approved drug, Lumacaftor (see Chapter “Cystic Fibrosis: a clinical view”), which improves folding and processing of F508del-CFTR to the plasma membrane, as well as Ivacaftor to improve channel activity. Another goal here would be to find ways of preventing the marked reduction in ATP levels in ductal (and acinar) cells [125, 183]. A potential clinical strategy would be to try and improve nutritional support at a very early stage in acute pancreatitis, although a recent trial looking at the benefits of nasoenteric feeding after ~20 h of admission did not show any improvement in outcome compared to those patients that had on-demand oral feeding commencing at 24 h [184].

Role of CFTR in sweat gland physiology

The human eccrine (or atrichal) sweat gland helps to maintain whole body temperature via the production of sweat in response to a hot environment, exercise, or emotional situations. An individual can secrete up to 4L of sweat in an hour to thermoregulate [185, 186]. Sweat consists primarily of water and salt, mostly NaCl but is hypotonic with respect to the interstitium. The sweat gets secreted onto the surface of the skin, where heat is lost from the body by the latent heat of evaporation of the sweat fluid. The ability to lose heat is affected by the prevailing outside temperature as well as humidity. However, it is

increasingly being recognised that sweat has a number of other important roles which include the production and secretion of a range of AMPs such as LL-37, lactoferrin and dermicidin, an AMP unique to skin [187–189], as well as compounds that maintain skin condition (barrier function) and skin lubricants [190]. Sweat glands also contain stem cells important for renewing skin after wounding and burns [186, 191]. A number of disorders of sweat glands exist, with most involving defects in electrolyte and fluid production. These include CF, where lack of functional CFTR prevents normal NaCl absorption and leads to excessive salt loss ([192]; reviewed by [193]) which is discussed in more detail below; idiopathic anhidrosis (decreased volume of secretion), which may be caused by defects in calcium signalling [194, 195] and hyperhidrosis (uncontrolled and excessive sweat secretion), that may be due to changes in calcium signalling and water transport by aquaporin 5 [196, 197]. In addition, lack of sufficient antimicrobials can lead to skin infections and atopic dermatitis [198], and problems with wound healing and re-epithelialisation may be linked to stem cell dysfunction [191, 199, 200].

Ecocrine sweat glands are derived from embryonic ectoderm (as is the exocrine pancreas), and it is estimated that there are 2–4 million glands dispersed over the surface of the body, with the highest density on the forehead, palms of the hands and soles of the feet. The human sweat gland is a simple, coiled tubular, exocrine gland, 2–5 mm in length that resides in the lower part of the dermis, and which connects to the surface of the skin by a straight absorptive duct [186, 193]. The gland is composed of two structurally and functionally different units

[201, 202] (Fig. 6). (1) The secretory coil (Fig. 6a) that makes up most of the coiled part of the gland and which is responsible for producing the primary secretion mainly in response to sympathetic cholinergic innervation, but beta-adrenergic stimulation also elicits a low volume secretion. The secretory coil is composed of three cell types. There is a single layer of epithelial cells, which consist of two morphologically and functional distinct types of cells known as clear (agranular) and dark (granular) cells which occur in equal proportions (Fig. 6a). These cells are responsible for producing the primary ‘sweat’ secretion, together with various glycoproteins and AMPs, respectively [185, 203]. Clear cells express CFTR [203, 204] as well as a CaCC, most probably TMEM16A (ANO1) [205]. The myoepithelial cells are the third type of cells (Fig. 6a), and are modified smooth muscle cells that contract during sweat secretion, and which provide structural integrity to the secretory coil preventing damage to the tubule during gland stimulation [206]. (2) The absorptive straight duct (Fig. 6b) that acts to absorb NaCl, but not water, producing the final, hypotonic, sweat secretion that flows out to the skin. The absorption of salt, but not water, helps minimise salt loss from the body, which, if not controlled, could lead to circulatory collapse. Structurally, the proximal part of the absorptive duct is composed of a double layer of epithelial cells, which are electrically connected by gap junctions, and physically joined by desmosomes, to form a syncytium [202]. Further up towards the skin exit, the distal part of the duct consists of multiple layers of cells and is not thought to be involved in electrolyte or fluid transport [185].




Fig. 6 Schematic representation of an eccrine sweat gland. The human sweat gland is a simple-coiled tubular exocrine gland that resides in the dermis and connects to the surface of the skin by a straight absorptive duct. **a** The secretory coil duct epithelium is

composed of clear (CC), dark (DC), and myoepithelial cells (MEC), and is responsible for producing the primary secretion. **b** The reabsorptive duct epithelium is composed of two layers of cuboidal cells which absorb salt but not water




Fig. 7 Ion transport in the sweat gland. **a** Secretory coil duct epithelial cells secrete NaCl and water mainly in response to cholinergic (acetylcholine, ACh) stimulation. **b** The reabsorptive duct epithelial cells express a constitutively active CFTR on both apical and basolateral membranes. Both Na^+ and Cl^- move

transcellularly and both ENaC and CFTR work together to regulate net transepithelial NaCl absorption. Na^+ is pumped out of the cell across the basolateral membrane to the interstitial fluid by the Na^+/K^+ -pump, generating a transepithelial electrical gradient favouring Cl^- absorption

Role of CFTR in sweat gland electrolyte and fluid transport

CFTR is expressed in both portions of the sweat gland and has been shown to be involved in the elaboration of the primary secretion from the secretory coil in response to beta-adrenergic stimulation, as well as in the absorption of NaCl in the absorptive duct [193]. However, in the secretory coil, the major physiological stimulus is acetylcholine (ACh), which is released from post-ganglionic sympathetic cholinergic fibres, and which stimulates copious fluid transport through a non-CFTR dependent pathway. Although not fully resolved, the most likely Cl^- exit pathway is via the TMEM16A Cl^- channel, which has been localised to secretory cells [205], and which is activated by increases in cytosolic calcium via cholinergic stimulation. However, recent studies suggest that dark cells may also contribute to sweat secretion, but employ a distinct anion channel known as best2 (see [186] for further discussion), which may have an additional role in sweat pH and fluid regulation. Furthermore, studies from isolated sweat glands from adult mouse foot pads have shown expression of the mRNA for the Chloride Intracellular Channel 6 (Clc6), as well as for the slc26a4 $\text{Cl}^-/\text{HCO}_3^-$ anion exchanger known as pendrin. The latter protein could play a role in HCO_3^- secretion [207], as discussed in the section on the airways. In the clear cells at least, the production of the primary secretion involves accumulation of Cl^- above electrochemical equilibrium by a basolateral NKCC1, followed by Cl^- exit through apical-located Cl^-

channels, CFTR or TMEM16A. This creates a lumen negative potential difference that drives paracellular transport of Na^+ , and water follows osmotically via aquaporins to produce an isotonic secretion, containing ~ 145 mM Na^+ , 115 mM Cl^- , with the remaining anions being lactate and HCO_3^- [185].

The duct cells express the highest levels of CFTR, and in contrast to all other CF-affected epithelial tissues, the channel is functionally present at both apical and basolateral membranes [193, 208], and appears to be constitutively active [209] via PKA activity. Similar to the airways, ENaC is also expressed with CFTR at the apical membrane. However, there are marked differences in the way the two tissues operate (Compare Figs. 3 and 7). In the sweat duct, both Na^+ and Cl^- move transcellularly and both channels work together to regulate net transepithelial NaCl absorption. Indeed, pioneering work from the Quinton lab showed that ENaC activity was dependent on a functional interaction with a phosphorylated, Cl^- transporting, CFTR [193, 210, 211], highlighting a completely different regulatory interaction between the two channels. Exactly how the activities of CFTR and ENaC are coordinated is still not fully established but may be due to changes in intracellular pH that accompany Na^+ absorption through ENaC [212]. In the duct cells, (as in the airways) Na^+ entry occurs through ENaC down a large electrochemical gradient. Intracellular Na^+ is then pumped out of the cell across the basolateral membrane to the interstitial fluid by the Na^+/K^+ -ATPase, generating a transepithelial electrical gradient favouring Cl^- absorption (Fig. 7b).

However, unlike the airways, the paracellular pathway in the absorptive duct has little intrinsic Cl^- permeability in both normal and CF glands [208, 213] and due to high anion conductance of the apical and basolateral membranes of the duct cells (due to active CFTR), Cl^- moves transcellularly. Indeed, the declining Na^+ concentration in the sweat fluid creates a driving force for passive Cl^- entry into the cell through active CFTR, and then, Cl^- exits via basolateral CFTR [185, 193]. Through this process NaCl concentration can fall to ~ 50 mM, before diffusion of Cl^- becomes limited by luminal Cl^- concentration [214], at least under normal flow rates. However, it has been observed that at low flow rates Cl^- levels can fall as low as 10–15 mM, and pH becomes very acidic [215], which must involve a different set of transporters. As discussed by Bovell [185], this could be due to a yet unidentified apical $\text{Cl}^-/\text{HCO}_3^-$ exchanger that is coupled to proton-secretion via a V-type ATPase [216, 217], leading to net Cl^- absorption. Because of the low water permeability of the duct epithelium, NaCl absorption does not lead to concurrent water movement, and therefore, the resulting modified fluid is hypotonic, thus conserving valuable salt for whole body salt and fluid homeostasis [193].

Sweat gland and cystic fibrosis

As discussed above, CFTR has an important role in electrolyte and fluid secretion as well as absorption. In people with CF, dysfunctional CFTR affects both processes, and this leads to two important clinical manifestations of the disease. (1) Although rates of cholinergic sweat secretion are very similar in non-CF and CF individuals, beta-adrenergic (cAMP) secretion is absent in CF [218]. (2) The inability of the sweat duct to absorb Cl^- prevents Na^+ transport, and therefore, salt absorption is markedly reduced and sweat NaCl levels rise, producing an abnormally ‘salty’ sweat, one of the hallmarks of the disease. Although, under normal conditions, this is not a major clinical problem, during hot and humid conditions, people with CF can lose excessive salt and fluid, causing dehydration and heat prostration as originally observed by Dorothy Anderson [219]. Indeed, the increase in sweat forms the basis of the Quantitative Pilocarpine Iontophoresis Test (QPIT) that is still used today for diagnostic purposes [220, 221]. This test measures the concentration of Cl^- in sweat and requires iontophoretic introduction of pilocarpine (a cholinergic agonist) to stimulate local sweating and then sufficient sweat is collected for analysis of Cl^- . Concentrations of Cl^- in excess of 60 mM in children are diagnostic of CF. Normal values are less than ~ 40 mM. More recently, four new assays have been introduced to assess both the secretory and reabsorptive capacity of sweat glands. These assays can

measure changes in bioelectric potentials, which are higher in CF glands compared to normal [222], as well as skin impedance and rates of secretion, thus improving overall sensitivity and diagnostic value in sweat gland pathologies (see [221]) for a recent summary of these new tests).

In summary, CFTR plays an important role in both salt and fluid secretion and absorption in the sweat gland, which is similar to the airways. However, in marked contrast to the airways, the ability of CFTR to regulate NaCl absorption relies on a positive interaction and regulation of ENaC; in other words, CFTR works with ENaC, and not against it! Exactly how two channels in the same membrane have completely different regulatory interactions is intriguing but poorly understood, and requires further research. Uniquely, CFTR is expressed in both apical and basolateral membranes of duct epithelial cells, although the underlying mechanism for this dual targeting has not been elucidated. Undoubtedly, cell/tissue specific regulatory interactions will have a role to play, but little is known in the sweat gland of these processes. It is intriguing that recent work from the CF pig has shown CFTR is not expressed in the plasma membrane of smooth muscle cells but is targeted to the sarcoplasmic reticulum [223], which illustrates the importance of cell context in relation to CFTR targeting. Defects in CFTR-mediated Cl^- transport lead to severe sweat gland dysfunction and excessive salt loss in CF. Although not a major clinical problem, there is no doubt that the sweat gland has been pivotal to our general understanding of the role of CFTR in CF which has been eloquently summarised in a recent review by Paul Quinton [193], and it still holds great promise as a diagnostic tool for evaluation of new therapies for CF in this age of personalised ($N = 1$) medicine [224].

Final summary

CFTR has a central role in coordinating electrolyte and fluid transport in a range of epithelial tissues including the airways, GI and reproductive tracts and secretory glands. It achieves this not only by acting as a conduit for anion transport (its ion channel function), but also through its varied and complex regulation of other, non-CFTR channels, and anion transporters (its transporter regulatory function), that also participate in salt and fluid transport. Through this activity, CFTR has a key role in maintaining epithelial integrity and defence of the body as a whole. Although we have only described the role of CFTR in three epithelial tissues, we chose these primarily to illustrate the varied functional roles of CFTR. Understanding the basis for these different activities of CFTR has not only helped us gain a better understanding of the role of CFTR in the physiology of the tissue, but importantly, has led to the

development of new and better strategies to help overcome CFTR-related pathologies. This is best exemplified by the development of a new generation of small molecule therapies being used to treat the basic defect in CF, which is described in more detail in Chapter “Cystic Fibrosis: a clinical view”. However, this approach is also being applied to CFTR-dependent secretory diarrhoeas, and may be relevant to other disease in which a link to CFTR malfunction is emerging, such as acute pancreatitis, COPD and asthma. However, we still lack precise details about the underlying molecular mechanisms that endow CFTR with the ability to modulate other transporters and defence mechanisms, so further research is clearly important and warranted.

Compliance with ethical standards

Funding Work from the authors is supported by a Strategic Research Centre Grant (INOVCF) from the CF Trust UK (SRC003).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Hong JH, Park S, Shcheynikov N, Muallem S (2014) Mechanism and synergism in epithelial fluid and electrolyte secretion. *Pflugers Arch* 466:1487–1499
- Angelow S, Kim KJ, Yu AS (2006) Claudin-8 modulates paracellular permeability to acidic and basic ions in MDCK II cells. *J Physiol* 571:15–26
- Elias BC, Mathew S, Srichai MB, Palamuttam R, Bulus N, Mernaugh G, Singh AB, Sanders CR, Harris RC, Pozzi A, Zent R (2014) The integrin beta1 subunit regulates paracellular permeability of kidney proximal tubule cells. *J Biol Chem* 289:8532–8544
- Hou J, Rajagopal M, Yu AS (2013) Claudins and the kidney. *Annu Rev Physiol* 75:479–501
- Gunzel D, Yu AS (2013) Claudins and the modulation of tight junction permeability. *Physiol Rev* 93:525–569
- Hou J, Renigunta A, Yang J, Waldegger S (2010) Claudin-4 forms paracellular chloride channel in the kidney and requires claudin-8 for tight junction localization. *Proc Natl Acad Sci USA* 107:18010–18015
- Anderson JM (2001) Molecular structure of tight junctions and their role in epithelial transport. *News Physiol Sci* 16:126–130
- Fromter E, Diamond J (1972) Route of passive ion permeation in epithelia. *Nat New Biol* 235:9–13
- Fischbarg J (2010) Fluid transport across leaky epithelia: central role of the tight junction and supporting role of aquaporins. *Physiol Rev* 90:1271–1290
- Shachar-Hill B, Hill AE (2002) Paracellular fluid transport by epithelia. *Int Rev Cytol* 215:319–350
- Verkman AS, Matthay MA, Song Y (2000) Aquaporin water channels and lung physiology. *Am J Physiol Lung Cell Mol Physiol* 278:L867–L879
- Shan J, Liao J, Huang J, Robert R, Palmer ML, Fahrenkrug SC, O’Grady SM, Hanrahan JW (2012) Bicarbonate-dependent chloride transport drives fluid secretion by the human airway epithelial cell line Calu-3. *J Physiol* 590:5273–5297
- Salvatore D, Buzzetti R, Baldo E, Forneris MP, Lucidi V, Manunza D, Marinelli I, Messori B, Neri AS, Raia V, Furnari ML, Mastella G (2011) An overview of international literature from cystic fibrosis registries. Part 3. Disease incidence, genotype/phenotype correlation, microbiology, pregnancy, clinical complications, lung transplantation, and miscellaneous. *J Cyst Fibros* 10:71–85
- Southern KW, Munck A, Pollitt R, Travert G, Zanolli L, Dankert-Roelse J, Castellani C, Group ECNSW (2007) A survey of newborn screening for cystic fibrosis in Europe. *J Cyst Fibros* 6:57–65
- De Boeck K, Zolin A, Cuppens H, Olesen HV, Viviani L (2014) The relative frequency of CFTR mutation classes in European patients with cystic fibrosis. *J Cyst Fibros* 13:403–409
- van der Doef HP, Kokke FT, Beek FJ, Woestenenk JW, Froeling SP, Houwen RH (2010) Constipation in pediatric cystic fibrosis patients: an underestimated medical condition. *J Cyst Fibros* 9:59–63
- Mugie SM, Benninga MA, Di Lorenzo C (2011) Epidemiology of constipation in children and adults: a systematic review. *Best Pract Res Clin Gastroenterol* 25:3–18
- Quinton PM (1999) Physiological basis of cystic fibrosis: a historical perspective. *Physiol Rev* 79:S3–S22
- Borowitz D (2015) CFTR, bicarbonate, and the pathophysiology of cystic fibrosis. *Pediatr Pulmonol* 50(Suppl 40):S24–S30
- Chan HC, Sun X (2014) SLC26 anion exchangers in uterine epithelial cells and spermatozoa: clues from the past and hints to the future. *Cell Biol Int* 38:1–7
- De Lisle RC, Borowitz D (2013) The cystic fibrosis intestine. *Cold Spring Harb Perspect Med* 3:a009753
- Liu Y, Wang DK, Chen LM (2012) The physiology of bicarbonate transporters in mammalian reproduction. *Biol Reprod* 86:99
- Seidler UE (2013) Gastrointestinal HCO₃⁻ transport and epithelial protection in the gut: new techniques, transport pathways and regulatory pathways. *Curr Opin Pharmacol* 13:900–908
- Verkman AS, Galletta LJ (2009) Chloride channels as drug targets. *Nat Rev Drug Discov* 8:153–171
- Meyrick B, Reid L (1970) Ultrastructure of cells in the human bronchial submucosal glands. *J Anat* 107:281–299
- Ballard ST, Trout L, Bebek Z, Sorscher EJ, Crews A (1999) CFTR involvement in chloride, bicarbonate, and liquid secretion by airway submucosal glands. *Am J Physiol* 277:L694–L699
- Widdicombe JH, Wine JJ (2015) Airway gland structure and function. *Physiol Rev* 95:1241–1319
- Knowles MR, Boucher RC (2002) Mucus clearance as a primary innate defense mechanism for mammalian airways. *J Clin Invest* 109:571–577
- Tarran R, Loewen ME, Paradiso AM, Olsen JC, Gray MA, Argent BE, Boucher RC, Gabriel SE (2002) Regulation of murine airway surface liquid volume by CFTR and Ca²⁺-activated Cl⁻ conductances. *J Gen Physiol* 120:407–418
- Joo NS, Irokawa T, Wu JV, Robbins RC, Whyte RI, Wine JJ (2002) Absent secretion to vasoactive intestinal peptide in cystic fibrosis airway glands. *J Biol Chem* 277:50710–50715
- Engelhardt JF, Yankaskas JR, Ernst SA, Yang Y, Marino CR, Boucher RC, Cohn JA, Wilson JM (1992) Submucosal glands are the predominant site of CFTR expression in the human bronchus. *Nat Genet* 2:240–248
- Inglis SK, Corboz MR, Ballard ST (1998) Effect of anion secretion inhibitors on mucin content of airway submucosal gland ducts. *Am J Physiol* 274:L762–L766

33. Abou Alaiwa MH, Reznikov LR, Gansemer ND, Sheets KA, Horswill AR, Stoltz DA, Zabner J, Welsh MJ (2014) pH modulates the activity and synergism of the airway surface liquid antimicrobials beta-defensin-3 and LL-37. *Proc Natl Acad Sci USA* 111:18703–18708
34. Pezzulo AA, Tang XX, Hoegger MJ, Alaiwa MH, Ramachandran S, Moninger TO, Karp PH, Wohlford-Lenane CL, Haagsman HP, van Eijk M, Banfi B, Horswill AR, Stoltz DA, McCray PB Jr, Welsh MJ, Zabner J (2012) Reduced airway surface pH impairs bacterial killing in the porcine cystic fibrosis lung. *Nature* 487:109–113
35. Hoegger MJ, Fischer AJ, McMenimen JD, Ostedgaard LS, Tucker AJ, Awadalla MA, Moninger TO, Michalski AS, Hoffman EA, Zabner J, Stoltz DA, Welsh MJ (2014) Impaired mucus detachment disrupts mucociliary transport in a piglet model of cystic fibrosis. *Science* 345:818–822
36. Garland AL, Walton WG, Coakley RD, Tan CD, Gilmore RC, Hobbs CA, Tripathy A, Clunes LA, Bencharit S, Stutts MJ, Betts L, Redinbo MR, Tarran R (2013) Molecular basis for pH-dependent mucosal dehydration in cystic fibrosis airways. *Proc Natl Acad Sci USA* 110:15973–15978
37. Huang J, Shan J, Kim D, Liao J, Evagelidis A, Alper SL, Hanrahan JW (2012) Basolateral chloride loading by the anion exchanger type 2: role in fluid secretion by the human airway epithelial cell line Calu-3. *J Physiol* 590:5299–5316
38. Gowen CW, Lawson EE, Gingras-Leatherman J, Gatzky JT, Boucher RC, Knowles MR (1986) Increased nasal potential difference and amiloride sensitivity in neonates with cystic fibrosis. *J Pediatr* 108:517–521
39. Rowe SM, Accurso F, Clancy JP (2007) Detection of cystic fibrosis transmembrane conductance regulator activity in early-phase clinical trials. *Proc Am Thorac Soc* 4:387–398
40. Mall M, Grubb BR, Harkema JR, O'Neal WK, Boucher RC (2004) Increased airway epithelial Na⁺ absorption produces cystic fibrosis-like lung disease in mice. *Nat Med* 10:487–493
41. Collawn JF, Lazrak A, Bebek Z, Matalon S (2012) The CFTR and ENaC debate: how important is ENaC in CF lung disease? *Am J Physiol Lung Cell Mol Physiol* 302:L1141–L1146
42. Grubb BR, O'Neal WK, Ostrowski LE, Kreda SM, Button B, Boucher RC (2012) Transgenic hCFTR expression fails to correct beta-ENaC mouse lung disease. *Am J Physiol Lung Cell Mol Physiol* 302:L238–L247
43. Lazrak A, Jurkuvenaite A, Chen L, Keeling KM, Collawn JF, Bedwell DM, Matalon S (2011) Enhancement of alveolar epithelial sodium channel activity with decreased cystic fibrosis transmembrane conductance regulator expression in mouse lung. *Am J Physiol Lung Cell Mol Physiol* 301:L557–L567
44. Itani OA, Chen JH, Karp PH, Ernst S, Keshavjee S, Parekh K, Klesney-Tait J, Zabner J, Welsh MJ (2011) Human cystic fibrosis airway epithelia have reduced Cl[−] conductance but not increased Na⁺ conductance. *Proc Natl Acad Sci USA* 108:10260–10265
45. Donaldson SH, Galiotta L (2013) New pulmonary therapies directed at targets other than CFTR. *Cold Spring Harb Perspect Med* 3:a009787
46. Alper SL, Sharma AK (2013) The SLC26 gene family of anion transporters and channels. *Mol Aspects Med* 34:494–515
47. Sheffield VC, Kraiem Z, Beck JC, Nishimura D, Stone EM, Salameh M, Sadeh O, Glaser B (1996) Pendred syndrome maps to chromosome 7q21-34 and is caused by an intrinsic defect in thyroid iodine organification. *Nat Genet* 12:424–426
48. Blackman SM, Commander CW, Watson C, Arcara KM, Strug LJ, Stonebraker JR, Wright FA, Rommens JM, Sun L, Pace RG, Norris SA, Durie PR, Drumm ML, Knowles MR, Cutting GR (2013) Genetic modifiers of cystic fibrosis-related diabetes. *Diabetes* 62:3627–3635
49. Sun L, Rommens JM, Corvol H, Li W, Li X, Chiang TA, Lin F, Dorfman R, Busson PF, Parekh RV, Zelenika D, Blackman SM, Corey M, Doshi VK, Henderson L, Naughton KM, O'Neal WK, Pace RG, Stonebraker JR, Wood SD, Wright FA, Zielenski J, Clement A, Drumm ML, Boelle PY, Cutting GR, Knowles MR, Durie PR, Strug LJ (2012) Multiple apical plasma membrane constituents are associated with susceptibility to meconium ileus in individuals with cystic fibrosis. *Nat Genet* 44:562–569
50. Lohi H, Kujala M, Makela S, Lehtonen E, Kestila M, Saarialho-Kere U, Markovich D, Kere J (2002) Functional characterization of three novel tissue-specific anion exchangers SLC26A7, -A8, and -A9. *J Biol Chem* 277:14246–14254
51. Anagnostopoulou P, Riederer B, Duerr J, Michel S, Binia A, Agrawal R, Liu X, Kalitzki K, Xiao F, Chen M, Schatterny J, Hartmann D, Thum T, Kabesch M, Soleimani M, Seidler U, Mall MA (2012) SLC26A9-mediated chloride secretion prevents mucus obstruction in airway inflammation. *J Clin Invest* 122:3629–3634
52. Lloriol C, Dulong S, Avella M, Gabillat N, Bouloukos K, Borgese F, Ehrenfeld J (2008) Characterization of SLC26A9, facilitation of Cl(−) transport by bicarbonate. *Cell Physiol Biochem* 22:15–30
53. Bertrand CA, Zhang R, Pilewski JM, Frizzell RA (2009) SLC26A9 is a constitutively active, CFTR-regulated anion conductance in human bronchial epithelia. *J Gen Physiol* 133:421–438
54. Chang MH, Plata C, Sindic A, Ranatunga WK, Chen AP, Zandi-Nejad K, Chan KW, Thompson J, Mount DB, Romero MF (2009) Slc26a9 is inhibited by the R-region of the cystic fibrosis transmembrane conductance regulator via the STAS domain. *J Biol Chem* 284:28306–28318
55. Clarke LL, Paradiso AM, Boucher RC (1992) Histamine-induced Cl[−] secretion in human nasal epithelium: responses of apical and basolateral membranes. *Am J Physiol* 263:C1190–C1199
56. Mall M, Gonska T, Thomas J, Schreiber R, Seydewitz HH, Kuehr J, Brandis M, Kunzelmann K (2003) Modulation of Ca²⁺-activated Cl[−] secretion by basolateral K⁺ channels in human normal and cystic fibrosis airway epithelia. *Pediatr Res* 53:608–618
57. Grubb BR, Vick RN, Boucher RC (1994) Hyperabsorption of Na⁺ and raised Ca(2+)-mediated Cl[−] secretion in nasal epithelia of CF mice. *Am J Physiol* 266:C1478–C1483
58. Caputo A, Caci E, Ferrera L, Pedemonte N, Barsanti C, Sondo E, Pfeiffer U, Ravazzolo R, Zegarar-Moran O, Galiotta LJ (2008) TMEM16A, a membrane protein associated with calcium-dependent chloride channel activity. *Science* 322:590–594
59. Schroeder BC, Cheng T, Jan YN, Jan LY (2008) Expression cloning of TMEM16A as a calcium-activated chloride channel subunit. *Cell* 134:1019–1029
60. Yang YD, Cho H, Koo JY, Tak MH, Cho Y, Shim WS, Park SP, Lee J, Lee B, Kim BM, Raouf R, Shin YK, Oh U (2008) TMEM16A confers receptor-activated calcium-dependent chloride conductance. *Nature* 455:1210–1215
61. Ousingsawat J, Kongsuphol P, Schreiber R, Kunzelmann K (2011) CFTR and TMEM16A are separate but functionally related Cl[−] channels. *Cell Physiol Biochem* 28:715–724
62. Schreiber R, Nitschke R, Greger R, Kunzelmann K (1999) The cystic fibrosis transmembrane conductance regulator activates aquaporin 3 in airway epithelial cells. *J Biol Chem* 274:11811–11816
63. Nilsson HE, Dragomir A, Lazorova L, Johannesson M, Roomans GM (2010) CFTR and tight junctions in cultured bronchial epithelial cells. *Exp Mol Pathol* 88:118–127
64. Weiser N, Molenda N, Urbanova K, Bahler M, Pieper U, Oberleithner H, Schillers H (2011) Paracellular permeability of

- bronchial epithelium is controlled by CFTR. *Cell Physiol Biochem* 28:289–296
65. Quinton PM (2008) Cystic fibrosis: impaired bicarbonate secretion and mucoviscidosis. *Lancet* 372:415–417
 66. Solymosi EA, Kaestle-Gembardt SM, Vadasz I, Wang L, Neye N, Chupin CJ, Rozowsky S, Ruehl R, Tabuchi A, Schulz H, Kapus A, Morty RE, Kuebler WM (2013) Chloride transport-driven alveolar fluid secretion is a major contributor to cardiogenic lung edema. *Proc Natl Acad Sci USA* 110:E2308–E2316
 67. Basset G, Crone C, Saumon G (1987) Fluid absorption by rat lung in situ: pathways for sodium entry in the luminal membrane of alveolar epithelium. *J Physiol* 384:325–345
 68. Basset G, Crone C, Saumon G (1987) Significance of active ion transport in transalveolar water absorption: a study on isolated rat lung. *J Physiol* 384:311–324
 69. Berthiaume Y, Staub NC, Matthay MA (1987) Beta-adrenergic agonists increase lung liquid clearance in anesthetized sheep. *J Clin Invest* 79:335–343
 70. O'Brodovich H, Hannam V, Seear M, Mullen JB (1990) Amiloride impairs lung water clearance in newborn guinea pigs. *J Appl Physiol* (1985) 68:1758–1762
 71. Fang X, Fukuda N, Barbry P, Sartori C, Verkman AS, Matthay MA (2002) Novel role for CFTR in fluid absorption from the distal airspaces of the lung. *J Gen Physiol* 119:199–207
 72. Fang X, Song Y, Hirsch J, Galletta LJ, Pedemonte N, Zemans RL, Dolganov G, Verkman AS, Matthay MA (2006) Contribution of CFTR to apical-basolateral fluid transport in cultured human alveolar epithelial type II cells. *Am J Physiol Lung Cell Mol Physiol* 290:L242–L249
 73. Korbmayer JP, Michel C, Neubauer D, Thompson K, Mizaikoff B, Frick M, Dietl P, Wittekindt OH (2014) Amiloride-sensitive fluid resorption in NCI-H441 lung epithelia depends on an apical Cl(–) conductance. *Physiol Rep* 2:e00201
 74. Quinton PM (2001) The neglected ion: HCO₃[–]. *Nat Med* 7:292–293
 75. Hunt JF, Fang K, Malik R, Snyder A, Malhotra N, Platts-Mills TA, Gaston B (2000) Endogenous airway acidification. Implications for asthma pathophysiology. *Am J Respir Crit Care Med* 161:694–699
 76. Kostikas K, Papatheodorou G, Ganas K, Psathakis K, Panagou P, Loukides S (2002) pH in expired breath condensate of patients with inflammatory airway diseases. *Am J Respir Crit Care Med* 165:1364–1370
 77. Gessner C, Hammerschmidt S, Kuhn H, Seyfarth HJ, Sack U, Engelmann L, Schauer J, Wirtz H (2003) Exhaled breath condensate acidification in acute lung injury. *Respir Med* 97:1188–1194
 78. Goodwin J, Spitalo N, Yaghi A, Dolovich M, Nair P (2012) Cystic fibrosis transmembrane conductance regulator gene abnormalities in patients with asthma and recurrent neutrophilic bronchitis. *Can Respir J* 19:46–48
 79. Maurya N, Awasthi S, Dixit P (2012) Association of CFTR gene mutation with bronchial asthma. *Indian J Med Res* 135:469–478
 80. Sanford A (2012) The role of CFTR mutations in asthma. *Can Respir J* 19:44–45
 81. Schroeder SA, Gaughan DM, Swift M (1995) Protection against bronchial asthma by CFTR delta F508 mutation: a heterozygote advantage in cystic fibrosis. *Nat Med* 1:703–705
 82. Welsh MJ (1983) Cigarette smoke inhibition of ion transport in canine tracheal epithelium. *J Clin Invest* 71:1614–1623
 83. Cantin AM, Hanrahan JW, Bilodeau G, Ellis L, Dupuis A, Liao J, Zielinski J, Durie P (2006) Cystic fibrosis transmembrane conductance regulator function is suppressed in cigarette smokers. *Am J Respir Crit Care Med* 173:1139–1144
 84. Clunes LA, Davies CM, Coakley RD, Aleksandrov AA, Henderson AG, Zeman KL, Worthington EN, Gentzsch M, Kreda SM, Cholon D, Bennett WD, Riordan JR, Boucher RC, Tarran R (2012) Cigarette smoke exposure induces CFTR internalization and insolubility, leading to airway surface liquid dehydration. *FASEB J* 26:533–545
 85. Coakley RD, Grubb BR, Paradiso AM, Gatzky JT, Johnson LG, Kreda SM, O'Neal WK, Boucher RC (2003) Abnormal surface liquid pH regulation by cultured cystic fibrosis bronchial epithelium. *Proc Natl Acad Sci USA* 100:16083–16088
 86. Song Y, Salinas D, Nielson DW, Verkman AS (2006) Hyperacidity of secreted fluid from submucosal glands in early cystic fibrosis. *Am J Physiol Cell Physiol* 290:C741–C749
 87. Chen EY, Yang N, Quinton PM, Chin WC (2010) A new role for bicarbonate in mucus formation. *Am J Physiol Lung Cell Mol Physiol* 299:L542–L549
 88. Gustafsson JK, Ermund A, Ambort D, Johansson ME, Nilsson HE, Thorell K, Hebert H, Sjoval H, Hansson GC (2012) Bicarbonate and functional CFTR channel are required for proper mucin secretion and link cystic fibrosis with its mucus phenotype. *J Exp Med* 209:1263–1272
 89. Tang XX, Ostedgaard LS, Hoegger MJ, Moninger TO, Karp PH, McMenimen JD, Choudhury B, Varki A, Stoltz DA, Welsh MJ (2016) Acidic pH increases airway surface liquid viscosity in cystic fibrosis. *J Clin Invest* 126:879–891
 90. Perez-Vilar J, Olsen JC, Chua M, Boucher RC (2005) pH-dependent intraluminal organization of mucin granules in live human mucous/goblet cells. *J Biol Chem* 280:16868–16881
 91. Verdugo P, Deyrup-Olsen I, Aitken M, Villalon M, Johnson D (1987) Molecular mechanism of mucin secretion: I. The role of intragranular charge shielding. *J Dent Res* 66:506–508
 92. Ishida A, Ohta N, Suzuki Y, Kakehata S, Okubo K, Ikeda H, Shiraishi H, Izuhara K (2012) Expression of pendrin and perlestin in allergic rhinitis and chronic rhinosinusitis. *Allergol Int* 61:589–595
 93. Seshadri S, Lu X, Purkey MR, Homma T, Choi AW, Carter R, Suh L, Norton J, Harris KE, Conley DB, Kato A, Avila PC, Czarnocka B, Kopp PA, Peters AT, Grammer LC, Chandra RK, Tan BK, Liu Z, Kern RC, Schleimer RP (2015) Increased expression of the epithelial anion transporter pendrin/SLC26A4 in nasal polyps of patients with chronic rhinosinusitis. *J Allergy Clin Immunol* 136:1548–1558
 94. Adams KM, Abraham V, Spielman D, Kolls JK, Rubenstein RC, Conner GE, Cohen NA, Kreindler JL (2014) IL-17A induces Pendrin expression and chloride-bicarbonate exchange in human bronchial epithelial cells. *PLoS One* 9:e103263
 95. Scanlon KM, Gau Y, Zhu J, Skerry C, Wall SM, Soleimani M, Carbonetti NH (2014) Epithelial anion transporter pendrin contributes to inflammatory lung pathology in mouse models of Bordetella pertussis infection. *Infect Immun* 82:4212–4221
 96. Garnett JP, Hickman E, Burrows R, Hegyi P, Tiszlavicz L, Cuthbert AW, Fong P, Gray MA (2011) Novel role for pendrin in orchestrating bicarbonate secretion in cystic fibrosis transmembrane conductance regulator (CFTR)-expressing airway serous cells. *J Biol Chem* 286:41069–41082
 97. Garcia-Caballero A, Rasmussen JE, Gaillard E, Watson MJ, Olsen JC, Donaldson SH, Stutts MJ, Tarran R (2009) SPLUNC1 regulates airway surface liquid volume by protecting ENaC from proteolytic cleavage. *Proc Natl Acad Sci USA* 106:11412–11417
 98. Ahmad S, Tyrrell J, Walton WG, Tripathy A, Redinbo MR, Tarran R (2016) SPLUNC1 has antimicrobial and antibiofilm activity against burkholderia cepacia complex. *Antimicrob Agents Chemother* 60:6003–6012
 99. Verhaeghe C, Remouchamps C, Hennuy B, Vanderplasschen A, Chariot A, Tabruyn SP, Oury C, Bours V (2007) Role of IKK and ERK pathways in intrinsic inflammation of cystic fibrosis airways. *Biochem Pharmacol* 73:1982–1994

100. Delplanque A, Coraux C, Tirouvanziam R, Khazaal I, Puchelle E, Ambros P, Gaillard D, Peault B (2000) Epithelial stem cell-mediated development of the human respiratory mucosa in SCID mice. *J Cell Sci* 113(Pt 5):767–778
101. Tirouvanziam R, Khazaal I, Peault B (2002) Primary inflammation in human cystic fibrosis small airways. *Am J Physiol Lung Cell Mol Physiol* 283:L445–L451
102. Wine JJ (2010) The development of lung disease in cystic fibrosis pigs. *Sci Transl Med* 2:29ps20
103. Bartlett JA, Ramachandran S, Wohlford-Lenane CL, Barker CK, Pezzulo AA, Zabner J, Welsh MJ, Meyerholz DK, Stoltz DA, McCray PB Jr (2016) Newborn cystic fibrosis pigs have a blunted early response to an inflammatory stimulus. *Am J Respir Crit Care Med* 194:845–854
104. Linsdell P, Hanrahan JW (1998) Glutathione permeability of CFTR. *Am J Physiol* 275:C323–C326
105. Moskwa P, Lorentzen D, Excoffon KJ, Zabner J, McCray PB Jr, Nauseef WM, Dupuy C, Banfi B (2007) A novel host defense system of airways is defective in cystic fibrosis. *Am J Respir Crit Care Med* 175:174–183
106. Roud JH, Buhl R, McElvaney NG, Borok Z, Crystal RG (1993) Systemic deficiency of glutathione in cystic fibrosis. *J Appl Physiol* (1985) 75:2419–2424
107. Pedemonte N, Caci E, Sondo E, Caputo A, Rhoden K, Pfeiffer U, Di Candia M, Bandettini R, Ravazzolo R, Zegar-Moran O, Galletta LJ (2007) Thiocyanate transport in resting and IL-4-stimulated human bronchial epithelial cells: role of pendrin and anion channels. *J Immunol* 178:5144–5153
108. Xu Y, Szepe S, Lu Z (2009) The antioxidant role of thiocyanate in the pathogenesis of cystic fibrosis and other inflammation-related diseases. *Proc Natl Acad Sci USA* 106:20515–20519
109. Escotte S, Catusse C, Coraux C, Puchelle E (2004) Reconstitution of human airway tissue in the humanized xenograft model. *J Cyst Fibros* 3(Suppl 2):63–65
110. Crespin S, Bacchetta M, Bou Saab J, Tantilipikorn P, Bellec J, Dudez T, Nguyen TH, Kwak BR, Lacroix JS, Huang S, Wisniewski L, Chanson M (2014) Cx26 regulates proliferation of repairing basal airway epithelial cells. *Int J Biochem Cell Biol* 52:152–160
111. Hajj R, Lesimple P, Nawrocki-Raby B, Birembaut P, Puchelle E, Coraux C (2007) Human airway surface epithelial regeneration is delayed and abnormal in cystic fibrosis. *J Pathol* 211:340–350
112. Trinh NT, Bardou O, Prive A, Maille E, Adam D, Lingee S, Ferraro P, Desrosiers MY, Coraux C, Brochiero E (2012) Improvement of defective cystic fibrosis airway epithelial wound repair after CFTR rescue. *Eur Respir J* 40:1390–1400
113. Argent BE, Gray MA, Steward MC, Case RM (2012) Cell physiology of pancreatic ducts. In: *Physiology of the gastrointestinal tract*, Elsevier Inc, pp 1399–1424
114. Crawford I, Maloney PC, Zeitlin PL, Guggino WB, Hyde SC, Turley H, Gatter KC, Harris A, Higgins CF (1991) Immunocytochemical localization of the cystic fibrosis gene product CFTR. *Proc Natl Acad Sci USA* 88:9262–9266
115. Marino CR, Matovic LM, Gorelick FS, Cohn JA (1991) Localization of the cystic fibrosis transmembrane conductance regulator in pancreas. *J Clin Invest* 88:712–716
116. Durie PR (1989) The pathophysiology of the pancreatic defect in cystic fibrosis. *Acta Paediatr Scand Suppl* 363:41–44
117. Kopelman H, Durie P, Gaskin K, Weizman Z, Forstner G (1985) Pancreatic fluid secretion and protein hyperconcentration in cystic fibrosis. *N Engl J Med* 312:329–334
118. Novak I, Haanes KA, Wang J (2013) Acid-base transport in pancreas—new challenges. *Front Physiol* 4:380
119. Williams JA, Yule, David I (2012) Stimulus-secretion coupling in pancreatic acinar cells. In: *Physiology of the gastrointestinal tract*. Elsevier Inc, pp 1361–1398
120. Behrendorff N, Floetenmeyer M, Schwiening C, Thorn P (2010) Protons released during pancreatic acinar cell secretion acidify the lumen and contribute to pancreatitis in mice. *Gastroenterology* 139:1711–1720
121. Steward MC, Ishiguro H, Case RM (2005) Mechanisms of bicarbonate secretion in the pancreatic duct. *Annu Rev Physiol* 67:377–409
122. Freedman SD, Kern HF, Scheele GA (2001) Pancreatic acinar cell dysfunction in CFTR(−/−) mice is associated with impairments in luminal pH and endocytosis. *Gastroenterology* 121:950–957
123. Bhoomagoud M, Jung T, Atladottir J, Kolodczek TR, Shugrue C, Chaudhuri A, Thrower EC, Gorelick FS (2009) Reducing extracellular pH sensitizes the acinar cell to secretagogue-induced pancreatitis responses in rats. *Gastroenterology* 141:2228–2239
124. Pallagi P, Venglovecz V, Rakonczay Z Jr, Borka K, Korompay A, Ozsvari B, Judak L, Sahin-Toth M, Geisz A, Schnur A, Maleth J, Takacs T, Gray MA, Argent BE, Mayerle J, Lerch MM, Wittmann T, Hegyi P (2011) Trypsin reduces pancreatic ductal bicarbonate secretion by inhibiting CFTR Cl(−) channels and luminal anion exchangers. *Gastroenterology* 141:2228–2239
125. Hegyi P, Wilschanski M, Muallem S, Lukacs GL, Sahin-Toth M, Uc A, Gray MA, Rakonczay Z Jr, Maleth J (2016) CFTR: a new horizon in the pathomechanism and treatment of pancreatitis. *Rev Physiol Biochem Pharmacol* 170:37–66
126. Gray MA, Greenwell JR, Argent BE (1988) Secretin-regulated chloride channel on the apical plasma membrane of pancreatic duct cells. *J Membr Biol* 105:131–142
127. Gray MA, Harris A, Coleman L, Greenwell JR, Argent BE (1989) Two types of chloride channel on duct cells cultured from human fetal pancreas. *Am J Physiol* 257:C240–C251
128. Gray MA, Plant S, Argent BE (1993) cAMP-regulated whole cell chloride currents in pancreatic duct cells. *Am J Physiol* 264:C591–C602
129. Gray MA, Pollard CE, Harris A, Coleman L, Greenwell JR, Argent BE (1990) Anion selectivity and block of the small-conductance chloride channel on pancreatic duct cells. *Am J Physiol* 259:C752–C761
130. Novak I, Greger R (1988) Properties of the luminal membrane of isolated perfused rat pancreatic ducts. Effect of cyclic AMP and blockers of chloride transport. *Pflugers Arch* 411:546–553
131. Novak I, Greger R (1988) Electrophysiological study of transport systems in isolated perfused pancreatic ducts: properties of the basolateral membrane. *Pflugers Arch* 411:58–68
132. Stuenkel EL, Machen TE, Williams JA (1988) pH regulatory mechanisms in rat pancreatic ductal cells. *Am J Physiol* 254:G925–G930
133. Villanger O, Veil T, Raeder MG (1995) Secretin causes H⁺/HCO₃[−] secretion from pig pancreatic ductules by vacuolar-type H⁺(+)-adenosine triphosphatase. *Gastroenterology* 108:850–859
134. Sohma Y, Gray MA, Imai Y, Argent BE (1996) A mathematical model of the pancreatic ductal epithelium. *J Membr Biol* 154:53–67
135. Sohma Y, Gray MA, Imai Y, Argent BE (2000) HCO₃[−] transport in a mathematical model of the pancreatic ductal epithelium. *J Membr Biol* 176:77–100
136. Lee MG, Ohana E, Park HW, Yang D, Muallem S (2012) Molecular mechanism of pancreatic and salivary gland fluid and HCO₃[−] secretion. *Physiol Rev* 92:39–74
137. Stewart AK, Shmukler BE, Vandorpe DH, Reimold F, Henehan JF, Nakakuki M, Akhavein A, Ko S, Ishiguro H, Alper SL (2011) SLC26 anion exchangers of guinea pig pancreatic duct: molecular cloning and functional characterization. *Am J Physiol Cell Physiol* 301:C289–C303
138. Stewart AK, Yamamoto A, Nakakuki M, Kondo T, Alper SL, Ishiguro H (2009) Functional coupling of apical Cl[−]/HCO₃[−]

- exchange with CFTR in stimulated HCO_3^- secretion by guinea pig interlobular pancreatic duct. *Am J Physiol Gastrointest Liver Physiol* 296:G1307–G1317
139. Shcheynikov N, Wang Y, Park M, Ko SB, Dorwart M, Naruse S, Thomas PJ, Muallem S (2006) Coupling modes and stoichiometry of $\text{Cl}^-/\text{HCO}_3^-$ exchange by *slc26a3* and *slc26a6*. *J Gen Physiol* 127:511–524
 140. Ishiguro H, Namkung W, Yamamoto A, Wang Z, Worrell RT, Xu J, Lee MG, Soleimani M (2007) Effect of *Slc26a6* deletion on apical $\text{Cl}^-/\text{HCO}_3^-$ exchanger activity and cAMP-stimulated bicarbonate secretion in pancreatic duct. *Am J Physiol Gastrointest Liver Physiol* 292:G447–G455
 141. Gray MA (2004) Bicarbonate secretion: it takes two to tango. *Nat Cell Biol* 6:292–294
 142. Ko SB, Shcheynikov N, Choi JY, Luo X, Ishibashi K, Thomas PJ, Kim JY, Kim KH, Lee MG, Naruse S, Muallem S (2002) A molecular mechanism for aberrant CFTR-dependent HCO_3^- transport in cystic fibrosis. *EMBO J* 21:5662–5672
 143. Ko SB, Zeng W, Dorwart MR, Luo X, Kim KH, Millen L, Goto H, Naruse S, Soyombo A, Thomas PJ, Muallem S (2004) Gating of CFTR by the STAS domain of SLC26 transporters. *Nat Cell Biol* 6:343–350
 144. Wang Y, Soyombo AA, Shcheynikov N, Zeng W, Dorwart M, Marino CR, Thomas PJ, Muallem S (2006) *Slc26a6* regulates CFTR activity in vivo to determine pancreatic duct HCO_3^- secretion: relevance to cystic fibrosis. *EMBO J* 25:5049–5057
 145. Lohi H, Lamprecht G, Markovich D, Heil A, Kujala M, Seidler U, Kere J (2003) Isoforms of SLC26A6 mediate anion transport and have functional PDZ interaction domains. *Am J Physiol Cell Physiol* 284:C769–C779
 146. Ishiguro H, Steward MC, Sohma Y, Kubota T, Kitagawa M, Kondo T, Case RM, Hayakawa T, Naruse S (2002) Membrane potential and bicarbonate secretion in isolated interlobular ducts from guinea-pig pancreas. *J Gen Physiol* 120:617–628
 147. Ishiguro H, Naruse S, Kitagawa M, Mabuchi T, Kondo T, Hayakawa T, Case RM, Steward MC (2002) Chloride transport in microperfused interlobular ducts isolated from guinea-pig pancreas. *J Physiol* 539:175–189
 148. Park HW, Nam JH, Kim JY, Namkung W, Yoon JS, Lee JS, Kim KS, Venglovecz V, Gray MA, Kim KH, Lee MG (2010) Dynamic regulation of CFTR bicarbonate permeability by $[\text{Cl}^-]_i$ and its role in pancreatic bicarbonate secretion. *Gastroenterology* 139:620–631
 149. Shcheynikov N, Kim KH, Kim KM, Dorwart MR, Ko SB, Goto H, Naruse S, Thomas PJ, Muallem S (2004) Dynamic control of cystic fibrosis transmembrane conductance regulator $\text{Cl}^-/\text{HCO}_3^-$ selectivity by external Cl^- . *J Biol Chem* 279:21857–21865
 150. Ishiguro H, Steward MC, Naruse S, Ko SB, Goto H, Case RM, Kondo T, Yamamoto A (2009) CFTR functions as a bicarbonate channel in pancreatic duct cells. *J Gen Physiol* 133:315–326
 151. Ishiguro H, Naruse S, Steward MC, Kitagawa M, Ko SB, Hayakawa T, Case RM (1998) Fluid secretion in interlobular ducts isolated from guinea-pig pancreas. *J Physiol* 511(Pt 2):407–422
 152. Satoh H, Moriyama N, Hara C, Yamada H, Horita S, Kunimi M, Tsukamoto K, Iso ON, Inatomi J, Kawakami H, Kudo A, Endou H, Igarashi T, Goto A, Fujita T, Seki G (2003) Localization of $\text{Na}^+/\text{HCO}_3^-$ cotransporter (NBC-1) variants in rat and human pancreas. *Am J Physiol Cell Physiol* 284:C729–C737
 153. Ishiguro H, Steward MC, Lindsay AR, Case RM (1996) Accumulation of intracellular HCO_3^- by $\text{Na}^+/\text{HCO}_3^-$ cotransport in interlobular ducts from guinea-pig pancreas. *J Physiol* 495(Pt 1):169–178
 154. Ishiguro H, Steward MC, Wilson RW, Case RM (1996) Bicarbonate secretion in interlobular ducts from guinea-pig pancreas. *J Physiol* 495(Pt 1):179–191
 155. Venglovecz V, Rakonczay Z Jr, Gray MA, Hegyi P (2015) Potassium channels in pancreatic duct epithelial cells: their role, function and pathophysiological relevance. *Pflugers Arch* 467:625–640
 156. Yang D, Shcheynikov N, Zeng W, Ohana E, So I, Ando H, Mizutani A, Mikoshiba K, Muallem S (2009) IRBIT coordinates epithelial fluid and HCO_3^- secretion by stimulating the transporters pNBC1 and CFTR in the murine pancreatic duct. *J Clin Invest* 119:193–202
 157. Park S, Shcheynikov N, Hong JH, Zheng C, Suh SH, Kawaai K, Ando H, Mizutani A, Abe T, Kiyonari H, Seki G, Yule D, Mikoshiba K, Muallem S (2013) Irbit mediates synergy between Ca^{2+} and cAMP signaling pathways during epithelial transport in mice. *Gastroenterology* 145:232–241
 158. Yang D, Li Q, So I, Huang CL, Ando H, Mizutani A, Seki G, Mikoshiba K, Thomas PJ, Muallem S (2011) IRBIT governs epithelial secretion in mice by antagonizing the WNK/SPAK kinase pathway. *J Clin Invest* 121:956–965
 159. Hegyi P, Gray MA, Argent BE (2003) Substance P inhibits bicarbonate secretion from guinea pig pancreatic ducts by modulating an anion exchanger. *Am J Physiol Cell Physiol* 285:C268–C276
 160. Hegyi P, Rakonczay Z Jr, Tiszlavicz L, Varro A, Toth A, Racz G, Varga G, Gray MA, Argent BE (2005) Protein kinase C mediates the inhibitory effect of substance P on HCO_3^- secretion from guinea pig pancreatic ducts. *Am J Physiol Cell Physiol* 288:C1030–C1041
 161. Kemeny LV, Hegyi P, Rakonczay Z Jr, Borka K, Korompay A, Gray MA, Argent BE, Venglovecz V (2011) Substance P inhibits pancreatic ductal bicarbonate secretion via neurokinin receptors 2 and 3 in the guinea pig exocrine pancreas. *Pancreas* 40:793–795
 162. Durie PR (1998) Pancreatitis and mutations of the cystic fibrosis gene. *N Engl J Med* 339:687–688
 163. Scheele GA, Fukuoka SI, Kern HF, Freedman SD (1996) Pancreatic dysfunction in cystic fibrosis occurs as a result of impairments in luminal pH, apical trafficking of zymogen granule membranes, and solubilization of secretory enzymes. *Pancreas* 12:1–9
 164. De Boeck K, Weren M, Proesmans M, Kerem E (2005) Pancreatitis among patients with cystic fibrosis: correlation with pancreatic status and genotype. *Pediatrics* 115:e463–e469
 165. Durno C, Corey M, Zielenski J, Tullis E, Tsui LC, Durie P (2002) Genotype and phenotype correlations in patients with cystic fibrosis and pancreatitis. *Gastroenterology* 123:1857–1864
 166. Zielenski J (2000) Genotype and phenotype in cystic fibrosis. *Respiration* 67:117–133
 167. Lee MG, Wigley WC, Zeng W, Noel LE, Marino CR, Thomas PJ, Muallem S (1999) Regulation of $\text{Cl}^-/\text{HCO}_3^-$ exchange by cystic fibrosis transmembrane conductance regulator expressed in NIH 3T3 and HEK 293 cells. *J Biol Chem* 274:3414–3421
 168. Choi JY, Muallem D, Kiselyov K, Lee MG, Thomas PJ, Muallem S (2001) Aberrant CFTR-dependent HCO_3^- transport in mutations associated with cystic fibrosis. *Nature* 410:94–97
 169. Rakonczay Z Jr, Hegyi P, Hasegawa M, Inoue M, You J, Iida A, Ignath I, Alton EW, Griesenbach U, Ovari G, Vag J, Da Paula AC, Crawford RM, Varga G, Amaral MD, Mehta A, Lonovics J, Argent BE, Gray MA (2008) CFTR gene transfer to human cystic fibrosis pancreatic duct cells using a Sendai virus vector. *J Cell Physiol* 214:442–455
 170. Ignath I, Hegyi P, Venglovecz V, Szekely CA, Carr G, Hasegawa M, Inoue M, Takacs T, Argent BE, Gray MA, Rakonczay Z Jr (2009) CFTR expression but not Cl^- transport is involved in the stimulatory effect of bile acids on apical $\text{Cl}^-/\text{HCO}_3^-$ exchange activity in human pancreatic duct cells. *Pancreas* 38:921–929

171. Venglovecz V, Hegyi P, Rakonczay Z Jr, Tiszlavicz L, Nardi A, Grunnet M, Gray MA (2011) Pathophysiological relevance of apical large-conductance $\text{Ca}(2)+$ -activated potassium channels in pancreatic duct epithelial cells. *Gut* 60:361–369
172. Venglovecz V, Rakonczay Z Jr, Ozsvari B, Takacs T, Lonovics J, Varro A, Gray MA, Argent BE, Hegyi P (2008) Effects of bile acids on pancreatic ductal bicarbonate secretion in guinea pig. *Gut* 57:1102–1112
173. Judak L, Hegyi P, Rakonczay Z Jr, Maleth J, Gray MA, Venglovecz V (2014) Ethanol and its non-oxidative metabolites profoundly inhibit CFTR function in pancreatic epithelial cells which is prevented by ATP supplementation. *Pflugers Arch* 466:549–562
174. Maleth J, Venglovecz V, Razga Z, Tiszlavicz L, Rakonczay Z Jr, Hegyi P (2011) Non-conjugated chenodeoxycholate induces severe mitochondrial damage and inhibits bicarbonate transport in pancreatic duct cells. *Gut* 60:136–138
175. Ko SB, Mizuno N, Yatabe Y, Yoshikawa T, Ishiguro H, Yamamoto A, Azuma S, Naruse S, Yamao K, Muallem S, Goto H (2010) Corticosteroids correct aberrant CFTR localization in the duct and regenerate acinar cells in autoimmune pancreatitis. *Gastroenterology* 138:1988–1996
176. Maleth J, Balazs A, Pallagi P, Balla Z, Kui B, Katona M, Judak L, Nemeth I, Kemeny LV, Rakonczay Z Jr, Venglovecz V, Foldesi I, Peto Z, Somoracz A, Borka K, Perdomo D, Lukacs GL, Gray MA, Monterisi S, Zaccolo M, Sendler M, Mayerle J, Kuhn JP, Lerch MM, Sahin-Toth M, Hegyi P (2015) Alcohol disrupts levels and function of the cystic fibrosis transmembrane conductance regulator to promote development of pancreatitis. *Gastroenterology* 148(427–439):e416
177. LaRusch J, Jung J, General JJ, Lewis MD, Park HW, Brand RE, Gelrud A, Anderson MA, Banks PA, Conwell D, Lawrence C, Romagnuolo J, Baillie J, Alkaade S, Cote G, Gardner TB, Amann ST, Slivka A, Sandhu B, Aloe A, Kienholz ML, Yadav D, Barmada MM, Bahar I, Lee MG, Whitcomb DC, North American Pancreatitis Study, G (2014) Mechanisms of CFTR functional variants that impair regulated bicarbonate permeation and increase risk for pancreatitis but not for cystic fibrosis. *PLoS Genet* 10:e1004376
178. Masson E, Chen JM, Audrezet MP, Cooper DN, Ferec C (2013) A conservative assessment of the major genetic causes of idiopathic chronic pancreatitis: data from a comprehensive analysis of PRSS1, SPINK1, CTFR and CFTR genes in 253 young French patients. *PLoS One* 8:e73522
179. Ooi CY, Durie PR (2012) Cystic fibrosis transmembrane conductance regulator (CFTR) gene mutations in pancreatitis. *J Cyst Fibros* 11:355–362
180. Davies JC, Cunningham S, Harris WT, Lapey A, Regelman WE, Sawicki GS, Southern KW, Robertson S, Green Y, Cooke J, Rosenfeld M, Group KS (2016) Safety, pharmacokinetics, and pharmacodynamics of ivacaftor in patients aged 2–5 years with cystic fibrosis and a CFTR gating mutation (KIWI): an open-label, single-arm study. *Lancet Respir Med* 4:107–115
181. Miller MR, Soave D, Li W, Gong J, Pace RG, Boelle PY, Cutting GR, Drumm ML, Knowles MR, Sun L, Rommens JM, Accurso F, Durie PR, Corvol H, Levy H, Sontag MK, Strug LJ (2015) Variants in solute carrier SLC26A9 modify prenatal exocrine pancreatic damage in cystic fibrosis. *J Pediatr* 166(1152–1157):e1156
182. Pallagi P, Balla Z, Singh AK, Dosa S, Ivanyi B, Kukor Z, Toth A, Riederer B, Liu Y, Engelhardt R, Jarmay K, Szabo A, Janovszky A, Perides G, Venglovecz V, Maleth J, Wittmann T, Takacs T, Gray MA, Gacser A, Hegyi P, Seidler U, Rakonczay Z Jr (2014) The role of pancreatic ductal secretion in protection against acute pancreatitis in mice*. *Crit Care Med* 42:e177–e188
183. Hegyi P, Petersen OH (2013) The exocrine pancreas: the acinar-ductal tango in physiology and pathophysiology. *Rev Physiol Biochem Pharmacol* 165:1–30
184. Bakker OJ, van Brunschot S, van Santvoort HC, Besselink MG, Bollen TL, Boermeester MA, Dejong CH, van Goor H, Bosscha K, Ahmed Ali U, Bouwense S, van Grevenstein WM, Heisterkamp J, Houdijk AP, Jansen JM, Karsten TM, Manusama ER, Nieuwenhuijs VB, Schaapherder AF, van der Schelling GP, Schwartz MP, Spanier BW, Tan A, Vecht J, Weusten BL, Witteman BJ, Akkermans LM, Bruno MJ, Dijkgraaf MG, van Ramshorst B, Gooszen HG, Dutch Pancreatitis Study, G (2014) Early versus on-demand nasoenteric tube feeding in acute pancreatitis. *N Engl J Med* 371:1983–1993
185. Bovell D (2015) The human eccrine sweat gland: structure, function and disorders. *J Local Global Health Sci* 5
186. Cui CY, Schlessinger D (2015) Eccrine sweat gland development and sweat secretion. *Exp Dermatol* 24:644–650
187. Murakami M, Ohtake T, Dorschner RA, Schitteck B, Garbe C, Gallo RL (2002) Cathelicidin anti-microbial peptide expression in sweat, an innate defense system for the skin. *J Invest Dermatol* 119:1090–1095
188. Park JH, Park GT, Cho IH, Sim SM, Yang JM, Lee DY (2011) An antimicrobial protein, lactoferrin exists in the sweat: proteomic analysis of sweat. *Exp Dermatol* 20:369–371
189. Rieg S, Garbe C, Sauer B, Kalbacher H, Schitteck B (2004) Dermcidin is constitutively produced by eccrine sweat glands and is not induced in epidermal cells under inflammatory skin conditions. *Br J Dermatol* 151:534–539
190. Peng Y, Cui X, Liu Y, Li Y, Liu J, Cheng B (2014) Systematic review focusing on the excretion and protection roles of sweat in the skin. *Dermatology* 228:115–120
191. Rittie L, Sachs DL, Orringer JS, Voorhees JJ, Fisher GJ (2013) Eccrine sweat glands are major contributors to reepithelialization of human wounds. *Am J Pathol* 182:163–171
192. Di Sant'Agnes PA, Darling RC, Perera GA, Shea E (1953) Abnormal electrolyte composition of sweat in cystic fibrosis of the pancreas; clinical significance and relationship to the disease. *Pediatrics* 12:549–563
193. Quinton PM (2007) Cystic fibrosis: lessons from the sweat gland. *Physiology (Bethesda)* 22:212–225
194. Cui CY, Childress V, Piao Y, Michel M, Johnson AA, Kunisada M, Ko MS, Kaestner KH, Marmorstein AD, Schlessinger D (2012) Forkhead transcription factor FoxA1 regulates sweat secretion through Bestrophin 2 anion channel and Na-K-Cl cotransporter 1. *Proc Natl Acad Sci USA* 109:1199–1203
195. Robertson J, Bovell D (2014) Pharmacological blockers of STIM1 inhibit increases in intracellular calcium in horse sweat gland cells. *FASEB J* 28:650.2
196. Bovell DL, MacDonald A, Meyer BA, Corbett AD, MacLaren WM, Holmes SL, Harker M (2011) The secretory clear cell of the eccrine sweat gland as the probable source of excess sweat production in hyperhidrosis. *Exp Dermatol* 20:1017–1020
197. Strutton DR, Kowalski JW, Glaser DA, Stang PE (2004) US prevalence of hyperhidrosis and impact on individuals with axillary hyperhidrosis: results from a national survey. *J Am Acad Dermatol* 51:241–248
198. Rieg S, Steffen H, Seeber S, Humeny A, Kalbacher H, Dietz K, Garbe C, Schitteck B (2005) Deficiency of dermcidin-derived antimicrobial peptides in sweat of patients with atopic dermatitis correlates with an impaired innate defense of human skin in vivo. *J Immunol* 174:8003–8010
199. Lu C, Fuchs E (2014) Sweat gland progenitors in development, homeostasis, and wound repair. *Cold Spring Harb Perspect Med* 4:a015222
200. Lu CP, Polak L, Rocha AS, Pasolli HA, Chen SC, Sharma N, Blanpain C, Fuchs E (2012) Identification of stem cell populations in sweat glands and ducts reveals roles in homeostasis and wound repair. *Cell* 150:136–150

201. Hashimoto K (1978) The eccrine gland. In: Jarrett A (ed) The physiology and pathophysiology of the skin. Academic Press Ltd, London
202. Hibbs RG (1958) The fine structure of human eccrine sweat glands. *Am J Anat* 103:201–217
203. Reddy MM, Bell CL, Quinton PM (1992) Evidence of two distinct epithelial cell types in primary cultures from human sweat gland secretory coil. *Am J Physiol* 262:C891–C898
204. Sato F, Sato K (2000) cAMP-dependent Cl(–) channel protein (CFTR) and its mRNA are expressed in the secretory portion of human eccrine sweat gland. *J Histochem Cytochem* 48:345–354
205. Ertongur-Fauth T, Hochheimer A, Buescher JM, Rapprich S, Krohn M (2014) A novel TMEM16A splice variant lacking the dimerization domain contributes to calcium-activated chloride secretion in human sweat gland epithelial cells. *Exp Dermatol* 23:825–831
206. Sato K, Nishiyama A, Kobayashi M (1979) Mechanical properties and functions of the myoepithelium in the eccrine sweat gland. *Am J Physiol* 237:C177–C184
207. Cui CY, Sima J, Yin M, Michel M, Kunisada M, Schlessinger D (2016) Identification of potassium and chloride channels in eccrine sweat glands. *J Dermatol Sci* 81:129–131
208. Reddy MM, Quinton PM (1989) Localization of Cl[–] conductance in normal and Cl[–] impermeability in cystic fibrosis sweat duct epithelium. *Am J Physiol* 257:C727–C735
209. Reddy MM, Quinton PM (2009) PKA mediates constitutive activation of CFTR in human sweat duct. *J Membr Biol* 231:65–78
210. Reddy MM, Light MJ, Quinton PM (1999) Activation of the epithelial Na⁺ channel (ENaC) requires CFTR Cl[–] channel function. *Nature* 402:301–304
211. Reddy MM, Quinton PM (2003) Functional interaction of CFTR and ENaC in sweat glands. *Pflugers Arch* 445:499–503
212. Reddy MM, Wang XF, Quinton PM (2008) Effect of cytosolic pH on epithelial Na⁺ channel in normal and cystic fibrosis sweat ducts. *J Membr Biol* 225:1–11
213. Bijman J, Quinton P (1987) Permeability properties of cell membranes and tight junctions of normal and cystic fibrosis sweat ducts. *Pflugers Arch* 408:505–510
214. Reddy MM, Quinton PM (1994) Intracellular Cl activity: evidence of dual mechanisms of cl absorption in sweat duct. *Am J Physiol* 267:C1136–C1144
215. Quinton PM, Reddy MM (1989) Cl[–] conductance and acid secretion in the human sweat duct. *Ann N Y Acad Sci* 574:438–446
216. Bovell DL, Clunes MT, Roussa E, Burry J, Elder HY (2000) Vacuolar-type H⁺-ATPase distribution in unstimulated and acetylcholine-activated isolated human eccrine sweat glands. *Histochem J* 32:409–413
217. Clunes MT, Lindsay SL, Roussa E, Quinton PM, Bovell DL (2004) Localisation of the vacuolar proton pump (V-H⁺-ATPase) and carbonic anhydrase II in the human eccrine sweat gland. *J Mol Histol* 35:339–345
218. Sato K, Sato F (1984) Defective beta adrenergic response of cystic fibrosis sweat glands in vivo and in vitro. *J Clin Invest* 73:1763–1771
219. Kessler WR, Andersen DH (1951) Heat prostration in fibrocystic disease of the pancreas and other conditions. *Pediatrics* 8:648–656
220. Gibson LE, Cooke RE (1959) A test for concentration of electrolytes in sweat in cystic fibrosis of the pancreas utilizing pilocarpine by iontophoresis. *Pediatrics* 23:545–549
221. Quinton PM (2014) A synopsis of methods of sweat tests in pathology. *Clin Biochem* 47:757–758
222. Quinton PM, Bijman J (1983) Higher bioelectric potentials due to decreased chloride absorption in the sweat glands of patients with cystic fibrosis. *N Engl J Med* 308:1185–1189
223. Cook DP, Rector MV, Bouzek DC, Michalski AS, Gansemer ND, Reznikov LR, Li X, Stroik MR, Ostedgaard LS, Abou Alaiwa MH, Thompson MA, Prakash YS, Krishnan R, Meyerholz DK, Seow CY, Stoltz DA (2016) Cystic fibrosis transmembrane conductance regulator in sarcoplasmic reticulum of airway smooth muscle. Implications for airway contractility. *Am J Respir Crit Care Med* 193:417–426
224. Brodlie M, Haq IJ, Roberts K, Elborn JS (2015) Targeted therapies to improve CFTR function in cystic fibrosis. *Genome Med* 7:101