



HAL
open science

Candida albicans: An emerging yeast model to study eukaryotic genome plasticity

Mélanie Legrand, Priya Jaitly, Adeline Feri, Christophe d'Enfert, Kaustuv Sanyal

► **To cite this version:**

Mélanie Legrand, Priya Jaitly, Adeline Feri, Christophe d'Enfert, Kaustuv Sanyal. *Candida albicans: An emerging yeast model to study eukaryotic genome plasticity*. Trends in Genetics, 2019, 35 (4), pp.292-307. 10.1016/j.tig.2019.01.005 . hal-02628092

HAL Id: hal-02628092

<https://hal.inrae.fr/hal-02628092v1>

Submitted on 28 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

***Candida albicans*: An Emerging Yeast Model to Study Eukaryotic Genome Plasticity**

Mélanie Legrand,¹ Priya Jaitly,² Adeline Feri,^{1,3,4} Christophe d'Enfert,^{1,*} and Kaustuv Sanyal^{2,*}

Saccharomyces cerevisiae and *Schizosaccharomyces pombe* have served as uncontested unicellular model organisms, as major discoveries made in the field of genome biology using yeast genetics have proved to be relevant from yeast to humans. The yeast *Candida albicans* has attracted much attention because of its ability to switch between a harmless commensal and a dreaded human pathogen. *C. albicans* bears unique features regarding its life cycle, genome structure, and dynamics, and their links to cell biology and adaptation to environmental challenges. Examples include a unique reproduction cycle with haploid, diploid, and tetraploid forms; a distinctive organisation of chromosome hallmarks; a highly dynamic genome, with extensive karyotypic variations, including aneuploidies, isochromosome formation, and loss-of-heterozygosity; and distinctive links between the response to DNA alterations and cell morphology. These features have made *C. albicans* emerge as a new and attractive unicellular model to study genome biology and dynamics in eukaryotes.

***Candida albicans*: A Model for Studying Genome Biology**

Maintenance of genome integrity and the accuracy of DNA replication are at the core of cell function, survival, and propagation. Thus, deciphering the molecular mechanisms that underlie genome biology is of crucial importance, especially as they have relevance in numerous human diseases such as cancer but also because they underlie species evolution and adaptation. For more than 50 years, *Saccharomyces cerevisiae* (see Glossary) and *Schizosaccharomyces pombe* have served as uncontested yeast models for molecular understanding of the processes underlying eukaryotic genome biology [1–6]. Major discoveries in this field have benefited from the ‘awesome power of yeast genetics’ and proven to be relevant across eukaryotes. In addition, the ascomycetous yeast *Candida albicans* – a distantly related cousin of *S. cerevisiae* (Box 1) – has attracted considerable interest because of its dominant importance as a human pathogen. While normally a commensal of humans, *C. albicans* is also responsible for superficial infections – thrush, oropharyngeal candidiasis, vaginal candidiasis – in healthy individuals as well as disseminated infections in hospitalised patients that receive broad-spectrum antibiotic treatment and have debilitated immunity [7]. The investigation of putative *C. albicans* virulence factors, in particular the ability to alternate between yeast and filamentous forms; the exploration of *C. albicans* interactions with the host; and the search for new antifungal targets have been accompanied by the development of a molecular toolkit that allows gene function to be accurately characterised in this species (Box 1) [8–16]. Notably, this toolkit has enabled exploring other aspects in depth, especially the genome biology of *C. albicans*. A number of features distinguish *C. albicans* from other yeast species – a unique reproduction cycle where haploid, diploid, and tetraploid forms are observed; a distinctive organisation of chromosome hallmarks; a highly dynamic genome, with extensive karyotypic

Highlights

C. albicans is a major human pathogen, with features that distinguish it from other yeast species.

Rather than meiosis, ploidy reduction upon mating occurs by a parasexual process of concerted chromosome loss.

Centromeres have unique organisation and are epigenetically regulated. Neither centromeres nor the DNA replication origins share any common defining DNA sequence.

Loss-of-heterozygosity events are frequent and widespread in the genome of *C. albicans*.

The DNA damage response is coupled to morphogenetic shifts.

C. albicans emerges as a new unicellular model to study eukaryotic genome biology and dynamics.

¹Fungal Biology and Pathogenicity Unit, Department of Mycology, Institut Pasteur, INRA, Paris, France

²Molecular Mycology Laboratory, Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore, India

³Université Paris Diderot, Sorbonne Paris Cité, Cellule Pasteur, Paris, France

⁴Current address: Pathoquest, BioPark, 11 rue Watt, 75013 Paris, France

*Correspondence: christophe.denfert@pasteur.fr (C. d'Enfert) and sanyal@jncasr.ac.in (K. Sanyal).

variations, including chromosomal loss and gain, as well as rearrangements, **isochromosome** formation, segmental duplication, and loss-of-heterozygosity (LOH); and distinctive links between the response to DNA alterations and cell morphology (Figure 1, Key Figure). In this review, we highlight these unique features that have helped *C. albicans* emerge as a new unicellular model to study genome biology and its evolution.

***Candida albicans*: A Yeast Species with a Cryptic Reproduction Cycle**

Sequencing of the *C. albicans* genome has revealed that this diploid species, long thought to be devoid of sex [17–20] is actually equipped with the majority of the genetic circuit required for sexual reproduction. Similar to mating type (**MAT**) loci in *S. cerevisiae*, mating type-like (**MTL**; Figure 2A) regions encoding the transcriptional regulators that control expression of mating specific genes [21] and elements of a functional pheromone response pathway exist in the *C. albicans* genome [22]. While many of the meiotic regulators from *S. cerevisiae* have counterparts in *C. albicans*, the latter lacks key meiotic components including **IME1**, the master switch for entry into the meiotic pathway in *S. cerevisiae*, and **SPO13**, which in *S. cerevisiae* is essential for proper execution of **meiosis I**. Existence of functional analogues of these genes in *C. albicans* is difficult to rule out as extensive sequence divergence may cause difficulty in their *in silico* identification. While **IME1** is essential for meiosis in a number of yeast species, strikingly, this gene is absent from many sexual species in the CTG clade, suggesting a possible clade-specific rewiring of the meiotic cycle (Box 1) [23]. In addition, the highly conserved pheromone response pathways were detected in nonmating species [23]. The apparent plasticity of mating and meiosis pathways in *Candida* species reinforces the fact that knowledge of gene products involved in sexual reproduction is insufficient to accurately predict the reproductive behaviour of an organism.

Hints from *in silico* data have fuelled the quest for evidence of mating and sexual reproduction in *C. albicans*. While engineered *C. albicans* strains homozygous for **MTL loci** are able to mate and form tetraploids both in laboratory conditions and in a mammalian host [24,25], meiosis remains to be demonstrated in *C. albicans*. Tetraploid cells revert to diploidy by undergoing a parasexual process of concerted chromosome loss [26–29]. The combination of mating and subsequent concerted chromosome loss that allows *C. albicans* to alternate between diploid and tetraploid (but also haploid and diploid) has been referred to as a **parasexual cycle** (Figure 2B). The apparent lack of conventional meiosis in *C. albicans* suggests that the function of meiotic genes may have diverged [22,23,30,31]. A classic example is Ndt80, the meiosis-specific transcription factor and a key modulator of progression of meiosis and sporulation in *S. cerevisiae*, that functionally diverged to participate in the biofilm pathway in *C. albicans* [32,33]. Similarly, Ume6, a key transcriptional regulator of early meiosis-specific genes in *S. cerevisiae*, has been rewired towards autophagy and hyphal growth regulation in *C. albicans* [30,34]. The latest example of such rewiring is Rme1; the function of which diverged from preventing meiosis by repressing **IME1** in *S. cerevisiae* to regulating chlamyospore formation in *C. albicans* (Hernandez-Cervantes *et al.*, personal communication).

Although we cannot rule out the possibility of *C. albicans* undergoing meiosis in conditions that have not been explored thus far, the hypothetical absence of meiosis does not prevent *C. albicans* from generating genetic and phenotypic diversity necessary for this opportunistic human pathogen to adapt to new environments. In addition to extensive shuffling of parental chromosomes resulting in new combinations of homologues, completion of the parasexual cycle often gives rise to aneuploid strains and is accompanied by recombination events between homologous chromosomes [35,36]. Unexpectedly, recombination events during the nonmeiotic parasexual cycle are dependent on the DNA double-strand break (DSB)

Glossary

***Candida albicans*:** this ascomycetous yeast, normally a commensal of humans, is responsible for superficial infections in healthy individuals as well as disseminated infections in immunocompromised patients.

Centromere (CEN): an essential chromosomal element that facilitates sister chromatid separation via kinetochore formation during mitosis.

CRISPR-Cas9: Cas9 is an RNA-guided DNA endonuclease enzyme that associate with the CRISPR (clustered regularly interspaced short palindromic repeats) to target and cut specific sites in genomes. CRISPR-Cas9 can be exploited for genome editing in *C. albicans*.

DNA replication origins (ORIs): genetic elements bound by the origin recognition complex and where DNA replication initiates. In *C. albicans*, ORIs are categorized into arm ORIs and centromere ORIs.

Experimental evolution: use of laboratory experiments to study evolutionary dynamics in controlled conditions imposed by the experimenter.

Gene flow: transfer of genetic variation from one population to another.

Heterozygosity: presence of different alleles at one or more loci on homologous chromosomes in a diploid organism.

Homologous recombination (HR): pathway that repairs DSBs in DNA using a type of genetic recombination in which nucleotide sequences are exchanged between two identical DNA molecules.

Homozygosity bias: when one haplotype or even a part of it is never found in the homozygous state.

IME1: master regulator of meiosis. The **IME1** gene is required for expression of meiosis specific genes and sporulation and in *S. cerevisiae*.

Isochromosome: abnormal chromosome whereby two identical chromosome arms flank a centromere.

Kinetochore: large multiprotein complex that assembles on centromere DNA and serves as the chromosomal attachment site of the spindle microtubules.

inducing **Spo11**; the meiosis-specific endonuclease that initiates meiotic recombination in *S. cerevisiae* [35].

The majority of *C. albicans* isolates are found in the diploid state and diploidy is considered the preferred ploidy state of *C. albicans*. Nevertheless, nondiploid isolates have also been reported [37]. Changes in ploidy including haploidy, tetraploidy, or aneuploidy (primarily monosomy or trisomy) are thought to provide *C. albicans* with a rapid response to changing environments within the host [38]. Deviations from diploidy seem to harbour a fitness cost in the long term, as **experimental evolution** experiments using clinical and laboratory haploid, diploid, and polyploid *C. albicans* strains in complete medium and under nutrient-limited conditions have revealed that the stabilised genome nearly always reaches diploidy, a phenomenon termed as **ploidy drive** [39].

Large genomic changes, similar to the ones observed in products of *in vitro* parasexual genome reduction or long-term evolution experiments, are well tolerated by *C. albicans* and have been associated with acquisition of new phenotypic traits, such as drug resistance. Although population genetics approaches have recently confirmed the predominance of clonal reproduction in the *C. albicans* population [17,40,41], the work by Ropars and colleagues on genomes of 182 *C. albicans* isolates from diverse origins revealed the occurrence of **gene flow** in this population [42] (Figure 2C). These lines of evidence highlight the fact that parasexuality (or possibly sexuality) also occurs in nature and significantly contributes to *C. albicans* genetic and phenotypic diversities.

The *Candida albicans* Genome: An Organisational View

The essential elements of a eukaryotic chromosome – namely **centromeres**, **DNA replication origins**, and **telomeres** – have been identified in *C. albicans* (Figure 3). The centromere (CEN) is an essential chromosomal element that facilitates sister chromatid separation via **kinetochore** formation during mitosis. The CEN DNA sequences of *C. albicans* are 3–5 kb long and are all unique; devoid of any common sequence motif or repeat [43] (Figure 3A). The absence of a CEN-specific DNA sequence and the inability of the exogenously introduced CEN DNA to function as a native CEN suggest epigenetic regulation of CEN identity in *C. albicans* [44]. However, upon CEN deletion, **neocentromeres** are formed efficiently in *C. albicans*, mostly proximal and rarely distal to the native CEN [45–47]. Similar to the native CENs, neocentromeres also cluster in 3D with other functional CENs [47]. Gene conversion (GC) at the CENs can interchange the deleted CEN with the native CEN [45] and possibly explains the low frequency of SNPs across *C. albicans* CENs. Besides CEN clustering, structural integrity of the kinetochore is required for CEN function in *C. albicans*. Depletion of an essential kinetochore protein disrupts the integrity of the kinetochore architecture [48,49] and results in delocalisation and degradation of CENP-A [48] that forms centromeric chromatin.

DNA replication initiates from multiple discrete genetic loci – the DNA replication origins (*ORI*). Based on the location, *ORIs* in *C. albicans* are of two types [50]: (i) arm *ORIs*, which are located on the chromosomal arms; and (ii) centromeric *ORIs*, which are present on [51] or close to the CENs [52] (Figure 3B). While a subset of arm *ORIs* are defined by a 15-bp AC-rich consensus motif and a nucleosome-depleted pattern, centromeric *ORIs* are defined by epigenetic mechanisms [50] and replicate earliest in the genome [51]. The centromeric *ORIs* together with the **homologous recombination** (HR) proteins, Rad51 and Rad52, play a key role in loading CENP-A onto the CENs [52]. While replicating CEN DNA, the moving replication forks from CEN-proximal *ORIs* stall at CEN due to the presence of the kinetochore acting as a physical barrier [52]. The fork stalling accumulates single-stranded DNA that attracts HR proteins Rad51

Loss-of-heterozygosity (LOH):

genetic event resulting in the loss of one of the haplotype information in heterozygous diploid organisms.

MAT loci: mating-type or *MAT* locus harbours genes that control sexual reproduction in fungi.

Major repeat sequences (MRS):

special feature of the *C. albicans* genome, consisting of a long tract of repetitive DNA, which is present on all chromosomes, except chromosome 3.

Meiosis: cell division that reduces the number of chromosomes by half.

MTL loci: mating type-like or *MTL* locus encodes transcription factors responsible for cellular identity in *C. albicans*.

Neocentromere: non-native centromere locus where a functional kinetochore assembles to allow chromosome segregation following disruption or inactivation of the native centromere.

Nondisjunction: failure of homologous chromosomes or sister chromatids to segregate equally in daughter cells during cell division.

Nonhomologous end joining

(NHEJ): pathway that repairs DSBs in DNA by directly ligating the break ends without requirement for a homologous template.

Parasexual cycle: combination of mating and subsequent concerted chromosome loss that allows *C. albicans* to alternate between diploid and tetraploid (but also haploid and diploid).

Ploidy drive: process that brings an organism to its base ploidy level.

Saccharomyces cerevisiae:

commonly known as baker's yeast or brewer's yeast, *S. cerevisiae* is a single-celled eukaryote that operates in a manner similar to a human cell and therefore is used as an important model organism in genetics and molecular biology.



Spo11: endonuclease that initiates meiotic recombination by catalysing the formation of double-strand breaks in DNA.

Telomeres: regions of repetitive nucleotide sequences located at the termini of a eukaryotic chromosome to ensure chromosome end replication and protection from degradation or end-to-end chromosome fusion.

Box 1. *Candida albicans*, an Alternative Yeast Model with an Extended and Adapted Molecular Genetics Toolkit

Because *C. albicans* decodes the CUG codon as serine instead of leucine and is predominantly a diploid, numerous tools have been adapted for genetic engineering of this species, allowing most molecular approaches to be developed in this species from insertional mutagenesis to gene tagging and two-hybrid screens as well as the production of mutant collections. The recent identification of stable haploids allows new approaches to be developed. CRISPR-cas9-based gene editing allows one-step generation of mutants in the diploid background, speeding up their construction [99–112].

Why is *C. albicans* an attractive emerging model organism?

(A) Phylogenetic location	(B) Genomic and genetic resources	(C) Deletion collection library	(D) Overexpression ORFeome
 <p><i>C. albicans</i> belongs to the <i>Saccharomycotina</i>, the ascomycete yeasts. More specifically, <i>C. albicans</i> belongs to the CTG clade, in which the CTG codon is predominantly translated as serine instead of leucine. Molecular reporters (GFP, luciferases, lacZ) have been adapted to this unusual CTG decoding. Of note, the <i>Candida</i> genus is polyphyletic and includes all <i>Saccharomycotina</i> species that have not demonstrated sex.</p> <p style="text-align: right;">References [99,100]</p>	<p>The <i>Candida</i> Genome Database (CGD)</p> <p>A free online database providing access to genomic sequence data and manually curated functional information about genes and proteins of <i>C. albicans</i> since 2004.</p> <p style="text-align: center;">Transformation markers</p> <ul style="list-style-type: none"> • Auxotrophic markers • Drug resistance markers • Recyclable markers <p style="text-align: center;">Controlled promoters</p> <ul style="list-style-type: none"> • Constitutive promoters (P_{TDH3}, P_{ACT1}) • Inducible promoters (P_{TET}, P_{PCK1}, P_{MET3}, P_{MAL2}) <p style="text-align: center;">Transposons</p> <ul style="list-style-type: none"> • Tn5/Tn7 mutagenesis in <i>E. coli</i> followed by recombination in <i>C. albicans</i> • Maize Transposable Elements Ac/Ds adapted to <i>C. albicans</i> <p style="text-align: right;">References [101–104]</p>	<ul style="list-style-type: none"> • Transposon-mediated (in both diploid and stable haploid isolates). The collection of haploid transposon mutants can be assessed using transposon sequencing (TnSeq). • Directed, barcoded deletions in a diploid isolate, allowing implementation of Signature-Tagged Mutagenesis (STM) screens • Induced deletions (obtained following transposon-mediated mutagenesis of a diploid isolate) • The gene replacement and controlled expression (GRACE) collection, whereby one allele is deleted and the other allele is placed under the control of a $P_{TET-OFF}$ promoter. Strains are barcoded allowing STM screens. • CRISPR-cas9 gene drive array allowing the generation of double/multiple-gene deletion mutant libraries using haploids and parasexuality. <ul style="list-style-type: none"> • Collections are available from the Fungal Genetics Stock Center <p style="text-align: right;">References [105–111]</p>	<p style="text-align: center;">The ORFeome</p> <ul style="list-style-type: none"> • 5099 ORFs cloned in a Gateway™ donor vector • 83% of the <i>C. albicans</i> genome • Full-length sequence validated • 4061 variation-free ORFs <p style="text-align: center;">Collection of overexpression plasmids</p>  <p style="text-align: center;">Gateway™-compatible two-hybrid vectors</p> <p style="text-align: right;">Reference [112]</p>


and Rad52, which are shown to interact with CENP-A in *C. albicans* [52]. As a consequence, CENP-A is deposited onto the CENs. Consistent with this, in a CEN-deleted strain, the neocentromere becomes the earliest replicating region [51].

A telomere, at the termini of a eukaryotic chromosome, ensures chromosome end replication and protects the chromosome from degradation or end-to-end chromosome fusion. *C. albicans* telomeres are unique in containing tandem copies of unusually long 23-bp repeating units [53] (Figure 3C). However, they are assembled into heterochromatin via the classical Sir2-mediated pathway [53,54]. The subtelomeric regions of *C. albicans* consist of the telomere-associated (TLO) family of genes, which encode for the subunits of the mediator complex; a crucial component for transcription initiation [55]. There are 15 *TLO* genes (including one pseudogene) in *C. albicans* but other non-*C. albicans* species have either one or two *TLO* genes [56]. In addition, overexpression of *TLO* genes in *C. albicans* influences many growth- and virulence-related properties [57]. The expansion in the number of *TLO* genes could thus explain the ability of *C. albicans* to adapt in various host niches.

A special feature of the *C. albicans* genome is the **major repeat sequence (MRS)**. The MRS is a long tract (10–100 kb) of repetitive DNA that is present on all chromosomes, except chromosome 3. Structurally, an MRS is composed of three subunits: the repetitive RPS

Key Figure

Major Defining Features of *Candida albicans* and Its Genome.



Trends in Genetics

Figure 1. *C. albicans* undergoes a unique haploid–diploid–tetraploid life cycle. A phenotypic switch from white to opaque form due to homozygosis of the *MTL* locus is the initial step of this cycle. Opaque cells of opposite mating-type then fuse together to form tetraploids. These tetraploids undergo a nonmeiotic parasexual cycle to return to the diploid state. Diploid
(Figure legend continued on the bottom of the next page.)


subunit flanked by nonrepetitive elements RB2 and HOK (Figure 3D). Chromosome 3 contains only the RB2 element without the RPS or HOK unit [58]. Surprisingly, the MRS, being a repetitive region, is not assembled into classical heterochromatin but carries marks of both euchromatin and heterochromatin [54]. The MRS covers about 3% of the total genome, yet its function remains elusive, except that it is considered to be a hotspot for genome rearrangements in *C. albicans* [59]. The MRS is a preferred site for chromosomal translocations [58], and the expansion and contraction of its RPS region give rise to chromosome length polymorphism [60]. Furthermore, the presence of the MRS affects the frequency of **nondisjunction**, whereby a homologue bearing a larger MRS is more likely to be lost at the time of chromosome segregation [59]. Thus, the MRS serves as an important means of generating karyotypic diversity in *C. albicans* and needs to be studied in greater detail for its function and origin.

***Candida albicans*, a Heterozygous Diploid with a Dynamic Genome**

Although genome variation has so far been explored primarily either in haploid or homozygous diploid genomes, it is now being tackled in diploid organisms having **heterozygosity**, with a hope of better understanding the genomics of adaptation in various environmental or host niches. Population genomics studies have revealed a number of aspects regarding heterozygosity in the *C. albicans* genome [19,42,61–63] (Figure 4A). Natural heterozygosity is observed across the *C. albicans* genome with an average rate of one SNP every 237–283 bases. By comparing a large number of strains, a higher level of heterozygosity is found to correlate with faster growth rates. These observations likely reflect the loss of alleles that influence fitness in strains that have undergone partial or complete homozygosis. Genome sequencing data have identified about 3600 open reading frames (ORFs) with high-confidence SNPs leading to changes in the amino acid sequence. In addition, SNPs found in regulatory regions, or even in ORFs, can also affect transcription levels and/or translation efficiency between the alleles [63]. Thus, extensive allelic differences may function to increase genetic and phenotypic diversity in an organism devoid of a true sexual cycle and contribute to the acquisition of new phenotypic traits such as drug resistance.

Population genomics studies in *C. albicans* have revealed that genome heterozygosity can vary from 48% to 89% – heterozygous and homozygous regions being defined as such based on the number of heterozygous SNPs within 5-kb windows [19,23,42,61]. The levels of heterozygosity are primarily influenced by large LOH events encompassing whole chromosomes or extending from a specific chromosomal locus to the telomere. These LOH events have been shown to be pervasive in *C. albicans* isolates and can be detected on all chromosomes. LOH can involve an entire chromosome upon chromosome loss due to chromosome nondisjunction during mitosis. Depletion of the centromeric histone H3, Cse4/CENP-A at centromeres has been reported in response to changes in ploidy and the environment, and is associated with chromosome instability [64]. The genome of *C. albicans* contains two homologous histone H2A-encoding genes, *HTA1* on chromosome 3 encoding Hta1p and *HTA2* on chromosome 1

C. albicans can become haploid or vice versa by chromosome loss and autodiploidisation, respectively. Several host factors or environmental stress can lead to karyotypic variations in *C. albicans*. The genome of *C. albicans* is remarkably plastic and can tolerate segmental aneuploidies, whole chromosome aneuploidies, and loss-of-heterozygosity events. Accumulation of such DNA alterations can be associated to polarised growth and drug resistance in *C. albicans*. Several chromosomal elements influence the organisation and stability of the *C. albicans* genome. These include centromeres (CEN), which are all unique and different in *C. albicans*; replication origins, which are of two types, arm origins and centromeric origins; telomeres, which are unusually long and associated with a family of telomere-associated genes (TLO); and the major repeat sequence (MRS), a feature unique to *C. albicans*.



Trends in Genetics


Figure 2. *Candida albicans* Mating Type, Parasexuality, and Population Structure. (A) Schematic of the *MTLa* and *MTLα* idiotypes. The genes encoding transcription factors responsible for cellular identity are shown in red, while the *OBP*, *PAP*, and *PIK* genes that are not involved in cellular identity but differ between the two idiotypes are shown in grey. (B) Schematic of the haploid–diploid–tetraploid life cycle of *C. albicans*. *C. albicans* is predominantly existing in the diploid (Figure legend continued on the bottom of the next page.)

encoding Hta2p. Unlike Hta2p, Hta1p has lost the conserved phosphorylation site for the Bub1 kinase, a key regulator of chromosome segregation, and has been shown to facilitate chromosome gain and loss events in *C. albicans* [64]. Another frequent cause of LOH is somatic HR used to repair DNA DSBs. DSBs are the consequence of DNA replication defects or R loop formation but can also arise due to external stresses [65]. The extent of the LOH event can reflect the molecular mechanisms involved in DSB repair in *C. albicans*. Short-range LOH can occur by GC without crossover and long-range LOH including whole chromosome arms can occur by either GC with crossover, break-induced replication (BIR), or mitotic crossover (MCO). If left unrepaired, a DSB may result in chromosome truncation or chromosome loss, characterized by LOH events that span an entire chromosome or an arm of it. Several tools have been developed to study LOH events in *C. albicans* (Box 2). They have in particular revealed that DNA DSBs are predominantly repaired by GC but other repair events such as BIR/MCO and GC with crossover (CO) can also be observed at a significant frequency [66]. They have also revealed that the frequency at which LOH events arise and the nature of these LOH are influenced by environmental parameters (see below).

A link between LOH, aneuploidy, and the elevation of antifungal resistance by studying drug-resistant *C. albicans* isolates has been precisely established (Figure 4B). Gain-of-function alleles of genes involved in the resistance to azole antifungals have been shown to be codominant with wild-type alleles and therefore high levels of resistance cannot be achieved in the presence of the wild-type allele. It is only upon homozygosity of the gain-of-function allele, as a result of LOH, that high levels of azole resistance can be achieved [67–69]. In addition, the appearance of aneuploidy and, in particular, the formation of an isochromosome composed of the two left arms of chromosome 5 is often associated with the acquisition of azole resistance [70]. Increased copy number of *ERG11* and *TAC1*, both located on the left arm of chromosome 5, accounts for the majority of drug resistance associated with the chromosome 5 isochromosome [71]. Large-scale genome changes have been characterized in *S. cerevisiae* as a means of adaptation in response to stress [72,73]. Similarly, *in vitro* exposure to oxidative stress, elevated temperature, and antifungal drugs [74,75], as well as passaging through an animal model of infection result in increased genomic rearrangements in *C. albicans* [76,77]. The large-scale changes described above provide *C. albicans* with the ability to rapidly generate genetic and phenotypic diversity within the host environment.

Although LOH events are frequent and widespread in the genome of *C. albicans*, several studies have observed that, for some chromosomes, one haplotype or even a part of it is never found in the homozygous state in the *C. albicans* laboratory strain SC5314. This homozygosity bias has been observed in haploids, in parasexual derivatives, and in *rad52* mutant derivatives [35,37,78]. These observations suggest that recessive lethal and deleterious alleles can be found in the heterozygous state in the genome of *C. albicans*. Several groups, including ours, have identified such recessive lethal or deleterious alleles on chromosome 3A and chromosome 4B [66,79]. Overall, consistent with clonal reproduction, *C. albicans* strains harbour

state with heterozygosity at the *MTL* locus. Homozygosity at the *MTL* locus allows occasional white–opaque phenotypic switching and more efficient mating between opaque cells. Transition from tetraploidy to diploidy or diploidy to haploidy is independent of meiosis and involves random concerted chromosome loss with the intermediate aneuploidy progeny cells. Haploids are shown in blue, diploids in red, and tetraploids in green. (C) Genome sequencing of 182 *C. albicans* isolates confirms a predominantly clonal population structure and reveals, for the first time, footprints of admixtures in two genetic clusters (green arrows), demonstrating the occurrence of (para)sexuality in the *C. albicans* natural environment. LOH (BIR/MCO and GC) is a major driver of intraclade evolution and major LOH events can result in the emergence of clusters with altered virulence/niche restriction, possibly due to pseudogenization [42]. Abbreviations: BIR, break-induced replication; GC, gene conversion; LOH, loss of heterozygosity; MCO, mitotic crossover.





Trends in Genetics

Figure 3. Schematic of Essential Chromosomal Elements in *Candida albicans*. (A) A 3–5-kb long CENP-A-rich centromere (CEN) that lacks any common DNA sequence elements or pericentric repeats. The AT% of CEN regions is not significantly different from the rest of the genome and the CEN DNA sequence does not show any DNA methylation. The CEN function is dependent on the chromosomal context rather than DNA sequence, and thus epigenetically regulated. (B) DNA replication origins (*OR*s), identified as ORC-bound regions, are categorised into arm *OR*s and centromere *OR*s. Like centromeres, origins do not show any strong common DNA sequence motif. (C) Telomere repeats in *C. albicans* are unusually long and subtelomeric regions have an unusually high number of *TLO* genes in this organism. The high copy number of *TLO* genes in *C. albicans* is expected to be the result of subtelomeric recombination, mediated positively by *TLO* recombination element and negatively by Sir2 [98]. (D) The MRS, which further consists of three sequence elements, namely, RB2, RPS, and HOK. The RB2 (~6 kb) and HOK (~8 kb) elements are nonrepetitive sequences that occur only once per MRS, flanking the RPS element. The repeated sequence (RPS) is a ~2-kb-long repetitive sequence whose number can vary in an MRS. Each RPS unit carries an *Sfi*I restriction enzyme site. *Sfi*I mapping of the *C. albicans* genome served as a valuable tool to study chromosomal rearrangements before the whole genome sequence was available.

recessive lethal and deleterious alleles that constrain the outcome of LOH events. Although **homozygosity** of some alleles has been linked to fitness advantage in a specific host niche, overall genome-wide heterozygosity remains prevalent in the *C. albicans* population. For these reasons, the fate of cells having undergone LOH still needs to be addressed in *C. albicans*.

Specificities of DNA Repair Mechanisms in *Candida albicans*

DNA DSBs, where both strands of the double helix are severed, are the most serious forms of DNA damage. Indeed, failure to repair a DSB leads to loss of the CEN-lacking chromosome fragment, while improper repair of a DSB can lead to gross chromosomal rearrangements such as translocations, inversions, and deletions. Two major pathways of DSB repair are known: HR and **nonhomologous end joining (NHEJ)**. Characterisation of deletion mutants of genes involved in HR (*RAD52*, *RAD51*, *RAD59*, *RAD54*, and *RDH54*) have shown that HR plays a crucial role in DNA damage repair in *C. albicans* [78,80–83]. In contrast, the characterisation of mutants impaired for NHEJ has suggested that this process is not efficient in *C. albicans* [78,82]. Recent results on the mode of repair of **CRISPR-cas9**-induced DSBs are in agreement with this observation but suggest that NHEJ could occur in *C. albicans*, albeit rarely. Indeed, it has been shown that HR is the predominant repair pathway of CRISPR-Cas9-induced DSBs in *S. cerevisiae* and *C. albicans*, in contrast to other yeast species such as




Trends in Genetics

Figure 4. Heterozygosity, LOH, Genome Dynamics, and Antifungal Resistance. (A) Heat map illustrating the density of heterozygous SNPs across chromosome 3 for seven sequenced isolates. Regions that have undergone LOH appear white and are most often extending towards the telomere, indicating that they are the result of either mitotic recombination, break-induced replication, or gene conversion with crossover. (B) Schematic view of the impact of environmental stresses on genome dynamics and adaptation in *C. albicans*. Abbreviation: LOH, loss-of-heterozygosity.

Box 2. A Molecular Toolkit to Study *Candida albicans* Genome Dynamics

Molecular tools have been developed and successfully used to identify and study genome dynamics in *C. albicans*. (A) Identification of LOH events using the *URA3* or *GAL1* markers. In strains that have been engineered to be heterozygous for the *URA3* or *GAL1* gene at a given locus, LOH at this locus leads to resistance to 5-fluoroorotic acid (5-FOA) or 2-deoxygalactose (2-DG), respectively. (B) Identification of LOH events using a combination of *GFP* (green fluorescent protein-coding gene) and *BFP* (blue fluorescent protein-coding gene). In strains that have been engineered to be heterozygous at a given locus through insertion of *GFP* on one chromosome and *BFP* on its homologue, LOH at this locus results in loss of one of the *FP* genes and therefore results in monofluorescence, which can be revealed by flow cytometry. The monofluorescent cells are localised in the side gates and the double fluorescent cells are found in the middle gate. (C) Genome editing systems have been developed in *C. albicans*. A DNA DSB-inducing system was developed through conditional expression of the *S. cerevisiae* I-SceI meganuclease in a *C. albicans* strain engineered to harbour a unique I-SceI cleavage site. Characterisation of repair events at this I-SceI site show that they almost always correspond to GC events but some instances of BIR/MCO or WCL were also observed, as well as combinations of independent events. A method based on the CRISPR/CRISPR-Cas9 system has emerged for genome editing in *C. albicans*. By combining several guiding RNAs, multiple sites can be targeted simultaneously allowing simultaneous editing of multiple chromosomal sites. (D) Live cell fluorescence imaging of chromosome segregation. CENP-A-GFP strains with Tub1-mCherry allows one to follow kinetochore integrity and spindle morphology [110,113–117].



Candida glabrata and *Naumovia castelli* [84]. Moreover, while HR-defective *S. cerevisiae* mutants could still repair a CRISPR-Cas9-induced DSB in the absence of a repair template, likely via NHEJ, the same is not true in *C. albicans*, consistent with limited efficiency of NHEJ in this species [84]. Nevertheless, NHEJ is likely to occur in *C. albicans* as scars typical of this repair process have been reported at repaired CRISPR-Cas9-induced DSBs [85]. Notably, all of these studies have been performed in diploid isolates of *C. albicans* that are heterozygous at the *MTL* locus. In *S. cerevisiae*, NHEJ is downregulated in a/α diploid cells when compared with homozygous diploid or haploid cells only expressing *MATa* or *MAT α* . This is shown to be accomplished by transcriptional repression of specific genes by the $a1/\alpha2$ repressor [86]. Therefore, NHEJ studies in *MTL* homozygous diploid cells or haploid cells should address the absence or presence of NHEJ in *C. albicans*.

Several studies have shown that treatment of *C. albicans* with DNA-damaging agents triggers polarized growth [87,88]. Mutants with altered expression of genes coding for proteins involved in DNA damage response and cell cycle regulation also display aberrant filamentous morphology [48,49,78,87,89–93] (Figure 5). Strikingly, these data suggest that genotoxic-stress-induced polarised growth involves but does not require the expression of hyphal-specific genes. It is possible that stress-induced polarised growth is different from standard hyphal growth. Another key aspect of DNA repair is the accessibility of the DNA lesion to the DNA repair machinery. Proteins involved in chromatin assembly and remodelling have been shown to be important for efficient DNA repair in *C. albicans* as they change chromatin structure and allow repair proteins to access damaged DNA through the acetylation of histone H4 [94]. Similarly, involvement of HR in the kinetochore assembly has been demonstrated in *C. albicans* [52]. In mammalian cells, acetylation of histone H4 has been shown to also play a critical role in directing changes both in chromatin organisation and in promoting recruitment of DSB repair proteins to sites of DNA damage [95]. Moreover, different chromatin signatures associated with HR or NHEJ repair have been recently defined [96]. It has also been demonstrated that, apart from rapidly accumulating DNA damage, *C. albicans* cells lacking the histone acetyltransferase Hat1 also switch from yeast-like to polarized growth [94]. Altogether, these data corroborate




Figure 5. DNA-Damage-Induced Morphogenesis in *Candida albicans*. Schematic of the link between accumulation of DNA damage induced by various genotoxic stresses and polarised growth in *C. albicans*.

that perturbations of cell-cycle progression, a direct consequence of DNA damage, induce filamentous growth in *C. albicans*, in a manner dependent on the DNA damage/replication checkpoint kinase Rad53.

Concluding Remarks

In this review, we have highlighted some of the distinctive features uncovered from recent studies on the genome biology of *C. albicans* that make this yeast a model complementary to *S. cerevisiae* and *S. pombe*; two model species that are at the two extremes of yeast evolution. *C. albicans* exhibits specificities with respect to: (i) its life cycle whereby haploid, diploid, and tetraploid alternate through means of a meiosis-independent, recombinogenic parasexual cycle; (ii) chromosomal landmarks, especially the CENs that have unique organisation and epigenetic properties; (iii) an unusual genome plasticity, that frequently generates aneuploidies and LOH events; (iv) an almost obligate usage of HR for DSB repair; and (v) a coupling of the DNA damage response to morphogenetic shifts. A unique combination of these attributes in one organism allows investigators to address a variety of questions in genome biology that are not generally studied using the conventional yeast models. For instance, as most studies on recombination in *S. cerevisiae* and *S. pombe* employ laboratory haploid or homozygous diploid strains, little is known about the impact of heterozygosity on the biology of yeasts. Results from studies in *C. albicans* clearly demonstrate that heterozygosity is often advantageous, yet allows the propagation of recessive deleterious or lethal alleles that are detrimental upon homozygosis.

Despite changes in epidemiology, *C. albicans* remains responsible for a majority of yeast infections and the treatment of these infections is still a challenge in specific cases, such as systemic or recurrent vulvovaginal candidiasis. One of the key questions is whether *C. albicans* genome plasticity and hallmarks indeed contribute to making this species such an important pathogen. There is no doubt that LOH, ploidy changes, and isochromosome formation contribute to the elevation of antifungal resistance and treatment failure. Moreover, exposure of *C. albicans* to a variety of stresses – antifungals, oxidative stress, high temperature – has been shown to promote recombination events that could help *C. albicans* permanently adapt to changes in environment. Residence of *C. albicans* in animals including in healthy humans, whether as a commensal or pathogen, is accompanied by mutation and recombination events, predominantly short-range LOH (76; Sitterlé *et al.*, personal communication), that could not only reflect adaptation to the host but also help in repairing DNA alterations caused by constant exposure to DNA damaging agents (e.g., reactive oxygen species produced by immune cells). A recent study reports that a combination of mutations and LOH events acquired during serial passaging in the gastrointestinal tract of mice, where *C. albicans* does not normally reside, allows *C. albicans* to shift from a poor gastrointestinal commensal and harmful bloodstream-borne pathogen to an efficient commensal and poor systemic pathogen [97]. Therefore, the high genomic plasticity of *C. albicans* can bear long-term impact on its biology and gaining further insights into the genome biology of this species will certainly impact not only on our understanding of eukaryotic genome biology in general but also the mechanisms that make *C. albicans* such a successful opportunistic pathogen (see Outstanding Questions).

Acknowledgements

M.L., C.D., and K.S. are supported by a grant from the Indo French Centre for the Promotion of Advanced Research (CEFIPRA, Project no. 5703-2). Work in the laboratory of C.D. has been supported by the French Government's Investissement d'Avenir program (Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases, ANR-10-LABX-62-IBED). A.F. was the recipient of a PhD. grant from INRA Jouy-en-Josas and Institut Pasteur. Work in the laboratory of K.S. is supported by grants from the Department of Biotechnology, Government of India (BT/PR13909/BRB/

Outstanding Questions

Is *C. albicans* completely devoid of meiosis?

How does Spo11 function during the parasexual cycle?

Can we utilise the parasexual cycle to understand recombination in organisms with cryptic meiosis?

Why is the MRS the hotspot for chromosomal rearrangements?

What is the contribution of the DNA sequence in replication origin and centromere function?

Is heterozygosity necessary for fitness of *C. albicans* in the host?

How is heterozygosity maintained in the largely clonal *C. albicans* population?

What are the mechanisms for tolerance of genome plasticity in *C. albicans*?

Is genome plasticity contributing to *C. albicans* success as a commensal and/or pathogen?

Is there an influence of the microbiota on *C. albicans* genome plasticity?

Is NHEJ active in *C. albicans*?

How is the link between DNA damage and cell polarity precisely established?

10/1432/2015; BT/PR14557/BRB/10/15529/2016; BT/HRD/35/01/03/2017) and a grant for Life Science Research, Education, and Training (BT/INF/22/SP27679/2018). P.J. and K.S. are also financially supported by an intramural Jawaharalal Nehru Centre for Advanced Scientific Research (JNCASR) grant.

References

1. Botstein, D. and Fink, G.R. (1988) Yeast: an experimental organism for modern biology. *Science* 240, 1439–1443
2. Cavanaugh, A.M. and Jaspersen, S.L. (2017) Big lessons from little yeast: budding and fission yeast centrosome structure, duplication, and function. *Annu. Rev. Genet.* 51, 361–383
3. Fair, B.J. and Pleiss, J.A. (2017) The power of fission: yeast as a tool for understanding complex splicing. *Curr. Genet.* 63, 375–380
4. Hinnebusch, A.G. and Johnston, M. (2011) YeastBook: an encyclopedia of the reference eukaryotic cell. *Genetics* 189, 683–684
5. Noma, K.I. (2017) The yeast genomes in three dimensions: mechanisms and functions. *Annu. Rev. Genet.* 51, 23–44
6. Nurse, P.M. (2002) Nobel Lecture: Cyclin dependent kinases and cell cycle control. *Biosci. Rep.* 22, 487–499
7. Brown, G.D. *et al.* (2012) Hidden killers: human fungal infections. *Sci. Transl. Med.* 4, 165rv113
8. Ene, I.V. and Bennett, R.J. (2014) The cryptic sexual strategies of human fungal pathogens. *Nat. Rev. Microbiol.* 12, 239–251
9. Erwig, L.P. and Gow, N.A. (2016) Interactions of fungal pathogens with phagocytes. *Nat. Rev. Microbiol.* 14, 163–176
10. Gow, N.A. *et al.* (2011) *Candida albicans* morphogenesis and host defence: discriminating invasion from colonization. *Nat. Rev. Microbiol.* 10, 112–122
11. Lohse, M.B. *et al.* (2018) Development and regulation of single- and multi-species *Candida albicans* biofilms. *Nat. Rev. Microbiol.* 16, 19–31
12. Netea, M.G. *et al.* (2015) Immune defence against *Candida* fungal infections. *Nat. Rev. Immunol.* 15, 630–642
13. Noble, S.M. *et al.* (2017) *Candida albicans* cell-type switching and functional plasticity in the mammalian host. *Nat. Rev. Microbiol.* 15, 96–108
14. Perfect, J.R. (2017) The antifungal pipeline: a reality check. *Nat. Rev. Drug Discov.* 16, 603–616
15. Shapiro, R.S. *et al.* (2011) Regulatory circuitry governing fungal development, drug resistance, and disease. *Microbiol. Mol. Biol. Rev.* 75, 213–267
16. Sudbery, P.E. (2011) Growth of *Candida albicans* hyphae. *Nat. Rev. Microbiol.* 9, 737–748
17. Bougnoux, M.E. *et al.* (2008) Mating is rare within as well as between clades of the human pathogen *Candida albicans*. *Fungal. Genet. Biol.* 45, 221–231
18. Graser, Y. *et al.* (1996) Molecular markers reveal that population structure of the human pathogen *Candida albicans* exhibits both clonality and recombination. *Proc. Natl. Acad. Sci. U. S. A.* 93, 12473–12477
19. Jones, T. *et al.* (2004) The diploid genome sequence of *Candida albicans*. *Proc. Natl. Acad. Sci. U. S. A.* 101, 7329–7334
20. Riggsby, W.S. *et al.* (1982) DNA content, kinetic complexity, and the ploidy question in *Candida albicans*. *Mol. Cell Biol.* 2, 853–862
21. Hull, C.M. and Johnson, A.D. (1999) Identification of a mating type-like locus in the asexual pathogenic yeast *Candida albicans*. *Science* 285, 1271–1275
22. Tzung, K.W. *et al.* (2001) Genomic evidence for a complete sexual cycle in *Candida albicans*. *Proc. Natl. Acad. Sci. U. S. A.* 98, 3249–3253
23. Butler, G. *et al.* (2009) Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. *Nature* 459, 657–662
24. Hull, C.M. *et al.* (2000) Evidence for mating of the asexual yeast *Candida albicans* in a mammalian host. *Science* 289, 307–310
25. Magee, B.B. and Magee, P.T. (2000) Induction of mating in *Candida albicans* by construction of MTL α and MTL α strains. *Science* 289, 310–313
26. Bennett, R.J. and Johnson, A.D. (2003) Completion of a parasexual cycle in *Candida albicans* by induced chromosome loss in tetraploid strains. *EMBO J.* 22, 2505–2515
27. Dumitru, R. *et al.* (2007) *In vivo* and *in vitro* anaerobic mating in *Candida albicans*. *Eukaryot. Cell* 6, 465–472
28. Lachke, S.A. *et al.* (2003) Skin facilitates *Candida albicans* mating. *Infect. Immun.* 71, 4970–4976
29. Miller, M.G. and Johnson, A.D. (2002) White-opaque switching in *Candida albicans* is controlled by mating-type locus homeodomain proteins and allows efficient mating. *Cell* 110, 293–302
30. Banerjee, M. *et al.* (2008) UME6, a novel filament-specific regulator of *Candida albicans* hyphal extension and virulence. *Mol. Biol. Cell* 19, 1354–1365
31. Chen, C.G. *et al.* (2004) CaNdt180 is involved in drug resistance in *Candida albicans* by regulating CDR1. *Antimicrob. Agents Chemother.* 48, 4505–4512
32. Nobile, C.J. *et al.* (2012) A recently evolved transcriptional network controls biofilm development in *Candida albicans*. *Cell* 148, 126–138
33. Nocedal, I. *et al.* (2017) Gene regulatory network plasticity predates a switch in function of a conserved transcription regulator. *eLife* 6, e23250
34. Bartholomew, C.R. *et al.* (2012) Ume6 transcription factor is part of a signaling cascade that regulates autophagy. *Proc. Natl. Acad. Sci. U. S. A.* 109, 11206–11210
35. Forche, A. *et al.* (2008) The parasexual cycle in *Candida albicans* provides an alternative pathway to meiosis for the formation of recombinant strains. *PLoS Biol.* 6, e110
36. Hickman, M.A. *et al.* (2015) Parasexual ploidy reduction drives population heterogeneity through random and transient aneuploidy in *Candida albicans*. *Genetics* 200, 781–794
37. Hickman, M.A. *et al.* (2013) The ‘obligate diploid’ *Candida albicans* forms mating-competent haploids. *Nature* 494, 55–59
38. Berman, J. (2016) Ploidy plasticity: a rapid and reversible strategy for adaptation to stress. *FEMS yeast research* 16,
39. Gerstein, A.C. *et al.* (2017) Ploidy tug-of-war: Evolutionary and genetic environments influence the rate of ploidy drive in a human fungal pathogen. *Evolution* 71, 1025–1038
40. Schmid, J. *et al.* (2016) Last hope for the doomed? Thoughts on the importance of a parasexual cycle for the yeast *Candida albicans*. *Curr. Genet.* 62, 81–85
41. Zhang, N. *et al.* (2015) Selective advantages of a parasexual cycle for the yeast *Candida albicans*. *Genetics* 200, 1117–1132
42. Ropars, J. *et al.* (2018) Gene flow contributes to diversification of the major fungal pathogen *Candida albicans*. *Nat. Commun.* 9, 2253
43. Sanyal, K. *et al.* (2004) Centromeric DNA sequences in the pathogenic yeast *Candida albicans* are all different and unique. *Proc. Natl. Acad. Sci. U. S. A.* 101, 11374–11379
44. Baum, M. *et al.* (2006) Formation of functional centromeric chromatin is specified epigenetically in *Candida albicans*. *Proc. Natl. Acad. Sci. U. S. A.* 103, 14877–14882
45. Thakur, J. and Sanyal, K. (2013) Efficient neocentromere formation is suppressed by gene conversion to maintain centromere function at native physical chromosomal loci in *Candida albicans*. *Genome Res.* 23, 638–652
46. Ketel, C. *et al.* (2009) Neocentromeres form efficiently at multiple possible loci in *Candida albicans*. *PLoS Genet.* 5, e1000400

47. Burrack, L.S. *et al.* (2016) Neocentromeres provide chromosome segregation accuracy and centromere clustering to multiple loci along a *Candida albicans* chromosome. *PLoS Genet.* 12, e1006317
48. Thakur, J. and Sanyal, K. (2012) A coordinated interdependent protein circuitry stabilizes the kinetochore ensemble to protect CENP-A in the human pathogenic yeast *Candida albicans*. *PLoS Genet.* 8, e1002661
49. Roy, B. *et al.* (2011) CaMtw1, a member of the evolutionarily conserved Mis12 kinetochore protein family, is required for efficient inner kinetochore assembly in the pathogenic yeast *Candida albicans*. *Mol. Microbiol.* 80, 14–32
50. Tsai, H.J. *et al.* (2014) Origin replication complex binding, nucleosome depletion patterns, and a primary sequence motif can predict origins of replication in a genome with epigenetic centromeres. *mBio* 5, e01703-01714
51. Koren, A. *et al.* (2010) Epigenetically-inherited centromere and neocentromere DNA replicates earliest in S-phase. *PLoS Genet.* 6, e1001068
52. Mitra, S. *et al.* (2014) Rad51–Rad52 mediated maintenance of centromeric chromatin in *Candida albicans*. *PLoS Genet.* 10, e1004344
53. McEachern, M.J. and Hicks, J.B. (1993) Unusually large telomeric repeats in the yeast *Candida albicans*. *Mol. Cell Biol.* 13, 551–560
54. Freire-Beneitez, V. *et al.* (2016) *Candida albicans* repetitive elements display epigenetic diversity and plasticity. *Sci Rep.* 6, 22989
55. Zhang, A. *et al.* (2012) The Tlo proteins are stoichiometric components of *Candida albicans* mediator anchored via the Med3 subunit. *Eukaryot. Cell* 11, 874–884
56. Jackson, A.P. *et al.* (2009) Comparative genomics of the fungal pathogens *Candida dubliniensis* and *Candida albicans*. *Genome Res.* 19, 2231–2244
57. Dunn, M.J. *et al.* (2018) Functional diversification accompanies gene family expansion of MED2 homologs in *Candida albicans*. *PLoS Genet.* 14, e1007326
58. Chibana, H. and Magee, P.T. (2009) The enigma of the major repeat sequence of *Candida albicans*. *Future Microbiol.* 4, 171–179
59. Lephart, P.R. *et al.* (2005) Effect of the major repeat sequence on chromosome loss in *Candida albicans*. *Eukaryot. Cell* 4, 733–741
60. Chibana, H. *et al.* (2000) Fine-resolution physical mapping of genomic diversity in *Candida albicans*. *Genome Res.* 10, 1865–1877
61. Hirakawa, M.P. *et al.* (2015) Genetic and phenotypic intra-species variation in *Candida albicans*. *Genome Res.* 25, 413–425
62. Muzzey, D. *et al.* (2013) Assembly of a phased diploid *Candida albicans* genome facilitates allele-specific measurements and provides a simple model for repeat and indel structure. *Genome Biol.* 14, R97
63. Muzzey, D. *et al.* (2014) Extensive and coordinated control of allele-specific expression by both transcription and translation in *Candida albicans*. *Genome Res.* 24, 963–973
64. Brimacombe, C.A. *et al.* (2018) Chromatin rewiring mediates programmed evolvability via aneuploidy. *bioRxiv* In: <https://doi.org/10.1101/407841>
65. Chatterjee, N. and Walker, G.C. (2017) Mechanisms of DNA damage, repair, and mutagenesis. *Environ. Mol. Mutagen.* 58, 235–263
66. Feri, A. *et al.* (2016) Analysis of repair mechanisms following an induced double-strand break uncovers recessive deleterious alleles in the *Candida albicans* diploid genome. *mBio* 7, <http://dx.doi.org/10.1128/mBio.01109-16>
67. Coste, A. *et al.* (2007) Genotypic evolution of azole resistance mechanisms in sequential *Candida albicans* isolates. *Eukaryot. Cell* 6, 1889–1904
68. Ford, C.B. *et al.* (2015) The evolution of drug resistance in clinical isolates of *Candida albicans*. *eLife* 4, e00662
69. Morschhauser, J. *et al.* (2007) The transcription factor Mrr1p controls expression of the MDR1 efflux pump and mediates multidrug resistance in *Candida albicans*. *PLoS Pathog.* 3, e164
70. Selmecki, A. *et al.* (2006) Aneuploidy and isochromosome formation in drug-resistant *Candida albicans*. *Science* 313, 367–370
71. Selmecki, A. *et al.* (2008) An isochromosome confers drug resistance *in vivo* by amplification of two genes, ERG11 and TAC1. *Mol. Microbiol.* 68, 624–641
72. Chang, S.L. *et al.* (2013) Dynamic large-scale chromosomal rearrangements fuel rapid adaptation in yeast populations. *PLoS Genet.* 9, e1003232
73. Yona, A.H. *et al.* (2012) Chromosomal duplication is a transient evolutionary solution to stress. *Proc. Natl. Acad. Sci. U. S. A.* 109, 21010–21015
74. Forche, A. *et al.* (2011) Stress alters rates and types of loss of heterozygosity in *Candida albicans*. *mBio* 2, <http://dx.doi.org/10.1128/mBio.00129-11>
75. Harrison, B.D. *et al.* (2014) A tetraploid intermediate precedes aneuploid formation in yeasts exposed to fluconazole. *PLoS Biol.* 12, e1001815
76. Ene, I.V. *et al.* (2018) Global analysis of mutations driving microevolution of a heterozygous diploid fungal pathogen. *Proc. Natl. Acad. Sci. U. S. A.* 115, E8688–E8697
77. Forche, A. *et al.* (2018) Rapid phenotypic and genotypic diversification after exposure to the oral host niche in *Candida albicans*. *Genetics* 209, 725–741
78. Andaluz, E. *et al.* (2011) Rad52 function prevents chromosome loss and truncation in *Candida albicans*. *Mol. Microbiol.* 79, 1462–1482
79. Ciudad, T. *et al.* (2016) Phenotypic consequences of a spontaneous loss of heterozygosity in a common laboratory strain of *Candida albicans*. *Genetics* 203, 1161–1176
80. Bellido, A. *et al.* (2015) Genetic interactions among homologous recombination mutants in *Candida albicans*. *Fungal. Genet. Biol.* 74, 10–20
81. Garcia-Prieto, F. *et al.* (2010) Role of the homologous recombination genes RAD51 and RAD59 in the resistance of *Candida albicans* to UV light, radiomimetic and anti-tumor compounds and oxidizing agents. *Fungal. Genet. Biol.* 47, 433–445
82. Legrand, M. *et al.* (2007) Role of DNA mismatch repair and double-strand break repair in genome stability and antifungal drug resistance in *Candida albicans*. *Eukaryot. Cell* 6, 2194–2205
83. Legrand, M. *et al.* (2011) The contribution of the S-phase checkpoint genes MEC1 and SGS1 to genome stability maintenance in *Candida albicans*. *Fungal. Genet. Biol.* 48, 823–830
84. Vyas, V.K. *et al.* (2018) New CRISPR mutagenesis strategies reveal variation in repair mechanisms among fungi. *mSphere* 3, In: <https://doi.org/10.1128/mSphere.00154-18>
85. Ng, H. and Dean, N. (2017) Dramatic improvement of CRISPR/Cas9 editing in *Candida albicans* by increased single guide RNA expression. *mSphere* 2, In: <https://doi.org/10.1128/mSphere.00385-16>
86. Valencia, M. *et al.* (2001) NEJ1 controls non-homologous end joining in *Saccharomyces cerevisiae*. *Nature* 414, 666–669
87. Bachewich, C. *et al.* (2005) Cell cycle arrest during S or M phase generates polarized growth via distinct signals in *Candida albicans*. *Mol. Microbiol.* 57, 942–959
88. da Silva Dantas, A. *et al.* (2010) Thioredoxin regulates multiple hydrogen peroxide-induced signaling pathways in *Candida albicans*. *Mol. Cell Biol.* 30, 4550–4563
89. Bachewich, C. and Whiteway, M. (2005) Cyclin Cln3p links G1 progression to hyphal and pseudohyphal development in *Candida albicans*. *Eukaryot. Cell* 4, 95–102
90. Bensen, E.S. *et al.* (2005) The mitotic cyclins Clb2p and Clb4p affect morphogenesis in *Candida albicans*. *Mol. Biol. Cell* 16, 3387–3400

91. Shi, Q.M. *et al.* (2007) Critical role of DNA checkpoints in mediating genotoxic-stress-induced filamentous growth in *Candida albicans*. *Mol. Biol. Cell* 18, 815–826
92. Thakur, J. and Sanyal, K. (2011) The essentiality of the fungus-specific Dam1 complex is correlated with a one-kinetochore-one-microtubule interaction present throughout the cell cycle, independent of the nature of a centromere. *Eukaryot. Cell* 10, 1295–1305
93. Loll-Kripplbeber, R. *et al.* (2014) A study of the DNA damage checkpoint in *Candida albicans*: uncoupling of the functions of Rad53 in DNA repair, cell cycle regulation and genotoxic stress-induced polarized growth. *Mol. Microbiol.* 91, 452–471
94. Tscherner, M. *et al.* (2012) The histone acetyltransferase Hat1 facilitates DNA damage repair and morphogenesis in *Candida albicans*. *Mol. Microbiol.* 86, 1197–1214
95. Dhar, S. *et al.* (2017) The tale of a tail: histone H4 acetylation and the repair of DNA breaks. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 372
96. Clouaire, T. *et al.* (2018) Comprehensive mapping of histone modifications at DNA double-strand breaks deciphers repair pathway chromatin signatures. *Mol. Cell.* 72, 250–262 e256
97. Tso, G.H.W. *et al.* (2018) Experimental evolution of a fungal pathogen into a gut symbiont. *Science* 362, 589–595
98. Freire-Beneitez, V. *et al.* (2016) Sir2 regulates stability of repetitive domains differentially in the human fungal pathogen *Candida albicans*. *Nucleic Acids Res.* 44, 9166–9179
99. Santos, M.A. *et al.* (2011) The genetic code of the fungal CTG clade. *C. R. Biol.* 334, 607–611
100. Shen, X.X. *et al.* (2016) Reconstructing the backbone of the Saccharomycotina yeast phylogeny using genome-scale data. *G3 (Bethesda)* 6, 3927–3939
101. Skrzypek, M.S. *et al.* (2018) Using the *Candida* genome database. *Methods Mol. Biol.* 1757, 31–47
102. Hernday, A.D. *et al.* (2010) Genetics and molecular biology in *Candida albicans*. *Methods Enzymol.* 470, 737–758
103. Xu, T. *et al.* (2011) Genome-wide transposon mutagenesis in *Saccharomyces cerevisiae* and *Candida albicans*. *Methods Mol. Biol.* 765, 207–224
104. Mielich, K. *et al.* (2018) Maize transposable elements Ac/Ds as insertion mutagenesis tools in *Candida albicans*. *G3 (Bethesda)* 8, 1139–1145
105. Roemer, T. *et al.* (2003) Large-scale essential gene identification in *Candida albicans* and applications to antifungal drug discovery. *Mol. Microbiol.* 50, 167–181
106. Xu, D. *et al.* (2007) Genome-wide fitness test and mechanism-of-action studies of inhibitory compounds in *Candida albicans*. *PLoS Pathog.* 3, e92
107. Nobile, C.J. and Mitchell, A.P. (2009) Large-scale gene disruption using the *UAU1* cassette. *Methods Mol. Biol.* 499, 175–194
108. Homann, O.R. *et al.* (2009) A phenotypic profile of the *Candida albicans* regulatory network. *PLoS Genet.* 5, e1000783
109. Noble, S.M. *et al.* (2010) Systematic screens of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and pathogenicity. *Nat. Genet.* 42, 590–598
110. Shapiro, R.S. *et al.* (2018) A CRISPR-Cas9- based gene drive platform for genetic interaction analysis in *Candida albicans*. *Nat. Microbiol.* 3, 73–82
111. Segal, E.S. *et al.* (2018) Gene essentiality analyzed by *in vivo* transposon mutagenesis 658 and machine learning in a stable haploid isolate of *Candida albicans*. *mBio* 9, In: <https://doi.org/10.1128/mBio.02048-18>
112. Legrand, M. *et al.* (2018) Generating genomic platforms to study *Candida albicans* pathogenesis. *Nucleic Acids Res.* 46, 6935–6949
113. Forche, A. *et al.* (2003) A system for studying genetic changes in *Candida albicans* during infection. *Fungal. Genet. Biol.* 39, 38–50
114. Loll-Kripplbeber, R. *et al.* (2015) A FACS-optimized screen identifies regulators of genome stability in *Candida albicans*. *Eukaryot. Cell* 14, 311–322
115. Vyas, V.K. *et al.* (2015) A *Candida albicans* CRISPR system permits genetic engineering of essential genes and gene families. *Science advances* 1, e1500248
116. Min, K. *et al.* (2016) *Candida albicans* gene deletion with a transient CRISPR-Cas9 system. *mSphere* 1, In: <https://doi.org/10.1128/mSphere.00130-16>
117. Nguyen, N. *et al.* (2017) An efficient, rapid, and recyclable system for CRISPR-mediated genome editing in *Candida albicans*. *mSphere* 2, In: <https://doi.org/10.1128/mSphereDirect.00149-17>