

Changes in soil carbon inputs and outputs along a tropical altitudinal gradient of volcanic soils under intensive agriculture

Jorge Sierra, François Causeret

► To cite this version:

Jorge Sierra, François Causeret. Changes in soil carbon inputs and outputs along a tropical altitudinal gradient of volcanic soils under intensive agriculture. Geoderma, 2018, 320, pp.95-104. 10.1016/j.geoderma.2018.01.025 . hal-02628286

HAL Id: hal-02628286 https://hal.inrae.fr/hal-02628286

Submitted on 11 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Changes in soil carbon inputs and outputs along a tropical altitudinal gradient of volcanic soils
2	under intensive agriculture
3	
4	Jorge Sierra*, François Causeret
5	
6	ASTRO Agrosystèmes Tropicaux, INRA, 97170, Petit-Bourg, Guadeloupe, France.
7	* Corresponding author at: ASTRO Agrosystèmes Tropicaux, INRA, 97170, Petit-Bourg, Guadeloupe,
8	France.
9	E-mail address: jorge.sierra@inra.fr (J. Sierra).
10	
11	
12	ABSTRACT
13	
14	Volcanic soils contain a large stock of soil organic carbon (SOC) which is highly vulnerable to changes
15	in land use and climate warming. In this study we examine the changes of SOC stocks along a tropical
16	elevation gradient (100 m-700 m; +437 mm yr ⁻¹ of rainfall and –0.7°C every 100 m), which is subject
17	to two intensive agricultural systems (banana monoculture and vegetable crops) characterised by the
18	heavy use of fertilisers and pesticides. We hypothesise that in these systems SOC is mainly controlled
19	by soil mineralogy, climatic factors and soil tillage. We used a process-based approach to determine
20	soil C inputs and outputs along the elevation gradient. The banana monoculture systems (198 plots)
21	are characterised by a temporal steady-state of SOC, and the vegetable crop systems (55 plots)
22	present high annual SOC losses. The banana systems were used to determine for the first time the
23	altitudinal change in the in situ rate constant of SOC mineralisation (k_{soc}). Under banana
24	monoculture, SOC stock increased by 218%, C input from crop residues decreased by 44%, and ${\sf k}_{\sf soc}$
25	decreased by 570% across the altitudinal gradient. These results indicated that the SOC gradient was
26	mainly induced by changes in C outputs. Allophane content increased with altitude (R^2 = 0.61; 8 g kg ⁻¹
27	every 100 m) and was positively correlated with SOC content ($R^2 = 0.51$) and negatively correlated
28	with k_{SOC} (R ² = 0.53). We hypothesise that the physical protection of SOC within amorphous
29	allophanic minerals was the main factor responsible for the k_{soc} and SOC stock gradients. Strictly, the
30	effects of allophane content and rainfall cannot be separated because soil mineralogy in the studied
31	area is mainly determined by the level of rainfall. Our process-based analysis indicated that changes
32	in temperature along the elevation gradient could only explain 22% of the observed change in k_{soc} .
33	SOC stocks under vegetable crops also increased with altitude; they were 57% lower than under
34	banana at low altitude but only 9% lower at high altitude. This reflected simultaneously the higher C
35	outputs under vegetable crops due to more intensive soil tillage, and the greater resilience to soil
36	disturbance of protected SOC in altitude. The results suggested that neither soil nutrient content nor

- 37 soil biological properties contributed to the SOC and k_{soc} gradients under the intensive cropping
- 38 systems analysed during this study. Overall, we conclude that SOC might be more vulnerable to soil
- 39 tillage and warming in low-altitude soils.
- 40

Keywords: allophane; banana; SOC decomposition; soil tillage; temperature sensitivity; vegetable
crops

- 43
- 44

45 1. Introduction

46

47 Although volcanic soils account for less than 1% of Earth's land area, they store about 5% of the global stock of soil organic carbon (SOC) (Dahlgren et al., 2004). These soils have peculiar physical 48 49 (e.g. very high water content at field capacity, high resilience to compaction and erosion), chemical 50 (e.g. high SOC and basic cations contents), and physico-chemical properties (e.g. high P fixation, high 51 anion exchange capacity), which are controlled by soil mineralogy, characterised by the presence of 52 amorphous minerals such as allophane (Clermont-Dauphin et al., 2004). These soils are mostly found 53 on the flanks of volcanic mountains and are occupied by forests and agricultural land, the latter 54 mainly in the inter-tropical zone where altitude moderates climate (Legros, 2012). Volcanic soils have 55 high agricultural productivity and are easy to work, but aluminium toxicity and P deficiency may 56 occur in those that are more acidic (Dahlgren et al., 2004). Because of the high SOC stocks in these 57 soils, conversion of the natural vegetation into farmland frequently induces SOC losses that can reach 58 2%-3% per year (e.g. Lemenih and Itanna, 2004; Sierra et al., 2015). However, some authors have 59 reported that the proper management of cropping systems, including diversified production and 60 reduced soil tillage, can be useful to preserving SOC stocks in volcanic soils (Segnini et al., 2011; 61 Sierra et al., 2017). This is of crucial concern in tropical mountain systems where warming is likely to 62 be particularly marked during coming decades and could contribute to increasing SOC losses 63 (Nottingham et al., 2016).

64 Most worldwide studies dealing with the dynamics of SOC in natural mountain systems have 65 observed positive correlations between SOC stocks and altitude (Tashi et al., 2016, for a metaanalysis on this topic). A few exceptions did not demonstrate any such relationship (e.g. Tan and 66 67 Wang, 2016) or an increase up to a given altitude and then a decrease (e.g. Djukic et al., 2010). 68 Generally as altitude increases so does annual rainfall, while air temperature decreases, thus causing marked changes in vegetation, soil type and soil microbial activity across the elevation gradient, 69 70 which in turn may affect the C cycle. Numerous factors and mechanisms have been proposed to 71 explain the increase of SOC in line with elevation in each particular system; e.g. an improved physical 72 protection of SOC because of the increase in the soil aggregate rate (Li et al., 2016), a reduction in

SOC decomposition linked to the decline of soil temperature (Dieleman et al., 2013), or the increase
of C humification associated with changes in vegetation and litter quality (Wang et al., 2016). This
general pattern of increasing SOC with altitude is frequently masked in areas where forests have
been replaced by agricultural systems which cause local decline in SOC stocks (Ozalp et al., 2016).

77 The high SOC stocks in mountain systems dominated by volcanic soils have generally been 78 attributed to the physical protection of organic matter provided by allophanic minerals, together 79 with a reduction in microbial activity breaking down SOC due to soil acidity, poor P availability and low soil temperature (Lemenih and Itanna, 2004; Naafs et al., 2004). As these factors co-vary with 80 81 altitude, confounding effects may occur when the relationship between them and SOC is evaluated 82 using correlation analysis (Körner, 2007). In this sense, although some authors have proposed that 83 mountain areas - with their elevation gradients of temperature and rainfall - may be useful to assess 84 the impact of climate changes similar to those observed across latitudes (Klimek et al., 2016; Wang et 85 al., 2016), Nottingham et al. (2015) pointed out that careful consideration should be given to the 86 effects of other soil and plant factors varying concomitantly with elevation. Undoubtedly, knowledge 87 of the interactions between the factors that induce changes in SOC is necessary to assess the impact 88 of projected warming in mountain areas. For example, if falling temperatures at higher altitudes is 89 the main factor that controls rises in SOC, then studies addressing the temperature sensitivity of SOC 90 decomposition (e.g. evaluation of the Q₁₀ coefficient) are essential to predict the impact of warming. 91 Indeed, several studies have focused on this topic (e.g. Zimmermann et al., 2009; Zimmermann and 92 Bird, 2012). By contrast, more complex studies will be necessary if warming simultaneously affects C 93 outputs (e.g. SOC decomposition) and C inputs (i.e. changes in vegetation and litter quality) 94 (Nottingham et al., 2016), or if SOC changes are primarily controlled by the physical protection 95 provided by soil structure, whose characteristics vary with altitude (Kramer and Chadwick, 2016).

96 The flanks of volcanic islands in tropical regions where the elevation gradient of climate is 97 one of the strongest in the world present a wide range of soil types, with andosols on the summit 98 and ferralsols, luvisols and nitisols on the footslopes (Legros, 2012). These regions therefore 99 represent an interesting edaphic and climatic setting to determine the effects of the factors 100 impacting SOC stocks. In this study we examine the changes of SOC stocks in an altitudinal gradient 101 of a tropical volcanic island where the land is subject to two intensive agricultural systems (banana 102 monoculture and vegetable crops), which are characterised by the heavy use of fertilisers and 103 pesticides. We hypothesise that in these systems SOC is mainly controlled by soil mineralogy, climatic 104 factors and soil tillage. We consider that the effects of other factors such as soil nutrient content and 105 soil biological properties are of minor importance in our intensive agricultural systems. The aim of 106 this study was therefore to assess the underlying mechanisms controlling SOC by focusing on the 107 effect of the gradients of soil mineralogy (e.g. allophanic mineral content), climate (e.g. temperature 108 and rainfall), and the intensity of soil tillage (low under banana, high under vegetable crops). The

109 banana cropping systems are characterised by a temporal steady-state of SOC stocks and the 110 vegetable crop systems present high annual SOC losses (Sierra et al., 2015). To reduce the impact of 111 confounding effects caused by factors that co-vary with altitude, we used a process-based approach to estimate C outputs and C inputs and to assess their effects on the SOC balance along the elevation 112 113 gradient. The study was carried out in Guadeloupe (French West Indies), which exhibits within a 114 small area nearly every physical landscape and cropping system found in the Caribbean region. This 115 study formed part of a larger project devoted to quantifying the impact of cropping systems, the 116 pedoclimate and warming on SOC stocks, using Guadeloupe as a case study of the Caribbean (Sierra et al., 2017). 117 118 119 120 2. Materials and methods 121 122 2.1. Study location, soils and climate 123 124 The study was carried out in the uplands of south Basse-Terre, the main island of the 125 Guadeloupe archipelago located in the eastern Caribbean (16° 05' N, 61° 40' W) (Fig. 1). Basse-Terre 126 Island is dominated by a mountain chain oriented northwest to southeast where the crest standing at 127 1456 m.a.s.l. (La Soufrière volcano). Although the land west of the crest slopes steeply toward the 128 Caribbean Sea (slopes of 20-30%), the eastern side of the chain slopes more gently toward the Atlantic Ocean (slopes of 10%). On both slopes, deep valleys trending at right angles to the mountain 129 130 chain are separated by sharp ridges. During the present study we focused on the land areas 131 dominated by andosols (FAO classification) (Fig. 1), which had developed on volcanic ash deposited 132 directly by successive volcanic eruptions throughout the Late Glacial and Holocene (Legros, 2012). 133 Because of the increase in the allophanic clay content with altitude, the hydrated microporous 134 structure of the soil, responsible for its high water content at field capacity, progressively expands 135 with increasing altitude, whereas bulk density decreases (Colmet-Daage and Lagache, 1965). The 136 zone of transition between the upland andosols and rich halloysite nitisols of the lowlands is placed 137 between 100 m and 200 m on the eastern slopes, and between 200 m and 300 m on the western 138 slopes (Fig. 1). In the studied area, the mean air temperature drops by 0.7°C and annual rainfall 139 increases by 437 mm yr⁻¹ per every 100 m (Fig. 2). The level of rainfall is lower from December to 140 May, which is slightly more pronounced in the lowlands (38% of the annual rainfall takes place in this

6-month period; an average of 170 mm per month) than in the uplands (45% of the annual rainfall;

- 142 an average of 380 mm per month).
- 143

141

144 2.2. Land use and farming practices

145

The agricultural land in the studied area is oriented toward the production of bananas for the 146 147 European market and vegetable crops for the local market (Sierra et al., 2015). Farms devoted to 148 banana production are situated from 100 m to 700 m and average 10 ha, while those focused on 149 vegetable crops are located from 100 m to 900 m and average 4 ha. The land above an altitude of 150 900 m is occupied by tropical rainforest; this is a protected area where agriculture and deforestation 151 have been banned. Only the agricultural land was included in the present study (Fig. 1). Sierra et al. 152 (2015) had previously reported that the SOC stored under banana cropping systems in this region is 153 likely to be close to, or at, a steady-state when crop residues are the only C input to the soil (i.e. 154 when no organic amendments are applied). By contrast, SOC stocks decrease by 2.0 ± 1.1 Mg C ha⁻¹ 155 yr⁻¹ in soils cultivated with vegetable crops (Sierra et al., 2017).

156 Banana is managed mainly as a pluriannual system and its cycle is 5-6 years. The growth 157 season is 9-12 months and varies among plants in the same plot; so, the initially homogeneous plant 158 population becoming heterogeneous after few years. This is important to the time of harvest 159 because each plant is harvested separately (Raphael et al., 2012). Harvesting is performed manually 160 and residues (leaves and stems) are cut and placed on the soil. During growth, each plant (mother 161 plant) produces a sucker (daughter plant) which grows after harvest of the mother plant. For this 162 reason, soil tillage is only applied before planting every 5-6 years. In plots with steep slopes where 163 mechanised tillage is not possible, the soil is not ploughed and replanting is ensured gradually by 164 hand as the old plants die. Such a system can be considered as being perennial (Clermont-Dauphin et al., 2004). 165

The principal vegetable crops in the studied area are tomato, cabbage, salad, eggplant and pepper. They are cultivated under monoculture systems; some farmers apply one year of fallow after 5-6 years of cultivation. Soil tillage is highly intensive with at least six tillage operations per year. Harvesting is manual and the residues (leaves and stems) are placed on the soil or buried at a depth of 0.1–0.2 m.

171 Most banana and vegetable crop systems are based on the heavy use of synthetic fertilisers and pesticides. For example, N fertilisers are applied at a rate of 400-600 kg N ha⁻¹ yr⁻¹ in banana and 172 173 150-200 kg N ha⁻¹ yr⁻¹ in vegetable crops. For K, the rates are 120 kg K ha⁻¹ yr⁻¹ for banana and 80 kg K 174 ha⁻¹ yr⁻¹ for vegetables. For P, the rates are 50 kg P ha⁻¹ yr⁻¹ for banana and 80 kg P ha⁻¹ yr⁻¹ for 175 vegetables. For banana, P and K fertilisers are applied after harvest of the mother plant, and N 176 fertiliser is split into 8-12 applications along the growth cycle. For vegetables, all the fertilisers are 177 applied at planting. No irrigation is applied in these cropping systems. In fact, the water requirement 178 for banana crops in the studied area is 120-150 mm per month (Lassoudière, 2007), which is fully 179 supplied by rainfall even during the drier season in the lowlands. About 50% of the plots under

banana and vegetable crop systems are limed every 5-10 years to raise soil pH and to reduce theimpact of aluminium toxicity on plant growth on the more acidic soils.

182

183 2.3. Soil and plant data

184

185 SOC stocks were evaluated using the soil database developed for the TropEmis Project 186 performed in Guadeloupe (Sierra et al., 2015). This database contains information on soils and farm 187 characteristics covering all agro-ecological regions of the archipelago. For the present study we 188 selected data corresponding to the studied area, which included 198 plots under banana 189 monoculture since about 1980, and 55 plots cultivated with vegetable crops since about 1990. These 190 plots had never received organic amendments. Information on the farms included their size, the 191 number and size of plots, location, altitude, soil type, cropping systems, and the management of 192 fertilizers, organic amendments, crop residues and soil tillage. The soil characteristics of the selected 193 plots included SOC (determined with a C analyser), organic N (Kjeldahl method), extractable P (Truog 194 method), cation exchangeable capacity and exchangeable cations (ammonium acetate method) and 195 pH (soil-water ratio of 1:5). Soil bulk density (core method; NRCS, 2014) was available for 89 plots (53 196 under banana monoculture and 36 under vegetable crops). Allophane content was available for 28 197 plots under banana monoculture and was estimated as proposed by Buurman et al. (2007):

198 Allophane = Si_o/0.14

acid method

(1)

where Si_o (g kg⁻¹) is Si in allophane determined using the ammonium oxalate-oxalic acid method
(Matus et al., 2008). We selected allophane content to assess soil mineralogy because it was well
correlated with SOC content in several studies carried out on volcanic soils (e.g. Matus et al., 2008;
Garrido and Matus, 2012).

203 Soil analyses were performed on the 0-0.25 m soil layer and all soil samples were collected by 204 the same team. Soil sampling was carried out on a systematic grid using a 0.06-m diameter auger and 205 collecting the equivalent of 30 sub-samples per hectare. Chemical analyses were performed on a 206 composite sample (~1 kg dry matter) at the SADEF Soil Testing Laboratory in France. Bulk density was 207 determined with five replications per plot. No significant differences between the two cropping 208 systems were observed for bulk density for plots located at the same altitude (Sierra et al., 2015). 209 Only yields that concern export crops are available in Guadeloupe. Banana yields in the studied area 210 were obtained from the Banana Producer Group of Guadeloupe and corresponded to the yields 211 observed in 2015 for 90 plots included in the soil database. It is important to point out that the 212 groups of banana plots with available information concerning bulk density, allophane content and 213 yields were not fully identical, but they were within the selected 198 plots under banana 214 monoculture.

215

216 2.4. Analytical approach

217

218 In this study, most of the factors affecting SOC present a high level of collinearity (e.g. Fig. 2), 219 which prevents them from being including together in a multiple regression model to predict SOC 220 content (de Oliveira Aparecido et al., 2017). We therefore adopted a mechanistic approach by 221 applying a process-based analysis to estimate C fluxes and SOC balance along the elevation gradient. 222 The analysis involved two steps. In the first step, we determined C inputs and outputs in order to 223 assess their relative influence on the SOC balance along the elevation gradient. Although C input was 224 represented by the humified crop residue C, C output corresponded to the in situ rate constant of 225 SOC mineralisation. This step was based on the results obtained in this study. In the second step we 226 investigated the relationships between these C fluxes and the soil (e.g. allophane and nutrient 227 contents, C/N ratio, microbial biomass, abundance of detritivore mesofauna) and climatic factors 228 (e.g. temperature and rainfall) affecting SOC turnover. This step was based on our results as well as 229 on those obtained in previous studies performed in the same area analysed here. The analysis was 230 applied to the plots under banana monoculture and calculations were made under the hypothesis of 231 a temporal steady-state of SOC stock. As cited above, this hypothesis was supported by the results 232 reported by Sierra et al. (2015), and also by the long period under banana monoculture without any 233 application of organic amendments. Because this hypothesis could not be applied to the plots 234 cultivated with vegetable crops (i.e. SOC stocks decrease over time), the rate constant of SOC 235 mineralisation was not calculated for these systems. SOC stocks under vegetable crop systems were 236 used to compare them with the altitudinal trend of SOC stocks observed for the banana systems. 237 We considered that in situ SOC mineralisation follows a first order kinetic (Saffih-Hdadi and

Mary, 2008). Therefore, under the hypothesis of a temporal steady-state of SOC stock at the plotscale it may be established that:

$$dSOC_{stock} / dt = 0 = (C_{res} \times h_{res}) - (SOC_{stock} \times k_{SOC})$$
(2)

241 and then:

242
$$k_{SOC} = (C_{res} \times h_{res}) / SOC_{stock}$$
 (3)

243 where SOC_{stock} (Mg C ha⁻¹) is the SOC stock and $dSOC_{stock}$ / dt is the annual change in SOC stock (Mg C

ha⁻¹ yr⁻¹), C_{res} (Mg C ha⁻¹ yr⁻¹) is the annual C input from crop residues (aboveground and roots), h_{res}

245 (unitless) is the humification coefficient of crop residues, and k_{SOC} (yr⁻¹) is the rate constant of

246 mineralisation of SOC_{stock}.

247 The characteristics of crop residues analysed in this study and the equations used to calculate 248 SOC_{stock}, C input from crop residues and the temperature sensitivity of k_{soc} (the Q₁₀ coefficient) are

- 249 presented as Supplementary Information in Appendix A (Table A.1 and Eqs. (A), respectively). Bulk 250 density was used to calculate SOC_{stock} (Eq. (A.1)), banana yields were used to calculate C inputs from 251 crop residues (Eqs. (A.2) and (A.3)), and temperature was used to calculate the Q₁₀ coefficient as 252 proposed by Hamdi et al. (2013) (Eqs. (A.4) and (A.5)). Although bulk density, banana yields and 253 temperature varied with altitude (see section 3 and Fig. 2), as mentioned above, these data were not 254 available for all of the selected banana plots. We therefore used the equations fitted to quantify the 255 altitudinal gradient of these three variables in order to estimate their values for each selected plot 256 based on knowledge of its altitude. All the fits were performed using the Newton solution of the Excel[™] solver. 257
- 258 259
- 260 **3. Results**
- 261

To simplify the presentation of the results, the altitudinal pattern of some soil properties are provided as Supplementary Information in Appendix A (Figs. A).

264 Only 15% of the selected banana plots were managed as perennial systems. These plots were 265 present from an altitude of about 300 m, and their SOC stocks were similar to those observed for the 266 plots managed as pluriannual systems located at the same altitude (Fig. A.1). In this way, all the 267 banana plots could be analysed as a single population regardless of their management. The SOC 268 content increased with altitude under both banana monoculture and vegetable crop systems (Fig. 3). 269 The intercept and the slope of the linear regression fitted to SOC data differed significantly between 270 the two cropping systems (P < 0.05); e.g. the intercept was higher and the slope was lower for the 271 banana monoculture. At an altitude of 100 m, the estimated SOC content was about 35 g C kg⁻¹ under 272 banana monoculture and 25 g C kg⁻¹ under vegetable crops; these values were respectively 75 g C kg⁻¹ and 65 g C kg⁻¹ at 700 m. Soil bulk density decreased with altitude from about 0.82 Mg m⁻³ at 100 m 273 274 to about 0.67 Mg m^{-3} at 700 m (Fig. 4a). Allophane content increased with altitude from about 25 g 275 kg⁻¹ at 100 m to 75 g kg⁻¹ at 700 m (Fig. 4b). SOC stocks were calculated from the SOC content of each 276 selected plot and the soil bulk density data obtained using the equation shown in Fig. 4a. The SOC 277 stocks displayed the same trend as that observed for SOC content (Fig. A.2). The differences between 278 cropping systems in terms of SOC stocks, in absolute and in relative terms, were higher at low 279 altitude (e.g. at 100 m, 55 Mg C ha⁻¹ for banana and 35 Mg C ha⁻¹ for vegetable crops; i.e. 57% higher 280 for banana) than at high altitude (e.g. at 700 m, 120 Mg C ha⁻¹ for banana and 110 Mg C ha⁻¹ for 281 vegetable crops; i.e. 9% higher for banana).

For both banana and vegetable crop systems, no changes with altitude were observed for cation exchangeable capacity (Figs. 5a and 5b) and soil pH (Figs. 5c and 5d). The same was observed for the C/N ratio (Fig. A.3), cations and base saturation (Fig. A.4), and extractable P (Fig. A.5). The average values of these soil properties are presented in Table 1. No differences were observed
between the two cropping systems except for extractable P (higher under vegetable crops) and
exchangeable K (higher under banana) (P < 0.05) (Table 1).

As the C/N ratio did not display any trend with altitude, the altitudinal gradient of organic N was very similar to that presented in Fig. 3 for SOC (Fig. A.6). Under banana monoculture, organic N increased from 2.7 g N kg⁻¹ at 100 m (5.6 Mg N ha⁻¹) to 7.3 g N kg⁻¹ at 700 m (12.3 Mg N ha⁻¹). For vegetable crops, organic N increased from 1.7 g N kg⁻¹ at 100 m (3.8 Mg N ha⁻¹) to 6.6 mg N kg⁻¹ at 700 m (10.9 Mg N ha⁻¹).

- 293 Banana yields decreased with altitude from about 45 Mg ha⁻¹ at 100 m to 20 Mg ha⁻¹ at 700 294 m (Fig. 6a). Yields displayed a relatively high variability throughout the altitudinal gradient. Using 295 these yield values, banana residues estimated from Eqs. (A.2) and (A.3) decreased from 15 Mg ha⁻¹ at 100 m to 7 Mg ha⁻¹ at 700 m. Therefore, the input of humified C derived from banana residues ($C_{res} \times$ 296 297 h_{res}, in Eq. (2)) decreased 2.2-fold between 100 m (1.6 Mg C ha⁻¹ yr⁻¹) and 700 m (0.7 Mg C ha⁻¹ yr⁻¹). 298 The in situ k_{soc} estimated from Eq. (3) for the selected plots of banana monoculture are 299 presented in Fig. 6b. k_{soc} decreased 6.7-fold between 100 m (3.0 × 10⁻² yr⁻¹) and 700 m of altitude 300 $(4.4 \times 10^{-3} \text{ yr}^{-1})$. With such a steep change, the Q₁₀ temperature coefficient estimated from Eq. (A.5) 301 was as great as 31, which is highly unrealistic, as will be discussed in section 4.2.2. For the 28 banana 302 plots used to determine allophane content, SOC content correlated positively and k_{soc} correlated 303 negatively with allophane content (P < 0.05) (Fig. 7).
- 304
- 305

306 4. Discussion

307

308 309

310 Our results confirmed that tropical volcanic soils under agriculture can store amounts of 311 organic matter equivalent to those observed in natural forest systems. For example, the SOC stocks 312 observed under banana monoculture were similar to those reported by Powers and Schlesinger 313 (2002) for a tropical rainforest in Costa Rica on volcanic soils, with comparable altitudinal gradient (e.g. 50 Mg C ha⁻¹ at 50 m to 140 Mg C ha⁻¹ at 800 m) and climatic conditions (e.g. temperature 314 315 ranging from 25°C to 20°C and rainfall ranging from 3500 mm yr⁻¹ to 4700 mm yr⁻¹). In addition, the 316 rise in SOC levels with altitude in our study tended to be linear as had previously been found by other 317 authors in natural ecosystems (e.g. Zimmermann et al., 2009; Dieleman et al., 2013). However, it is likely that altitude did not exert a direct impact on the SOC gradient, but an indirect effect involving 318 319 soil (e.g. mineralogy), climatic (e.g. temperature and rainfall) and vegetation factors (e.g. amount of 320 crop residues) that varied across the elevation gradient. Because SOC stocks result from a balance of

4.1. Relative impact of C inputs and outputs on the SOC balance under banana monoculture

321 C inputs and outputs, such factors might induce the SOC gradient due to their impacts on C input and322 output fluxes.

323 In our agricultural systems, C input was dependent on the quantity of crop residues and the 324 humification coefficient (Eq. (2)). In fact, the plots included in this study were selected because no 325 organic amendments were applied so that C from crop residues was the only C input into the soil. 326 The decline of banana yields at increasing altitudes, and consequently the reduction of the amount of 327 crop residues, revealed that changes in C input could not be responsible for the increase in SOC with 328 altitude, but might weaken it. Banana yields drop with altitude because lower temperatures reduce 329 the growth rate and extend the growing season, which together diminish plant biomass and annual 330 fruit production (Brisson et al., 1998). In contrast with this finding in our agricultural systems, several 331 authors have found that in natural forest systems, C inputs rise with altitude because of an increase 332 in the amount of litter associated with marked changes in the composition and density of vegetation 333 (e.g. Powers and Schlesinger, 2002; Tashi et al., 2016).

334 Although the temperature gradient may also affect the decomposition rate of crop residues, 335 the humification coefficient of crop residues is generally considered to be dependent solely on 336 residue quality (e.g. Justes et al., 2009). We thus considered that h_{res} in Eq. (2) was not affected by 337 the temperature gradient, in which case C input only varied as a function of the amount of crop 338 residues (i.e. Eqs. (A.2) and (A.3)). Some authors have reported that the humification coefficient in 339 natural forest systems varies in line with changes in vegetation and litter quality along the altitudinal 340 gradient; e.g. humification of litter C increased in an elevation gradient in Taiwan (Wang et al., 2016) and decreased in an elevation gradient in Peru (Segnini et al., 2011). This was not the case during the 341 342 present study because vegetation and crop residue quality were the same across the altitudinal 343 gradient. This is consistent with the results obtained in a recent study carried out in the same area analysed here, which indicated that nutrient content of banana plants and residues was not affected 344 345 by the elevation gradient (G. Loranger-Merciris, unpublished).

346 On the basis of our results it appears that the increase in SOC stocks with altitude was 347 primarily controlled by the reduction of C outputs, as suggested by the steep decline of k_{soc} with 348 altitude. Indeed, the banana monoculture systems are characterised by a steady-state of SOC stock 349 at each altitude, where the level of SOC at equilibrium increases with altitude because C outputs 350 decrease faster than C inputs. Therefore, the assessment of the underlying mechanisms that induce 351 the SOC gradient should be based on the evaluation of the most important factors affecting the 352 gradient of k_{soc} .

353

4.2. Factors affecting the altitudinal gradient of k_{SOC}

355

In our process-based approach, k_{soc} represents the soil biological activity linked to SOC mineralisation and integrates the impact of the climatic and soil factors that affect mineralisation in the field. On the basis of our results, we hypothesise that the k_{soc} gradient was firstly controlled by soil mineralogy and rainfall and secondly by temperature. We consider that other factors such as soil nutrient content, soil C/N ratio and soil biological properties (e.g. microbial biomass and abundance of detritivore mesofauna) were of minor importance in this study. These factors are analysed below.

362

363 4.2.1. Soil mineralogy and rainfall

364

365 These factors cannot be analysed separately because soil mineralogy in the studied area is 366 mainly determined by the level of rainfall, which controls the intensity of the weathering processes 367 (Colmet-Daage and Lagache, 1965; Legros, 2012). In this way, soils located in the wet uplands are 368 characterised by chemically reactive allophanic materials, which incorporate organic compounds to 369 develop amorphous gels (Clermont-Dauphin et al., 2004). The content in allophanic materials 370 diminishes as the altitude and rainfall decrease, and some gel minerals reorganise progressively into 371 crystalline 1:1 clay minerals (e.g. halloysite) and aluminium and iron hydrous oxides, which adsorb 372 organic matter on their surfaces but do not incorporate it into the mineral matrix (Kramer and 373 Chadwick, 2016). The crystallisation of 1:1 minerals occurs during the drier season and is then more 374 rapid in lowlands with a lower level of rainfall in that season (Sedov et al., 2003). The increase of 375 allophane content and the decrease of bulk density with altitude observed in the present study 376 reflected this change in soil mineralogy, which was in line with the results reported by other studies 377 on volcanic soils (Powers and Schlesinger, 2002; Li et al., 2016).

378 Several authors reported that SOC content was positively correlated with allophane content 379 in elevation gradients of volcanic soils (e.g. Powers and Schlesinger, 2002; Matus et al., 2008), and 380 the same was observed in the present study (Fig. 7a). Moreover, the range of values of allophane 381 content observed in this study was similar to that reported by Matus et al. (2008) and Garrido and 382 Matus (2012) for volcanic soils in Chile. Buurman et al. (2007) proposed that high rainfall in the 383 uplands favours the presence of anoxic conditions inside the micropores of the allophanic gels, 384 reducing SOC decay and inducing the increase in SOC content with altitude. On the other hand, 385 Garrido and Matus (2012) proposed that SOC stabilisation in andosols is associated with the low 386 physical accessibility for microbial decomposers of SOC inside the micropores of allophanic clay 387 minerals. Although the mechanisms of SOC stabilisation in andosols are not fully elucidated at this 388 time (Filimonova et al., 2016), it appears that this stabilisation would be the major responsible for the increase in the mean residence time and the content of SOC with altitude in elevation gradients 389 390 of volcanic soils (Kramer and Chadwick, 2016). We hypothesise that the steep drop observed in k_{soc} 391 along the altitudinal gradient reflected the increase in the mean residence time of SOC. This

hypothesis is supported by the negative correlation observed between k_{soc} and allophane content, which indicated that allophane content explained about 53% of the variability of k_{soc} . (Fig. 7b). Since SOC is more protected against microbial degradation within allophanic materials than on the surfaces of secondary minerals, k_{soc} decreases and SOC stocks increase with altitude.

396 It is interesting to point out that k_{SOC} was higher at low altitudes despite a lower level of 397 rainfall, mainly in the drier season. However, an effect of rainfall on k_{soc} in lowlands cannot be 398 discarded because it could affect the activity of SOC decomposers. Although under the prevailing 399 rainfall regime in the uplands the soils are at field capacity much of the year (Clermont-Dauphin et 400 al., 2004), a modelling study carried out in the lowlands of the studied area indicated that water 401 content in the top soil layer can be lower than that at field capacity during a part of the drier season 402 (Brisson et al., 1998). Therefore, rainfall could weaken the k_{soc} gradient by reducing the k_{soc} values in 403 the lowlands. While it is difficult to separate the influence of rainfall on microbial activity and on soil 404 mineralogy in lowlands, it seems clear that the latter was a key factor explaining the decrease of k_{soc} 405 with increasing altitude.

406

407 4.2.2. Temperature

408

409 The decrease of temperature with altitude is frequently cited as the main factor inducing an 410 increase in SOC content in elevation gradients under natural forest systems. For example, Luo et al. 411 (2014) found that in situ soil respiration decreased by 35% in the altitudinal gradient of a subalpine 412 secondary forest in China, which was attributed to a drop in soil temperature from 11°C to 8°C along 413 the elevation gradient. Using Eqs. (A.4) and (A.5), we calculated that the Q_{10} coefficient of soil 414 respiration corresponding to the results of Luo et al. (2014) would be 2.7, which is within the range of 415 values cited by several authors (e.g. Nottingham et al., 2015). In our study, k_{soc} decayed 6.7-fold from 416 100 m to 700 m for a temperature gradient of only 4.4°C. Indeed, this sharp decay of k_{soc} gave rise to 417 a highly unrealistic Q_{10} value (31). The kinetic theory on which the estimate of Q_{10} is based predicts a value of around 2 for typical ambient temperatures of 0°C-20°C (Davidson and Janssens, 2006), which 418 419 increases for the low ranges of temperature and for organic compounds of greater chemical 420 complexity (Nottingham et al., 2015). For example, in laboratory studies carried out on soils collected 421 from altitudinal gradients, the Q₁₀ of SOC mineralisation was about 1.5 in the Changbai Mountains of 422 China (Wang et al., 2016), and ranged from 2 to 3 in the Western Carpathians of Poland (Klimek et 423 al., 2016). Nottingham et al. (2015) pointed out that the higher Q_{10} values frequently observed in situ 424 experiments (e.g. up to 7 in their study), when compared with the lower values obtained from 425 laboratory studies, might be associated with complex interactions between factors varying 426 simultaneously with elevation. We think this was also the case in our study, where the unrealistic Q_{10} 427 might have been an artefact induced by other factors changing with altitude such as soil mineralogy.

Using Eqs. (A.4) and (A.5), we calculated that for the temperature gradient observed in the present study, k_{soc} would decrease 1.4-fold from 100 m to 700 m for a Q_{10} value of 2. The decrease would be 1.6-fold for a Q_{10} of 3. This implies that changes in temperature in our elevation gradient only could explain between 21% and 24% (i.e. 1.4/6.7 and 1.6/6.7, respectively) of the observed change in k_{soc} in the studied area.

433

434 4.2.3. Soil nutrient content and C/N ratio

435

436 Natural systems in tropical uplands are frequently characterised by gradients of nutrient 437 limitation that affect plants, microbial activity and SOC decomposition, mainly with respect to N and 438 P (Buurman et al., 2007; Nottingham et al., 2015). This does not appear to be the case in our study 439 because of the heavy use of NPK fertilisers and lime in banana cropping systems, which could explain 440 the lack of altitudinal gradients for P and cation contents. In a study carried out on tropical forest 441 soils in Costa Rica, Cleveland et al. (2004) reported that available soil P was a limiting factor for SOC 442 and litter decomposition. However, the contents of extractable P found by these authors were 10-15 443 times smaller than those observed under banana monoculture in the present study (Table 1). 444 Similarly, the contents of exchangeable Ca, Mg and K observed in our soils were about two-fold 445 higher than the values reported for other elevation gradients of volcanic soils under natural 446 vegetation (e.g. Kramer and Chadwick, 2016).

447 Concerning N availability, even if organic N content was lower in the lowlands than in the 448 uplands (Fig. A.6), it is unlikely that N could limit k_{soc} at low altitudes because of the high rates of N fertiliser applied in banana monoculture along the growth cycle (400-600 kg N ha⁻¹ yr⁻¹, split into 449 450 several applications). Moreover, Dorel et al. (2005) found that the amount of mineralisable N 451 measured in a laboratory experiment using andosols of the same area analysed here was much 452 higher than that observed in other tropical soils and was not correlated with either altitude or SOC 453 content. For an elevation gradient in the Western Carpathians, Klimek et al. (2016) observed a 454 reduction in the soil C/N ratio that was associated with an increase at high altitudes of the more 455 chemically recalcitrant fraction of SOC, which could decrease the rate of SOC decomposition in the 456 uplands. In contrast to this, our results showed that the C/N ratio did not exhibit any trend with 457 altitude; i.e. there was no evidence of a change in SOC recalcitrance with altitude. Overall, these 458 results suggest that neither soil nutrient content nor soil C/N ratio contributed to the SOC and k_{SOC} 459 gradients.

460

461 4.2.4. Microbial biomass and detritivore mesofauna

462

Several authors observed that microbial biomass and the structure of bacterial and fungal communities change with altitude, which may affect SOC turnover. For example, Pabst et al. (2013) found that microbial biomass increased in an elevation gradient of Mount Kilimanjaro (Tanzania) under natural forest and savannah ecosystems. Similar results were obtained by Chang et al. (2016) for an elevation gradient in Taiwan dominated by bamboo plantations. In addition, these authors reported that the fungi-bacteria ratio also increased with altitude, which was related to the increase in SOC content, mainly in the amount of the labile SOC fraction.

470 In the present study, we did not focus on the biological properties of the soils because 471 previous studies performed in the same area indicated that they were affected by the management 472 of the banana cropping system but not by altitude. For example, Clermont-Dauphin et al. (2004) and 473 Dorel et al. (2005) found that microbial biomass was higher in less intensive cropping systems, 474 regardless of their altitude. These systems included low rates of fertilisers and pesticides and the use 475 of organic amendments. In addition, the number and the biomass of earthworms were higher under 476 these less intensive systems (Clermont-Dauphin et al., 2004), and the same was found for the 477 abundance of detritivore mesofauna such as ants and termites (G. Loranger-Merciris, unpublished). 478 These studies indicated that microbial biomass and the abundance of mesofauna did not display any 479 trend with altitude in the more intensive systems involved in the present study. As discussed above, 480 banana systems including the use of organic amendments were not analysed in our study because 481 the hypothesis of a steady-state of SOC could not be applied to these systems (Sierra et al., 2015). On the basis of these results, we consider that the SOC and k_{SOC} gradients observed in our intensive 482 483 banana systems cannot be attributed to changes in microbial biomass or in the size of the mesofauna 484 population with altitude.

485

486 4.3. Differences in SOC stocks between banana and vegetable cropping systems

487

488 Several authors reported that SOC stocks decreased dramatically following the conversion of 489 natural vegetation into arable land in elevation gradients (e.g. Lemenih and Itanna, 2004; Pabst et al., 490 2013). In our long-term agricultural systems, where deforestation had not been applied during the 491 past five decades, SOC stocks were lower under vegetable crop systems than under banana 492 monoculture throughout the elevation gradient. This could be due to two factors: a lower C input 493 because of smaller quantities of crop residues from vegetable crops, and a higher C output because 494 of the more intensive soil tillage applied in these systems. Crop residues from vegetable crops vary 495 from 10.3 Mg dry matter ha⁻¹ yr⁻¹ for eggplant to 2.2 Mg ha⁻¹ yr⁻¹ for salad (Table A.1). The coefficient of humification for vegetable crop residues averaged 0.34, which is close to that of banana residues 496 497 (i.e. 0.31; Table A.1), which suggests that residue humification was of minor importance when trying 498 to explain the differences between the two cropping systems. By considering the mean quantity of

499 residues from vegetable crops (i.e. 6 Mg ha⁻¹ yr⁻¹) and the mean humification coefficient (Table A.1), 500 the mean input of humified C derived from vegetable crop residues would be about 0.9 Mg C ha⁻¹ yr⁻¹ 501 ¹, which is 45% lower than that observed for banana monoculture at 100 m (i.e. 1.6 Mg C ha⁻¹ yr⁻¹) 502 and 30% higher than that observed at 700 m (i.e. 0.7 Mg C ha⁻¹ yr⁻¹). Although in Guadeloupe there 503 are not data on the changes of vegetable crop yields and their crop residues with altitude, it seems 504 that on average the C input from crop residues was probably lower under vegetable crops than under 505 banana monoculture, which may partially explain their lower SOC stocks. Concerning C outputs, 506 Sierra et al. (2015) found that SOC mineralisation in the region studied during this work was about 507 110% higher in soils under vegetable crops compared to those under banana monoculture, linked to 508 differences between the systems in terms of the intensity of soil tillage. It therefore seems that 509 differences between the two cropping systems in C outputs were much higher than their differences 510 in C inputs. We concluded that differences in SOC stocks between the two cropping systems along 511 the elevation gradient were mainly associated with C outputs and secondarily with C inputs. 512 Although differences between the two systems regarding extractable P (higher under vegetable 513 crops) and exchangeable K (higher under banana) contents reflected the specific fertiliser practices of 514 each cropping system, the level of most soil properties was similar in both cropping systems (Table 515 1). We therefore consider that these properties were of minor importance to explain the differences 516 in SOC stocks between vegetable and banana cropping systems.

517 A more interesting finding from this study was that the differences in SOC stocks between 518 the two cropping systems decreased with altitude. This was consistent with the hypothesis that changes in soil mineralogy played a major role in SOC protection and decomposition. Thus SOC 519 520 occluded within allophanic minerals at high altitude would be less responsive to differences in soil 521 tillage between the cropping systems, and its vulnerability would increase in line with the reduction 522 of allophane content at lower altitudes. This is consistent with the kinetic theory of SOC 523 decomposition (Nottingham et al., 2016), and agrees with the results reported by Nottingham et al. 524 (2015) concerning the greater sensitivity of unprotected SOC to environmental changes along an 525 elevation gradient in the Peruvian Andes.

526 The results of this study may provide some insights concerning the impact of climate 527 warming on SOC stocks in our tropical altitudinal gradient. Climate models predict that the mean 528 temperature in the Guadeloupe archipelago will rise 0.7°C in the next three decades (Sierra et al., 529 2010). Assuming a Q_{10} value of 2 for SOC mineralisation, this rise in temperature would induce a 5% 530 increase in the current k_{SOC} values. The increase would be 8% for a Q_{10} of 3. Even though warming 531 and Q_{10} may vary along the altitudinal gradient, it appears that the magnitude of the change in k_{soc} due to warming might be much smaller than that observed for k_{soc} between the lowlands and 532 533 uplands in this study (i.e. 570%). It follows that SOC decomposition and C emissions will continue to 534 be higher in lowlands under a climate warming scenario, mainly under vegetable cropping systems. Indeed, this hypothesis needs to be tested during future laboratory and field studies designed to
determine Q₁₀ values across the gradient of soil mineralogy.

- 537
- 538

539 5. Conclusions

540

541 Our results confirm that any assessment of the underlying mechanisms controlling SOC changes in altitudinal gradients needs to consider the confounding influence of the various edaphic, 542 543 climatic and vegetation factors that co-vary with elevation. For this, the process-based approach 544 applied in this study to analyse the banana cropping systems, characterised by a steady-state of SOC 545 stock, was useful to assess the factors affecting the SOC gradient. This approach made it possible to 546 establish that the level of SOC at the steady-state increases with altitude because C outputs decrease 547 faster than C inputs. In other words, the increase in SOC with altitude was primarily controlled by the 548 reduction in C outputs, represented by the in situ rate constant of SOC mineralisation (k_{soc}). Changes in allophane content with altitude explained 53% of the observed variability of k_{SOC} , which probably 549 550 reflected the effect of the physical protection of SOC within amorphous allophanic minerals. 551 Although changes in temperature only could explain 22% of such variability, the unrealistic Q_{10} value 552 of k_{soc} estimated for the entire altitudinal gradient suggests that the temperature sensitivity of SOC 553 needs to be determined at a local level (e.g. for each soil mineralogical type defined by the allophane 554 content).

555 In contrast to reports on altitudinal gradients under natural ecosystems, it appears that 556 neither soil nutrient content nor biological soil properties contributed to the SOC and k_{soc} gradients 557 under our agricultural systems characterised by a heavy use of fertilisers and pesticides. In addition, 558 SOC vulnerability to climate warming would be less in uplands because of the physical protection 559 provided by allophanic amorphous minerals. These results suggest that SOC under vegetable crop 560 systems may be more vulnerable to warming than that under banana monoculture, which would be 561 linked to an increase in the content of unprotected SOC following the intensive soil tillage applied in 562 the former case. Further work is now necessary to assess the combined effects of warming and soil 563 mineralogy on SOC stocks under contrasting tropical agricultural systems.

564

565

566 Acknowledgements

567

568 This study formed part of the TropEmis Project funded by the Reacctif Programme of the 569 French Agency for the Environment and Energy Management (ADEME) (Grant 410-00159), the 570 European Regional Development Fund (FEDER) (Grant 410-00160) and the Regional Council of

571	Guadeloupe (Grant 410-00161), and of the RIVAGE Project funded by the FEDER (Grant 410-00242)
572	and the Regional Council of Guadeloupe (Grant 410-00325). We are grateful to Marcus Hery of the
573	Banana Producer Group of Guadeloupe for providing us with the database on banana yield, and to
574	the 253 farmers who were interviewed for this study. We would like to Victoria Hawken for
575	reviewing the English manuscript. We thank the anonymous reviewers for their constructive
576	comments, which helped us to improve the manuscript.
577	
578	
579	Appendix A. Supplementary Information
580	
581	Supplementary information linked to this article can be found at:
582	
583	
584	References
585	
586	Brisson, N., Ozier-Lafontaine, H., Dorel, M., 1998. Effects of soil management and water regime on
587	banana growth between planting and flowering. Simulation using the STICS model. Acta
588	Hortic. 490: 229-238.
589	Buurman, P., Peterse, F., Almendros Martin, G., 2007. Soil organic matter chemistry in allophanic
590	soils: a pyrolysis-GC/MS study of a Costa Rican Andosol catena. Eur. J. Soil Sci. 58: 1330-1347.
591	Chang, E.H., Chen, T.H., Tian, G., Chiu, C.Y., 2016. The effect of altitudinal gradient on soil microbial
592	community activity and structure in moso bamboo plantations. Appl. Soil Ecol. 98: 213-220.
593	Clermont-Dauphin, C., Cabidoche, Y.M., Meynard, J.M., 2004. Effects of intensive monocropping of
594	bananas on properties of volcanic soils in the uplands of the French West Indies. Soil Use
595	Manag. 20: 105-113.
596	Cleveland, C.C., Townsend, A.R., Constance, B.C., 2004. Soil microbial dynamics in Costa Rica:
597	seasonal and biogeochemical constraints. Biotropica 36: 184-195.
598	Colmet-Daage, F., Lagache, P., 1965. Caractéristiques de quelques groupes de sols dérivés de roches
599	volcaniques aux Antilles françaises. Cahiers ORSTOM Série Pédologie 3: 91-121 (in French).
600	Dahlgren, R. A., Saigusa, M., Ugolini, F.C., 2004. The nature, properties and management of volcanic
601	soils. Adv. Agron. 82: 114-183.
602	Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and
603	feedbacks to climate change. Nature 440: 165-173.
604	de Oliveira Aparecido, L.E., de Souza Rolim, G., Camargo Lamparelli, R.A., de Souza, P.S., dos Santos,
605	E.R., 2017. Agrometeorological models for forecasting coffee yield. Agron. J. 109: 249-258.

- Dieleman, W.I.J., Venter, M., Ramachandra, A., Krockenberger, A.K., Bird, M.I., 2013. Soil carbon
 stocks vary predictably with altitude in tropical forests: Implications for soil carbon storage.
 Geoderma 204-205: 59-67.
- Djukic, I., Zehetner, F., Tatzber, M., Gerzabek, M.H., 2010. Soil organic-matter stocks and
 characteristics along an Alpine elevation gradient. J. Plant Nutr. Soil Sci. 173: 30-38.
- Dorel, M., Lombard, K., Oliver, R., 2005. Mineralisable nitrogen and organic status of andosols in
 Guadeloupe: Climatic and edaphic determinants, influence of soil management. Étude et
 Gestion des Sols 12: 267-280 (in French).
- Filimonova, S., Kaufhold, S., Wagner, F.E., Häusler, W., Kögel-Knabner, I., 2016. The role of allophane
 nano-structure and Fe oxide speciation for hosting soil organic matter in an allophanic
 Andosol. Geochim. Cosmochim. Ac. 180: 284-302.
- 617 Garrido, E., Matus, F., 2012. Are organo-mineral complexes and allophane content determinant
 618 factors for the carbon level in Chilean volcanic soils? Catena 92: 106-112.
- Hamdi, S., Moyano, F., Sall, S., Bernoux, M., Chevallier, T., 2013. Synthesis analysis of the
 temperature sensitivity of soil respiration from laboratory studies in relation to incubation

621 methods and soil conditions. Soil Biol. Biochem. 58: 115-126.

- Justes, E., Mary, B., Nicolardot, B., 2009. Quantifying and modelling C and N mineralization kinetics of
 catch crop residues in soil: parameterization of the residue decomposition module of STICS
 model for mature and non mature residues. Plant Soil 325: 171-185.
- Klimek, B., Jelonkiewicz, Ł., Niklińska, M., 2016. Drivers of temperature sensitivity of decomposition
 of soil organic matter along a mountain altitudinal gradient in the Western Carpathians. Ecol.
 Res. 31: 609-615.
- Körner, C., 2007. The use of "altitude" in ecological research. Trends Ecol. Evol. 22: 569-574.
- Kramer, M.G., Chadwick, O.A., 2016. Controls on carbon storage and weathering in volcanic soils
 across a high-elevation climate gradient on Mauna Kea, Hawaii. Ecology 97: 2384-2395.
- 631 Lassoudière, A., 2007. Le bananier et sa culture. Editions Quae, Paris, France (in French).
- Legros, J.P., 2012. Major soil groups of the world: ecology, genesis, properties and classification. CRC
 Press-Taylor & Francis Group, Boca Raton, FL, USA.
- Lemenih, M., Itanna, F., 2004. Soil carbon stocks and turnovers in various vegetation types and arable
 lands along an elevation gradient in southern Ethiopia. Geoderma 123: 177-188.
- Li, L., Vogel, J., He, Z., Zou, X., Ruan, H., Huang, W., Wang, J., Bianchi, T.S., 2016. Association of soil
 aggregation with the distribution and quality of organic carbon in soil along an elevation
- 638 gradient on Wuyi Mountain in China. PLoS ONE 11(3): e0150898.
- 639 doi:10.1371/journal.pone.0150898.
- Luo, S., Liu, G., Li, Z., Hu, C., Gong, L., Wang, M., Hu, H., 2014. Soil respiration along an altitudinal
 gradient in a subalpine secondary forest in China. iForest 8: 526-532.

- Matus, F., Garrido, E., Sepúlveda, N., Cárcamo, I., Panichini, M., Zagal, E., 2008. Relationship between
 extractable Al and organic C in volcanic soils of Chile. Geoderma 148: 180-188.
- Naafs, D.F.W., van Bergen, P.F., Boogert, S.J., de Leeuw, J.W., 2004. Solvent-extractable lipids in an
 acid andic forest soil; variations with depth and season. Soil Biol. Biochem. 36: 297-308.
- 646 Nottingham, A.T., Turner, B.L., Whitaker, J., Ostle, N., Bardgett, R.D., McNamara, N.P., Salinas, N.,
- 647 Meir, P., 2016. Temperature sensitivity of soil enzymes along an elevation gradient in the
 648 Peruvian Andes. Biogeochemistry 127: 217-230.
- Nottingham, A.T., Whitaker, J., Turner, B.L., Salinas, N., Zimmermann, M., Malhi, Y., Meir, P., 2015.
 Climate warming and soil carbon in tropical forests: insights from an elevation gradient in the
 Peruvian Andes. BioScience 65: 906-921.
- NRCS, 2014. Kellogg soil survey laboratory methods manual. Soil Survey Investigations Report No. 42
 Version 5.0. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb1253872.pdf
 (accessed 03.05.17).
- Ozalp, M., Yuksel, E.E., Yuksek, T., 2016. Soil property changes after conversion from forest to
 pasture in mount Sacinka, Artvin, Turkey. Land Degrad. Develop. 27: 1007-1017.
- Pabst, H., Gerschlauer, F., Kiese, R., Kuzyakov, Y., 2013. Land use and precipitation affect organic and
 microbial carbon stocks and the specific metabolic quotient in soils of eleven ecosystems of
 Mt. Kilimanjaro, Tanzania. Land Degrad. Develop. 27: 592-602.
- Powers, J.S., Schlesinger, W.H., 2002. Relationships among soil carbon distributions and biophysical
 factors at nested spatial scales in rain forests of northeastern Costa Rica. Geoderma 109:
 165-190.
- Raphael, L., Sierra, J., Recous, S., Ozier Lafontaine, H., Desfontaines, L., 2012. Soil turnover of crop
 residues from the banana (Musa AAA cv. Petite-Naine) mother plant and simultaneous
 uptake by the daughter plant of released nitrogen. Eur. J. Agron. 38: 117-123.
- Saffih-Hdadi, K., Mary, B., 2008. Modeling consequences of straw residues export on soil organic
 carbon. Soil Biol. Biochem. 40: 594-607.
- Sedov, S.N., Solleiro-Rebolledo, E., Gama-Castro, J.E., 2003. Andosol to Luvisol evolution in Central
 Mexico: timing, mechanisms and environmental setting. Catena 54: 495-513.
- Segnini, A., Posadas, A., Quiroz, R., Milori, D.M.B.P., Vaz, C.M.P., Martin-Neto, L., 2011. Soil carbon
 stocks and stability across an altitudinal gradient in southern Peru. J. Soil Water Conserv. 66:
 213-220.

Sierra, J., Brisson, N., Ripoche, D., Déqué, M., 2010. Modelling the impact of thermal adaptation of soil microorganisms and crop system on the dynamics of organic matter in a tropical soil under a climate change scenario. Ecol. Model. 221: 2850-2858.

676 Sierra, J., Causeret, F., Chopin, P., 2017. A framework coupling farm typology and biophysical 677 modelling to assess the impact of vegetable crop-based systems on soil carbon stocks. 678 Application in the Caribbean. Agr. Syst. 153: 172-180. 679 Sierra, J., Causeret, F., Diman, J.L., Publicol, M., Desfontaines, L., Cavalier, A., Chopin, P., 2015. 680 Observed and predicted changes in soil carbon stocks under export and diversified 681 agriculture in the Caribbean. The case study of Guadeloupe. Agric. Ecosyst. Environ. 213: 252-682 264. 683 Tan, Q., Wang, G., 2016. Decoupling of nutrient element cycles in soil and plants across an altitude 684 gradient. Sci. Rep. 6: 34875. 685 Tashi, S., Singh, B., Keitel, C., Adams, M., 2016. Soil carbon and nitrogen stocks in forests along an 686 altitudinal gradient in the eastern Himalayas and a meta-analysis of global data. Glob. Change 687 Biol. 22: 2255-2268. 688 Wang, H.C., Chou, C.Y., Chiou, C.R., Tian, G., Chiu, C.Y., 2016. Humic acid composition and 689 characteristics of soil organic matter in relation to the elevation gradient of Moso Bamboo 690 plantations. PLoS ONE 11(9): e0162193. doi:10.1371/journal.pone.0162193. 691 Zimmermann, M., Bird, M.I., 2012. Temperature sensitivity of tropical forest soil respiration increase 692 along an altitudinal gradient with ongoing decomposition. Geoderma 187-188: 8-15. 693 Zimmermann, M., Meir, P., Bird, M.I., Malhi, Y., Ccahuana, A.J.Q., 2009. Climate dependence of 694 heterotrophic soil respiration from a soil-translocation experiment along a 3000-m tropical 695 forest altitudinal gradient. Eur. J. Soil Sci. 60: 895-906. 696 697 698

699	Figure captions
700	
701	Fig. 1. Location of the studied area in the southern part of Basse-Terre Island in the Guadeloupe
702	archipelago. The land occupied by andosols out of the area included in this study corresponds to the
703	rainforest.
704	
705	Fig. 2. Mean air temperature and rainfall across the altitudinal gradient. Vertical bars indicate the
706	standard deviation.
707	
708	Fig. 3. Soil organic C content across the altitudinal gradient under (a) banana monoculture and (b)
709	vegetable cropping systems.
710	
711	Fig. 4. Altitudinal gradient of (a) soil bulk density and (b) allophane content.
712	
713	Fig. 5. Changes across the altitudinal gradient in (a)(b) cation exchangeable capacity, and (c)(d) soil
714	pH, for banana monoculture and vegetable crops, respectively.
715	
716	Fig. 6. Altitudinal gradients of (a) banana yields and (b) the in situ rate constant of soil organic C
717	mineralisation (k_{soc}). k_{soc} was estimated for plots under banana monoculture.
718	
719	Fig. 7. Relationships between allophane content and (a) soil organic C and (b) the in situ rate
720	constant of soil organic C mineralisation (k _{soc}).
721	
722	



Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



Fig. 6.



Fig. 7.

Table 1

Mean values of the soil properties displaying no trend across the altitudinal gradient. CEC: cation exchangeable capacity. The values in brackets refer to the coefficient of variation expressed in %. For each property, mean values followed by different letters are significantly different at P <0.05.

Cropping system	рН	C/N	Extractable P	Exchangeable CEC				CEC	Base saturation
				Са	Mg	К	Na		
			mg kg⁻¹		cmol kg ⁻¹				%
Banana monoculture	5.8 (9)a	9.7 (10)a	30 (128)a	7.2 (72)a	1.9 (46)a	1.9 (61)b	0.1 (8)a	35 (20)a	32 (51)a
Vegetable crops	6.0 (9)a	9.9 (9)a	51 (52)b	8.4 (52)a	1.8 (47)a	1.1 (78)a	0.1 (6)a	34 (23)a	35 (47)a