



HAL
open science

Effect of natural antioxidants from grape seed and chestnut in combination with hydroxytyrosol, as sodium nitrite substitutes in Cinta Senese dry-fermented sausages

Chiara Aquilani, Francesco Sirtori, Mónica Flores, Riccardo Bozzi, Bénédicte Lebret, Carolina Pugliese

► To cite this version:

Chiara Aquilani, Francesco Sirtori, Mónica Flores, Riccardo Bozzi, Bénédicte Lebret, et al.. Effect of natural antioxidants from grape seed and chestnut in combination with hydroxytyrosol, as sodium nitrite substitutes in Cinta Senese dry-fermented sausages. *Meat Science*, 2018, 145, pp.389-398. 10.1016/j.meatsci.2018.07.019 . hal-02629320

HAL Id: hal-02629320

<https://hal.inrae.fr/hal-02629320v1>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Effect of natural antioxidants from grape seed and chestnut in combination with hydroxytyrosol, as sodium nitrite substitutes in Cinta Senese dry-fermented sausages

Chiara Aquilani, Francesco Sirtori, Monica Flores, Riccardo Bozzi, Benedicte Lebret, Carolina Pugliese



PII: S0309-1740(18)30453-4
DOI: doi:[10.1016/j.meatsci.2018.07.019](https://doi.org/10.1016/j.meatsci.2018.07.019)
Reference: MESC 7633
To appear in: *Meat Science*
Received date: 26 April 2018
Revised date: 13 July 2018
Accepted date: 13 July 2018

Please cite this article as: Chiara Aquilani, Francesco Sirtori, Monica Flores, Riccardo Bozzi, Benedicte Lebret, Carolina Pugliese, Effect of natural antioxidants from grape seed and chestnut in combination with hydroxytyrosol, as sodium nitrite substitutes in Cinta Senese dry-fermented sausages. *Mesc* (2018), doi:[10.1016/j.meatsci.2018.07.019](https://doi.org/10.1016/j.meatsci.2018.07.019)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comment citer ce document :

Aquilani, C., Sirtori, F., Flores, M., Bozzi, R., Lebret, B., Pugliese, C. (2018). Effect of natural antioxidants from grape seed and chestnut in combination with hydroxytyrosol, as sodium nitrite substitutes in Cinta Senese dry-fermented sausages. *Meat Science*, 145, 389-398. , DOI : [10.1016/j.meatsci.2018.07.019](https://doi.org/10.1016/j.meatsci.2018.07.019)

Effect of natural antioxidants from grape seed and chestnut in combination with hydroxytyrosol, as sodium nitrite substitutes in Cinta Senese dry-fermented sausages

Chiara Aquilani^{a,*} chiara.aquilani@unifi.it, Francesco Sirtori^a, Monica Flores^b, Riccardo Bozzi^a, Benedicte Lebret^c, Carolina Pugliese^a

^aDISPAA - Department of Agrifood Production and Environmental Sciences, Animal Science Section, Universita' degli Studi di Firenze, 50144 Firenze, Italy

^bIATA - Institute of Agrochemistry and Food Technology (IATA-CSIC), Paterna, Valencia, Spain

^cPEGASE, INRA, Agrocampus-Ouest, 35042 Rennes, France

*Corresponding author.

Abstract:

Dry-fermented pork sausages, from Cinta Senese local breed, were manufactured replacing sodium nitrite (NIT) with two mixtures of natural antioxidants consisting of: i) grape seed extract and olive pomace hydroxytyrosol (GSE); ii) chestnut extract and olive pomace hydroxytyrosol (CHE). The effects on physical-chemical, aromatic and sensory traits, as well as the microbiological safety, were tested. Nitrite replacement lowered the pH in GSE and CHE samples and resulted in several differences in physical traits between CHE and NIT samples. *Listeria monocytogenes*, *Salmonella* and *Clostridium botulinum* were not found in any samples. GSE and CHE mixtures showed a slightly lower antioxidant activity. Volatile profile showed a similar aromatic profile among the three treatments with differences mainly to abundance of the single compounds, indicating that replacement of nitrite by natural antioxidants did not affect the overall aroma profile, as outlined by olfactometry results. In addition, the replacement did not affect the overall acceptability, except for color-related traits, underscored in GSE and CHE products.

Keywords: Volatile compounds; meat quality; GC-olfactometry; local pig breed; lipid oxidation; pork products

1. Introduction

Dry cured meat products are typical of the Mediterranean area and they represent a high-value production in European countries, considering that curing process allows extension of meat shelf-life (Marco et al., 2006) and leads to typical pork products with specific eating quality and regional identity (Pugliese and Sirtori, 2012). In Southern Europe, salami and dry-fermented sausages, are generally characterized by slowly air-drying and mold-ripening (Flores, 1997). This curing process leads to peculiar characteristics and flavors that are widely appreciated by consumers; but is also related to longer curing times that may cause higher lipid oxidation levels. Moreover, natural fermentation, avoiding the addition of lactic acid-producing starter cultures, is more susceptible to the growth of harmful bacteria, such as *Listeria monocytogenes* or *Clostridium botulinum* (Lücke, 2000). Thus, to avoid a severe deterioration of nutritive and organoleptic attributes, as well as to ensure food safety, several synthetic food preservatives are commonly included. Among them, the most used are nitrites and nitrates (Hammes, 2012). Nitrite positively affects color, inhibits the growth of pathogenic bacteria, contributes to the development of typical cured meat flavor and delays oxidative rancidity (Marco et al., 2006). Despite their effectiveness as curing agents, the nitrite/nitrate intake represents a risk to human health, i.e. the formation of carcinogenic nitrosamines is one of the most current concerns (De Mey et al., 2017). Several studies have focused on nitrate/nitrite reduction or substitution (Purriños et al., 2013; Özvural and Vural, 2014; Pateiro et al., 2015), but the main issue remains finding an alternative able to address the multiple activities they perform. Up until now, most of the alternatives proposed are plant extracts, largely obtained from agricultural by-products. These compounds are very rich in polyphenols, flavonoids and terpenoids and are able to perform a double antioxidant-antimicrobial functions (Falowo et al., 2014; Hygreeva et al., 2014; Shah et al., 2014). These compounds might also constitute a great opportunity to exploit agricultural by-products, which otherwise would be wasted. The aim of this study was to assess the feasibility of producing dry-fermented sausages by replacing sodium nitrite

with natural antioxidants while trying to maintain quality traits. Grape seed extract, chestnut extract and hydroxytyrosol (extracted from defatted olive pomace), were chosen due to their great availability as by-products of important Tuscan agricultural products. Moreover, among the investigated plant extracts, they have shown an interesting potential both for antioxidant activity and microbial inhibition. This innovation also aimed to valorize Cinta Senese, a local pig breed strongly linked to the Tuscan region.

2. Materials and methods

2.1. Antioxidant mixtures

The natural antioxidants employed in the present studies were provided by Phytolab (Sesto Fiorentino, Florence, Italy). They consisted of grape seed and chestnut extracts, tocopherol and hydroxytyrosol (extracted by defatted olive pomace). The manufacturer provided the phenolic profile (Table 1), total phenolic content and antiradical scavenging activity (EC_{50}) (Table 2) of each extract. The grape seed and chestnut extracts were combined with the same amount of hydroxytyrosol and tocopherol to form two different mixtures; grape seed (GSE) and chestnut (CHE) mixtures.

2.2. Sausages manufacturing

In an industrial plant (Azienda Agricola Savigni, Pistoia, Italy), 24kg of pork lean and 6kg of subcutaneous backfat from Cinta Senese pig breed were minced and equally divided in three batches. Salt (23g/kg), sucrose (35g/kg) and black pepper (0.2g/kg) were added to each batch following the recipe traditionally used by the manufacturer. Thirty ppm of sodium nitrite (E250) were added to the first batch to constitute the control (NIT). In second batch, 10g/kg of GSE mixture were used to replace sodium nitrite, while 10g/kg of CHE were added to the third batch. Sausages were weighed, dried at 28°C and RH 85% for 4 days and then ripened 21 days (T 13°C, RH 70%). Once ripened, six samples of each batch were collected, pH, color, and processing loss were immediately measured. Samples were vacuum packed and stored at -80°C for physical,

chemical and aromatic analysis. Another 3 samples of each batch were stored at 4°C to be employed for sensory analysis the following day. This design was replicated to have two totally independent batches for each treatment.

2.3. Physical, chemical and microbiological parameters

At the end of ripening, physical parameters were assessed on 12 samples of each batch (6 for each replication). Sausage pH was measured at room temperature (20 °C) using a pH meter Crison GLP21 (Barcelona, Spain), the instrument was introduced in a sausage portion. Color (L^* , a^* and b^*) was determined by a Minolta Chromameter CR-200 (Tokyo, Japan) immediately after slicing. a_w was measured following the method ISO 21807:2004. Two 10 mm-thick and 10 mm-width slices of each sample, were cut and immediately analyzed at room temperature (22 °C), using a Zwick Roell Z2.5 apparatus (Ulm, Germany) with a loading cell of 1 kN at the crosshead speed of 1 mm/sec. Texture profile analysis (TPA) was performed assessing the following parameters: hardness, cohesiveness, gumminess, springiness and chewiness. Moisture was determined by lyophilizing to constant weight 40g of sample, according to AOAC methods (1990). Weight loss was measured as the difference between weight at time zero and end of ripening (after 24 days). Total protein, fat and ash contents were determined following AOAC (1990) methods. Lipid oxidation was determined according Vyncke (1970), using a PerkinElmer Lambda EZ150 spectrophotometer (Waltham, MA, USA). Results were expressed as mg of malondialdehyde (MDA)/kg of samples. Fatty acids were determined using a Varian GC-430 apparatus equipped with a flame ionization detector (FID) (Palo Alto, CA, USA) as reported by Sirtori et al. (2015). The individual methyl esters were identified by their retention time using an analytical standard (F.A.M.E. Mix, C8-C22 Supelco 18920-1AMP). Response factors based on the internal standard (C19:0) were used for quantification and results were expressed as mg/100g of sample. The fatty acid content was reported as saturated (SFA), monounsaturated (MUFA) and polyunsaturated (PUFA) fatty acids. Microbiological analyses were carried out in an external accredited laboratory

to determine the products' safety. The following bacteria were investigated: *Escherichia coli* (ISO 16649-2:2001), *Listeria monocytogenes* (UNI EN ISO 11290-1:2005), coagulase positive *Staphylococcus* spp. (UNI EN ISO 6888-1:2004), *Clostridium botulinum* (ISO 15213:2003) and *Salmonella* spp. (UNI EN ISO 6579:2008).

2.4. Volatile compounds analysis

2.4.1. Gas chromatography-mass spectrometry analysis (GC-MS)

Solid-phase microextraction (SPME) and GC-MS analysis were performed following the method described by Corral, Salvador, & Flores (2013) using a 85 μm Carboxen/Polydimethylsiloxane (CAR/PDMS) fiber (Supelco, Bellefonte, PA) installed in a Gerstel MPS2 multipurpose sampler (Gerstel, Germany) and an Agilent HP 7890 series II GC with an HP 5975C mass selective detector (Hewlett-Packard Palo Alto, CA, USA). The volatile compounds (VOCs) detected were identified by comparison with mass-spectra from the library database (Nist'05), linear retention index (van Den Dool and Dec. Kratz, 1963) and by comparison with authentic standards. The quantification of volatile compounds was done in SCAN mode using either total or extracted ion chromatogram (TIC or EIC) on an arbitrary scale.

2.4.2. Gas chromatography-olfactometry analysis (GC-O)

A gas chromatograph (Agilent 6890, USA) equipped with an FID detector and sniffing port (ODP3, Gerstel, Mülheim an der Ruhr, Germany) was used to analyze aroma compounds extracted by SPME as described by Corral et al. (2013). The detection frequency (DF) method was used to estimate the aromatic impact of each volatile and each assessment was carried out according to Olivares, Navarro, & Flores (2011). Four trained panelists evaluated the odors from the GC-effluent. Each assessor evaluated 3 sausages for a total of 12 assessments, the final DF was obtained by summing the 12 sniffings. The detection of an odor by less than three assessors was considered noise. Compounds were identified by comparison with mass spectra, with linear retention indices

of authentic standards injected in the GC-MS and GC-O, and by coincidence of the assessors' descriptors with those reported by Burdock (2010).

2.5. Sensory analysis

Sensory analysis was carried out in an equipped laboratory by 8 trained panelists using a quantitative-descriptive analysis method. Fourteen attributes (grease appearance, abnormal colors, firmness, color uniformity, redness, cured meat flavor, off odor, salty, rancid, off flavor, hardness, juiciness, aftertaste, general acceptability) were evaluated, each attribute was scored in a 10 cm non-structured line (Pugliese et al., 2010). Select subjects underwent an introductory session, where the testing procedures and the chosen sensory traits were discussed using two types of comparable commercial products. During three sessions, panelists evaluated a total of 9 sausages (3 samples x 3 treatments) identified by an alphanumeric code. The sausages were divided in 0.5cm-thick x 2cm-diameter slices and two slices of each samples were randomly served to judges at room temperature (20°C). Panelists were invited to eat a cracker and drink a glass of water between samples.

2.6. Statistical analysis

Data were analyzed by SAS software. Two-way ANOVAs were performed on physical and chemical data according to the following model:

$$Y_{ijk} = \mu + T_i + B_j + \varepsilon_{ijk}$$

Where μ is the mean, T is the i^{th} treatment, B is the j^{th} batch and ε is the error. For sensory data, effect of panelist was included in the previous model. The interaction between Treatment and Batch factors was tested but being not significant, it was not included in the model.

Volatile compounds data were also analyzed by a multivariate approach to determine the presence of characteristic compounds able to be allocated to samples among different treatments. A stepwise discriminant analysis (SDA) was first used to reduce the space-variables, selecting the subset of variables that better discriminated groups. Canonical discriminant analysis (CDA) were performed

using SDA selected variables, resulting in 2 new variables, called canonical functions (CAN1, CAN2). They consisted of a series of canonical coefficients (CC) that indicate the partial contribution of each variable in composing the CANs. The greater the CC, the more the variable contributes to CAN composition.

3. Results and discussion

3.1. Physical, chemical and microbiological parameters

The major foodborne pathogens (*Escherichia coli*, *Listeria monocytogenes*, *Staphylococcus* spp., *Clostridium* spp and *Salmonella* spp.) were absent or below the limit required (Reg CE/2073/05) in all samples (Table 3). Several studies on plant phenolic suggest these components have antimicrobial activity (Kao et al., 2010; Mujić et al., 2014; Fasolato et al., 2016). Further studies are, however, required to determine the effectiveness of the studied antioxidants against the development of the main foodborne pathogens. The a_w values (Table 4) recorded for GSE and CHE, being below 0.89, contributed to control pathogenic organisms development (Toldrá and Flores, 2014). Moisture, fat, protein and ash contents and weight loss were not affected by treatment and they were in line with values reported for dry-cured sausages (Olivares, Navarro, & Flores, 2015; Ribas-Agustí et al., 2014; Škrlep, Čandek-Potokar, Tomažin, Batorek Lukač, & Flores, 2017), except for weight loss, which was slightly greater in the present study; likely the smaller diameter of samples could have enhanced the water loss during ripening. The pH observed was in the range reported for natural fermented meat products at a similar curing time (Özvural and Vural, 2014; Škrlep et al., 2017); indeed, they were characterized by a higher pH compared to commercial products (Hospital et al., 2015; Montanari et al., 2018). Moreover, significant differences among treatments were found for pH, with lower values observed for GSE and CHE samples, suggesting that *Lactobacillus* (LAB) growth, that takes place during the first fermentation phase, could be slightly promoted in these products. Indeed, low or nitrites-free sausages showed an increased presence of LAB (Hospital et al., 2015). Growth of LAB was not, however, assessed in the present

experiment, and further studies will be required to assess effects of GSE and CHE. Concerning color attributes, L^* was not affected by treatment, a^* showed significant greater values in GSE and NIT samples than in CHE ones, while b^* was significant higher in NIT compared to the modified products. A change in a^* was expected considering the role nitrites play in nitrosomyoglobin formation, the characteristic red curing pigment (Hammes, 2012). Since neither chemical composition nor oxidation resulted in significant differences among groups, a different pathway for red color formation in GSE samples should be considered. A stable red color compound called Zn-protoporphyrin, derived from the substitution of heme iron with zinc, has been observed in Parma ham, an Italian nitrite/nitrate-free ham (Wakamatsu et al., 2004a). Up to now, mechanisms leading to its formation are not well-known (Hammes, 2012), but the absence of nitrites, low levels of oxygen, meat endogenous enzymes as well as microorganism, are all factors that may contribute to its formation (Wakamatsu et al., 2004b). Apart from the absence of nitrites, some compounds contained in grape seed extract may have promoted the Zn-protoporphyrin formation for the GSE group, while no formation occurred in the CHE samples. These results are partially in agreement with those reported by Lorenzo, González-Rodríguez, Sánchez, Amado, & Franco (2013), although their study was conducted on chorizo where pigmentation due to added paprika may have interfered.

According to TPA results (Table 5), cohesiveness, springiness and chewiness were affected by nitrite replacement, being highest in CHE samples, lowest in NIT, whereas GSE samples were similar to both. Since no differences in moisture and weight loss were found among the treatments, the results obtained were attributable basically to the differences in pH, which, declining, causes the aggregation of myofibrillar proteins and leads to gel formation (Lücke, 2000). The higher pH of NIT samples likely inhibited this phenomenon thus reducing the sausage cohesiveness and chewiness. The results reported are partially in agreement with Lorenzo et al., (2013), who also noticed the highest chewiness values for chorizo with added chestnut extract and ripened for 19

days, compared to the same product manufactured with GSE or synthetic antioxidant (BHT in this case).

The groups did not differ in SFA, MUFA and PUFA contents. Their relative amounts reflect those of fresh pork composition, slightly richer in MUFA than SFA, with PUFA being approximately a third of either SFA or MUFA categories (Škrlep et al., 2017). PUFA can also be considered an indicator of meat oxidative status, due to their double bonds being preferred substrates for oxidative reactions (Pateiro et al., 2015). Results suggest that the natural antioxidants employed were as effective as nitrites in control lipid oxidation during manufacturing and ripening. This is supported by TBARS results, showing no significant differences among treatments, however, further studies will be required to evaluate the antioxidant activity during the shelf-life. Nitrites exert their antioxidant activity in cured meat by forming the myoglobin-stable compounds and making the iron inaccessible for oxidation (Riazi et al., 2018); phenolic compounds instead, follow different pathways, acting as hydrogen donors. The phenolic hydroxyl groups intercept the free radicals to form stable end-products, interrupting and avoiding further lipid oxidation, especially of unsaturated FAs (Jayaprakasha et al., 2003). To the best of our knowledge, few data are available about dry-fermented pork sausages with added natural extracts, but the great variability of these traditional products makes comparisons difficult. The efficacy of hydroxytyrosol in preventing lipid oxidation was reported by Cofrades et al., (2011) for n-3 enriched frankfurters, while Lorenzo et al., (2013) observed comparable TBARS values in Spanish chorizo with added BHT, grape seed extract or chestnut extract.

3.2. Volatile profile and olfactometry

Ninety-one VOCs were identified by HS-SPME-GS-MS (Table 6). The most abundant groups originated from spices (51-61%) and carbohydrate fermentation (30-39%), followed by amino acid degradation (6-7%), while VOCs derived from lipid β -oxidation and lipid oxidation processes represented 1% of total extracted areas.

Among the 14 VOCs related to lipid auto-oxidation, 5 showed significant differences with NIT resulting in the lowest, while GSE and CHE products showed intermediate or higher abundances. These VOCs originate by autocatalytic fat oxidation and involves mostly unsaturated fatty acids. Among the identified VOCs, hexanal is of key-importance to better outline the products' oxidation status. The correlation between this compound and lipid oxidation is well-known and its low perception threshold makes hexanal an important contributor to overall aroma (Marco et al., 2006). The higher hexanal content in GSE and CHE samples than NIT is consistent with TBARs results, suggesting greater PUFA oxidation, even if the extent was limited and did not affect the parameters previously examined. The total lipid auto-oxidation values confirmed these differences, with NIT being the lowest, GSE the highest and CHE similar to both, likely related to a higher EC₅₀ for GSE. Hence, even though the phenolic extracts used contributed to maintain lipid oxidation below the perception threshold of rancid flavor, they appeared less effective than nitrites in controlling lipid oxidation. Partially in agreement with this, Purriños et al. (2013) reported grape seed was a less effective antioxidant in chorizo, but on the contrary, chestnut extract was found to have higher antioxidant activity than BHT.

Spice derived VOCs were the major group, due to the use of black pepper. Indeed, limonene, a compound particularly abundant in pepper (Moretti et al., 2004) represented approximately half of the total amount for each treatment. As the same amount of pepper was added to all treatments, differences among the 3 groups might be due to several external factors such as an irregular distribution of the ground pepper in the raw matrix (Montanari et al., 2018), as well as a heterogeneity of the spices themselves caused by different grinding techniques and/or storage times that could have led to differential oxidative status, odorant losses and sensory attribute changes (Orav et al., 2004; Liu et al., 2013).

Microbial enzymes degrade free fatty acids through β -oxidation reactions, generating methyl ketones as final products (Flores & Olivares, 2007). Two VOCs belonging to this group were

identified, but significant differences were found only for total abundance, which was highest for NIT, the lowest in GSE and CHE samples were similar to both. Since the main microbial populations were not examined in this study, the differences in volatile development due to microbial fermentation cannot be explained directly. However, the role of genus staphylococcus in incomplete β -oxidation are well-known, so the presence of 2-pentanone and 3-octanone were likely related to the presence of these bacteria (Chen et al., 2017).

Contrarily to Chen et al., (2017) and Marco et al. (2006), only one ester was observed. However, they worked on fermented sausages manufactured with starter cultures containing staphylococci strains. Staphylococci promote the formation of esters with staphylococci esterase activity being one of the main factors leading to ester formation in dry-fermented sausages (Wang et al., 2016).

Among VOCs generated by bacterial metabolism, products of carbohydrate fermentation form an important group, consisting in 13 identified compounds. Acetaldehyde, 2,3-butanedione and 2,3-butanediol were higher in GSE samples than in CHE and NIT; ethyl alcohol and butanoic acid were lowest in NIT samples, while 3-hydroxy-2-butanone was lowest in CHE samples. VOCs from carbohydrate metabolism were generally the highest for GSE. The great abundance of 2-butanone is remarkable, considering that this compound is known as a by-product LAB metabolism (Montanari et al., 2018), originating from 2,3-butanedione (significantly higher in GSE). Then, 2-butanone is reduced to 2-butanol and 3-hydroxy-2-butanone, that again, was significantly higher in GSE samples and preferentially formed in small diameter sausages (Montanari et al., 2018).

The last group of VOCs related to bacterial metabolism consisted of 19 VOCs from amino acid degradation. 2-methylpropanal and ethylbenzene were higher in CHE samples, while the lowest amounts of 3-methylbutanol and 2-acetyl-1-pyrroline were observed for natural antioxidant products and benzeneacetaldehyde and benzylalcohol were higher in GSE samples than for CHE and NIT. The highest total amino acid degradation products were observed in NIT samples, followed by CHE and then GSE. This was likely due to 3-methylbutanol, whose content almost

doubled in NIT samples while toluene, the most abundant compound, was similar for the three groups. The compounds observed are characteristic of dry-fermented sausages, being reported by several authors (Marco et al., 2006; Corral et al., 2013; Škrlep et al., 2017).

Despite the variability within each group of VOCs, total amounts of microbial metabolites suggest a greater development of microflora in NIT samples that might be related to the antimicrobial activity of phenolic extracts during ripening. Several studies have reported phenolic compounds diffuse into bacterial cells walls and interact with cytoplasmatic proteins, affecting Gram positive bacteria and, particularly, Gram positive cocci (Jayaprakasha et al., 2003; Fasolato et al., 2016; Riazi et al., 2018). It is worth noting that the main populations involved in sausage fermentation processes, LAB and Staphylococci, are both Gram positive bacteria.

The role each compound plays in defining the aromatic profile strongly depends on its abundance and on its perception threshold (Olivares et al., 2015). During the GC-O sessions, 31 aroma notes were perceived by trained assessors (Table 7). Seven aroma compounds were associated with spices, 1 to lipid beta-oxidation, 4 to carbohydrate fermentation, 1 to esterase activity, 11 to amino acid degradation, 4 to lipid auto-oxidation, 2 to unknown compounds (not identified with any of the VOCs identified by GC-MS) and 1 to contaminants. Considering their DF value as an aroma impact index, 11 VOCs had a DF higher than 8. Despite the differences outlined by SPME-GC-MS analysis, panelists did not detect any differences in the olfactometric profile of the three groups. This is likely because all the identified VOCs were observed in the three groups and the main differences among GSE, CHE and NIT samples were attributable only to differences in the single compounds abundances. As a consequence, GC-O data were displayed combined in a single aromatic profile (Table 7). Most of the identified compounds were previously observed as recurrent in dry-fermented sausages (Flores & Olivares, 2015; Söllner and Schieberle, 2009; Schmidt and Berger, 1998). Amino acid degradation compounds have a key-role in flavor development, contributing with malty, fruity, sweaty flavors and ripened aroma (Hospital et al., 2015; Chen et al.,

2017). Indeed, more than half of SPME-GC-MS identified VOCs were also observed in GC-O sessions as odor active compounds in Cinta Senese dry-fermented sausages. They accounted for one third of the compounds forming the GC-O profile and had high DF values. Among them, 2 acetyl-1-pyrroline and methional are considered as the most potent odor active compounds in dry-fermented sausages (Corral, Leitner, Siegmund, & Flores, 2016; Söllner and Schieberle, 2009). Also 3-methylbutanoic acid is considered a potent aroma contributor, giving cheesy, lactic and fatty notes (Flores & Olivares 2015), while 2,5-dimethylpyrazine is related to meaty and cooked potatoes notes.

The second most represented VOC group were spice-derived, especially α -terpinene, β -myrcene and terpinolene, which were previously reported as odor active compounds (Olivares et al., 2015; Schmidt & Berger, 1998). Another important group was composed of lipid oxidation VOCs, among them hexanal was the most potent odorant (Marco, Navarro, & Flores, 2007; Olivares et al., 2015; Söllner and Schieberle, 2009; Schmidt and Berger, 1998). Indeed, hexanal produces fresh and green notes (Table 7), but it turns to rancid notes as its abundance increases. Lastly, also 2,3-butanedione, derived from carbohydrate fermentation, due to its low threshold (about 4 μ g/l) (Hospital et al., 2015), was an important aroma contributor, characterized by buttery-sweet notes and a DF of 10.

Figure 1 displays the 19 compounds identified by SDA. The selected compounds were able to discriminate the three treatments. Can1 accounted for a great part of variance, separating CHE from GSE and NIT, while Can 2 sharply divided GSE and CHE from NIT. Multivariate analysis showed how lipid auto-oxidation compounds, comprising half of the compounds identified by SDA, were central in differentiating the three groups. Tetradecane, ethylbenzene and 3-octanone having the greatest negative Can1 scores, were considered mainly responsible for separating GSE and NIT from CHE, in agreement with ANOVA results. Concerning Can2, compounds with higher weighing were decane and octane for GSE and CHE, while 3-methylbutanol seems to characterize NIT samples. Among the 19 compounds identified, 8 were also perceived by GC-O panelists. However,

considering their DF and CCs together, only hexanal and 2.5-dimethylpyrazine might have an effective potential in discriminating the groups also from a sensory point of view.

3.3. Sensory analysis

Figure 2 shows sensory results. Abnormal colors, off-flavors, off-odors and rancid were scored as 0 (not present) and were not shown. As expected, the most affected traits were color related, with both color uniformity ($P < 0.01$) and redness ($P < 0.01$) scoring lower for GSE and CHE samples compared to NIT ones, while firmness was scored highest for CHE samples ($P < 0.05$), agreeing with TPA results. Likely, the lower moisture and fat contents observed in CHE could have affected the firmness, even if neither moisture nor fat, significantly differed among treatments. Despite the differences in lipid auto-oxidation VOCs, no perceivable rancid notes were detected by panelists, as the level of MDA found in the samples did not exceed the organoleptic perception of lipid oxidation (Campo et al., 2006). Effects of adding grape seed and chestnut extracts to dry-fermented sausages have not, however, always been positive. Ribas-Agustí et al. (2014) reported that panelists discarded grape seed extract added products, as there were judged to be abnormal compared to control samples. Similarly, Özvural and Vural (2011) observed a decrease in overall acceptability of frankfurters with grape seed extract added, even when products manufactured with concentrations lower than 0.05% resulted in scores similar to control.

4. Conclusions

The results on VOCs profiles suggested a greater antimicrobial activity of natural antioxidant mixtures (GSE and CHE) compared to sodium nitrite (NIT), likely due to their phenolic constituents; further none of the main foodborne pathogens were found in any sample. No significant differences among treatments were found for lipid oxidation, even if lipid auto-oxidation VOCs suggested a slightly lower antioxidant activity of GSE and CHE compared to sodium nitrite. Despite the differences in single VOCs abundances, the replacement did not affect the overall aroma profile, as outlined by GC-O results and sensory analysis. Some differences in instrumental

color and texture negatively affected GSE and CHE products, but the overall acceptability was not influenced. GSE and CHE effects on microbiota in dry-fermented sausages should be studied in depth, however, the results so far indicated that tested antioxidants are valid alternatives to sodium nitrite in Cinta Senese dry-fermented sausages.

5. Acknowledgments

The research was carried out with funds from the European Union's Horizon 2020 research and innovation program under grant agreement No 634476 (acronym TREASURE). The content of this work reflects only the author's view and the European Union Agency is not responsible for any use that may be made of the information it contains. We acknowledge Prof. Romani and Phytolab for providing the antioxidant mixtures. Authors also acknowledge the financial support of AGL2015-64673-R (MINEICO, Spain) and FEDER funds.

6. References

- Burdock GA 2010. Fenaroli's handbook of flavor ingredients.
- Campo MM, Nute GR, Hughes SI, Enser M, Wood JD and Richardson RI 2006. Flavour perception of oxidation in beef. *Meat Science* 72, 303–311.
- Chen Q, Kong B, Han Q, Xia X and Xu L 2017. The role of bacterial fermentation in lipolysis and lipid oxidation in Harbin dry sausages and its flavour development. *LWT - Food Science and Technology* 77, 389–396.
- Cofrades S, Salcedo Sandoval L, Delgado-Pando G, López-López I, Ruiz-Capillas C and Jiménez-Colmenero F 2011. Antioxidant activity of hydroxytyrosol in frankfurters enriched with n-3 polyunsaturated fatty acids. *Food Chemistry* 129, 429–436.
- Corral S, Leitner E, Siegmund B and Flores M 2016. Determination of sulfur and nitrogen compounds during the processing of dry fermented sausages and their relation to amino acid

generation. *Food Chemistry* 190, 657–664.

Corral S, Salvador A and Flores M 2013. Salt reduction in slow fermented sausages affects the generation of aroma active compounds. *Meat Science* 93, 776–785.

van Den Dool H and Dec. Kratz P 1963. A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *Journal of Chromatography A* 11, 463–471.

Falowo AB, Fayemi PO and Muchenje V 2014. Natural antioxidants against lipid-protein oxidative deterioration in meat and meat products: A review. *Food Research International* 64, 171–181.

Fasolato L, Carraro L, Facco P, Cardazzo B, Balzan S, Taticchi A, Andreani NA, Montemurro F, Martino ME, Di Lecce G, Toschi TG and Novelli E 2016. Agricultural by-products with bioactive effects: A multivariate approach to evaluate microbial and physicochemical changes in a fresh pork sausage enriched with phenolic compounds from olive vegetation water. *International Journal of Food Microbiology* 228, 34–43.

Flores J 1997. Mediterranean vs northern European meat products. Processing technologies and main differences. *Food Chemistry* 59, 505–510.

Flores M and Olivares A 2015. 25 25.1 Flavors.

Flores M, Olivares A, Chen M, Tu R and Wang S 2015. *Handbook of Fermented Meat and Poultry*. Blackwell Publishing Ltd, Oxford, UK.

Hammes WP 2012. Metabolism of nitrate in fermented meats: The characteristic feature of a specific group of fermented foods. *Food Microbiology* 29, 151–156.

Hospital XF, Carballo J, Fernández M, Arnau J, Gratacós M and Hierro E 2015. Technological implications of reducing nitrate and nitrite levels in dry-fermented sausages: Typical microbiota, residual nitrate and nitrite and volatile profile. *Food Control* 57, 275–281.

- Hygreeva D, Pandey MC and Radhakrishna K 2014. Potential applications of plant based derivatives as fat replacers, antioxidants and antimicrobials in fresh and processed meat products. *Meat Science* 98, 47–57.
- Jayaprakasha GK, Selvi T and Sakariah KK 2003. Antibacterial and antioxidant activities of grape (*Vitis vinifera*) seed extracts. *Food Research International* 36, 117–122.
- Kao TT, Tu HC, Chang WN, Chen BH, Shi YY, Chang TC and Fu TF 2010. Grape seed extract inhibits the growth and pathogenicity of *Staphylococcus aureus* by interfering with dihydrofolate reductase activity and folate-mediated one-carbon metabolism. *International Journal of Food Microbiology* 141, 17–27.
- Liu H, Zeng F, Wang Q, Ou S, Tan L and Gu F 2013. The effect of cryogenic grinding and hammer milling on the flavour quality of ground pepper (*Piper nigrum* L.). *Food Chemistry* 141, 3402–3408.
- Lorenzo JM, González-Rodríguez RM, Sánchez M, Amado IR and Franco D 2013. Effects of natural (grape seed and chestnut extract) and synthetic antioxidants (butylatedhydroxytoluene, BHT) on the physical, chemical, microbiological and sensory characteristics of dry cured sausage ‘chorizo’. *Food Research International* 54, 611–620.
- Lücke FK 2000. Utilization of microbes to process and preserve meat. *Meat Science* 56, 105–115.
- Marco A, Navarro JL and Flores M 2006. The influence of nitrite and nitrate on microbial, chemical and sensory parameters of slow dry fermented sausage. *Meat Science* 73, 660–673.
- Marco A, Navarro JL and Flores M 2007. Quantitation of selected odor-active constituents in dry fermented sausages prepared with different curing salts. *Journal of Agricultural and Food Chemistry* 55, 3058–3065.
- De Mey E, De Maere H, Paelinck H and Fraeye I 2017. Volatile N-nitrosamines in meat products:

Potential precursors, influence of processing, and mitigation strategies. *Critical Reviews in Food Science and Nutrition* 57, 2909–2923.

Montanari C, Gatto V, Torriani S, Barbieri F, Bargossi E, Lanciotti R, Grazia L, Magnani R, Tabanelli G and Gardini F 2018. Effects of the diameter on physico-chemical, microbiological and volatile profile in dry fermented sausages produced with two different starter cultures. *Food Bioscience* 22, 9–18.

Moretti VM, Madonia G, Diaferia C, Mentasti T, Paleari MA, Panseri S, Pirone G and Gandini G 2004. Chemical and microbiological parameters and sensory attributes of a typical Sicilian salami ripened in different conditions. *Meat Science* 66, 845–854.

Mujić I, Rudić D, Živković J, Jukić H, Jug T, Nikolić G and Trutić N 2014. Antioxidant and antibacterial properties of castanea sativa mill. Catkins extracts.

Olivares A, Navarro JL and Flores M 2011. Effect of fat content on aroma generation during processing of dry fermented sausages. *Meat Science* 87, 264–273.

Olivares A, Navarro JL and Flores M 2015. Characterization of volatile compounds responsible for the aroma in naturally fermented sausages by gas chromatography-olfactometry. *Food Science and Technology International* 21, 110–123.

Orav A, Stulova I, Kailas T and Müürisepp M 2004. Effect of Storage on the Essential Oil Composition of *Piper nigrum* L. Fruits of Different Ripening States. *Journal of Agricultural and Food Chemistry* 52, 2582–2586.

Özvural EB and Vural H 2011. Grape seed flour is a viable ingredient to improve the nutritional profile and reduce lipid oxidation of frankfurters. *Meat Science* 88, 179–183.

Özvural EB and Vural H 2014. Which is the best grape seed additive for frankfurters: Extract, oil or flour? *Journal of the Science of Food and Agriculture* 94, 792–797.

- Pateiro M, Bermúdez R, Lorenzo J and Franco D 2015. Effect of Addition of Natural Antioxidants on the Shelf-Life of ‘Chorizo’, a Spanish Dry-Cured Sausage. *Antioxidants* 4, 42–67.
- Pugliese C and Sirtori F 2012. Quality of meat and meat products produced from southern European pig breeds. *Meat Science*.
- Pugliese C, Sirtori F, Adorante SD, Parenti S, Rey A, Lopez-bote C and Franci O 2010. Effect of pasture in oak and chestnut groves on chemical and sensorial traits of cured lard of Cinta Senese pigs. *Italian Journal Of Animal Science* 8, 131–142.
- Purriños L, García Fontán MC, Carballo J and Lorenzo JM 2013. Study of the counts, species and characteristics of the yeast population during the manufacture of dry-cured ‘lacón’ Effect of salt level. *Food Microbiology* 34, 12–18.
- Riazi F, Zeynali F, Hoseini E, Behmadi H and Savadkoobi S 2018. Corrigendum to ‘Oxidation phenomena and color properties of grape pomace on nitrite-reduced meat emulsion systems’ [*Meat Sci.* (121) (2016) 350–358], (S0309174016302182), (10.1016/j.meatsci.2016.07.008). *Meat Science* 135, 189.
- Ribas-Agustí A, Gratacós-Cubarsí M, Sárraga C, Guàrdia MD, García-Regueiro JA and Castellari M 2014. Stability of phenolic compounds in dry fermented sausages added with cocoa and grape seed extracts. *LWT - Food Science and Technology* 57, 329–336.
- Schmidt S and Berger RG 1998. Aroma Compounds in Fermented Sausages of Different Origins. *LWT - Food Science and Technology* 31, 559–567.
- Shah MA, Bosco SJD and Mir SA 2014. Plant extracts as natural antioxidants in meat and meat products. *Meat Science* 98, 21–33.
- Sirtori F, Crovetto A, Acciaioli A, Bonelli A, Pugliese C, Bozzi R, Campodoni G and Franci O 2015. Effect of Replacing a Soy Diet with *Vicia Faba* and *Pisum Sativum* on Performance, Meat

and Fat Traits of Cinta Senese Pigs. *Italian Journal of Animal Science* 14, 3659.

Škrlep M, Čandek-Potokar M, Tomažin U, Batorek Lukač N and Flores M 2017. Properties and aromatic profile of dry-fermented sausages produced from Krškopolje pigs reared under organic and conventional rearing regime. *Animal*, 1–8.

Söllner K and Schieberle P 2009. Decoding the key aroma compounds of a Hungarian-type salami by molecular sensory science approaches. *Journal of Agricultural and Food Chemistry* 57, 4319–4327.

Toldrá F and Flores M 2014. SAUSAGES, TYPES OF | Dry and Semidry. In *Encyclopedia of Meat Sciences*, pp. 248–255. Elsevier.

Vyncke W 1970. Direct Determination of the Thiobarbituric Acid Value in Trichloroacetic Acid Extracts of Fish as a Measure of Oxidative Rancidity. *Fette, Seifen, Anstrichmittel* 72, 1084–1087.

Wakamatsu J, Nishimura T and Hattori A 2004a. A Zn-porphyrin complex contributes to bright red color in Parma ham. *Meat Science* 67, 95–100.

Wakamatsu J, Okui J, Ikeda Y, Nishimura T and Hattori A 2004b. Establishment of a model experiment system to elucidate the mechanism by which Zn-protoporphyrin IX is formed in nitrite-free dry-cured ham. *Meat Science* 68, 313–317.

Wang Y, Li Y, Yang J, Ruan J and Sun C 2016. Microbial volatile organic compounds and their application in microorganism identification in foodstuff. *TrAC - Trends in Analytical Chemistry* 78, 1–16.

Table 1. Phenolic profile of olive pomace and defatted grape seed and chestnut extracts.

	Olive Pomace (g/L) (hydroxytyrosol)	Grape Seed (mg/g)	Chestnut (mg/g)
Hydroxytyrosol	11.65	Gallic acid 0.01	Vescalin 9.34
Tyrosol & hydroxytyrosol derived compounds	15.13	Catechin B3 (dimers) 2.22	Castalin 8.99

Verbascosid	5.84	Catechin	11.07	Pedunculagin I	3.88
		Catechin (trimers)	3.21	Monogalloil glucose I	3.58
		Catechin B6 (dimers)	2.61	Gallic acid	18.50
		Catechin B2 (dimers)	5.37	Monogalloil glucose II	2.73
		Epicatechin	13.62	Roburin D	10.51
		Catechina trimer	3.71	Vescalagin	32.15
		Epicatechin gallate (PM 730)	6.65	C-glucoside tergallic dehydrate	2.73
		Epicatechin gallate (PM 442)	6.10	Castalagin	31.03
		Oligomers (tetramers)	54.88	Digalloil glucose I	10.03
		Epicatechin gallate (PM 882)	180.65	Digalloil glucose II	2.09
		Epicatechin gallate oligomers (trimers)	382.97	Hydrolyzable tannin m/z 1085	8.05
		Epicatechin gallate oligomers (trimers)	149.66	Trigalloil glucose I	4.61
				Trigalloil glucose II	6.74
				Tetragalloil glucose	2.05
				Ellagic acid	4.08

Table 2. Total phenolic content and radical scavenging activity of natural antioxidant constituting the mixtures

	Total phenolic content	Antiradical scavenging activity (EC ₅₀)
Grape seed extract	822.709 (mg/g)	0.147
Chestnut extract	161.091 (mg/g)	0.085
Olive pomace (hydroxytyrosol)	32.62 (g/l)	0.196
α -tocopherol	-	0.184

Table 3. Microbiological safety parameters on Cinta Sense dry-fermented sausages manufactured with natural antioxidant (GSE=grape seed extract; CHE=chestnut extract) as replacement of sodium nitrite (NIT).

	GSE	CHE	NIT
<i>Escherichia coli</i>	<10	<10	<10
<i>Listeria monocytogenes</i>	-	-	-
Coagulase positive <i>Staphilococcus spp.</i>	<10	<10	<10
Sulfite-reducing bacteria (<i>Clostridium botulinum</i>)	<10	1.2 10 ²	0.7 10 ²
<i>Salmonella spp.</i>	-	-	-

Results are expressed as ufc/g; the symbol “-“ indicates that the organism was not present

Table 4. Physical and chemical parameters on Cinta Sense dry-fermented sausages manufactured with natural antioxidant as replacement of sodium nitrite.

	GSE	CHE	NIT	SEM ^a	P ^b
--	-----	-----	-----	------------------	----------------

pH 0 days	6.13	6.40	6.35	0.05	n.s.
pH 24 days	6.02b	6.04b	6.10a	0.04	**
a_w	0.89	0.88	0.90	0.01	n.s.
L*	43.21	41.67	42.37	0.67	n.s.
a*	17.22a	15.92b	18.06a	0.41	**
b*	5.31b	4.87b	6.48a	0.21	**
Weight loss (%)	40.23	43.30	42.92	1.35	n.s.
Moisture (%)	30.43	29.97	30.94	0.66	n.s.
Fat (g/100 g dm)	41.82	41.01	41.32	0.33	n.s.
Protein (g/100 g dm)	48.88	49.74	49.14	0.31	n.s.
Ash (g/100g dm)	8.46	8.69	8.80	0.12	n.s.
TBARs (mg MDA/kg)	1.05	0.98	0.93	0.05	n.s.
SFA (mg/100g)	8.05	7.57	7.71	0.23	n.s.
MUFA (mg/100g)	9.72	9.17	9.10	0.25	n.s.
PUFA (mg/100g)	3.32	3.38	3.47	0.10	n.s.

Values are reported as means of the two replications within the same treatment, where GSE is grape seed extract added group, CHE the chestnut extract added group and NIT the control group added with sodium nitrite

^a Standard error

^b P value of natural antioxidant effect ** p<0.01, * p<0.05, different letters in the same row indicate significant differences at p<0.05.

Table 5. Texture traits of Cinta Sense dry-fermented sausages manufactured with natural antioxidant as replacement of sodium nitrite.

	GSE	CHE	NIT	SEM ^a	P ^b
Hardness (N)	104.93	102.82	102.68	6.38	n.s.
Cohesiveness	0.38ab	0.42a	0.35b	0.01	**
Gumminess	39.57	42.93	35.99	2.11	n.s.
Springiness	3.04ab	3.26a	2.70b	0.13	*
Chewiness (N)	120.67ab	139.55a	96.79b	8.36	**

Values are reported as means of the two replications within the same treatment, where GSE is grape seed extract added group, CHE the chestnut extract added group and NIT the control group added with sodium nitrite

^aStandard error

^bP value of natural antioxidant effect ** p<0.01, * p<0.05, different letters in the same row indicate significant differences at p<0.05.

Table 6. Volatile compounds in Cinta Sense dry-fermented sausages manufactured with natural antioxidant as replacement of sodium nitrite.

Compound	LRI ^a	RI ^b	GSE	CHE	NIT	SEM ^c	P ^d
Lipid auto-oxidation							
Pentane	500	a	0.59	0.60	0.56	0.19	
Propanal	524	a	0.19	0.13	0.13	0.06	
Isopropanol (45)	539	-	1.20	0.79	1.13	0.40	
Hexane	600	a	0.76	0.74	0.56	0.22	

1-Propanol	613	a	4.90	4.90	3.72	2.37	
Octane	800	a	4.20	3.498	3.90	0.98	
Propanoic acid (74)	814	a	0.14a	0.05b	0.02b	0.05	**
1-Pentanol	827	a	0.34ab	0.43a	0.22b	0.16	*
Hexanal	841	a	2.62a	2.27 a	1.36 b	0.27	*
1-Hexanol	924	a	0.79	0.75	0.88	0.16	
Decane	1000	a	1.10a	1.26a	0.48b	0.18	**
Dodecane	1200	a	0.96	1.14	0.92	0.37	
Tridecane	1300	a	0.36	0.44	0.34	0.10	
Tetradecane	1400	a	0.18b	0.87a	0.22b	0.16	**
Total			19.79a	16.19ab	14.57b	0.96	**
<hr/>							
<i>Spices</i>							
α -Thujene	934	b	24.40	26.07	27.29	6.22	
α -Pinene	940	a	17.84	19.00	17.33	3.68	
Sabinene+ β -pinene	986	a	133.28	149.70	145.97	19.83	
β -Myrcene	1003	a	37.19b	53.47a	39.01b	9.25	**
α -Phellandrene (93)	1019	b	8.71b	12.39a	9.41b	2.41	**
3-Carene	1023	a	73.20	86.24	77.49	15.77	
α -Terpinene	1035	a	3.66b	4.54a	3.28b	0.78	*
Unknown (57)	1042	b	0.17	0.16	0.14	0.11	
Limonene	1046	a	307.31b	401.19a	289.74b	60.75	**
β -Phellandrene (93)	1051	b	13.98b	19.28a	14.42b	3.74	*
p-Cymene (119)	1052	b	8.49	9.078	8.98	2.73	
β -Ocimene	1067	b	1.18b	1.87a	1.15b	0.44	**

4-Carene	1073	b	0.21	0.27	0.22	0.12	
Unkwon	1075	b	3.21b	4.28a	3.16b	0.93	*
Styrene	1091	a	2.20	2.43	2.34	0.31	
Terpene	1099	b	0.95	1.46	1.35	0.31	
Terpinolene	1101	b	4.64b	6.48a	4.46b	1.20	**
Unknown (93)	1120	b	0.51	0.55	0.56	0.11	
Linalool (93)	1150	a	0.28b	0.33a	0.32ab	0.04	*
β -Terpinene/ γ -Terpinene	1158	b	1.4	1.37	1.43	0.24	
4-Terpineol	1231	a	2.14	2.31	2.12	0.29	
Estragole	1249	a	1.28	1.33	1.20	0.21	
α -terpineol	1256	a	0.82	0.91	0.88	0.13	
δ -Elemene	1342	b	3.58a	4.19a	2.74b	0.66	**
α -Cubebene	1348	b	0.85a	0.92a	0.68b	0.12	**
Cyclosativene (161)	1402	b	0.02b	0.04a	0.02b	0.00	*
Copaene (161)	1406	b	1.50a	1.42a	1.16b	0.28	*
β -Cubabene (161)	1421	b	0.20a	0.19a	0.16b	0.04	*
β -Elemene (93)	1423	b	0.11b	0.13a	0.09b	0.04	**
α -Bergamotene	1433	b	0.79a	0.91a	0.69c	0.13	**
trans- α -Bergamotene	1450	b	0.67	0.84	0.76	0.18	
β -Caryophyllene	1455	b	107.97a	113.01a	91.99b	13.04	**
α -Caryophyllene	1486	b	4.87a	4.99a	4.19b	0.68	*
Isocaryophyllene	1498	b	0.65	0.84	0.57	0.26	
Isolongifolene	1510	b	10.64b	11.05b	25.91a	10.05	**
Valencene	1513	b	0.00b	5.31a	0.00b	1.78	**
γ -Cadinene	1518	b	1.72b	1.89b	2.92a	0.56	**

δ -Cadinene	1529	b	0.96b	1.04b	1.29a	0.20	**
Total			783.55b	984.05a	785.83b	134.67	**

Bacterial metabolism**Lipid β -oxidation**

2-Pentanone	734	a	5.64	5.96	6.92	1.79	
3-Octanone	1032	a	2.23	2.51	2.63	0.65	
Total			7.87b	8.47ab	9.55a	0.63	*

Esterase activity

3-Methyl-1-butanol acetate	907	a	0.59	0.49	0.51	0.12	
----------------------------	-----	---	------	------	------	------	--

Carbohydrate**fermentation**

Acetaldehyde	462	a	1.26a	0.86b	0.89b	0.25	**
Ethyl alcohol	507	a	6.79a	5.65ab	3.68b	2.23	*
Acetone	529	a	72.92	76.47	72.72	20.58	
2,3-Butanedione	627	a	9.76a	7.47b	7.83b	2.01	*
2-Butanone	632	a	241.96	287.47	332.22	91.70	
2-Butanol	644	a	19.99	20.52	29.19	11.02	
Acetic acid (60)	720	a	11.99	12.19	7.08	5.34	
1-Butanol (56)	727	a	0.28	0.18	0.71	0.67	
2-Pentanol	756	a	0.49	0.43	0.38	0.15	
3-Hydroxy-2-butanone	782	a	124.14a	89.35b	136.91a	31.84	**

2,3-Butanediol (45)	884	a	6.20a	1.50b	3.76b	2.92	**
2,3-Butanediol (45)	892	a	4.17	2.30	2.36	1.98	
Butanoic acid (60)	894	a	0.52a	0.49a	0.18b	0.254808	*
Total			499.97b	489.85b	597.61a	23.69	**

Amino acid degradation

2-Methylpropanal	595	a	0.54b	0.66a	0.39b	0.17	**
3-Methylbutanal	691	a	11.34	12.43	15.34	4.53	
2-Methylbutanal	701	a	6.18	7.53	8.85	2.88	
Toluene	789	a	51.99	56.94	59.18	6.88	
3-Methylbutanol	795	a	7.39b	8.46b	15.13a	5.92	*
2-Methylbutanol	797	a	1.27	1.44	1.82	5.92	
Pyrrrole	845	a	0.31	0.30	0.25	0.05	
2-Methylpropanoic acid	868	a	2.01	1.70	2.70	1.06	
Ethylbenzene (91)	884	a	0.09b	0.18a	0.09b	0.05	**
3-Methylbutanoic acid (60)	942	a	2.52	2.25	2.27	1.05	
2,5- Dimethylpirazine (108)	943	a	0.34	0.34	0.36	0.09	
2-Methylbutanoic acid	948	a	2.32	2.73	2.98	1.25	
2-Acetyl-1-pyrroline	961	a	0.72b	0.91b	1.37a	0.37	**
Methional	986	a	0.16	0.18	0.15	0.02	
Benzaldehyde (106)	1020	a	1.14	1.40	1.27	0.33	
Benzeneacetaldehyde	1110	a	6.61a	4.14b	2.72b	3.17	*
Tetramethylpyrazine	1118	a	0.03	0.02	0.03	0.01	
Benzylalcohol (79)	1122	a	0.27a	0.18b	0.19b	0.06	**
Phenylethyl alcohol (91)	1195	a	0.53	0.64	0.81	0.35	

Total			96.59b	103.09ab	115.63a	4.55	*
Total microbial metabolism			605.04b	601.91b	723.03a	24.32	**
Unknown or contaminant compounds							
Carbon disulfide (76)	537	a	5.80	6.42	6.70	1.89	
p-Xylene (91)	892	a	0.09b	0.31a	0.11a	0.09	**
2-Butoxyethanol	953	a	2.45	2.23	2.33	0.33	
4-Methylphenol (108)	1199	a	0.10	0.06	0.08	0.03	
Total			8.45	8.98	9.23	0.55	

Values are reported as means of the two replications within the same treatment, where GSE is grape seed extract added group, CHE the chestnut extract added group and NIT the control group added with sodium nitrite.

Abundance expressed as $AU \times 10^{-6}$ (AU: abundance unit, expressed as total ion chromatogram (TIC) or area of the target ion shown in parenthesis).

^aLinear retention indices (LRI) of the compounds eluted from the GC-MS using a DB-624 capillary column.

^bReliability of identification: a, identification by mass-spectrum and by coincidence with the LRI of an authentic standard; b, tentatively identification by mass-spectrum.

^cStandard error

^dP value of natural antioxidant effect ** $p < 0.01$, * $p < 0.05$, different letters in the same row indicate significant differences at $p < 0.05$.

Table 7. Odor active compounds identified by GC-O in Cinta Sense dry-fermented sausages manufactured with natural antioxidant as replacement of sodium nitrite.

Compound	GC-MS ¹		GC-O ²		RI ³	Odour Description	DF ₄
	LRI	LRI std	LRI initial	LRI final			
Lipid auto-oxidation							
1-Propanol	613	611	613	618	a	vegetal, green, pungent, fresh, floral,	5
Propanoic Acid	814	806	800	810	a	tasty, fresh, green, cheese, cured, pungent	4
1-Pentanol	827	823	821	827	a	vegetable, pungent, unpleasant, cabbage, acid	6
Hexanal	841	839	834	844	a	green, grass, vegetable, fresh	8
Spices							
Linalool	1150	1145	1141	1149	a	fresh, floral, cabbage, unpleasant, soap	7
β -Terpinene/ γ -Terpinene	1158		1160	1167	b	cooked, cooked vegetable, floral, pungent, resin	8
β -Myrcene	1003	1003	1000	1004	a	irritating, spicy, pepper, green, leaves, earthy	10
α -Terpinene	1035	1035	1030	1034	a	mushrooms, wetness, burnt, unpleasant, pungent, pine, woody, earthy	12
Unkown terpene	1075		1076	1080	c	earthy, green, vegetable, fresh, fruity, cologne	5
Terpinolene	1101	1106	1107	1113	a	floral, rose, grass, green	11

α -Thujene	934		929	933	c	sour, vinager, unpleasant, fruity,	5
Lipid β-oxidation							
2-Pentanone	735	731	725	733	a	floral, green, fresh, oxidized fat, cheese	8
Esterase activity							
3-Methyl-1-butanol acetate	907	905	905	911	a	sweet, fresh, floral	7
Carbohydrate fermentation							
2-Butanone	632	629	631	642	a	sweet, slightly unpleasant, green, grass	5
2-Butanol	644	643	650	661	a	sweet, caramel, malt, unpleasant,	5
Acetic acid	720	718	714	722	a	grass, vegetable, fresh, wine, green	5
Amino acid degradation							
2-Methylpropanal	595	590	602	609	a	acid, floral, green, weak	4
3-Methylbutanal	691	687	690	697	a	caramel, chocolate, grass, fresh	6
2-Methylbutanal	702	699	691	700	a	sweet, floral, fruity, toasted,	6
3-Methylbutanol	795	793	791	793	a	sweet, spicy, toasted, floral	4
2-Methylpropanoic acid	868	864	869	872	a	cheese, roasted, cured, green, slightly sweet, unpleasant	6

Ethylbenzene	884	881	884	891	a	earthy, fresh, green, mushroom	4
3-Methylbutanoic acid	942	941	922	926	a	cheese, rancid, oxidized fat	8
2,5-Dimethylpyrazine	943	943	936	943	a	meaty, cooked potatoes, sweet, buttery	7
2-Acetyl-1-pyrroline	961	960	960	964	a	roasted, nuts, bread, pop-corn, biscuits, fried potatoes	12
Methional	986	964	966	969	a	mashed potato, cooked onion, roasted meat	9
Tetramethylpyrazine	1118	1118	1115	1121	a	earthy, green, grass, wetness, fresh	7
Unknown or contaminant compounds							
Carbon Disulfide	537	537	531	543	a	weak, burnt, malt	4
Unknown		776	762	766	c	cured, meat, acid, fresh, acid	6
Unknown	1190	1182	1176	1182	c	roasted, fried nuts, biscuits	11

¹Linear retention index (LRI) of the compounds eluted from the GC-MS and LRI of standard compound

²Initial and end linear retention index of aroma compound in GC-FID-O

³Reliability of identification (RI): a, identification by mass spectrum, coincidence with LRI of an authentic standard and by coincidence of the assessor's descriptors with those described in the Fenaroli's handbook of flavor ingredients (Burdock, 2002); b, tentatively identification by mass spectrum; c, unknown compounds.

⁴Detection frequency value.

Figure 1. Scores of Canonical discriminant analysis for GSE (●), CHE (●) and NIT (●) samples and loadings of Canonical discriminant analysis for volatile compounds identified by Stepwise discriminant analysis.

Figure 2. Sensorial traits of Cinta Sense dry-fermented sausages manufactured with natural antioxidant as replacement of sodium nitrite




Figure 1




Figure 2