



HAL
open science

Viticulture in Portugal: A review of recent trends and climate change projections

Helder Fraga, Inaki Garcia de Cortazar Atauri, Aureliano C. Malheiro, Jose Moutinho-Pereira, João A. Santos

► **To cite this version:**

Helder Fraga, Inaki Garcia de Cortazar Atauri, Aureliano C. Malheiro, Jose Moutinho-Pereira, João A. Santos. Viticulture in Portugal: A review of recent trends and climate change projections. *OENO One*, 2017, 51 (2), pp.61 - 69. 10.20870/oeno-one.2016.0.0.1621 . hal-02629331

HAL Id: hal-02629331

<https://hal.inrae.fr/hal-02629331>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Viticulture in Portugal: A review of recent trends and climate change projections

Helder Fraga^{1*}, Iñaki García de Cortázar Atauri², Aureliano C. Malheiro¹,
José Moutinho-Pereira¹ and João A. Santos¹

¹ Centre for the Research and Technology of Agro-Environmental and Biological Sciences,
Universidade de Trás-os-Montes e Alto Douro, 5000-801 Vila Real, Portugal

² French National Institute for Agricultural Research, INRA, US1116 AgroClim, F-84914 Avignon, France

*This article is published in cooperation
with the ClimWine international conference held in Bordeaux 11-13 April 2016.
Guest editor: Nathalie Ollat*

Abstract

Aim: The winemaking sector in Portugal is of major socio-economic relevance, significantly contributing to the national exports and sustaining many wine-related activities, including oenotourism. Portuguese viticultural regions present a wide range of edaphoclimatic conditions with remarkable regional specificities, thus contributing to the individuality of their wines. Owing to the strong influence of climate and weather factors on grapevines, climate change may drive significant impacts on Portuguese viticulture.

Methods and results: Climatic projections for the next decades in Portugal highlight an overall warming and drying trend of the grapevine growing season, potentially resulting in modifications in phenology, growth, development, yields and eventually wine characteristics and typicity. Furthermore, the current viticultural suitability of each region is projected to undergo significant changes, suggesting a reshaping of the optimal conditions for viticulture throughout the country. In order to sustain high quality levels and affordable yield regularity, cost-effective, appropriate and timely adaptation measures must be implemented by the sector.

Conclusion: The most recent scientific studies covering the potential impacts of climate change on Portuguese viticulture are herein presented.

Significance and impact of the study: Possible adaptation measures against these threats are also discussed, foreseeing their integration into decision support systems by stakeholders and decision-makers.

Keywords: *Vitis vinifera*, viticulture, wine regions, Portugal, climate change

*Received : 10 July 2016; Accepted : 18 October 2016
DOI: 10.20870/oeno-one.2016.0.0.1621*

Viticulture and winemaking in Portugal

1. Socio-economic context

In Portugal, winemaking is historically one of the most relevant socio-economic activities. In the context of the overall agricultural sector (e.g. cereals, vegetables), this industry represents roughly 14% of the total planted area and 6% of the total productions (INE, 2016). With an average vineyard area of over 200 thousand hectares and a yearly wine production of about 6 million hectolitres (IVV, 2013), the national production has shown a slight decrease over the past decade (-2%/yr), which can be mostly attributed to the gradual decrease in vineyard area (-1%/yr). Nonetheless, Portugal is currently the 11th wine producer and the 10th exporter in the world (OIV, 2013), which is a remarkable outcome taking into account the size of the country. Approximately half of the total annual wine production is currently being exported. In absolute terms, this contributes to the national exports with over 700 million €/yr, which corresponds to nearly 2% of total national exports. A major factor for this success is the wide recognition of Portuguese wines in foreign markets, just to mention the renowned Port wine.




Figure 1. Wine regions in mainland Portugal. The Azores and Madeira archipelagos are not shown.

2. The winemaking regions

Portugal comprises a total of 14 wine regions (mainland Portugal, Azores and Madeira archipelagos) (Figure 1), which include 31 Protected Denominations of Origin. In the north, the Douro Demarcated Region, with almost 1.5 Mhl of total wine production and 45,000 ha of vineyards, is the oldest and one of the most important wine regions of the country. This region, famous for its Port wine, is responsible for one fourth of all wine produced in Portugal (IVV, 2013), and its vineyard landscape is also considered World Heritage by the UNESCO since 2001. In the northwest, the most maritime area of Portugal, the Minho wine region produces mostly white wines, distinguishable for their typical freshness and slightly higher acidity. In the south, the Alentejo wine region, with typical Mediterranean climates, has undergone remarkable growth rates over the recent decades and is currently the leading region in terms of non-fortified wine production. In general, from the sparkling wines of the Beira-Atlântico region to the fortified Madeira wine, many other regions in Portugal present unique wines resulting from their specific *terroirs*. The wines of Portugal are thereby valuable national brands, increasingly recognised worldwide.

3. The climate and the soils

Overall, the wine regions in mainland Portugal present Mediterranean-like climatic conditions, with warm dry summers and mild wet autumns-winters. In the northern/coastal areas (i.e. Minho and Beira-Atlântico), the Atlantic influence is strong, resulting in relatively high precipitation totals (>1,000 mm). In general, temperatures are higher in the south (e.g. Alentejo) and lower in the north (e.g. Minho and Trás-os-Montes). In the inner areas, however, summertime low water availability critically limits grapevine development (inner Alentejo and Douro). Winter temperatures tend to be milder in the southern regions, such as in Lisboa and Algarve (January mean temperature of 10–12°C), though late spring frost is common in some northern regions (e.g. Terras do Dão), which may lead to important damages in vineyards.

Growing degree-day (GDD (Winkler, 1974); April–October, 1950–2000) climatologies over Portugal (Figure 2a) indicate that the northern regions typically show temperate climates, with cool areas at higher elevations and warm/temperate-warm areas at lower elevations, especially in the Douro region. Almost all of the southern part of the country shows a warm climate, but with some inner areas already in

the very-warm class. Regarding dryness conditions (Dryness Index (Tonietto and Carbonneau, 2004); April–September, 1950–2000), the Atlantic influence is noticeable in the northern coastal areas, with sub-humid or even humid climates (Figure 2b), whereas the rest of the country depicts moderate dryness.

There are two main soil types in Portugal: cambisols in the centre-north and lithosols/luvisols in the south (FAO, 2006). The most important exceptions are the Douro region in the north, with lithosols, and the Peninsula-de-Setubal and Tejo regions, with podsols. Loam is the predominant soil texture in Portugal,




Figure 2. a) Growing degree-days and b) Dryness Index calculated for the growing season (April–September, 1950–2000) over Portugal.

while sand and clay are less frequent (Fraga *et al.*, 2014a). With respect to physiography, the most mountainous areas are located in inner northern and central areas, northwards of the Tagus River, while flatlands prevail in coastal and southern areas.

4. The varieties and practices

Portuguese vineyards preserve a large number of autochthonous and international varieties, with over 300 authorized varieties. Aragonez or Tinta-Roriz (also known as Tempranillo, red) is the most planted variety in Portugal, followed by Touriga-Franca (red), Castelão (red), Fernão-Pires (white) and Touriga-Nacional (red). These varieties are present in nearly all regions. Other varieties are more region-specific, such as Alvarinho (white) in Minho or Baga (red) in Beira-Atlântico. All these varieties present unique agronomic and oenological characteristics that ultimately result in distinctive wines, either mono-varietal or blended, though blended wines are more traditional, including the Port wine. The cordon (unilateral or bilateral) is the most used training system, though pergola (in Minho) and gobelet (e.g. Trás-os-Montes) can also be found. Phenological timings (budburst, flowering, veraison and maturation), although varietal-dependent, tend to occur earlier in the southern/warmest part of the country. Budburst occurs from March to April and flowering in May–June, while harvest is typically carried out from late August to early October.

Climate change

1. Climate change projections

Atmospheric conditions are one of the most important controlling factors for the growth and productivity of grapevines (Keller, 2010). In fact, grapevine physiology and berry composition are highly influenced by air temperatures during the growth cycle (Keller, 2010). Furthermore, winter chilling is also very important for bud dormancy (Bates *et al.*, 2002; Field *et al.*, 2009). Consequently, a base temperature of ca. 10°C is required to break this dormancy period and to onset the growing/vegetative cycle (Amerine and Winkler, 1944; Winkler, 1974). Despite its high adaptability to different climatic conditions and its resilience to moderate water and heat stresses, this crop can be severely affected by stresses derived from extreme weather events. Extremely low negative temperatures in spring may significantly damage grapevine development (Branas, 1974). Grapevines are also very sensitive to frost and hail during their vegetative period (e.g. Spellman, 1999). Heat waves may also

considerably affect physiology and yields (Kliewer, 1977; Mullins *et al.*, 1992).

According to the International Panel on Climate Change, global temperature is expected to rise from 2 to 5°C by 2100 (IPCC, 2013). For Portugal, temperature projections are in agreement with these changes, while precipitation is expected to decrease, particularly in the south-innermost areas (SIAM2, 2006). The recent climatic trends over Portugal are already in line with these climate projections (Fraga *et al.*, 2012).

For the future, an overall warming and drying of the grapevine growing season is indeed anticipated (Fraga *et al.*, 2012; Fraga *et al.*, 2014b). GDD projections for 2041-2070 under the IPCC A1B scenario suggest large increases in accumulated temperatures (Figure 3a), particularly in the innermost regions, reaching values above 2700 °C (excessively hot class). In addition to the overall warming, the drying trend will lead to changes in the DI patterns. Severe dryness is likely to occur in the future, particularly in the innermost southern regions (Figure 3b). The expected decrease in rainfall in spring and summer will enhance water requirements and may trigger severe water stress in vineyards. Moreover, updated projections following the IPCC RCP scenarios are in close agreement with these outcomes (Fraga *et al.*, 2016a). Given the key role played by atmospheric factors on viticulture, climate change is expected to bring new challenges to this crop.

2. Impacts on phenology

Grapevines will be particularly affected by the projected higher temperatures during the growing season. As temperatures are a major driver of the grapevine development stages (Parker *et al.*, 2013), significant warmings are expected to lead to earlier onsets (Bock *et al.*, 2011; Chuine *et al.*, 2004; Dalla Marta *et al.*, 2010; Daux *et al.*, 2011; Jones *et al.*, 2005a; Molitor *et al.*, 2014; Sadras and Petrie, 2011; Webb *et al.*, 2011). Recent studies for Portugal isolated future projections for the phenological stages of 16 native varieties under RCP4.5 and 8.5 (Fraga *et al.*, 2015; Fraga *et al.*, 2016c; Malheiro *et al.*, 2013). The results hint at earlier onsets of 2–5 days for budburst and flowering, and of 7–15 days for veraison until 2070, depending on the selected future scenario and variety.

Earlier phenological timings will bring heterogeneous outcomes. Earlier budburst and flowering may result in substantial increases in the risks of frost damages. Given the projected increase

of grapevine-related pests/diseases (Francesca *et al.*, 2006; Valero *et al.*, 2003; Van Niekerk *et al.*, 2011), this may also entail increased risks and cause a strong impact on management practices. Extreme heat during this period may abruptly reduce vine metabolism, affecting the aroma and colour of wines. Still, higher sugar concentrations and lower acidity are expected, which may potentially increase the risk of organoleptic degradation or even spoilage (Orduna, 2010), threatening the production of well-balanced wines.

3. Impacts on yield

Under future climates, the potential impacts on grapevine yield can be very diverse. The interaction between negative (higher heat and water stresses) and positive (enhanced CO₂ effect on plant physiology) climate change effects on yields are expected to lead to different outcomes (Bindi *et al.*, 1996; Fraga *et al.*, 2014c). Basically, the overall effect on production will depend on CO₂ concentrations, temperature, solar radiation, precipitation and many other factors. As an example, for the Douro region, several studies suggest higher grapevine yields and wine productions in future climates (Gouveia *et al.*, 2011; Santos *et al.*, 2011; Santos *et al.*, 2013). Nonetheless, these studies were conducted considering the more humid part of the region (Baixo-Corgo), while projections for the driest areas hint at yield decreases (Cima-Corgo and Douro-Superior). Furthermore, yield decreases are also projected to occur in the Alentejo region (Coelho *et al.*, 2013), which shows a much warmer and drier climate. Although projections for yield are largely heterogeneous and site-specific, most studies agree regarding the projections for the annual yield irregularity. The expected increase in the frequency and intensity of weather extremes (Andrade *et al.*, 2014) will lead to higher inter-annual yield variability, which may affect the whole winemaking sector (Jones *et al.*, 2005a; Schultz, 2000).

4. Impacts on wine quality

Under future warmer climates, extremely high temperatures may inhibit the formation of anthocyanins (Buttrose *et al.*, 1971), impacting berry colour and aroma (Bureau *et al.*, 2000; Downey *et al.*, 2006). Higher sugar concentrations and lower acidity are also expected under future warming. For regions already presenting warm climates (e.g. Alentejo, Douro), climate change may thus endanger the balanced ripening of grapes and the sustainability of the existing varieties and wine styles (Fraga *et al.*, 2016b; Jones and Alves, 2012). However, future warming in the cooler climate regions (e.g. Minho,

Beira-Atlântico) may improve suitability for the production of high quality wines. As such, although a modification of the currently established wine types may occur, the socio-economic impacts of climate change on wine quality can be quite diverse.

Adaptation measures

As a protection strategy against the unwanted impacts of climate change, adaptation measures should be




Figure 3. a) Growing degree-days and b) Dryness Index calculated for the growing season (April–September) over Portugal for 2041-2070 and the A1B scenario.

considered, focusing on specific problems. Adequate and timely planning of these measures needs to be adopted by the winemaking sector. The readiness to define and implement adaptation strategies to climate change will result in lower detrimental impacts to the sector (Barbeau *et al.*, 2014; Battaglini *et al.*, 2009; Olesen *et al.*, 2011). These strategies may include short-term measures, such as changes in viticultural management practices, grown varieties or even oenological practices (Neethling *et al.*, 2016). Additionally, long-term adaptation measures should also be considered, as some regions may become excessively warm and dry. These different types of adaptation measures are discussed in the following subsections.

1. Short-term adaptation measures

Taking into account the warming trend projected for the future, selecting varieties with higher thermal requirements and higher stress resistance may strengthen the overall resilience of the Portuguese viticulture under future climates (Fraga *et al.*, 2016b). Therefore, preserving the existing biodiversity is critical, as some varieties may thrive in the future and may thereby be part of the solution. Furthermore, oenological practices may also play a key role in maintaining regional wine typicity and quality.

Regions under severe dryness, such as Alentejo and Douro, should promote higher water use efficiencies (Flexas *et al.*, 2010) by adopting training systems that promote shorter trunks and lower total leaf areas, such as gobelet. The selection of more drought tolerant rootstocks must also be regarded as a possible adaptation measure (Harbertson and Keller, 2012; Keller *et al.*, 2011). Changes in tillage systems and soil management should also be considered (Bahar and Yasasin, 2010; Kvaternjak *et al.*, 2008). One of the most controversial (in Portugal) measures is the application of water by irrigation, but smart irrigation strategies can promote a balanced compromise solution between environment, economy and plant water requirements (Chaves *et al.*, 2007; Chaves *et al.*, 2010; dos Santos *et al.*, 2003; Ferreira *et al.*, 2012).

Excessive solar radiation can also be damaging for grapevines, already under water and heat stresses. Shading materials, either natural (e.g. olive trees) or artificial, can help overcome this shortcoming (Greer *et al.*, 2011; Shahak *et al.*, 2008). Furthermore, the use of chemical sunscreens for leaf protection against sunburns may also represent an important alternative (Dinis *et al.*, 2016). Another option to consider is the

adjustment of the implemented training system (Pieri and Gaudillere, 2003). Changing canopy geometry or orientation can also significantly influence light interception (Grifoni *et al.*, 2008; Intrieri *et al.*, 1998).

2. Long-term adaptation measures

Long-term adaptation measures should also be considered, although their extent and application may bring significant socio-economic implications. These measures include changing vineyard location, as some regions may become excessively warm and dry (Fraga *et al.*, 2016b; Moriondo *et al.*, 2013). Relocations of vineyards to cooler sites, such as higher elevations, coastal zones or simply areas with lower solar exposures, are possible measures.

Conclusions

Although climate change is expected to drive significant changes on Portuguese viticulture, large uncertainties still remain regarding the true extent of its impacts. The expected warming and drying trends throughout Portugal may bring some additional challenges for grapevine production (Santos *et al.*, 2011). Increases in the growing-season mean temperatures are indeed expected not only in all of the Portuguese winemaking regions, but also in other regions worldwide (Duchene and Schneider, 2005; Jones *et al.*, 2005b; Neumann and Matzarakis, 2011). This will lead to earlier phenological timings, with potential detrimental impacts (Bock *et al.*, 2011; Chuine *et al.*, 2004; Dalla Marta *et al.*, 2010; Webb *et al.*, 2008). Some southern regions are projected to become excessively dry to grapevine production using the currently established viticultural practices and varieties. Additionally, enhanced risks of pests and diseases in vineyards can be an additional threat (Francesca *et al.*, 2006; Valero *et al.*, 2003; Van Niekerk *et al.*, 2011).

The implementation of adaptation measures is urging, as scientific confidence for significant climate change in the upcoming decades is growing. Appropriate measures need to be addressed by the wine industry to face climate change impacts, mainly by developing suitable strategies at regional scales (Metzger *et al.*, 2008). Winegrape growers are becoming progressively more aware of this problem (Battaglini *et al.*, 2009), since timely strategic planning will provide competitive advantages. Nevertheless, in order to effectively cope with the projected changes, continuous research is needed as climate change is progressing. As such, it is up to the decision-makers and stakeholders from the winemaking sector to implement actions against

climate change. These actions will critically contribute to the future economic and environmental sustainability of the Portuguese viticulture.

Acknowledgments : This work was supported by: R&D project ModelVitiDouro (PA 53774), funded by the Agricultural and Rural Development Fund (EAFRD) and the Portuguese Government (Measure 4.1 – Cooperation for Innovation PRODER programme – Rural Development Programme); R&D project INNOVINE&WINE – Vineyard and Wine Innovation Platform, (NORTE-01-0145-FEDER-000038), co-funded by the European Regional Development Fund through the programme NORTE 2020 (Programa Operacional Regional do Norte 2014/2020); European Investment Funds (FEDER/COMPETE/POCI – Operacional Competitiveness and Internacionalization Programme) under Project POCI-01-0145-FEDER-006958 and National Funds (FCT – Portuguese Foundation for Science and Technology) under Project UID/AGR/04033/2013. The postdoctoral fellowship (SFRH/BPD/119461/2016) awarded to the first author is appreciated.

References

- Amerine, M.A. and Winkler, A.J., 1944. Composition and quality of musts and wines of California grapes. *Hilgardia*, 15(6): 493-675.
- Andrade, C., Fraga, H. and Santos, J.A., 2014. Climate change multi-model projections for temperature extremes in Portugal. *Atmos Sci Lett*, 15(2): 149-156.
- Bahar, E. and Yasasin, A.S., 2010. The yield and berry quality under different soil tillage and clusters thinning treatments in grape (*Vitis vinifera* L.) cv. Cabernet-Sauvignon. *Afr J Agric Res*, 5(21): 2986-2993.
- Barbeau, G., Goulet, E., Neethling, E., Ollat, N. and Touzard, J.M., 2014. Les méthodes d'adaptation au changement climatique. In «*Changement climatique et terroirs viticoles*» (H. Quénel), Lavoisier, pp. 347-376.
- Bates, T.R., Dunst, R.M. and Joy, P., 2002. Seasonal dry matter, starch, and nutrient distribution in 'Concord' grapevine roots. *HortScience*, 37(2): 313-316.
- Battaglini, A., Barbeau, G., Bindi, M. and Badeck, F.W., 2009. European winegrowers' perceptions of climate change impact and options for adaptation. *Reg Environ Change*, 9(2): 61-73.
- Bindi, M., Fibbi, L., Gozzini, B., Orlandini, S. and Miglietta, F., 1996. Modelling the impact of future climate scenarios on yield and yield variability of grapevine. *Clim Res*, 7(3): 213-224.
- Bock, A., Sparks, T., Estrella, N. and Menzel, A., 2011. Changes in the phenology and composition of wine from Franconia, Germany. *Clim Res*, 50(1): 69-81.
- Branas, J., 1974. Viticulture. Dehan, Montpellier, France, 990 pp.

- Bureau, S.M., Razungles, A.J. and Baumes, R.L., 2000. The aroma of Muscat of Frontignan grapes: effect of the light environment of vine or bunch on volatiles and glycoconjugates. *J Sci Food Agric*, 80(14): 2012-2020.
- Buttrose, M.S., Hale, C.R. and Kliewer, W.M., 1971. Effect of temperature on the composition of Cabernet Sauvignon berries. *Am J Enol Vitic*, 22(2): 71-75.
- Chaves, M.M., Santos, T.P., Souza, C.R., Ortuno, M.F., Rodrigues, M.L., Lopes, C.M., Maroco, J.P. and Pereira, J.S., 2007. Deficit irrigation in grapevine improves water-use efficiency while controlling vigour and production quality. *Ann Appl Biol*, 150(2): 237-252.
- Chaves, M.M., Zarrouk, O., Francisco, R., Costa, J.M., Santos, T., Regalado, A.P., Rodrigues, M.L. and Lopes, C.M., 2010. Grapevine under deficit irrigation: hints from physiological and molecular data. *Ann Bot*, 105(5): 661-676.
- Chuine, I., Yiou, P., Viovy, N., Seguin, B., Daux, V. and Le Roy Ladurie, E., 2004. Historical phenology: grape ripening as a past climate indicator. *Nature*, 432(7015): 289-290.
- Coelho, J.C., Lopes, C.M., Braga, R., Pinto, P.A. and Egipto, R.J.L., 2013. Avaliação do impacte das alterações climáticas na sustentabilidade económica da cultura da vinha no Alentejo. VII APDEA Congress - ESADR 2013, Évora, P15: 4015-4039.
- Dalla Marta, A., Grifoni, D., Mancini, M., Storchi, P., Zipoli, G. and Orlandini, S., 2010. Analysis of the relationships between climate variability and grapevine phenology in the Nobile di Montepulciano wine production area. *J Agric Sci*, 148(6): 657-666.
- Daux, V., Garcia de Cortazar-Atauri, I., Yiou, P., Chuine, I., Garnier, E., Le Roy Ladurie, E., Mestre, O. and Tardaguila, J., 2011. An open-database of grape harvest dates for climate research: data description and quality assessment. *Clim Past Discuss*, 7(6): 3823-3858.
- Dinis, L.-T., Ferreira, H., Pinto, G., Bernardo, S., Correia, C.M. and Moutinho-Pereira, J., 2016. Kaolin-based, foliar reflective film protects photosystem II structure and function in grapevine leaves exposed to heat and high solar radiation. *Photosynthetica*, 54(1): 47-55.
- dos Santos, T.P., Lopes, C.M., Rodrigues, M.L., de Souza, C.R., Maroco, J.P., Pereira, J.S., Ricardo-da-Silva, J. and Chaves, M.M., 2003. Partial rootzone drying: effects on growth and fruit quality of field-grown grapevines (*Vitis vinifera*). *Funct Plant Biol*, 30(6): 663-671.
- Downey, M.O., Dokoozlian, N.K. and Krstic, M.P., 2006. Cultural practice and environmental impacts on the flavonoid composition of grapes and wine: a review of recent research. *Am J Enol Vitic*, 57(3): 257-268.
- Duchene, E. and Schneider, C., 2005. Grapevine and climatic changes: a glance at the situation in Alsace. *Agron Sustain Dev*, 25(1): 93-99.
- FAO, 2006. World reference base for soil resources 2006, a framework for international classification, correlation and communication. World soil resources reports 103, *Food and Agriculture Organization of the United Nations*, Rome.
- Ferreira, M.I., Silvestre, J., Conceição, N. and Malheiro, A.C., 2012. Crop and stress coefficients in rainfed and deficit irrigation vineyards using sap flow techniques. *Irrig Sci*, 30(5): 433-447.
- Field, S.K., Smith, J.P., Holzapfel, B.P., Hardie, W.J. and Emery, R.J.N., 2009. Grapevine response to soil temperature: xylem cytokinins and carbohydrate reserve mobilization from budbreak to anthesis. *Am J Enol Vitic*, 60(2): 164-172.
- Flexas, J., Galmes, J., Galle, A., Gulias, J., Pou, A., Ribas-Carbo, M., Tomas, M. and Medrano, H., 2010. Improving water use efficiency in grapevines: potential physiological targets for biotechnological improvement. *Aust J Grape Wine Res*, 16(s1): 106-121.
- Fraga, H., Santos, J.A., Malheiro, A.C. and Moutinho-Pereira, J., 2012. Climate change projections for the Portuguese viticulture using a multi-model ensemble. *Ciência Téc Vitiv*, 27(1): 39-48.
- Fraga, H., Malheiro, A.C., Moutinho-Pereira, J., Cardoso, R.M., Soares, P.M.M., Cancela, J.J., Pinto, J.G. and Santos, J.A., 2014a. Integrated analysis of climate, soil, topography and vegetative growth in Iberian viticultural regions. *PLoS One*, 9(9): e108078.
- Fraga, H., Malheiro, A.C., Moutinho-Pereira, J., Jones, G.V., Alves, F., Pinto, J.G. and Santos, J.A., 2014b. Very high resolution bioclimatic zoning of Portuguese wine regions: present and future scenarios. *Reg Environ Change*, 14(1): 295-306.
- Fraga, H., Malheiro, A.C., Moutinho-Pereira, J. and Santos, J.A., 2014c. Climate factors driving wine production in the Portuguese Minho region. *Agric For Meteorol*, 185: 26-36.
- Fraga, H., Costa, R., Moutinho-Pereira, J., Correia, C.M., Dinis, L.-T., Gonçalves, I., Silvestre, J., Eiras-Dias, J., Malheiro, A.C. and Santos, J.A., 2015. Modeling phenology, water status, and yield components of three Portuguese grapevines using the STICS crop model. *Am J Enol Vitic*, 66(4): 482-491.
- Fraga, H., García de Cortázar Atauri, I., Malheiro, A.C. and Santos, J.A., 2016a. Modelling climate change impacts on viticultural yield, phenology and stress conditions in Europe. *Glob Change Biol*, 22(11): 3774-3788.
- Fraga, H., Santos, J.A., Malheiro, A.C., Oliveira, A.A., Moutinho-Pereira, J. and Jones, G.V., 2016b. Climatic suitability of Portuguese grapevine varieties

- and climate change adaptation. *Int J Climatol*, 36(1): 1-12.
- Fraga, H., Santos, J.A., Moutinho-Pereira, J., Carlos, C., Silvestre, J., Eiras-Dias, J., Mota, T. and Malheiro, A.C., 2016c. Statistical modelling of grapevine phenology in Portuguese wine regions: observed trends and climate change projections. *J Agric Sci*, 154(5): 795-811.
- Francesca, S., Simona, G., Francesco Nicola, T., Andrea, R., Vittorio, R., Federico, S., Cynthia, R. and Maria Lodovica, G., 2006. Downy mildew (*Plasmopara viticola*) epidemics on grapevine under climate change. *Glob Change Biol*, 12(7): 1299-1307.
- Gouveia, C., Liberato, M.L.R., DaCamara, C.C., Trigo, R.M. and Ramos, A.M., 2011. Modelling past and future wine production in the Portuguese Douro Valley. *Clim Res*, 48(2): 349-362.
- Greer, D.H., Weedon, M.M. and Weston, C., 2011. Reductions in biomass accumulation, photosynthesis in situ and net carbon balance are the costs of protecting *Vitis vinifera* 'Semillon' grapevines from heat stress with shade covering. *AoB Plants*, 2011: plr023.
- Grifoni, D., Carreras, G., Zipoli, G., Sabatini, F., Dalla Marta, A. and Orlandini, S., 2008. Row orientation effect on UV-B, UV-A and PAR solar irradiation components in vineyards at Tuscany, Italy. *Int J Biometeorol*, 52(8): 755-763.
- Harbertson, J.F. and Keller, M., 2012. Rootstock effects on deficit-irrigated winegrapes in a dry climate: grape and wine composition. *Am J Enol Vitic*, 63(1): 40-48.
- INE, 2016. Instituto nacional de estatística - <http://www.ine.pt/>.
- Intrieri, C., Poni, S., Rebucci, B. and Magnanini, E., 1998. Row orientation effects on whole-canopy gas exchange of potted and field-grown grapevines. *Vitis*, 37(4): 147-154.
- IPCC, 2013. Climate change 2013: the physical science basis. Summary for policymakers. Working group I contribution to the IPCC fifth assessment report.
- IVV, 2013. Vinhos e Aguardentes de Portugal, Anuário 2013. Ministério da Agricultura, do Desenvolvimento Rural e das Pescas: Instituto da Vinha e do Vinho, Lisboa: 236.
- Jones, G.V., Duchêne, E., Tomasi, D., Yuste, J., Braslavská, O., Schultz, H.R., Martínez, C., Boso, S., Langellier, F., Perruchot, C. and Guimberteau, G., 2005a. Changes in European winegrape phenology and relationships with climate. In *Proceedings of the XIV GESCO Viticulture Symposium*, Geisenheim, Germany, pp. 54-61.
- Jones, G.V., White, M.A., Cooper, O.R. and Storchmann, K., 2005b. Climate change and global wine quality. *Clim Change*, 73(3): 319-343.
- Jones, G.V. and Alves, F., 2012. Impact of climate change on wine production: a global overview and regional assessment in the Douro Valley of Portugal. *Int J Global Warm*, 4(3/4): 383-406.
- Keller, M., 2010. *The science of grapevines: anatomy and physiology*. Elsevier, 400 pp.
- Keller, M., Mills, L.J. and Harbertson, J.F., 2011. Rootstock effects on scion vigor and fruit and wine composition in a dry climate. *Am J Enol Vitic*, 62(3): 388a-388a.
- Kliewer, W.M., 1977. Effect of high temperatures during the bloom-set period on fruit-set, ovule fertility, and berry growth of several grape cultivars. *Am J Enol Vitic*, 28(4): 215-222.
- Kvaternjak, I., Kisić, I., Birkas, M., Sajko, K. and Simunić, I., 2008. Soil tillage as influenced by climate change. *Cereal Res Commun*, 36: 1203-1206.
- Malheiro, A.C., Campos, R., Fraga, H., Eiras-Dias, J., Silvestre, J. and Santos, J.A., 2013. Winegrape phenology and temperature relationships in the Lisbon wine region, Portugal. *J Int Sci Vigne Vin*, 47(4): 287-299.
- Metzger, M.J., Schroter, D., Leemans, R. and Cramer, W., 2008. A spatially explicit and quantitative vulnerability assessment of ecosystem service change in Europe. *Reg Environ Change*, 8(3): 91-107.
- Molitor, D., Junk, J., Evers, D., Hoffmann, L. and Beyer, M., 2014. A high-resolution cumulative degree day-based model to simulate phenological development of grapevine. *Am J Enol Vitic*, 65(1): 72-80.
- Moriondo, M., Jones, G.V., Bois, B., Dibari, C., Ferrise, R., Trombi, G. and Bindi, M., 2013. Projected shifts of wine regions in response to climate change. *Clim Change*, 119(3-4): 825-839.
- Mullins, M.G., Bouquet, A. and Williams, L.E., 1992. *Biology of the grapevine*. Cambridge University Press, Cambridge, UK.
- Neethling, E., Petitjean, T., Quéno, H. and Barbeau, G., 2016. Assessing local climate vulnerability and winegrowers' adaptive processes in the context of climate change. *Mitig Adapt Strateg Glob Change*, doi: 10.1007/s11027-015-9698-0.
- Neumann, P.A. and Matzarakis, A., 2011. Viticulture in southwest Germany under climate change conditions. *Clim Res*, 47(3): 161-169.
- OIV, 2013. *Statistical report on world vitiviniculture*. OIV, Paris, France, 32 pp.
- Olesen, J.E., Trnka, M., Kersebaum, K.C., Skjelvåg, A.O., Seguin, B., Peltonen-Sainio, P., Rossi, F., Kozyra, J. and Micale, F., 2011. Impacts and adaptation of European crop production systems to climate change. *Eur J Agron*, 34(2): 96-112.

- Orduna, R.M., 2010. Climate change associated effects on grape and wine quality and production. *Food Res Int*, 43(7): 1844-1855.
- Parker, A., García de Cortázar-Atauri, I., Chuine, I., Barbeau, G., Bois, B., Boursiquot, J.-M., Cahurel, J.-Y., Claverie, M., Dufourcq, T., Gény, L., Guimberteau, G., Hofmann, R.W., Jacquet, O., Lacombe, T., Monamy, C., Ojeda, H., Panigai, L., Payan, J.-C., Lovelle, B.R., Rouchaud, E., Schneider, C., Spring, J.-L., Storchi, P., Tomasi, D., Trambouze, W., Trought, M. and van Leeuwen, C., 2013. Classification of varieties for their timing of flowering and veraison using a modelling approach: a case study for the grapevine species *Vitis vinifera* L. *Agric For Meteorol*, 180: 249-264.
- Pieri, P. and Gaudillère, J.P., 2003. Sensitivity to training system parameters and soil surface albedo of solar radiation intercepted by vine rows. *Vitis*, 42(2): 77-82.
- Sadras, V.O. and Petrie, P.R., 2011. Climate shifts in south-eastern Australia: early maturity of Chardonnay, Shiraz and Cabernet Sauvignon is associated with early onset rather than faster ripening. *Aust J Grape Wine Res*, 17(2): 199-205.
- Santos, J.A., Malheiro, A.C., Karremann, M.K. and Pinto, J.G., 2011. Statistical modelling of grapevine yield in the Port Wine region under present and future climate conditions. *Int J Biometeorol*, 55(2): 119-131.
- Santos, J.A., Grätsch, S.D., Karremann, M.K., Jones, G.V. and Pinto, J.G., 2013. Ensemble projections for wine production in the Douro Valley of Portugal. *Clim Change*, 117(1-2): 211-225.
- Schultz, H., 2000. Climate change and viticulture: a European perspective on climatology, carbon dioxide and UV-B effects. *Aust J Grape Wine Res*, 6(1): 2-12.
- Shahak, Y., Ratner, K., Giller, Y.E., Zur, N., Or, E., Gussakovsky, E.E., Stern, R., Sarig, P., Raban, E., Harcavi, E., Doron, I. and Greenblat-Avron, Y., 2008. Improving solar energy utilization, productivity and fruit quality in orchards and vineyards by photosensitive netting. *Acta Hort*, 772: 65-72.
- SIAM2, 2006. Alterações climáticas em Portugal: cenários, impactos e medidas de adaptação. *Gradiva*, Lisboa, 506 pp.
- Spellman, G., 1999. Wine, weather and climate. *Weather*, 54: 230-239.
- Tonietto, J. and Carbonneau, A., 2004. A multicriteria climatic classification system for grape-growing regions worldwide. *Agric For Meteorol*, 124(1-2): 81-97.
- Valero, M., Ibanez, A. and Morte, A., 2003. Effects of high vineyard temperatures on the grapevine leafroll associated virus elimination from *Vitis vinifera* L. cv. Napoleon tissue cultures. *Sci Hort*, 97(3-4): 289-296.
- Van Niekerk, J.M., Bester, W., Halleen, F., Crous, P.W. and Fourie, P.H., 2011. The distribution and symptomatology of grapevine trunk disease pathogens are influenced by climate. *Phytopathol Mediterr*, 50: S98-S111.
- Webb, L.B., Whetton, P.H. and Barlow, E.W.R., 2008. Climate change and winegrape quality in Australia. *Clim Res*, 36: 99-111.
- Webb, L.B., Whetton, P.H. and Barlow, E.W.R., 2011. Observed trends in winegrape maturity in Australia. *Glob Change Biol*, 17(8): 2707-2719.
- Winkler, A.J., 1974. *General viticulture*. University of California Press, California, USA.