



Genome Sequencing of *Ralstonia solanacearum* Biovar 3, Phylotype I, Strains Rs-09-161 and Rs-10-244, Isolated from Eggplant and Chili in India.

Raman Ramesh, Sapna Gaitonde, Gauri Achari, Trupti Asolkar, Narendra Pratap Singh, Sebastien Carrere, Stéphane Genin, Nemo Peeters

► To cite this version:

Raman Ramesh, Sapna Gaitonde, Gauri Achari, Trupti Asolkar, Narendra Pratap Singh, et al.. Genome Sequencing of *Ralstonia solanacearum* Biovar 3, Phylotype I, Strains Rs-09-161 and Rs-10-244, Isolated from Eggplant and Chili in India.. *Genome Announcements*, 2014, 2 (3), 10.1128/genomeA.00323-14 . hal-02629436

HAL Id: hal-02629436

<https://hal.inrae.fr/hal-02629436>

Submitted on 22 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Genome Sequencing of *Ralstonia solanacearum* Biovar 3, Phylotype I, Strains Rs-09-161 and Rs-10-244, Isolated from Eggplant and Chili in India

Raman Ramesh,^a Sapna Gaitonde,^a Gauri Achari,^a Trupti Asolkar,^a Narendra Pratap Singh,^a Sebastien Carrere,^{b,c} Stephane Genin,^{b,c} Nemo Peeters^{b,c}

^aICAR Research Complex for Goa, Old Goa, Goa, India^a; INRA, Laboratoire des Interactions Plantes-Microorganismes (LIPM), Castanet-Tolosan, France^b; CNRS, Laboratoire des Interactions Plantes-Microorganismes (LIPM), Castanet-Tolosan, France^c

***Ralstonia solanacearum* Indian strains Rs-09-161 and Rs-10-244 were isolated from the coastal region of Goa and from the Andaman Islands. We report the draft genome sequences of these representative isolates infecting solanaceous vegetables in India.**

Received 27 March 2014 Accepted 1 May 2014 Published 29 May 2014

Citation Ramesh R, Gaitonde S, Achari G, Asolkar T, Singh NP, Carrere S, Genin S, Peeters N. 2014. Genome sequencing of *Ralstonia solanacearum* biovar 3, phylotype I, strains Rs-09-161 and Rs-10-244, isolated from eggplant and chili in India. *Genome Announc.* 2(3):e00323-14. doi:10.1128/genomeA.00323-14.

Copyright © 2014 Ramesh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/nd/3.0/).

Address correspondence to Raman Ramesh, rameshicar@yahoo.co.in.

Ralstonia solanacearum is a major plant-pathogenic bacterium that causes bacterial wilt disease in over 400 plant species (1). *R. solanacearum* strains from Africa, South America, North America, and the Caribbean have been sequenced (2). Phylotype I *R. solanacearum* strains from China have already been sequenced (3–5). Though bacterial wilt is severe in India (6), no genomic sequence is available as of yet. We sequenced the *R. solanacearum* strains Rs-09-161 and Rs-10-244, respectively isolated from eggplant (*Solanum melongena*) grown in the west coast region of India and chili (*Capsicum annuum*) grown in the Andaman Islands, India. Both strains belong to biovar 3, phylotype I, historically classified as race 1. Endoglucanase (*egl*) gene sequence analysis of *R. solanacearum* isolates collected from India indicated that Rs-09-161 belongs to the major group found within the phylotype I strains and hence represents the *R. solanacearum* population infecting solanaceous vegetables from India. Based on the *egl* gene sequence analysis, Rs-10-244 is the representative *R. solanacearum* isolate from the other subgroup.

Using Illumina HiSeq 2000, we obtained the nucleotide sequences of the Rs-09-161 and Rs-10-244 genomes from a paired-end library with an average insert size of 300 to 500 bp. The trimmed reads were assembled into contigs by using SOAP *de novo* (1.05) and Velvet (1.2.07). Meta-assembly was done using progressiveMauve (2.3.1), with *R. solanacearum* strain GMI1000 as the reference genome. Structural annotation was carried out using EuGene-P (7), with *R. solanacearum* strain GMI1000 as the reference proteome.

The overall G+C content of the Rs-09-161 genome is 66.82%. It contains 5,215 predicted genes, similar to the species complex average of 5,213. The draft genome sequence consists of 5,185 protein-coding genes with a mean gene length of 974 bp and 30 non-protein-coding genes with a mean gene length of 534 bp. The overall G+C content of the Rs-10-244 genome is 66.98%. It contains 5,202 predicted genes. The draft genome sequence consists of

5,171 protein-coding genes with a mean gene length of 969 bp and 31 non-protein-coding genes with a mean gene length of 446 bp. Compared with the published *R. solanacearum* genomes, Rs-09-161 and Rs-10-244 are most closely related to phylotype I strains GMI 1000 and FQY_4.

Further, T3E genes present in these genomes are identified according to the latest nomenclature (2). 71 T3E genes are present in Rs-09-161, which includes two putative candidate T3E genes, and 76 T3E genes are present in Rs-10-244.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at GenBank under the accession numbers **JHBO00000000** for strain Rs-09-161 and **JHAM00000000** for strain Rs-10-244. The versions described in this paper are the first versions.

ACKNOWLEDGMENTS

We acknowledge financial support from the Indian Council of Agricultural Research, New Delhi, India, through the “Outreach project on *Phytophthora*, *Fusarium* and *Ralstonia* diseases of horticultural and field crops” (*PhytoFuRa*).

REFERENCES

- Wicker E, Grassart L, Coranson-Beaudu R, Mian D, Guilbaud C, Fegan M, Prior P. 2007. *Ralstonia solanacearum* strains from Martinique (French West Indies) exhibiting a new pathogenic potential. *Appl. Environ. Microbiol.* 73:6790–6801. <http://dx.doi.org/10.1128/AEM.00841-07>.
- Peeters N, Carrère S, Anisimova M, Plener L, Cazalé AC, Genin S. 2013. Repertoire, unified nomenclature and evolution of the type III effector gene set in the *Ralstonia solanacearum* species complex. *BMC Genomics* 14:859. <http://dx.doi.org/10.1186/1471-2164-14-859>.
- Li Z, Wu S, Bai X, Liu Y, Lu J, Liu Y, Xiao B, Lu X, Fan L. 2011. Genome sequence of the tobacco bacterial wilt pathogen *Ralstonia solanacearum*. *J. Bacteriol.* 193:6088–6089. <http://dx.doi.org/10.1128/JB.06009-11>.
- Cao Y, Tian B, Liu Y, Cai L, Wang H, Lu N, Wang M, Shang S, Luo Z, Shi J. 2013. Genome sequencing of *Ralstonia solanacearum* FQY_4, isolated from a bacterial wilt nursery used for breeding crop resistance.

- Genome Announc. 1(3):e00125-13. <http://dx.doi.org/10.1128/genomeA.00125-13>.
5. Shan W, Yang X, Ma W, Yang Y, Guo X, Guo J, Zheng H, Li G, Xie B. 2013. Draft genome sequence of *Ralstonia solanacearum* race 4 biovar 4 strain SD54. Genome Announc. 1(6):e00890-13. <http://dx.doi.org/10.1128/genomeA.00890-13>.
6. Ramesh R, Phadke GS. 2012. Rhizosphere and endophytic bacteria for the suppression of eggplant wilt caused by *Ralstonia solanacearum*. Crop Protect. 37:35–41. <http://dx.doi.org/10.1016/j.cropro.2012.02.008>.
7. Sallet E, Roux B, Sauviac L, Jardinaud MF, Carrère S, Faraut T, De Carvalho-Niebel F, Gouzy J, Gamas P, Capela D, Bruand C, Schiex T. 2013. Next-generation annotation of prokaryotic genomes with EuGene-P: application to *Sinorhizobium meliloti* 2011. DNA Res. 20:339–354. <http://dx.doi.org/10.1093/dnares/dst014>.