



**HAL**  
open science

## Data in support of peptidomic analysis of spermatozoa during epididymal maturation

Valérie Labas, Lucie Spina, Clémence Belleannée, Ana-Paula Teixeira-Gomes, Audrey Gargaros, Françoise Dacheux, Jean-Louis Dacheux

### ► To cite this version:

Valérie Labas, Lucie Spina, Clémence Belleannée, Ana-Paula Teixeira-Gomes, Audrey Gargaros, et al.. Data in support of peptidomic analysis of spermatozoa during epididymal maturation. Data in Brief, 2014, 1, pp.79-84. 10.1016/j.dib.2014.10.003 . hal-02629498

**HAL Id: hal-02629498**

**<https://hal.inrae.fr/hal-02629498>**

Submitted on 27 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

## Data in support of peptidomic analysis of spermatozoa during epididymal maturation



Valérie Labas<sup>a,b,c,d,e,\*</sup>, Lucie Spina<sup>a,b,c,d,e</sup>,  
Clémence Belleannee<sup>b,c,d,e,1</sup>, Ana-Paula Teixeira-Gomes<sup>a,f,g</sup>,  
Audrey Gargaros<sup>a,b,c,d,e</sup>, Françoise Dacheux<sup>b,c,d,e</sup>,  
Jean-Louis Dacheux<sup>b,c,d,e</sup>

<sup>a</sup> INRA, Plateforme d'Analyse Intégrative des Biomolécules, Laboratoire de Spectrométrie de Masse, F-37380 Nouzilly, France

<sup>b</sup> INRA, UMR85 Physiologie de la Reproduction et des Comportements, F-37380 Nouzilly, France

<sup>c</sup> CNRS, UMR7247, F-37380 Nouzilly, France

<sup>d</sup> Université François Rabelais de Tours, F-37000 Tours, France

<sup>e</sup> IFCE, F-37380 Nouzilly, France

<sup>f</sup> INRA, UMR1282 Infectiologie et Santé Publique, F-37380 Nouzilly, France

<sup>g</sup> Université François Rabelais de Tours, UMR1282 Infectiologie et Santé Publique, F-37000 Tours, France

## ARTICLE INFO

## Article history:

Received 6 October 2014

Accepted 7 October 2014

Available online 6 November 2014

## Keywords:

Peptidome  
Spermatozoa  
Maturation  
Epididymis

## ABSTRACT

The final differentiation of the male germ cell occurs in the epididymal duct where the spermatozoa develop the ability to be motile and fertilize an ovum. Understanding of these biological processes is the key to understanding and controlling male fertility. Comparative studies between several epididymal maturation states could be an informative approach to finding sperm modifications related to maturation and fertility. Here we show the data from differential peptidomic/proteomic analyses on spermatozoa isolated from 4 epididymal regions (immature to mature stage) using a profiling approach based on MALDI-TOF mass spectrometry and, combined to top-down MS in order to identify peptidofoms and proteofoms. By this way, 172m/z peaks ranging between 2 and 20 kDa were found to be

DOI of original article: <http://dx.doi.org/10.1016/j.jprot.2014.09.031>

\* Corresponding author.

E-mail address: [valerie.labas@tours.inra.fr](mailto:valerie.labas@tours.inra.fr) (V. Labas).

<sup>1</sup> Present address: Centre de Recherche du Centre Hospitalier de l'Université Laval (CHUQ), Département d'Obstétrique-Gynécologie, Faculté de Médecine, Université Laval, Québec, Canada.

<http://dx.doi.org/10.1016/j.dib.2014.10.003>

2352-3409/© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

modified during maturation of sperm. A total of 62  $m/z$  were identified corresponding to 32 different molecular species. The interpretation of these data can be found in the research article published by Labas and colleagues in the Journal of Proteomics in 2014 [1].

© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

## Specifications table

Subject area	Reproductive biology
More specific subject area	Peptidomic analysis of boar epididymal spermatozoa and molecular phenotypes at different maturation stages
Type of data	Top down raw data and.csv tables with identified proteins
How data was acquired	Experiments performed on a LTQ Orbitrap Velos Mass Spectrometer (Thermo Fisher Scientific, Bremen, Germany)
Data format	Raw data .csv results tables of top-down identification
Experimental factors	Boar spermatozoa from 4 different regions of the epididymis
Experimental features	Epididymal spermatozoa peptidome ( <i>Sus scrofa</i> ) and investigation of the boar sperm maturation process
Data source location	N/A
Data accessibility	Data are available here and via the PRIDE partner repository with the dataset identifier PXD001303.

## Value of the data

- First description of peptidome/degradome of epididymal spermatozoa from boar (*Sus scrofa*).
- Molecular phenotypes distinguishing degrees of maturation of spermatozoa during the epididymal transit.
- First description of protease activities involved in maturation of epididymal spermatozoa.

## 1. Data

Mean peak values obtained from intact cells (IC), detergent-soluble extracts (SD) and detergent-insoluble extracts (ID) associated with immature (IC2, SD2, ID2) and mature epididymal spermatozoa (IC9, SD9, ID9) are shown in [Supplementary Table 1](#). Only 172  $m/z$  peak values for which the fold-changes were  $>2$  and which presented at least one significant difference between epididymal samples ( $p < 0.05$ ) are reported in this table. Some of them are identified. The variation index was calculated as the product of the absolute difference between the maximum (max) and minimum (min) peak intensity values multiplied by the fold-change (fold) between the two extreme values. The variations are expressed as linear decrease (LD) or linear increase (LI) when all the mean values for the four epididymal samples were significantly different from each other ( $p < 0.05$ ), expressed as increase (I) or decrease (D) when at least one of the mean values was not significantly different from the others, and intermediate (inter.) when at least one value was different. In the first column,  $m/z$  values in green correspond to  $m/z$  peaks observed only in IC analysis (a total of 135  $m/z$ ). Rows with color in IC, SD and ID columns correspond to specific peaks for the sample preparation.

A list of endogenous biomolecules identified by top-down MS is shown in [Supplementary Table 2](#). This table shows the raw file name associated to the NCBI nr accession number, the gene name, the protein description, the location, the characterized post-translational modifications, the number of *b* ions, *y* ions and total ions, the delta mass (Da and ppm) between the ProSight theoretical mass *M* and the mass *M* observed by nano-ESI-FTMS, the mass *M* (Da) observed by nano-ESI-FTMS, the ProSight theoretical mass *M* (Da), the ProSight PDE score, the *E*-value identification probability, the *p* score, the precursor *m/z* and mass type with corresponding charge state (*z*), the fragmentation mode, the database used for identification, the signal/noise, the ProSight search type mode, the precursor mass type, the precursor mass tolerance (Da or ppm), the fragment mass type, the fragment mass tolerance (ppm), the delta *M* mode and disulfide activation/deactivation, the minimum of matching fragment between observed and theoretical fragmentation mass spectra, the include modified forms activation/deactivation, the sequence, the calculated theoretical average and monoisotopic mass  $[M+H]^+$  (Da) and the mass  $[M+H]^+$  (Da) observed previously by MALDI-MS.

## 2. Materials and methods: organ sampling and sperm preparation

The epididymides were collected from four one-year-old adult boars. The luminal contents of the tubules of four epididymal regions (E2, E4, E6 and E9) were collected by micro perfusion [2]. Spermatozoa were isolated by centrifugation. The pellets were washed once with PBS (140 mM NaCl, 15 mM KCl, 7 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.5 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4), centrifuged and resuspended in PBS. To remove protein contamination, the spermatozoa were centrifuged on 40% Percoll in PBS. The pellets were washed again with PBS and resuspended in 20 mM Tris-HCl, pH 6.8, and 260 mM sucrose (Tris-sucrose buffer (TS)) (Fig. 1).




Fig. 1. Experimental design.

### 3. Whole and fractionated cell preparations

For whole cell analysis, about  $2 \times 10^6$  spermatozoa were spotted for each epididymal region onto the MALDI sample probe and immediately mixed with a sinapinic acid matrix solution.

For fractionated cell preparations, sperm suspensions in TS buffer were centrifuged and about  $1 \times 10^7$  spermatozoa were resuspended in 1 volume of 2% octyl- $\beta$ -D-1-thioglucoopyranoside in 20 mM Tris-HCl, pH 7, with 2 mM protease inhibitors (2 mM 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride, 0.3  $\mu$ M aprotinin, 130  $\mu$ M bestatine hydrochloride, 14  $\mu$ M E-64, 1 mM EDTA, 0.9  $\mu$ M leupeptine hemisulfate and 1 mM phenylmethanesulfonyl fluoride). The mixture was agitated at 4 °C for 1 h and centrifuged. Supernatants corresponding to enriched membrane and cytosolic proteins (detergent-soluble extract (SD)) were dried and resuspended in TFA 0.1%. The pellet (detergent-insoluble extract (ID)), corresponding to nuclei, mitochondria and cytoskeleton, was washed with TS and centrifuged.

### 4. MALDI-TOF profiling of whole sperm cells and sub-cellular fractions

All samples were analyzed by a MALDI-TOF mass spectrometer (Waters Corporation, Micromass Ltd., Manchester, UK) operating in positive linear mode as previously described [1]. Spectral profiles were collected in the 2000–20,000  $m/z$  mass range. Data processing was performed using MassLynx™ 4.0 software. To increase mass accuracy, internal calibration was performed. Thus, the major unknown 6797  $m/z$  constant was used as “lock mass” for all spectra.

Intact spermatozoa and corresponding detergent-soluble and -insoluble extracts obtained from 4 epididymal regions (E2, E4, E6 and E9) of both epididymes from 4 animals were analyzed by MALDI-TOF MS, with 8 replicates for each of the three sample preparations. A total of 768 spectra were generated in this study. The spectra were analyzed by Progenesis MALDI™ 1.2 software (NonLinear Dynamics) as previously described [1]. After alignment of all spectra, a total of 253  $m/z$  peaks were detected (Supplementary Table 1). A total of 135  $m/z$  molecular species were characterized by Intact Cell MALDI-MS and 118  $m/z$  were newly observed by MALDI MS from SD and ID fractions.

### 5. Quantitative analysis linked to the sperm maturation process

In order to characterize peak differences between epididymal spermatozoa, the intensity of each normalized peak was subjected to one way analysis of variance with Progenesis (factor epididymal regions) and to three-way analysis of variance with R software (factors being animals, epididymis, epididymal regions, the replicate interactions being the residuals), as previously described [1]. All peak signals with a differential fold-change greater than two average values of normalized intensity and a  $p$  value  $< 0.05$  were selected (Supplementary Table 1). Thus 89  $m/z$  peaks were retained for intact cells (IC), 112  $m/z$  peaks for detergent-soluble extracts (SD) and 59  $m/z$  peaks for detergent-insoluble extracts (ID) for a total of 172 unique  $m/z$  peaks (Supplementary Table 1).

### 6. Top-down mass spectrometry

Identification of peptidofoms and proteofoms (endogenous species) was achieved by acquiring nano-ESI tandem high resolution mass spectrometry (MS and MS/MS). Molecular species were previously extracted with 1% and 5% formic acid from IC (region 2 or 9) or from ID fraction (region 9), desalted and concentrated using ZipTip C4 (Millipore Corporation, Billerica, MA). Eluted peptides and proteins were directly analyzed using a LTQ Orbitrap Velos mass spectrometer (Thermo Fisher Scientific, Germany) operating in positive mode, as previously described (1). Data were acquired

using Xcalibur software v2.1 (Thermo Fisher Scientific, San Jose, CA). All analyses were performed manually using a high-high strategy, meaning that a MS spectrum in the 400–2000 $m/z$  mass range was followed by a MS/MS spectrum obtained by High energy Collisional Dissociation (HCD). Thus 412 $m/z$  corresponding to 217 non-redundant molecular ions were selected to induce HCD fragmentation.

Identification and structural characterization were performed using ProSight PC software 2.0 (Thermo Scientific, San Jose). Raw data files were processed by THRASH (signal/noise: 2–3), and data were compared to a simple annotated “*Sus scrofa*” house database generated from NCBI using Proteome Discoverer (Thermo Fisher Scientific). Automated searches were performed using the “Absolute Mass and Biomarker” search options. The mass tolerances were set at 5 ppm for the monoisotopic precursors, 5 Da for the average precursor and 15 ppm for fragment ions mass tolerance. Disulfide modifications and N-terminal post-translation modifications (acetylation and initial methionine cleavage) were activated. Post-translational modifications such as phosphorylation and disulfide bridges were confirmed using the manual Single Protein mode. Proposed sequences with  $E$ -value  $< 1 \times 10^{-2}$  were considered positively identified with a minimum of 10 matching fragment ions. The data were deposited with the ProteomeXchange Consortium (<http://proteomecentral.proteomexchange.org>) via the PRIDE partner repository [3,4] with the dataset identifier PXD001303.

A total of 62 $m/z$  were identified and attributed to 32 different molecular species corresponding to three intact and whole proteins and 58 peptides from 29 proteins (Supplementary Table 2). Forty-five of these peptides presented tryptic or semi-tryptic cleavages, suggesting protease activities by trypsin-like or kallikrein enzymes (Supplementary Table 2).

Gene symbols were mapped for peptidofoms and proteofoms identified by top-down MS, and analyzed using the online PANTHER classification system (database version 9.0; <http://www.pantherdb.org/>) [5] (Fig. 2). Go terms from the Biological Process and the Molecular Function domains were considered. The background dataset for the analysis was the *Homo sapiens* and *Sus scrofa* genomes.

### BIOLOGICAL PROCESS



- **Biological regulation** (AKAP3, AKAP4, GATA1, PRKAR2A)
- **Cellular component organization or biogenesis** (TPPP2)
- **Cellular process** (AKAP3, AKAP4, CALM1, LOC100626147, PRKAR2A, TPPP2)
- **Developmental process** (GATA1, TPPP2)
- **Immune system process** (TrxR1)
- **Localization** (AKAP3, AKAP4, ATP5E, MTMR11)
- **Metabolic process** (ACR, AKAP3, AKAP4, ATP5E, COX7C, GAPDHS, GAT1, GATA1, LOC100523928, MTMR11, PRKAR2A, TrxR1, UBC)
- **Reproduction** (MTMR11)

### MOLECULAR FUNCTION



- **Binding** (AKAP3, AKAP4, CALM1, GATA1, PRKAR2A, TPPP2, UBC)
- **Catalytic activity** (ACR, AKAP3, AKAP4, ATP5E, COX7C, GAPDHS, GATA1, LOC100523928, PRKAR2A, TrxR1)
- **Enzyme regulator activity** (AKAP3, AKAP4, PRKAR2A)
- **Nucleic acid binding transcription factor activity** (GATA1)
- **Receptor activity** (LOC100626147)
- **Structural molecule activity** (TPPP2, UBC)
- **Transporter activity** (ATP5E)

**Fig. 2.** Classification of identified molecular species based on biological process and molecular function using PANTHER classification system.

### Conflict of interest

The authors declare that they have no conflicts of interest.

## Appendix A. Supplementary information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2014.10.003>.

## References

- [1] V. Labas, L. Spina, C. Bellannée, A.P. Teixeira-Gomes, A. Gargaros, F. Dacheux, et al., Analysis of epididymal sperm maturation by MALDI profiling and top-down mass spectrometry, *J. Proteomics* 113 (2015) 226–243, <http://dx.doi.org/10.1016/j.jprot.2014.09.031>.
- [2] J. Dacheux, An in vitro luminal perfusion technique to study epididymal secretion, *IRCS Med* 8 (1980) 137.
- [3] J.A. Vizcaino, R. Cote, F. Reisinger, H. Barsnes, J.M. Foster, J. Rameseder, et al., The proteomics Identifications database: 2010 update, *Nucleic Acids Res.* 38 (2010) D736–D742.
- [4] J.A. Vizcaino, R.G. Cote, A. Csordas, J.A. Dienes, A. Fabregat, J.M. Foster, et al., The PRoteomics IDentifications (PRIDE) database and associated tools: status in 2013, *Nucleic Acids Res.* 41 (2013) D1063–D1069.
- [5] H. Mi, A. Muruganujan, P.D. Thomas, PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees, *Nucleic Acids Res.* 41 (2013) D377–D386.