



HAL
open science

Proteome data to explore the impact of pBClin15 on Bacillus cereus ATCC 14579

Jean-Paul Madeira, Beatrice Alpha-Bazin, J. Armengaud, Helene Omer,
Catherine Duport

► **To cite this version:**

Jean-Paul Madeira, Beatrice Alpha-Bazin, J. Armengaud, Helene Omer, Catherine Duport. Proteome data to explore the impact of pBClin15 on Bacillus cereus ATCC 14579. *Data in Brief*, 2016, 8, pp.1243-1246. 10.1016/j.dib.2016.07.042 . hal-02629907

HAL Id: hal-02629907

<https://hal.inrae.fr/hal-02629907>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Proteome data to explore the impact of pBClin15 on *Bacillus cereus* ATCC 14579Jean-Paul Madeira^{a,b}, Béatrice Alpha-Bazin^b,
Jean Armengaud^b, H el ene Omer^a, Catherine Duport^{a,*}^a SQPOV, UMR0408, Avignon Universit e, INRA, F-84914 Avignon, France^b CEA, DSV, IBItec-S, SPI, Li2D, Laboratory "Innovative technologies for Detection and Diagnostics", Bagnols-sur-C eze F-30200, France

ARTICLE INFO

Article history:

Received 17 June 2016

Received in revised form

13 July 2016

Accepted 19 July 2016

Available online 26 July 2016

ABSTRACT

This data article reports changes in the cellular and exoproteome of *B. cereus* cured from pBClin15. Time-course changes of proteins were assessed by high-throughput nanoLC-MS/MS. We report all the peptides and proteins identified and quantified in *B. cereus* with and without pBClin15. Proteins were classified into functional groups using the information available in the KEGG classification and we reported their abundance in term of normalized spectral abundance factor. The repertoire of experimentally confirmed proteins of *B. cereus* presented here is the largest ever reported, and provides new insights into the interplay between pBClin15 and its host *B. cereus* ATCC 14579. The data reported here is related to a published shotgun proteomics analysis regarding the role of pBClin15, "Deciphering the interactions between the *Bacillus cereus* linear plasmid, pBClin15, and its host by high-throughput comparative proteomics" Madeira et al. [1]. All the associated mass spectrometry data have been deposited in the ProteomeXchange Consortium (<http://proteomecentral.proteomexchange.org>) via the PRIDE partner repository (<http://www.ebi.ac.uk/pride/>), with the dataset identifier PRIDE: PXD001568, PRIDE: PXD002788 and PRIDE: PXD002789.

  2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.jprot.2016.06.022>

* Correspondence to: UMR SQPOV -INRA PACA, 228, route de l'A eroport, CS 40509, Domaine Saint Paul-Site Agroparc, 84914 Avignon Cedex 9, France.

E-mail address: catherine.duport@univ-avignon.fr (C. Duport).

<http://dx.doi.org/10.1016/j.dib.2016.07.042>

2352-3409/  2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Proteomics
More specific subject area	Microbial proteomics
Type of data	Figure, Tables
How data was acquired	NanoLC-MS/MS using an LTQ-Orbitrap XL hybrid mass spectrometer (ThermoFisher) coupled to an Ultimate 3000 nRSLC system (Dionex, ThermoFisher).
Data format	Analyzed
Experimental factors	<i>Bacillus cereus</i> cells with and without pBclin15 plasmid
Experimental features	Proteins were extracted from bacterial cultures harvested at the early exponential, late exponential and stationary growth phases. The extracellular proteins were obtained by trichloroacetic acid precipitation of culture supernatant. The cellular proteins were obtained after disruption of bacteria with a Precellys 24 disruptor (Bertin Technologies).
Data source location	France
Data accessibility	Analyzed datasets are within this article and raw data are available via the PRIDE partner repository (http://www.ebi.ac.uk/pride) with the dataset identifiers, PRIDE: PXD001568, PRIDE: PXD002788 and PRIDE: PXD002789.

Value of the data

- This is the first proteomic study that assesses the influence of plasmid curation on a bacterium.
- This large proteomic dataset on *B. cereus* is a valuable resource for understanding the relationships between linear plasmids and bacterial cells.
- The data are presented as a reference for other investigators who like to check other functional changes of *B. cereus* whole-cell proteome or exoproteome.

1. Data

In this paper we provide the list of the 44 proteins of *B. cereus* that we found previously by proteogenomics (unpublished work) and the lists of proteins identified in the cellular proteome and exoproteome of the Δ pBclin15 and wild-type ATCC 14579 strains [1]. These files comprise label-free quantitation of the proteins based on spectral counts estimated for the three biological replicates in early exponential, late exponential and stationary growth phase. The average \log_2 -abundance level for each protein across all samples, *t*-statistics, *p*-values, *p*-value adjusted for multiple testing, and *B*-statistics are provided [2].

2. Experimental design, materials and methods

B. cereus cells (with and without pBclin15, [3,4]) were cultured in batches as described previously [5] with three independent cultures for each strain. Fig. 1 depicts the collection points depending on the time-growth curve, *i.e.* during early exponential growth phase (EE), at the late exponential growth phase (LE) signifying the transition between exponential and stationary phases, and during the stationary phase (S). At each time-point, cells and culture supernatant were collected resulting into two fractions: cellular soluble proteome and exoproteome, respectively.




Fig. 1. Time course of Δ pBclin15 and its parental strain, ATCC 14579 (WT). The strains were cultured in MOD medium supplemented with 30 mM glucose under aerobiosis. Samples from Δ pBclin15 and WT were isolated from the early exponential (EE), late exponential (LE) and stationary (S) growth phases as indicated by the arrows.

2.1. Sample preparation and shotgun tandem mass spectrometry

Soluble proteins were extracted and processed as described previously [5–7]. Briefly, total protein samples were loaded onto denaturing NuPAGE 4–12% Bis-Tris gels (Invitrogen) for a 3 mm electrophoretic migration. Proteins were then subjected to proteolysis with sequencing grade trypsin (Roche) using ProteaseMAX surfactant (Promega) at 0.01% [7,8]. The resulting peptides were analyzed by tandem mass spectrometry as described previously [5,9]. The peptides from the extracellular digests and those from cellular digests were resolved by reverse chromatography using a 90-min gradient, or a 180-min gradient, respectively, from 4 to 40% solvent B (0.01% HCOOH, 100% CH₃CN) with solvent A being 0.01% HCOOH, 100% H₂O.

2.2. Database searching and criteria

Tandem mass spectrometry raw data were assigned to peptide sequences with the MASCOT search engine (version 2.3.02) from Matrix Science, the following parameters: full-trypsin specificity, a mass tolerance of 5 ppm on the parent ion and 0.5 Da on the MS/MS, carboxyamidomethylated Cys (+57.0215) as a fixed modification and oxidized methionine (+15.9949) as a variable modification, and a maximum of two missed cleavages. All peptide matches with a score below a *p*-value of 0.05 were filtered by the IRMa 1.28.0 parser [10]. A protein was considered validated when at least two different peptides were identified when considering all the samples. In terms of protein identification, the false-positive rate estimated using the appropriate decoy database resulted below 0.1%.

2.3. Data analysis

Spectral counts corresponding to the number of MS/MS spectra per protein were extracted in the three different nanoLC-MS/MS biological replicates for each growth phase. The normalized spectral abundance factor (NSAF) for each protein was calculated by dividing the number of spectral count (SC) by the protein length (L), divided by the sum of SC/L for all N proteins in the experiment [11]. Analyses of abundance level change of proteins were performed with the LIMMA package by the LIMMA Voom method [12]. The data were normalized with the trimmed mean of M-values (TMM) normalization. Then, quantitative proteomics data were analyzed by empirical Bayes moderation of the standard errors towards a common value.

Transparency document. Supporting information

Transparency data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.07.042>.

Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.07.042>.

References

- [1] J.P. Madeira, H. Omer, B. Alpha-Bazin, J. Armengaud, C. Duport, Deciphering the interactions between the *Bacillus cereus* linear plasmid, pBClin15, and its host by high-throughput comparative proteomics, *J. Proteom.* 146 (2016) 25–33.
- [2] G.K. Smyth, Linear models and empirical bayes methods for assessing differential expression in microarray experiments, *Stat. Appl. Genet. Mol. Biol.* 3 (2004), Article3.
- [3] N. Ivanova, A. Sorokin, I. Anderson, N. Galleron, B. Candelon, V. Kapatral, et al., Genome sequence of *Bacillus cereus* and comparative analysis with *Bacillus anthracis*, *Nature* 423 (2003) 87–91.
- [4] H. Omer, B. Alpha-Bazin, J.L. Brunet, J. Armengaud, C. Duport, Proteomics identifies *Bacillus cereus* EntD as a pivotal protein for the production of numerous virulence factors, *Front. Microbiol.* 6 (2015) 1004.
- [5] J.P. Madeira, B. Alpha-Bazin, J. Armengaud, C. Duport, Time dynamics of the *Bacillus cereus* exoproteome are shaped by cellular oxidation, *Front Microbiol.* 6 (2015) 342.
- [6] G. Clair, J. Armengaud, C. Duport, Restricting fermentative potential by proteome remodeling: an adaptative strategy evidenced in *Bacillus cereus*, *Mol. Cell Proteom.* 11 (M111) (2012) 013102.
- [7] G. Clair, S. Roussi, J. Armengaud, C. Duport, Expanding the known repertoire of virulence factors produced by *Bacillus cereus* through early secretome profiling in three redox conditions, *Mol. Cell Proteom.* 9 (2010) 1486–1498.
- [8] A. de Groot, R. Dulermo, P. Ortet, L. Blanchard, P. Guerin, B. Fernandez, et al., Alliance of proteomics and genomics to unravel the specificities of Sahara bacterium *Deinococcus deserti*, *PLoS Genet.* 5 (2009) e1000434.
- [9] A. Dedieu, J.C. Gaillard, T. Pourcher, E. Darrouzet, J. Armengaud, Revisiting iodination sites in thyroglobulin with an organ-oriented shotgun strategy, *J Biol. Chem.* 286 (2011) 259–269.
- [10] V. Dupierris, C. Masselon, M. Court, S. Kieffer-Jaquinod, C. Bruley, A toolbox for validation of mass spectrometry peptides identification and generation of database: IRMa, *Bioinformatics* 25 (2009) 1980–1981.
- [11] B.L. Zybaylov, L. Florens, M.P. Washburn, Quantitative shotgun proteomics using a protease with broad specificity and normalized spectral abundance factors, *Mol. Biosyst.* 3 (2007) 354–360.
- [12] G.K. Smyth, J. Michaud, H.S. Scott, Use of within-array replicate spots for assessing differential expression in microarray experiments, *Bioinformatics* 21 (2005) 2067–2075.