



HAL
open science

High-Quality Draft Genome Sequences of Two Xanthomonas Pathotype Strains Infecting Aroid Plants

I. Robène, Stéphanie S. Bolot, Olivier Pruvost, Matthieu Arlat, Laurent Noel, Sébastien Carrere, Marie Agnes Jacques, R. Koebnik, Lionel Gagnevin

► **To cite this version:**

I. Robène, Stéphanie S. Bolot, Olivier Pruvost, Matthieu Arlat, Laurent Noel, et al.. High-Quality Draft Genome Sequences of Two Xanthomonas Pathotype Strains Infecting Aroid Plants. *Genome Announcements*, 2016, 4 (5), pp.e00902-16. 10.1128/genomeA.00902-16 . hal-02629928

HAL Id: hal-02629928

<https://hal.inrae.fr/hal-02629928v1>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High-Quality Draft Genome Sequences of Two *Xanthomonas* Pathotype Strains Infecting Aroid Plants

I. Robène,^a S. Bolot,^{b,c} O. Pruvost,^a M. Arlat,^{b,c,d} L. D. Noël,^{b,c} S. Carrère,^{b,c} M.-A. Jacques,^e  R. Koebnik,^f  L. Gagnevin^g

CIRAD, UMR CIRAD-UR Peuplements Végétaux et Bioagresseurs en Milieu Tropical (PVBMT), Saint-Pierre, La Réunion, France^a; INRA, Laboratoire des Interactions Plantes Micro-Organismes (LIPM), UMR 441, Castanet-Tolosan, France^b; CNRS, Laboratoire des Interactions Plantes Micro-Organismes (LIPM), UMR 2594, Castanet-Tolosan, France^c; Université Paul Sabatier, Toulouse, France^d; INRA, Institut de Recherche en Horticulture et Semences (IRHS), UMR 1345 SFR 4207 QUASAV, Beaucauzé, France^e; IRD, UMR IRD-CIRAD-UM2 Interactions Plantes-Microorganismes-Environnement (IPME), Montpellier, France^f; CIRAD, UMR IRD-CIRAD-UM2 Interactions Plantes-Microorganismes-Environnement (IPME), Montpellier, France^g

We present here the draft genome sequences of bacterial pathogens of the Araceae family, *Xanthomonas axonopodis* pv. *dieffenbachiae* LMG 695 and *Xanthomonas campestris* pv. *syngonii* LMG 9055, differing in host range. A comparison between genome sequences will help understand the mechanisms involved in tissue specificity and adaptation to host plants.

Received 12 July 2016 Accepted 18 July 2016 Published 1 September 2016

Citation Robène I, Bolot S, Pruvost O, Arlat M, Noël LD, Carrère S, Jacques M-A, Koebnik R, Gagnevin L. 2016. High-quality draft genome sequences of two *Xanthomonas* pathotype strains infecting aroid plants. *Genome Announc* 4(5):e00902-16. doi:10.1128/genomeA.00902-16.

Copyright © 2016 Robène et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to I. Robène, isabelle.robene@cirad.fr.

Among the different members of the aroids, the tropical flower anthurium (*Anthurium andreanum* Linden ex André) is an economically important crop cultivated in tropical and temperate areas. Production of anthurium in the world is threatened by *Anthurium* bacterial blight (ABB), which is caused by *Xanthomonas axonopodis* pv. *dieffenbachiae* (1). Strains of *X. axonopodis* pv. *dieffenbachiae* isolated from anthurium can infect other aroids, and particularly different ornamental species of *Dieffenbachia*, *Caladium*, *Philodendron*, *Syngonium*, or edible aroids, such as *Colocasia* species. Other *Xanthomonas* strains, first classified as *Xanthomonas campestris* pv. *syngonii* (2), have a narrower host range and infect *Syngonium* plants causing the bacterial leaf blight of *Syngonium*. Based on rep-PCR and multilocus sequence analysis (MLSA), the two strains LMG 695 and LMG 9055 grouped in the *X. axonopodis* 9.4 cluster (3, 4). Hybridization values (<70%) between LMG 695 or LMG 9055 and the type strain of *X. axonopodis* LMG 982 did not support their inclusion in *X. axonopodis* (5). A polyphasic taxonomic approach based on MLSA, DNA/DNA hybridization, average nucleotide identity (ANI) values, and phenotypic analyses confirmed these results and proposed to reclassify this cluster as *Xanthomonas phaseoli* (6). In addition to differences in pathogenicity, these data also supported the separation of the two categories of strains represented by LMG 695 and LMG 9055 in distinct pathovars named *dieffenbachiae* and *syngonii*, respectively.

The genomes of both strains were sequenced using the Illumina HiSeq 2000 platform (GATC Biotech, Germany). Shotgun sequencing yielded 25,320,564 read pairs (12,846,409 100-bp paired-end reads with an insert size of 250 bp, and 12,474,156 50-bp mate-pair reads with an insert size of 3 kb) and 30,521,960 read pairs (17,529,832 100-bp paired-end reads with an insert size of 250 bp and 12,992,128 50-bp mate-pair reads with an insert size of 3 kb) for LMG 695 and LMG 9055, respectively.

A combination of Velvet (7), SOAPdenovo, and SOAP GapCloser (8) yielded 10 contigs >500 bp (N_{50} , 1,343,781 bp), with

the largest contig being 1,543,007 bp, for a total assembly size of 5,035,943 bp for strain LMG 695, and 33 contigs >500 bp (N_{50} , 414,753 bp), with the largest contig being 751,399 bp, for a total assembly size of 5,000,894 bp for strain LMG 9055. Genomic contigs were annotated using the EuGene-P annotation pipeline to identify RNAs and protein-coding genes (9). The draft genomes were predicted to contain 4,423 and 4,665 coding sequences (CDSs) for strains LMG 695 and LMG 9055, respectively.

The two genomes have an average nucleotide identity (ANI) of 97.33% (10). The two genomes share >3,700 CDSs, and each contains approximately 200 CDSs, with no orthologs in other *Xanthomonas* species (11). Nineteen and 22 predicted type 3 effector (T3E) genes are present in the LMG 695 and LMG 9055 genomes, respectively. At least 13 of these T3E genes are shared by the two strains.

Accession number(s). These whole-genome shotgun projects have been deposited in GenBank under the accession no. CP014347 for strain LMG 695 and LSLD00000000 for strain LMG 9055. The versions described in this paper are the first versions.

ACKNOWLEDGMENTS

This research was funded by the French Agence Nationale de la Recherche, the European Regional Development Fund 2007–2014 (grant 2007FR161PO004), the Conseil Régional de la Réunion, and CIRAD.

FUNDING INFORMATION

This work, including the efforts of Matthieu Arlat, Marie-Agnès Jacques, Ralf Koebnik, and Lionel Gagnevin, was funded by Agence Nationale de la Recherche (ANR) (ANR-2010-GENM-013 XANTHOMIX).

REFERENCES

- Alvarez AM, Toves PJ, Vowell TS. 2006. Bacterial blight of anthuriums: Hawaii's experience with a global disease. *APSNet Features* <http://dx.doi.org/10.1094/APSnetFeature-2006-0206>.
- Dickey RS, Zumoff CH. 1987. Bacterial leaf blight of *Syngonium* caused

- by a pathovar of *Xanthomonas campestris*. *Phytopathology* 77:1257–1262. <http://dx.doi.org/10.1094/Phyto-77-1257>.
3. Rademaker JL, Louws FJ, Schultz MH, Rossbach U, Vauterin L, Swings J, De Bruijn FJ. 2005. A comprehensive species to strain taxonomic framework for *Xanthomonas*. *Phytopathology* 95:1098–1111. <http://dx.doi.org/10.1094/PHYTO-95-1098>.
 4. Bui Thi Ngoc L, Vernière C, Jouen E, Ah-You N, Lefeuvre P, Chiroleu F, Gagnevin L, Pruvost O. 2010. Amplified fragment length polymorphism and multilocus sequence analysis-based genotypic relatedness among pathogenic variants of *Xanthomonas citri* pv. *citri* and *Xanthomonas campestris* pv. *bilvae*. *Int J Syst Evol Microbiol* 60:515–525. <http://dx.doi.org/10.1099/ijs.0.009514-0>.
 5. Donahoo RS, Jones JB, Lacy GH, Stromberg VK, Norman DJ. 2013. Genetic analyses of *Xanthomonas axonopodis* pv. *dieffenbachiae* strains reveal distinct phylogenetic groups. *Phytopathology* 103:237–244. <http://dx.doi.org/10.1094/PHYTO-08-12-0191-R>.
 6. Constantin EC, Cleenwerck I, Maes M, Baeyen S, Van Malderghem C, De Vos P, Cottyn B. 2015. Genetic characterisation of strains named as *Xanthomonas axonopodis* pv. *dieffenbachiae* leads to a taxonomic revision of the *X. axonopodis* species complex. *Plant Pathol* 65:792–806. <http://dx.doi.org/10.1111/ppa.12461>.
 7. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 8. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu SM, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam TW, Wang J. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read *de novo* assembler. *GigaScience* 1:18. <http://dx.doi.org/10.1186/2047-217X-1-18>.
 9. Sallet E, Roux B, Sauviac L, Jardinaud MF, Carrère S, Faraut T, de Carvalho-Niebel F, Gouzy J, Gamas P, Capela D, Bruand C, Schiex T. 2013. Next-generation annotation of prokaryotic genomes with EuGene-P: application to *Sinorhizobium meliloti* 2011. *DNA Res* 20: 339–354. <http://dx.doi.org/10.1093/dnares/dst014>.
 10. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57: 81–91. <http://dx.doi.org/10.1099/ijs.0.64483-0>.
 11. Blom J, Albaum SP, Doppmeier D, Pühler A, Vorhölter FJ, Zakrzewski M, Goesmann A. 2009. Edgar: a software framework for the comparative analysis of prokaryotic genomes. *BMC Bioinformatics* 10:154. <http://dx.doi.org/10.1186/1471-2105-10-154>.