



HAL
open science

Molecular methods routinely used to detect *Coxiella burnetii* in ticks cross-react with *Coxiella*-like bacteria

Elsa Jourdain, Olivier Duron, Séverine Barry, Daniel González-Acuña, Karim Sidi-Boumedine

► To cite this version:

Elsa Jourdain, Olivier Duron, Séverine Barry, Daniel González-Acuña, Karim Sidi-Boumedine. Molecular methods routinely used to detect *Coxiella burnetii* in ticks cross-react with *Coxiella*-like bacteria. *Infection Ecology & Epidemiology*, 2015, 5 (1), pp.29230. 10.3402/iee.v5.29230 . hal-02630180

HAL Id: hal-02630180

<https://hal.inrae.fr/hal-02630180v1>

Submitted on 15 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORIGINAL RESEARCH ARTICLE

Molecular methods routinely used to detect *Coxiella burnetii* in ticks cross-react with *Coxiella*-like bacteria

Jourdain Elsa, PhD, DVM^{1*}, Olivier Duron, PhD², Barry Séverine¹,
Daniel González-Acuña, PhD, DVM³ and Karim Sidi-Boumedine, PhD, DVM⁴

¹INRA, UR0346 Animal Epidemiology Unit, Saint Genès Champanelle, France; ²Laboratoire MIVEGEC (Maladies Infectieuses et Vecteurs: Ecologie, Génétique, Evolution et Contrôle), Centre National de la Recherche Scientifique (CNRS) Unité Mixte de Recherche UMR 5290, Université Montpellier 1 – Université Montpellier 2 – Institut pour la Recherche et le Développement, Unité de Recherche UR 224, Montpellier, France; ³Veterinary School, University of Concepción, Chillán, Chile; ⁴Anses, Sophia-Antipolis Laboratory, Animal Q fever Unit, Sophia-Antipolis, France

Background: Q fever is a widespread zoonotic disease caused by *Coxiella burnetii*. Ticks may act as vectors, and many epidemiological studies aim to assess *C. burnetii* prevalence in ticks. Because ticks may also be infected with *Coxiella*-like bacteria, screening tools that differentiate between *C. burnetii* and *Coxiella*-like bacteria are essential.

Methods: In this study, we screened tick specimens from 10 species (*Ornithodoros rostratus*, *O. peruvianus*, *O. capensis*, *Ixodes ricinus*, *Rhipicephalus annulatus*, *R. decoloratus*, *R. geigy*, *O. sonrai*, *O. occidentalis*, and *Amblyomma cajennense*) known to harbor specific *Coxiella*-like bacteria, by using quantitative PCR primers usually considered to be specific for *C. burnetii* and targeting, respectively, the *IS1111*, *icd*, *scvA*, *p1*, and *GroEL/htpB* genes.

Results: We found that some *Coxiella*-like bacteria, belonging to clades A and C, yield positive PCR results when screened with primers initially believed to be *C. burnetii*-specific.

Conclusions: These results suggest that PCR-based surveys that aim to detect *C. burnetii* in ticks by using currently available methods must be interpreted with caution if the amplified products cannot be sequenced. Future molecular methods that aim at detecting *C. burnetii* need to take into account the possibility that cross-reactions may exist with *Coxiella*-like bacteria.

Keywords: Q fever; tick-borne diseases; tick endosymbiont; false positive; surveillance; PCR primers

Responsible Editor: Mare Löhmus, Institute for Environmental Medicine, Karolinska Institute, Sweden.

*Correspondence to: Jourdain Elsa, UR0346 Animal Epidemiology Unit, INRA, FR-63122 Saint Genès Champanelle, France, Email: elsa.jourdain@clermont.inra.fr

Received: 22 July 2015; Revised: 22 October 2015; Accepted: 27 October 2015; Published: 24 November 2015

Q fever is a worldwide zoonosis caused by *Coxiella burnetii*, a ubiquitous intracellular bacterium that infects humans and a variety of animals. Livestock, especially small ruminants, are the main sources of human infections (1–3). In domestic ruminants, Q fever's major clinical manifestations are abortions and stillbirths, whose occurrence may translate into significant economic losses (1–3). In humans, *C. burnetii* infections range from asymptomatic to severe. Acute forms of the disease may result in high fevers and severe pneumonia or hepatitis, and chronic forms are strongly debilitating and may be fatal when endocarditis develops in patients with underlying heart disease (3, 4). Both animals and humans essentially become infected through the inhalation of airborne particles contaminated with *C. burnetii* (2, 5).

Ticks are historically known to be potential vectors for Q fever. Indeed, the first strain of *C. burnetii* was isolated from a *Dermacentor andersoni* tick in the 1930s (6–8). First assigned to the genus *Rickettsia* (in which infection by arthropods is the rule), it was later recognized as the Q fever etiologic agent and is now considered as *C. burnetii* reference strain (9). Because *C. burnetii* is frequently detected in field-sampled ticks (10–13) and because laboratory experiments have revealed that at least some tick species are competent vectors (6, 14, 15), it is currently considered that ticks may act as vectors and help transmit the bacterium among wildlife and, on occasion, domestic ruminants (2, 16).

Interestingly, ticks also frequently carry *Coxiella*-like bacteria that are likely involved in mutualistic symbioses

with their arthropod hosts (16–19). To date, there has been no indication that these tick-carried *Coxiella*-like bacteria are transmitted to vertebrates, although *Coxiella*-like bacteria have sporadically been detected in pet birds (19, 20). Recent investigations based on multilocus phylogenetic analyses and whole genome sequencing data revealed that all known *C. burnetii* strains originate within the vast group of *Coxiella*-like endosymbionts and are the descendants of a *Coxiella*-like progenitor hosted by ticks (17).

Because epidemiologic studies that aim at assessing *C. burnetii* prevalence in ticks frequently rely on DNA detection by polymerase chain reaction (PCR), it is important to make sure that these screening methods are specific for *C. burnetii*. The objective of this study was to determine whether five molecular methods frequently used to detect or characterize *C. burnetii* cross-react with *Coxiella*-like bacteria present in ticks.

Materials and methods

Selection of a panel of 20 ticks infected with specific *Coxiella*-like bacteria

Ten tick species, previously shown to harbor specific *Coxiella*-like bacteria (17), were investigated.

They were selected with the aim to represent the four clades (A–D) currently described for the *Coxiella* genus (17) (Fig. 1): clade A (*Ornithodoros rostratus*, *O. peruvianus*, *O. capensis*), clade B (*Ixodes ricinus*), clade C (*Rhipicephalus annulatus*, *R. decoloratus*, *R. geigy*, *O. sonrai*, *O. occidentalis*), and clade D (*Amblyomma*

cajennense). This panel included three tick species from which *C. burnetii* had previously been reported, namely *O. sonrai* (12), *I. ricinus* (21–23), and *R. annulatus* (24).

Two tick specimens were examined for each species. They were either obtained from breeding colonies or sampled from their host species or habitats and they were processed as previously described (17). Briefly, the ticks were first washed with sterile water to avoid external bacterial contamination. Then, DNA was individually extracted using the DNeasy Blood & Tissue Kit (Qiagen) following manufacturer’s instructions. DNA template quality was verified via PCR amplification of 18S ribosomal RNA or cytochrome oxidase 1 arthropod primers. Nested PCR assays were conducted using primers designed to amplify bacteria from the Coxiellaceae family (i.e. *Coxiella* and its sister genus *Rickettsiella*) and to target the *rpoB* (DNA-directed RNA polymerase beta) gene and the *GroEL/htpB* (60 kDa chaperone heat shock protein B) gene as described elsewhere (17, 25). Sequencing of the PCR products obtained showed that each tick species was infected by a specific *Coxiella*-like bacterium that was genetically related to, but distinct from, *C. burnetii*. None of these tick DNA templates was found infected with *C. burnetii* on the basis of multilocus DNA sequencing (17).

Selection of qPCR primers thought to be specific for *C. burnetii*

The 20 tick specimens were tested using quantitative PCR (qPCR) methods using primers that are usually considered to be specific for *C. burnetii* (Table 1). We used TaqMan




Fig. 1. Genetic relatedness of the 10 tick species used in this study using as reference the phylogenetic network published by Duron *et al.* (17) with concatenated *16S rRNA*, *23S rRNA*, *GroEL/htpB*, *rpoB*, and *dnaK* sequences for 71 tick-borne *Coxiella* strains, 15 *C. burnetii* reference strains, and several bacterial outgroups.

Table 1. Details about the qPCR methods used in the study

Gene	Function	Primer designation	Primers and probe sequences (5'–3')	Fragment length (bp)	Reference	% covering with the endosymbiont of <i>R. turanicus</i> ^b	% covering with the endosymbiont of <i>A. americanum</i> ^c
<i>IS1111</i>	Insertion sequence	Forward primer	Confidential ^a	76	(1)	58	63 ^d
		Reverse primer	Confidential ^a			0	0
		Probe	Confidential ^a			0	63 ^d
<i>icd</i>	Isocitrate dehydrogenase	Forward primer	GACCGACCCATTATCCCT	139	(2)	84	0
		Reverse primer	CGGCGTAGATCTCCATCCA			0	0
		Probe	CGCCCGTCATGAAAAACGTGGTC			0	0
<i>p1</i>	Porine	Qp1-F	CGGCGATTGGCGTTTC	68	(3)	0	0
		Qp1-R	GGTTGCGGTAATGCCGTAA			12 ^d	0
		Probe	AACTGTTCAAATCCGAAACGAGTCGCA			50 ^d	0
<i>scvA</i>	Chromatin condensation	QscvA-F	TGGAAAGACAAAATGTCCAACAA	69	(3)	52 ^d	0
		QscvA-R	GGTTAGAAGCACCCGGTCGT			0	0
		Probe	ACGTGGAAAAGACCAACG			67 ^d	0
<i>GroEL/htpB</i>	Heat shock protein	HtpB-1	TGGCTCAAGCGATTTTGTT	82	(4)	65 ^d	0
		HtpB-2	TTATCAATACCCCGTTTCAAATCC			92 ^d	0
		Probe	AAAGCCGTTATTGCTGGAATGAACCCC			70 ^d	0

^aThe detailed protocol used for the amplification of *IS1111* will be soon published by Sidi-Boumedine *et al.* (in preparation) and remains meanwhile confidential; ^bGenBank accession number: CP011126; ^cGenBank accession number: CP007541; ^dsequence positions are distant from each other on the endosymbiont complete genome.

Universal PCR Master Mix (UMM 2 ×) following the amplification protocol: 1 cycle at 50°C for 2 min and 1 cycle at 95°C for 10 min, followed by 40 PCR cycles of 95°C for 15 s and 60°C for 1 min. Two of the targeted markers – the multicopy *IS1111* insertion sequence (26, 27) and the *icd* (isocitrate dehydrogenase) housekeeping gene (26) – are frequently used in surveys that aim to estimate the prevalence of *C. burnetii* infection in ticks (16). The following genes were also targeted: *scvA* (small-cell-variant protein A), which is likely involved in chromatin condensation when the bacterium is ‘sporulating’, and *porine p1*, which encodes an outer membrane protein (28). Finally, we focused on a specific region of the *GroEL/htpB*, distinct from the region targeted to amplify the genome of *Coxiella*-like bacteria (17) and considered to be specific to *C. burnetii* (29). Nine Mile phase II genomic DNA (RSA 493 isolate) was used as a reference. *In silico* comparisons of the primers and probes with currently published sequences of *R. turanicus* (GenBank accession number: CP011126) and *A. americanum* (GenBank accession number: CP007541) suggested that mismatches with these symbionts were unlikely (Table 1).

Results

We found that some *Coxiella*-like bacteria, belonging to clades A and C, yield positive PCR results when screened with primers initially believed to be *C. burnetii*-specific (Table 2). Overall, DNA was amplified for at least one marker in 6 of the 10 tick species studied. The most frequently amplified marker was *IS1111*, which was detected in five different species, whereas *GroEL/htpB* and *scvA* were amplified from three species. *Porine p1* was solely amplified from a *R. geigy* specimen, which was also positive for *htpB*, *scvA*, and *IS1111*, and displayed a particularly low C_t value ($C_t = 33$) for *IS1111*. Conversely, *icd* was not detected in any of our samples. Interestingly, we observed intraspecific variation: one of the *Coxiella*-like

endosymbiont from *R. decoloratus* was positive for *IS1111*, whereas the other was positive for *scvA*. Unfortunately, because all PCR products were poorly concentrated, sequencing was unsuccessful.

Discussion

The marker we most frequently detected in ticks infected with *Coxiella*-like endosymbionts was the *IS1111* transposable element, which is routinely targeted during epidemiological surveys examining *C. burnetii* prevalence in ticks (16). We thus showed that *C. burnetii* detection assays based only on *IS1111* may lead to misidentification with *Coxiella*-like endosymbionts. The recent work of Duron (30) corroborates this finding: several genetically divergent *IS1111* copies were found widespread in many *Coxiella*-like endosymbionts, therefore showing that *IS1111* can no longer be considered specific to *C. burnetii*. These findings may explain why surveys based on *IS1111* screening occasionally report prevalence levels > 10% (23, 31–33).

Our results also showed that the use of a combination of primers targeting different markers, as performed in some studies (11, 21, 34–36), is not sufficient to guarantee the specificity of *C. burnetii* detection. Indeed, up to four of our markers were detected in a same *Coxiella*-like endosymbiont. Interestingly, *icd*, which is frequently used as a PCR target in epidemiological studies (16, 34), was not amplified from our panel of *Coxiella*-like infected ticks. However, Reeves et al. (37) were able to amplify a 612-bp *icd* fragment, displaying 93% homology with *C. burnetii*, from a *Coxiella*-like bacterium that infects ticks from the *O. capensis* complex in South Carolina, USA. This result contrasts with our observation that *icd* was not amplified from the endosymbiont of *O. capensis* ticks sampled from Cape Verde and highlights the fact that the amplification of a specific genetic marker strongly depends on the PCR method (PCR, nested PCR, or qPCR) and the primer sequences used.

Table 2. C_t values obtained using qPCR for both specimens of the 10 tick species tested

<i>Coxiella</i> -like clade	Tick species	<i>IS1111</i>	<i>icd</i>	<i>GroEL/htpB</i>	<i>p1</i>	<i>scvA</i>
A	<i>O. rostratus</i>	–/–	–/–	–/–	–/–	–/–
	<i>O. peruvianus</i>	39/37	–/–	30/31	–/–	–/–
	<i>O. capensis</i>	–/–	–/–	35/35	–/–	–/38
B	<i>I. ricinus</i>	–/–	–/–	–/–	–/–	–/–
C	<i>R. annulatus</i>	37/38	–/–	–/–	–/–	–/–
	<i>R. decoloratus</i>	–/36	–/–	–/–	–/–	39/–
	<i>R. geigy</i>	37/33	–/–	–/38	–/35	–/38
	<i>O. sonrai</i>	39/36	–/–	–/–	–/–	–/–
	<i>O. occidentalis</i>	–/–	–/–	–/–	–/–	–/–
D	<i>A. cajennense</i>	–/–	–/–	–/–	–/–	–/–

The sign ‘/’ is used to separate the results obtained for the first and the second tick specimen; ‘–’ indicates that no amplification was observed.

More generally, negative results may be due to low detection sensitivity, which is supported by high detection thresholds for most of the genes. In particular, intraspecific variation may be due to the individual ticks having a low bacterial burden, sex- or stage-specific differences, and the presence of PCR inhibitors. In our study, it is possible that the *IS1111*-based PCR method was the most sensitive of the tests used, because several copies of this gene are likely present in the genome of *Coxiella*-like bacteria, as is the case for *C. burnetii*. This hypothesis is supported by the observation that *GroEL*/*htpB*, *porine p1*, and *sevA* were detected in the endosymbiont of a *R. geigy* specimen that displayed a low C_t value for *IS1111*.

Standardizing methodology across laboratories is essential to allow comparisons among studies. Although remarkable progress has recently been made in designing new PCR-based techniques to detect *C. burnetii*, these advances have overlooked that an important genetic diversity actually exists within the *Coxiella* genus (16). In this context, it may not be surprising that the PCR primers routinely used to target *C. burnetii* actually cross-react with *Coxiella*-like bacteria. Therefore, in the future, molecular methods aiming at detecting *C. burnetii* should make sure that no cross-reaction exists not only with other abortive agents but also with *Coxiella*-like organisms. Recent full-genome sequencing data indeed not only highlighted obvious genetic similarities of *C. burnetii* with *Coxiella*-like bacteria but also revealed some mutations specific to *Coxiella*-like bacteria (18, 38). This pattern likely explains why PCR cross-reactions with *Coxiella*-like bacteria are partial and variable between markers. Interestingly, identical *IS1111* copies were found in *C. burnetii* and some *Coxiella*-like bacteria (30), suggesting that the risk of detecting *Coxiella*-like bacteria with *IS1111* primers designed to detect *C. burnetii* is very high and must not be underestimated.

PCR-based surveys that aim to detect *C. burnetii* in ticks must be interpreted with caution if the amplified DNA products are not sequenced. Unfortunately, the ratio of bacterial DNA to tick DNA is frequently low, which makes it challenging to obtain PCR products concentrated enough for direct sequencing via conventional PCR. Additionally, currently available qPCR methods, such as those used in this study, often yield very short DNA fragments that are difficult to concentrate for sequencing purposes and that correspond to rather uninformative sequences. Therefore, there is an urgent need to develop a multiplex qPCR or microchip method that would make it possible to directly differentiate *Coxiella*-like bacteria from *C. burnetii* in tick samples and to detect co-infections. Pending development of such a test, useful alternative methods include the sequencing of the *16S rRNA*, *rpoB*, and *GroEL* genes of *Coxiella* bacteria, after amplification by nested PCR, as previously described (17, 25).

Acknowledgements

We thank Elodie Rousset for reading and providing comments on the manuscript; Aurélien Joulie, Sébastien Masegla, and Elise Yang for their involvement in the molecular analyses; and Karen McCoy, Christine Chevillon, Abel Biguezoton, Marcelo Labruna, Patrick Durand, and François Renaud for collecting the ticks.

Conflict of interest and funding

This work was supported by the LABEX ECOFECT (LEGOXiNET project, ANR-11-LABX-0048) of Université de Lyon, which is part of the French Stimulus Initiative program (Investissements d'Avenir; ANR-11-IDEX-0007), operated by the French National Research Agency (ANR) and the FONDECYT project (project number 1130948).

References

- Rodolakis A. Q fever in dairy animals. Rickettsiology and rickettsial diseases – Fifth International Conference. Ann N Y Acad Sci 2009; 1166: 90–3.
- EFSA. Scientific Opinion on Q fever. EFSA J 2010; 8: 1595.
- European Centre for Disease Prevention and Control (2010). Risk assessment on Q fever. Panel with representatives from the Netherlands, France, Germany, UK, United States. Asher, A., Bernard, H., Coutino, R., Durat, G., De Valk, H., Desenclos, J.-C., Holmberg, J., Kirkbridge, H., More, S., Scheenberger, P., van der Hoek, W., van der Poel, C., van Steenberg, J., Villanueva, S., Coulombier, D., Forland, F., Giesecke, J., Jansen, A., Nilsson, M., Guichard, C., Mailles, A., Pouchol, E., Rousset, E. ECDC Technical report, 40 pp. doi: <http://dx.doi.org/10.2900/28860>. Available from: <http://www.ecdc.europa.eu> [cited 13 November 2011].
- Anderson A, Bijlmer H, Fournier PE, Graves S, Hartzell J, Kersh GJ, et al. Diagnosis and management of Q Fever – United States, 2013: recommendations from CDC and the Q Fever Working Group. MMWR Recomm Rep 2013; 62: 1–28.
- Tissot-Dupont H, Amadei MA, Nezri M, Raoult D. Wind in November, Q fever in December. Emerg Infect Dis 2004; 10: 1264–9.
- Davis GE, Cox HR. A filter-passing infectious agent isolated from ticks. I. Isolation from *Dermacentor andersoni*, reactions in animals, and filtration experiments. Public Health Rep 1938; 53: 2259–67.
- Dyer RE. A filter-passing infectious agent isolated from ticks. III. Human infection. Public Health Rep 1938; 53: 2277–82.
- McDade JE. Historical aspects of Q fever. In: Marrie TJ ed. Q fever, Volume 1: the disease. Boca Raton, Florida: CRC Press; 1990, pp. 5–22.
- Hechemy KE. History and prospects of *Coxiella burnetii* research. In: Toman R, Heinzen RA, Samuel JE, Mege J-L eds. *Coxiella burnetii*: recent advances and new perspectives in research of the Q Fever bacterium. Dordrecht: Springer; 2012, pp. 1–11.
- Toma L, Mancini F, Di Luca M, Cecere JG, Bianchi R, Khoury C, et al. Detection of microbial agents in ticks collected from migratory birds in central Italy. Vector-Borne Zoonotic Dis 2014; 14: 199–205.
- Cooper A, Stephens J, Ketheesan N, Govan B. Detection of *Coxiella burnetii* DNA in wildlife and ticks in Northern Queensland, Australia. Vector-Borne Zoonotic Dis 2013; 13: 12–16.

12. Mediannikov O, Fenollar F, Socolovschi C, Diatta G, Bassene H, Molez JF, et al. *Coxiella burnetii* in humans and ticks in rural Senegal. *PLoS Negl Trop Dis* 2010; 4: 8.
13. Ioannou I, Sandalakis V, Kassinis N, Chochlakis D, Papadopoulos B, Loukaides F, et al. Tick-borne bacteria in mouflons and their ectoparasites in cyprus. *J Wildl Dis* 2011; 47: 300–6.
14. Siroky P, Kubelova M, Modry D, Erhart J, Literak I, Spitalska E, et al. Tortoise tick *Hyalomma aegyptium* as long term carrier of Q fever agent *Coxiella burnetii* – evidence from experimental infection. *Parasitol Res* 2010; 107: 1515–20.
15. Daitey AB. Transovarial and transspermal transmission of *Coxiella burnetii* by the tick *Hyalomma asiaticum* and its role in Q-rickettsiosis ecology. *Parazitologiya* 1977; 11: 403–11.
16. Duron O, Sidi-Boumedine K, Rousset E, Moutailler S, Jourdain E. The importance of ticks in Q fever transmission: what has (and has not) been demonstrated? *Trends Parasitol* 2015; 31: 536–52.
17. Duron O, Noël V, McCoy KD, Bonazzi M, Sidi-Boumedine K, Morel O, et al. The recent evolution of a maternally-inherited endosymbiont of ticks led to the emergence of the Q fever pathogen, *Coxiella burnetii*. *PLoS Pathog* 2015; 11: e1004892.
18. Smith TA, Driscoll T, Gillespie JJ, Raghavan R. A *Coxiella*-like endosymbiont is a potential vitamin source for the lone star Tick. *Genome Biol Evol* 2015; 7: 831–8.
19. Zhong J. *Coxiella*-like endosymbionts. In: Toman R, Heinzen RA, Samuel JE, Mege J-L eds. *Coxiella burnetii*: recent advances and new perspectives in research of the Q fever bacterium. Dordrecht: Springer; 2012, pp. 365–79.
20. Shivaprasad HL, Cadenas MB, Diab SS, Nordhausen R, Bradway D, Crespo R, et al. *Coxiella*-like infection in psittacines and a toucan. *Avian Dis* 2008; 52: 426–32.
21. Sprong H, Tjisse-Klasen E, Langelaar M, De Bruin A, Fonville M, Gassner F, et al. Prevalence of *Coxiella burnetii* in ticks after a large outbreak of Q fever. *Zoonoses Public Health* 2012; 59: 69–75.
22. Reye AL, Stegny V, Mishaeva NP, Velhin S, Hubschen JM, Ignatyev G, et al. Prevalence of tick-borne pathogens in *Ixodes ricinus* and *Dermacentor reticulatus* ticks from different geographical locations in Belarus. *PLoS One* 2013; 8: e54476.
23. Bonnet S, de la Fuente J, Nicollet P, Liu X, Madani N, Blanchard B, et al. Prevalence of tick-borne pathogens in adult *Dermacentor* spp. ticks from nine collection sites in France. *Vector-Borne Zoonotic Dis* 2013; 13: 226–36.
24. Reye AL, Arinola OG, Hubschen JM, Muller CP. Pathogen prevalence in ticks collected from the vegetation and livestock in Nigeria. *Appl Environ Microbiol* 2012; 78: 2562–8.
25. Duron O, Jourdain E, McCoy KD. Diversity and global distribution of the *Coxiella* intracellular bacterium in seabird ticks. *Ticks Tick Borne Dis* 2014; 5: 557–63.
26. de Bruin A, de Groot A, de Heer L, Bok J, Wielinga PR, Hamans M, et al. Detection of *Coxiella burnetii* in complex matrices by using multiplex quantitative PCR during a major Q fever outbreak in the Netherlands. *Appl Environ Microbiol* 2011; 77: 6516–23.
27. Joulie A, Laroucau K, Bailly X, Prigent M, Gasqui P, Lepetitcolin E, et al. Circulation of *Coxiella burnetii* in a naturally infected flock of dairy sheep: shedding dynamics, environmental contamination, and genotype diversity. *Appl Environ Microbiol* 2015; 81: 7253–60.
28. Coleman SA, Fischer ER, Howe D, Mead DJ, Heinzen RA. Temporal analysis of *Coxiella burnetii* morphological differentiation. *J Bacteriol* 2004; 186: 7344–52.
29. Fernandes I, Rousset E, Dufour P, Sidi-Boumedine K, Cupo A, Thiery R, et al. Evaluation of the recombinant heat shock protein B (HspB) of *Coxiella burnetii* as a potential antigen for immunodiagnostic of Q fever in goats. *Vet Microbiol* 2009; 134: 300–4.
30. Duron O. The *IS1111* insertion sequence used for detection of *Coxiella burnetii* is widespread in *Coxiella*-like endosymbionts of ticks. *FEMS Microbiol Lett* 2015; 362: fnv132. Available from: <http://femsle.oxfordjournals.org/content/362/17/fnv132>
31. Szymanska-Czerwinska M, Galinska EM, Niemczuk K, Zasepa M. Prevalence of *Coxiella burnetii* infection in foresters and ticks in south-eastern Poland and comparison of diagnostic methods. *Ann Agric Environ Med* 2013; 20: 699–704.
32. Pastiu AI, Matei IA, Mihalca AD, D'Amico G, Dumitrache MO, Kalmar Z, et al. Zoonotic pathogens associated with *Hyalomma aegyptium* in endangered tortoises: evidence for host-switching behaviour in ticks? *Parasit Vectors* 2012; 5: 301.
33. Loftis AD, Reeves WK, Szumlas DE, Abbassy MM, Helmy IM, Moriarity JR, et al. Population survey of Egyptian arthropods for rickettsial agents. *Ann N Y Acad Sci* 2006; 1078: 364–7.
34. Michelet L, Delannoy S, Devillers E, Umhang G, Aspan A, Juremalm M, et al. High-throughput screening of tick-borne pathogens in Europe. *Front Cell Infect Microbiol* 2014; 4: 103.
35. Lee JH, Park HS, Jang WJ, Koh SE, Park TK, Kang SS, et al. Identification of the *Coxiella* sp. detected from *Haemaphysalis longicornis* ticks in Korea. *Microbiol Immunol* 2004; 48: 125–30.
36. Tozer SJ, Lambert SB, Strong CL, Field HE, Sloots TP, Nissen MD. Potential animal and environmental sources of Q fever infection for humans in Queensland. *Zoonoses Public Health* 2014; 61: 105–12.
37. Reeves WK, Loftis AD, Sanders F, Spinks MD, Wills W, Denison AM, et al. *Borrelia*, *Coxiella*, and *Rickettsia* in *Carios capensis* (Acari: Argasidae) from a brown pelican (*Pelecanus occidentalis*) rookery in South Carolina, USA. *Exp Appl Acarol* 2006; 39: 321–9.
38. Gottlieb Y, Lalar I, Klasson L. Distinctive genome reduction rates revealed by genomic analyses of two *Coxiella*-Like endosymbionts in ticks. *Genome Biol Evol* 2015; 7: 1779–96.