

Development, anatomy, and genetic control of some teratological phenotypes of ranunculaceae flowers

Florian Jabbour, Felipe Espinosa, Thierry Deroin, Domenica Manicacci, Sophie Nadot, Catherine Damerval

► **To cite this version:**

Florian Jabbour, Felipe Espinosa, Thierry Deroin, Domenica Manicacci, Sophie Nadot, et al.. Development, anatomy, and genetic control of some teratological phenotypes of ranunculaceae flowers. *Modern Phytomorphology*, 2016, 9, pp.75-75. 10.5281/zenodo.159704 . hal-02633677

HAL Id: hal-02633677


<https://hal.inrae.fr/hal-02633677>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





DEVELOPMENT, ANATOMY, AND GENETIC CONTROL OF SOME TERATOLOGICAL PHENOTYPES OF RANUNCULACEAE FLOWERS

FLORIAN JABBOUR ^{1*}, FELIPE ESPINOSA ¹, THIERRY DEROIN ¹, DOMENICA MANICACCI ²,
SOPHIE NADOT ³, CATHERINE DAMERVAL ²

Key words: Ranunculaceae, evolution, flower, morpho-anatomy, phylogeny, teratology

¹ Institut de Systématique, Evolution, Biodiversité, ISYEB – UMR 7205 – CNRS, MNHN, UPMC, EPHE, Muséum national d'Histoire naturelle, Sorbonne Universités, 57 rue Cuvier, CP39, F-75005 Paris, France; * fjabbour@mnhn.fr

² UMR Génétique Quantitative et Evolution – Le Moulon, INRA-Univ Paris-Sud – CNRS- AgroParisTech, 91190 Gif-sur-Yvette, France

³ Laboratoire Ecologie, Systématique, Evolution, Univ. Paris-Sud, AgroParisTech, CNRS UMR 8079, 91405 Orsay, France

Teratological organisms originate from developmental anomalies, and exhibit structures and a body organization that deviate from the species standard. These monsters give essential clues about the formation and evolutionary significance of the wild-type groundplan. We focus on flower terata, which can be affected in their sterile and/or fertile organs, with special emphasis on the Ranunculaceae. The diversity of perianth shapes and organizations in flowers of this family is huge, and is even increased when anomalies occur during organo- and/or morphogenesis.

To begin with, we synthesize the observations and research conducted on the Ranunculacean floral terata, following the most

recent phylogenetic framework published in 2016 by our team.

Then, we report results regarding the morphology of developing meristems, the anatomy of buds, and the genetic control of selected teratological phenotypes of Ranunculaceae flowers. We focus on species and horticultural varieties belonging to the genera *Aquilegia*, *Delphinium*, and *Nigella*. Wild-type flowers of these species are actinomorphic (*Aquilegia*, *Nigella*) or zygomorphic (*Delphinium*), spurred (*Aquilegia*, *Delphinium*) or with pocket-like petals (*Nigella*).

Last, we discuss the evolutionary potential of such teratological phenotypes when they occur in the wild.