



HAL
open science

Effects of neonatal surgical castration and immunocastration in male pigs on blood T lymphocytes and health markers

Caroline Leclercq, Armelle Prunier, Elodie Merlot

► **To cite this version:**

Caroline Leclercq, Armelle Prunier, Elodie Merlot. Effects of neonatal surgical castration and immunocastration in male pigs on blood T lymphocytes and health markers. *Animal*, 2014, 8 (5), pp.836-843. 10.1017/S1751731114000445 . hal-02634076

HAL Id: hal-02634076

<https://hal.inrae.fr/hal-02634076v1>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of neonatal surgical castration and immunocastration in male pigs on blood T lymphocytes and health markers

C. Leclercq^{1,2}, A. Prunier^{1,2} and E. Merlot^{1,2†}

¹INRA, UMR1348 PEGASE, F-35590 Saint-Gilles, France; ²Agrocampus Ouest, UMR1348 PEGASE, F-35000 Rennes, France

(Received 24 September 2013; Accepted 14 February 2014)

Surgical castration in pig husbandry is criticized for welfare reasons. Thus, it is necessary to evaluate alternative ways of rearing male pigs, such as entire or immunocastrated animals. Immunocastration is a vaccination directed against gonadotropin-releasing hormone (GnRH) to suppress the production of sexual hormones. This study aimed at investigating the effects of these two methods of castration in comparison with intact male pigs on blood T-lymphocyte subsets and function, the immunoglobulin (Ig) response to an influenza vaccine and health markers during sexual development. A total of 70 animals were allocated to three experimental groups: entire (E), surgically castrated at 5 to 6 days of age (SC), and immunized against GnRH at 3 and 4 months of age (IC). Blood samples were collected at 3, 4 and 5 months. At slaughter, global health status and body and spleen weights were measured. Results showed that SC male pigs had fewer blood lymphocytes than E pigs at 4 and 5 months ($P < 0.05$), whereas IC pigs did not differ significantly from E pigs. The percentages of $CD3^+$, $CD3^+CD4^+$ and $CD3^+CD8^+$ lymphocytes were not altered by treatment ($P > 0.1$). Compared with E pigs, the SC pigs had a higher percentage of $CD3^+CD4^+CD8^+$ cells at 4 months, whereas the IC pigs had a higher percentage at 5 months ($P < 0.05$). Regarding $\gamma\delta T$ cells, SC pigs had a lower percentage than E pigs at 4 and 5 months ($P < 0.05$), whereas IC pigs did not differ significantly from E pigs at any age. However, there were no consequences on T-lymphocyte proliferation and total IgG or anti-influenza Ig. At slaughter, relative spleen weight was decreased in IC pigs, whereas pneumonia score was decreased in SC pigs relatively to E pigs. Overall, no clear functional consequences of either method on commercial pig immune abilities were demonstrated, but more investigations are required to ascertain this conclusion.

Keywords: pig, surgical castration, immunocastration, testosterone, T cells

Implications

Surgical castration of male pigs is commonly performed in European farms to avoid unappreciated sexual odors (boar taint) during cooking. Nevertheless, this method is criticized for welfare reasons. One alternative is rearing entire male pigs, provided that the problem of boar taint is solved. A second option is immunocastration that allows boar taint to be reduced. Because testicular hormones exert immunoregulatory functions, the influence of castration and immunocastration on immunity and health was evaluated using blood markers. Overall, the lack of clear negative functional consequences supports the generalization of these new practices.

Introduction

Sex steroids play a significant role in the regulation of the immune function (Bouman *et al.*, 2005) and hence influence

the resistance to disease. Surgical castration performed on rodents during adulthood reverses the age-related thymic atrophy and restores the thymic production of lymphocytes (Sutherland *et al.*, 2005). This leads, in secondary lymphoid organs, to an increase in the frequency of both $CD8^+$ and $CD4^+$ naïve T cells, in the T $CD4/T$ $CD8$ ratio and in the capacity of proliferation of T cells, restoring a juvenile phenotype (Sutherland *et al.*, 2005). Surgical castration of adult rodents also increases the production of B cells in the bone marrow, leading to an increased number of B cells in the spleen and a subsequent splenic enlargement (Viselli *et al.*, 1995). A study carried out on monkeys, using a gonadotropin-releasing hormone (GnRH) antagonist to suppress the neonatal surge of testosterone, showed that neonatal androgens have a long-term influence on the immune development that could be different from the effect of post-pubertal androgens (Gould *et al.*, 1998). Neonatal treatment with an antagonist of GnRH induced a diminution in white blood cells and more so in T- and B-cell numbers during both infancy and adulthood. Very few data describe the effects of testicular hormone deprivation in

† E-mail: elodie.merlot@rennes.inra.fr

pigs on blood T cells. We showed that neonatal surgical castration decreased the number of blood lymphocytes at 4 and 5 months, as well as the thymus weight at 5 months, suggesting that castration might affect T-cell production (Merlot *et al.*, 2013). However, in this study, functional tests involving T cells such as the specific response to vaccination against KLH and the mitogen-induced proliferation of peripheral blood cells were unaltered.

Sex steroid levels vary along the pubertal development. A first peak of androgen secretion occurs during the first few weeks after birth (Colenbrander *et al.*, 1978). Thereafter, not only testosterone but also estradiol levels progressively increase during pubertal development of male pigs, from about 2 until 6 months of age (Allrich *et al.*, 1982; Brunius *et al.*, 2011). Immunocastration is based on vaccination against GnRH resulting in the suppression of testicular activity. The primary immunization is usually carried out around 3 months of age. The second injection is carried out some few weeks later and leads to a rapid drop (within a couple of days) in LH and testosterone (Claus *et al.*, 2007). Thus, surgical castration suppresses the gonadal steroids from the neonatal period, whereas GnRH and LH are preserved and should even be increased owing to the lack of negative feedback of the testicular hormones on the hypothalamus and pituitary (Colenbrander *et al.*, 1987). Immunocastration allows natural activity of the hypothalamic–pituitary–gonadal (HPG) axis until the second injection of the vaccine, followed by a suppression of both pituitary and testicular hormones. Because of the differences in the timing and nature of the suppressed hormones, the two methods of castration may have different effects on the immune system.

The present experiment was carried out to investigate the consequences of two methods of castration that are used in pigs, surgical castration and immunocastration, on peripheral T cells. We hypothesized that the frequency and function of T cells would be similar in immunocastrated and entire male pigs until the second vaccine injection, whereas they could differ between surgically castrated and entire pigs from 3 months of age. The effects of the different treatments were observed from 3 to 5 months of age, as fattening pigs are usually slaughtered at around 5 months. We determined white blood cell counts, proportions of blood T-cell subsets, proliferation of T lymphocytes, markers relevant to evaluating the health status (total IgG, haptoglobin levels and clinical measures at slaughter), and the immune response to a vaccination against influenza.

Material and methods

The experiment was conducted following French guidelines for animal care and use (<http://ethique.ipbs.fr/sdv/charteexpeanimale.pdf>) in the experimental unit of INRA – Saint Gilles (35590 France). People involved in the experiment were holders of an animal experiment agreement delivered by the veterinary services of the French Ministry of Agriculture.

Animals, experimental design and sampling procedures

Two experiments were conducted on pigs born from Large White × Landrace sows inseminated with Pietrain semen.

Experiment A involved three experimental groups: 18 entire (E), 16 immunocastrated (IC) and 14 surgically castrated (SC) males. This experiment was run in four repetitions undertaken from December 2010 to July 2011. For each replicate, four litters were selected at farrowing, in which a third of the male piglets were surgically castrated at 5 to 6 days of age following the standard procedure used in pig farming. After weaning at 26 to 28 days of age, animals were always housed with pen-mates of the same gonadal status (entire and surgically castrated). At 68 to 71 days of age (29 ± 5 kg live weight), two to five male pigs were chosen in each of the four litters (depending on the number of male pigs available per litter) and assigned to experimental groups (SC, IC and E) on the basis of their BW and gonadal status to equilibrate, as well as possible, BW and litters of origin among treatments. They were transferred into the experimental unit, where they were housed in groups of two or three pigs from the same treatment. Pigs received *ad libitum* standard growth diet. Water was provided *ad libitum*. The primary and the booster immunizations of IC pigs against GnRH (Improvac[®]; Pfizer, Belgium, SC route) were carried out at 87 to 90 and 115 to 118 days of age. All the pigs were vaccinated against influenza (Gripovac; Merial, Lyon, France) at 129 to 132 and 143 to 146 days of age.

Blood samples were drawn by venipuncture in the jugular vein on EDTA-containing (hormonal assay and white blood cell counts) and heparinized (flow cytometry and lymphocyte proliferation) tubes at 87 to 90, 115 to 118 (before each injection of Improvac) and 150 to 153 days of age. At the second and third blood samplings, samples were also drawn in tubes without anti-coagulant for total immunoglobulin (Ig) G and anti-influenza Ig determinations. Plasma and sera were stored at -20°C until use. At 151 to 154 days of age, pigs were transferred to the slaughterhouse and were euthanized the following day by exsanguination after electronarcosis.

Experiment B was run to increase the number of pigs regarding white blood cell counts and lymphocyte subset analyses, and involved 10 entire male pigs and 10 surgically castrated male pigs originating from six different litters. In each litter, half of the male pigs were neonatally castrated, and at 68 to 71 days of age male piglets were chosen (one SC and one E, or two SC and two E per litter, depending of the number of male pigs available in the litter) and assigned to the E and SC experimental groups. This experiment was conducted from February to July 2012 in one replicate in the same rearing conditions as experiment A, except that animals were not vaccinated against influenza. Blood samples were drawn on EDTA-containing (white blood cell counts) and heparinized (flow cytometric analysis) tubes at 115 to 118 and 150 to 153 days of age.

White blood cell count and immunoassays

Total numbers of leukocytes, lymphocytes and granulocytes were measured with a hematology automatic cell counter

(MS-9; MeletSchloesing Laboratories, Osny, France). The coulter allows discriminating particles according to their electric impedance that is proportional to their volume, and has been calibrated for growing pigs by the manufacturer. The lymphocyte and granulocyte counts carried out by the coulter and by conventional microscopy in a veterinary laboratory gave correlations with a R^2 of 0.92 and 0.94, respectively. Total IgG levels were assessed by ELISA (Bethyl E100-104, pig IgG Fc Elisa quantitation; Interchim, Montluçon, France), as described previously (Courret *et al.*, 2009). Anti-influenza Ig levels were assessed by ELISA (Swine influenza Virus Antibody Test Kit – H1N1, IDEXX Europe, Hoofddorp, The Netherlands). The relative level of antibody in the sample was determined by calculating the sample to positive standard ratio of optical densities. Plasma testosterone was measured by the AIA 360 automaton (Tosoh Europe, Tessenderlo, Belgium) using a commercial kit (ST AIA – testosterone pack; Tosoh Europe), and haptoglobin was assessed using a commercial kit (Tridelta Ltd., AbCys s.a, Paris, France).

Proliferation assay and flow cytometry

All products were purchased from SIGMA Aldrich (St Louis, Missouri, USA). Peripheral blood mononuclear cells (PBMC) were isolated from heparinized blood by gradient centrifugation on histopaque 1077. They were adjusted at a concentration of 10×10^6 cells/ml RPMI 1640 supplemented with 10% fetal bovine serum 1% penicillin/streptomycin and 1% L-glutamine as described previously (Courret *et al.*, 2009). A viability analyzer using trypan-blue exclusion test (Vi-Cell XR; Beckman Coulter, Paris, France) was used to determine cell numbers and viability, which was always higher than 95%.

A total of 0.5×10^6 cells per well in a final volume of 200 μ l were cultured in triplicate in the presence or absence of the T-cell mitogen Concanavalin A (ConA 6.25 μ g/ml) and, after 48 h of incubation, were pulsed with tritiated thymidine (1 μ Ci per well) for 24 h. Plates were stored at -20°C , and then filtrated on filter glass paper (MultiScreen Filter Plates; Millipore, Molsheim, France). Radioactivity was counted (counts per minute = c.p.m.) in a microplate scintillation counter (TopCount from Perkin Elmer, Courtaboeuf, France). Mitogenic responsiveness was expressed as a proliferation index: PI = (c.p.m. of stimulated cells)/(c.p.m. of unstimulated cells).

The monoclonal antibodies (mAb) used to detect cell surface markers were SpectralRedTM (SPRD)-conjugated anti-swine CD3 (clone PPT3, IgG1), FITC-conjugated anti-swine CD4 (clone 74-12-4, IgG2b) and PE-conjugated anti-swine CD8 α (clone 76-2-1, IgG2a) that were purchased from Southern Biotech (Birmingham, Alabama). Anti-swine $\gamma\delta$ T-cell receptor (TCR) antibody (clone PGBL22A, IgG1, VMRD; Pullman, WA, USA) was used with an anti-mouse IgG secondary antibody conjugated with Alexa Fluor 633 (Invitrogen, Carlsbad, CA, USA). Appropriate isotype control antibodies were used (Southern Biotech). The direct labeling (CD3/CD4/CD8) was performed on 0.5×10^6 cells incubated with saturating

concentrations of anti CD3, anti CD4 and anti CD8 mAb (1/100 dilutions) in a final volume of 50 μ l. The indirect labeling consisted of successive incubations of 1×10^6 cells with anti $\gamma\delta$ TCR (1/100 dilution), anti-mouse IgG (1/1000 dilution) and anti-CD3 (1/100 dilution) antibodies in a final volume of 100 μ l. All incubations lasted 30 min and were carried out at 4°C . After each staining step, cells were washed twice in 250 μ l PBS. Cells were suspended in 400 μ l of fluorescence-activated cell sorter buffer before analysis. Analysis of PBMC was performed using a Macs Quant Analyzer (Miltenyi Biotec, Bergisch Gladbach, Germany) with MacsQuant Quantify software. The strategy of gating for the analysis is presented in Figure 1.

Clinical measurements at slaughter

In experiment A, the lungs were inspected for pericarditis and pleurisy at slaughter. Pneumonia lesions were scored by summing the score attributed to each of the seven lobes according to the consolidation surface area (from 0 for no lesion to 4 for total pulmonary consolidation), leading to an individual severity score ranging from 0 to 28 (Madec and Kobish, 1982). Rhinitis was also estimated by summing the shrinking score attributed to each of the four turbinate bones inside the snout (from 0 for intact to 4 for total atrophy) and the distortion score attributed to the nasal wall (from 0 for no deviation to 2 for severe deviation), leading to an individual severity score on a scale from 0 to 18. Stomach lesions were scored between 0 and 7 according to the severity or absence of ulcer (Henry *et al.*, 1970). The testes and spleen were weighed.

Statistical analysis

Analysis of variance was performed using the mixed procedure of SAS (version 8.1, 2000, SAS Institute Inc., Cary, NC, USA). The fixed factors of the analysis were the treatment and replicate. Normality of the data was checked on residuals with the Shapiro–Wilk test. For blood leukocyte counts and cytometry, data from experiments A and B were analyzed together with experiment B declared as a 5th replicate. Because the disequilibrium owing to the lack of IC pigs in experiment B could have biased the analysis, those results were compared with those of an analysis, taking into account only the repetitions from experiment A. The means that were statistically different when pooling experiment A + B were also significant (% CD3⁺CD4⁺CD8⁺ at 5 months of age, blood lymphocyte numbers at 4 months), close to statistical difference (% $\gamma\delta$ T cells at 5 months) or at least in the same numerical order (blood total leukocyte and lymphocyte counts at 5 months, $\gamma\delta$ T and CD3⁺CD4⁺CD8⁺% at 4 months) in experiment A alone. Thus, pooling experiments A and B increased the statistical power without inducing any bias. The Bonferroni test was used for post-hoc comparisons when the treatment effect in the ANOVA had a P -value < 0.05. Numeric data are presented as least square means \pm standard errors of the mean (s.e.) calculated by SAS for the lowest group size.




Figure 1 Strategy of analysis of lymphocyte subpopulations by flow cytometry. A first gate of lymphoid cells was designed on the forward – side scatter plot (gate P1, 10 000 events acquired). Among lymphoid cells, CD3⁺ T lymphocytes were selected (gate P2 included in P1). The percentages of CD3⁺CD4⁺CD8⁻, CD3⁺CD4⁻CD8⁺ and CD3⁺CD4⁺CD8⁺ cells (a) and of CD3⁺ $\gamma\delta$ TCR⁺ cells (b) were determined among the CD3⁺ population.

Results

Concentration of testosterone from 3 to 5 months of age
E pigs had higher concentration of testosterone at 4 and 5 months of age than SC pigs ($P < 0.001$ and < 0.001 , respectively, Figure 2). Until the second vaccination (at 4 months), testosterone concentration was not different in the E and IC pigs ($P > 0.1$) but higher in E than in SC pigs ($P < 0.001$). At 5 months, the concentration in IC pigs fell to levels significantly lower than those of E pigs ($P < 0.001$) and comparable to those of SC pigs (Figure 2).

Blood immune measurements

White blood cell counts. Compared with E male pigs, SC male pigs had a lower number of total leukocytes at 5 months ($P = 0.03$) as well as a lower number of lymphocytes at 4 and 5 months ($P < 0.01$ and $P = 0.02$, respectively) but a similar number of granulocytes at the three ages (Table 1). When the number of cells differed significantly




Figure 2 Levels of testosterone in entire (E, $n = 18$), immunocastrated (IC, $n = 16$ to 18) and surgically castrated (SC, $n = 14$) pigs from experiment A. ^{a,b}Within each age, means with different letters differ with P -values < 0.001 .

between the SC and E male pigs, it was intermediate in the IC male pigs and never differed significantly from one group or the other.

Table 1 Numbers of total leukocytes, lymphocytes and granulocytes in entire (E), immunocastrated (IC) and surgically castrated (SC) pigs

Age	Treatment groups			s.e.	P-values
	E	IC	SC		
Leukocytes (1000 cells/mm ³)					
3 months	33.8	30.3	31.5	1.4	ns
4 months	29.2	28.8	27.0	1.9	ns
5 months	26.0 ^b	24.1 ^{ab}	22.5 ^a	1.2	0.03
Lymphocytes (1000 cells/mm ³)					
3 months	16.5	14.5	14.4	0.8	0.07
4 months	16.1 ^b	15.7 ^{ab}	13.6 ^a	0.7	< 0.01
5 months	16.6 ^b	15.3 ^{ab}	13.7 ^a	0.9	0.02
Granulocytes (1000 cells/mm ³)					
3 months	16.5	15.0	16.3	1.0	ns
4 months	12.4	12.4	12.7	1.4	ns
5 months	8.9	8.2	8.3	0.5	ns

Data include the results from experiment A at 3 months of age (E: *n* = 18, IC: *n* = 16 to 18, SC: *n* = 14) and from experiment A + B at 4 and 5 months of age (E: *n* = 28, IC: *n* = 16 to 18, SC: *n* = 24).

ns: *P*-value of the treatment effect is > 0.10.

^{a,b}Within a row, means with different letters differ at *P* < 0.05.

PBMC subsets and proliferation. There was no effect of treatment on the percentages of CD3⁺, CD3⁺CD4⁺ and CD3⁺CD4⁺CD8⁺ cells (Table 2). E pigs had a percentage of CD3⁺CD4⁺CD8⁺ cells that was lower than in SC pigs at 4 months of age (*P* < 0.01), and lower than in IC pigs at 5 months of age (*P* = 0.03). At 4 and 5 months, E pigs had a higher percentage of TCRγδ⁺ T lymphocytes than the SC male pigs (*P* = 0.02 and 0.03, respectively), whereas the IC pigs had an intermediate percentage not differing from that observed in the E and SC pigs (*P* > 0.1).

At 5 months, the E male pigs tended to have a higher basal proliferation of T cells than the SC and IC male pigs (*P* = 0.05). T-cell proliferation in response to ConA was not influenced by treatment.

Plasma immunoglobulins and haptoglobin

Treatments did not influence anti-influenza Ig or haptoglobin levels (Table 3). At 4 months of age, gonadal status tended to influence total IgG level (*P* = 0.06), but the effect was no longer present at 5 months (*P* > 0.1).

Clinical measurements at slaughter

Relative weight of the spleen tended to be influenced by treatment (*P* = 0.06) and that of the liver was higher in IC pigs than in E and SC pigs (*P* < 0.001, Table 4). Lung exams revealed that E pigs had higher scores of pneumonia than SC pigs, but there was no significant difference for stomach and nose lesions.

Discussion

This experiment showed that the consequences of surgical castration and immunocastration on peripheral immune cells were moderate. Long-term suppression of testicular

Table 2 T-lymphocyte subsets in entire (E), immunocastrated (IC) and surgically castrated (SC) pigs

Age	Treatment groups			s.e.	P-values
	E	IC	SC		
% CD3 ⁺ among lymphoid cells					
3 months	80	83	81	2	NS
4 months	82	81	82	1	NS
5 months	86	84	85	1	NS
% CD3 ⁺ CD4 ⁺ CD8 ⁻ among CD3 ⁺ cells					
3 months	19	20	21	2	NS
4 months	17	16	19	2	NS
5 months	17	19	18	1	NS
% CD3 ⁺ CD4 ⁻ CD8 ⁺ among CD3 ⁺ cells					
3 months	24	23	21	2	NS
4 months	22	24	22	1	NS
5 months	18	19	18	1	NS
% CD3 ⁺ CD4 ⁺ CD8 ⁺ among CD3 ⁺ cells					
3 months	6.0	6.9	7.6	1.1	NS
4 months	5.7 ^a	7.1 ^{ab}	8.8 ^b	0.8	< 0.01
5 months	4.2 ^a	6.4 ^b	4.8 ^{ab}	0.6	0.03
% CD3 ⁺ TCRγδ ⁺ among CD3 ⁺ cells					
3 months	48	45	42	3	NS
4 months	46 ^b	45 ^{ab}	39 ^a	2	0.02
5 months	60 ^a	53 ^{ab}	53 ^b	2	0.03

Data include the results from experiment A at 3 months of age (E: *n* = 18, IC: *n* = 16 to 18, SC: *n* = 14) and from experiment A + B at 4 and 5 months of age (E: *n* = 28, IC: *n* = 16 to 18, SC: *n* = 24).

ns: *P* value of the treatment effect is > 0.10.

^{a,b}Within a row, means with different letters differ at *P* < 0.05.

hormones by surgical castration seemed to have a greater impact than late immunocastration.

Effect of neonatal deprivation of testicular hormones by surgical castration

In our study, SC pigs presented lower numbers of lymphoid cells from 3 to 5 months relatively to entire pigs. Our finding is in agreement with previous studies where castrated male pigs at slaughter age had fewer peripheral lymphoid cells compared with entire male pigs (Odink *et al.*, 1990; Merlot *et al.*, 2013). The blood lymphoid fraction is composed of T, B and NK cells. As our results showed that surgical castration did not decrease the percentage of T (CD3⁺) lymphocytes among lymphoid cells, we can hypothesize that the haematopoiesis of T cells, on the one hand, and B plus NK cells on the other was reduced in a similar proportion in SC pigs.

These differences between E and SC pigs in blood cell numbers could result from a direct effect of neonatal and prepubertal androgens on the haematopoiesis of the lymphoid lineage. This would be in contradiction with studies on rodents, showing that both pre- and post-pubertal surgical castration increase the weight of the thymus and spleen and the peripheral pool of both B and T cells during adulthood (Viselli *et al.*, 1995; Ellis *et al.*, 2001; Radojevic *et al.*, 2007). The divergence between studies on pigs and rodents might result from the fact that the influence of

androgens on immune cells depends on the age of the animals at the observational time. They might stimulate the development of the immune system during the neonatal period and infancy, and have an opposite action after puberty. For example, Shioya *et al.* (2000) observed that in rats the stimulatory influence of neonatal castration on thymus weight does not appear before 31 days of age, that is, after the theoretical age of puberty. Overall, in the rodent studies (Viselli *et al.*, 1995; Ellis *et al.*, 2001;

Radojevic *et al.*, 2007), the measurements were carried out in sexually mature animals, whereas observations in pigs were carried out during pubertal development. However, another hypothesis to explain the higher number of circulating lymphoid cells in E pigs is that entire pigs are more aggressive than castrates, leading to more injuries when animals are group-housed. These lesions could favor skin microbial infections and thus stimulate the immune system, resulting in an increased number of blood leukocytes. This hypothesis would be in agreement with the tendency for a higher spontaneous proliferation of blood lymphocytes compared with SC pigs, which might also suggest a higher immune stimulation in E than in SC pigs. However, bacterial challenges are usually also associated with increased numbers of neutrophils (Odink *et al.*, 1990), which was not the case in the present experiment. Furthermore, although aggressiveness and skin lesions were not measured in this experiment, our group has recently observed in another study that there was nearly no difference in the frequency of agonistic interactions and the number of skin lesions between groups of E and SC pigs at 4 and 5 months of age, when the pigs are kept in stable social groups (Tallet *et al.*, 2011 and 2013), which was the case in the present study. For these two reasons, we defend the hypothesis of a direct effect of androgens on immune cell production.

Surgical castration not only decreased the number of total blood lymphocytes but also modified the frequency of several subsets of T cells. It increased the frequency of the CD3⁺CD4⁺CD8⁺ lymphocyte subset at 4 months. In pigs, these cells are known to be a subset of memory T cells (Pescovitz *et al.*, 1994; Zuckermann and Husmann, 1996). The increase in these memory cells unlikely results from a greater immune stimulation of the immune system of surgically castrated male pigs (see the discussion above). Furthermore, this result must be considered cautiously because it could be observed only transiently at 4 but not 3 and 5 months of age.

The frequency of $\gamma\delta$ T cells was also influenced by treatment. These cells were less frequent in SC compared

Table 3 Blood titres of anti-influenza Ig, concentration of total IgG and haptoglobin and proliferation of peripheral blood mononuclear cells of entire (E, n = 18), immunocastrated (IC, n = 16 to 18) and surgically castrated (SC, n = 14) pigs from experiment A

	Age	Treatment groups			s.e.	P-values
		E	IC	SC		
Total IgG (mg/ml)						
	3 months	nd	nd	nd		
	4 months	15.0	17.0	19.7	1.5	0.06
	5 months	12.5	12.1	14.4	0.9	ns
Anti-influenza Ig titres (arbitrary unit)						
	3 months	nd	nd	nd		
	4 months	0.14	0.13	0.14	0.03	ns
	5 months	0.89	0.84	0.83	0.06	ns
Haptoglobin (mg/ml)						
	3 months	1.26	2.11	1.71	0.30	ns
	4 months	0.68	1.04	0.66	0.27	ns
	5 months	0.87	1.12	0.89	0.27	ns
Proliferation without ConA (cpm)						
	3 months	527	913	254	273	ns
	4 months	797	541	675	309	ns
	5 months	1601	750	803	282	0.05
Proliferation in response to ConA (proliferation index)						
	3 months	172	131	139	33	ns
	4 months	125	125	143	28	ns
	5 months	72	76	57	21	ns

nd = not determined.

ns: P-value of the treatment effect is > 0.10.

Table 4 Body and organ weights and health measures in entire (E, n = 18), immunocastrated (IM, n = 16 to 18) and surgically castrated (SC, n = 14) pigs from experiment A

	Age	Treatment groups			s.e.	P-values
		E	IC	SC		
BW (kg)	3 months	48	45	45	2	ns
	4 months	73	71	69	2	ns
	5 months	112	116	107	3	0.08
Spleen/BW ratio (g/kg)	5 months	1.49	1.31	1.42	0.05	0.06
Liver/BW ratio (g/kg)	5 months	17.9 ^a	19.9 ^b	17.7 ^b	0.4	< 0.01
Ulcer score (0 to 7)	5 months	1.8	1.9	2.2	0.3	ns
Pneumonia score (0 to 28)	5 months	5.6 ^b	4.6 ^{ab}	2.6 ^a	0.9	0.04
Rhinitis score (0 to 18)	5 months	1.1	2.3	2.2	0.6	ns

For stomach ulcer, pneumonia and rhinitis, intact organs received a score of zero.

ns: P-value of the treatment effect is > 0.10.

^{a,b}Means with different letters differ at P < 0.05.

with E male pigs at 4 and 5 months of age. In humans, testosterone during adulthood is suspected of inhibiting the production of a subset of $\gamma\delta$ T cells, the V γ 9/V δ 2 T cells (Caccamo *et al.*, 2006). However, our results suggest that androgens during the neonatal periods and/or infancy might favor the production of $\gamma\delta$ T cells. To the best of our knowledge, our study is the first to provide results concerning the effect of sex steroids on the production of $\gamma\delta$ T cells in sexually immature mammals.

The influence of neonatal surgical castration that we observed throughout the present study could be attributed, at least partly, to a long-term effect of the suppression of neonatal androgens (lack of imprinting effect). Indeed, plasma androgens are elevated during the first month of age in pigs (Colenbrander *et al.*, 1978). The surgical castration that was carried out at 5 to 6 days of age shortened this neonatal peak, and may have altered the process of masculinization of the immune system, which has been shown in rodents to be programmed by the neonatal surge of androgens (Leposavic *et al.*, 2009). It may also be hypothesized that the influence of surgical castration could be attributed to the suppression of androgens during pubertal development. Indeed, measurements of plasma testosterone from the pigs of experiment A confirmed that concentrations at 3 months of age were much higher in entire than in surgically castrated male pigs and that they increased between 3 and 5 months of age.

From a functional point of view, minor alterations were observed in SC pigs. Basal proliferation of T cells tended to be higher in entire than in castrated pigs at 5 months of age; however, the proliferation index in response to a polyclonal stimulation with Con A was equivalent. The possible effect of neonatal castration on the production of Ig was not clear as total IgG concentrations tended to increase at 4 but not at 5 months, and the titre of specific anti-influenza antibodies measured at 5 months of age did not differ between SC and E pigs. Nevertheless, increases in total IgG levels were observed in castrated pigs at 5 months of age in a previous study (Merlot *et al.*, 2013). The increase in IgG concentration might result from the absence of testosterone that favors the production of antibodies by B cells (Kanda *et al.*, 1996). Moreover, in adult rodents, androgen deprivation induces an increase in the pool of B cells (Viselli *et al.*, 1995).

Effect of late deprivation of testicular hormones by immunocastration

We expected different immune effects of immunocastration and surgical castration for at least two reasons. First, deprivation of the testicular hormones did not occur at the same period of sexual development, being effective only after the second injection, that is, after 4 months of age in immunocastrated male pigs. Second, the two methods are very different from the endocrine point of view. Surgical castration suppresses only testicular hormones, whereas immunization against GnRH blocks the action of GnRH and suppresses the secretion of FSH, LH and gonadal steroids (Claus *et al.*, 2007). GnRH itself is known to have an

immunomodulatory role. Indeed, GnRH is secreted in the thymus (Azad *et al.*, 1998), and GnRH receptors are expressed in the thymus and spleen (Jacobson *et al.*, 1998) as well as in blood lymphocytes (Chen *et al.*, 2002). In particular, GnRH is involved in the homing of T cells to specific organs such as the spleen (Chen *et al.*, 2002). Therefore, GnRH may affect the immune system independently of sex steroids (Jacobson and Ansari, 2004).

In agreement with our hypothesis, the effect of immunocastration was different from that of surgical castration. Furthermore, this effect was marginal. It reduced basal proliferation of T cells at 5 months of age and enhanced the distribution of blood CD3⁺CD4⁺CD8⁺ memory T cells compared with entire male pigs at 5 months. The increase of CD3⁺CD4⁺CD8⁺ cells in IC pigs 1 month after the second vaccine injection might result from the process of vaccination itself, which is likely to stimulate the production of memory lymphocytes. It is possible that nearly no consequences of immunocastration were detected because the last measures were taken only 1 month after the second vaccine injection. Indeed, it seems that immune alterations after manipulation of the gonadotropic axis in pubertal animals develop in several weeks. For instance, Hirakata *et al.* (2010) measured the histological modifications of the thymus after suppression of the HPG axis in miniature pigs and showed that thymic rejuvenation appeared 2 months after the beginning of the treatment. Their experiment was conducted on sexually mature pigs (22 and 29 months) and does not prove that such an effect is present in peripubertal animals.

Conclusion

Our data demonstrate that the two methods of castration had different immune consequences. Neonatal surgical castration seemed to influence blood lymphocyte numbers and the frequencies of the different T cell subpopulations until slaughter age, suggesting that immune development and particularly thymic function were altered. Immunocastration at puberty had virtually no effect on the immune system. Overall, no clear functional consequences of either method on commercial pig immune abilities were demonstrated, but more investigations are required to ascertain such a conclusion.

Acknowledgments

The authors thank Françoise Thomas, Raphael Comte, Michel Lefebvre, Rachel Verdet and Sarah Poirot from the PEGASE research unit for their help in the sample analysis, and Josselin Delamarre and Patrice Roger from the experimental unit for their technical assistance. This study was funded by INRA and the French National Research Agency (L'Agence Nationale de la Recherche, ANP-09-BLAN-0083 ANDROPIG). Proliferation plates were kindly processed by Xénoblis (Saint-Grégoire, France).

References

Allrich RD, Christenson RK, Ford JJ and Zimmerman DR 1982. Pubertal development of the boar -testosterone, estradiol-17-beta, cortisol and LH

- concentrations before and after castration at various ages. *Journal of Animal Science* 55, 1139–1146.
- Azad N, LaPaglia N, Agrawal L, Steiner J, Uddin S, Williams DW, Lawrence AM and Emanuele NV 1998. The role of gonadectomy and testosterone replacement on thymic luteinizing hormone-releasing hormone production. *Journal of Endocrinology* 158, 229–235.
- Bouman A, Heineman MJ and Faas MM 2005. Sex hormones and the immune response in humans. *Human Reproduction Update* 11, 411–423.
- Brunius C, Zamaratskaia G, Andersson K, Chen G, Norrby M, Madej A and Lundstrom K 2011. Early immunocastration of male pigs with Improvac (R) – effect on boar taint, hormones and reproductive organs. *Vaccine* 29, 9514–9520.
- Caccamo N, Dieli F, Wesch D, Jomaa H and Eberl M 2006. Sex-specific phenotypical and functional differences in peripheral human V gamma 9/V delta 2 T cells. *Journal of Leukocyte Biology* 79, 663–666.
- Chen A, Ganor Y, Rahimpour S, Ben-Aroya N, Koch Y and Levite M 2002. The neuropeptides GnRH-II and GnRH-I are produced by human T cells and trigger laminin receptor gene expression, adhesion, chemotaxis and homing to specific organs. *Nature Medicine* 8, 1421–1426.
- Claus R, Lacorn M, Danowski K, Pearce MC and Bauer A 2007. Short-term endocrine and metabolic reactions before and after second immunization against GnRH in boars. *Vaccine* 25, 4689–4696.
- Colenbrander B, Dejong FH and Wensing CJG 1978. Changes in serum testosterone concentrations in male pig during development. *Journal of Reproduction and Fertility* 53, 377–380.
- Colenbrander B, Meijer JC, Macdonald AA, Vandewiel DFM, Engel B and de Jong FH 1987. Feedback-regulation of gonadotropic hormone secretion in neonatal pigs. *Biology of Reproduction* 36, 871–877.
- Couret D, Jamin A, Kuntz-Simon G, Prunier A and Merlot E 2009. Maternal stress during late gestation has moderate but long-lasting effects on the immune system of the piglets. *Veterinary Immunology Immunopathology* 131, 17–24.
- Ellis TM, Moser MT, Le PT, Flanigan RC and Kwon ED 2001. Alterations in peripheral B cells and B cell progenitors following androgen ablation in mice. *Internal Immunology* 13, 553–558.
- Gould KG, Akinbami MA and Mann DR 1998. Effect of neonatal treatment with a gonadotropin releasing hormone antagonist on developmental changes in circulating lymphocyte subsets: a longitudinal study in male rhesus monkeys. *Developmental and Comparative Immunology* 22, 457–467.
- Henry Y, Dauloudet C, Kerisit R, Lefeuvre P, Gaye A, Denechaud MF and Bourdon D 1970. Effets nutritionnels de l'incorporation de cellulose purifiée dans le régime du porc en croissance-finition. III-Incidence sur le développement des ulcères gastro-oesophagiens. *Annales de Zootechnie* 19, 117–141.
- Hirakata A, Okumi M, Griesemer AD, Shimizu A, Nobori S, Tena A, Moran S, Arn S, Boyd RL, Sachs DH and Yamada K 2010. Reversal of age-related thymic involution by an LHRH agonist in miniature swine. *Transplant Immunology* 24, 76–81.
- Jacobson JD and Ansari MA 2004. Immunomodulatory actions of gonadal steroids may be mediated by gonadotropin-releasing hormone. *Endocrinology* 145, 330–336.
- Jacobson JD, Crofford LJ, Sun LH and Wilder RL 1998. Cyclical expression of GnRH and GnRH receptor mRNA in lymphoid organs. *Neuroendocrinology* 67, 117–125.
- Kanda N, Tsuchida T and Tamaki K 1996. Testosterone inhibits immunoglobulin production by human peripheral blood mononuclear cells. *Clinical and Experimental Immunology* 106, 410–415.
- Leposavic G, Perisic M, Kosec D, Arsenovic-Ranin N, Radojevic K, Stojic-Vukanic Z and Pilipovic I 2009. Neonatal testosterone imprinting affects thymus development and leads to phenotypic rejuvenation and masculinization of the peripheral blood T-cell compartment in adult female rats. *Brain, Behavior, and Immunity* 23, 294–304.
- Madec F and Kobish M 1982. A survey of pulmonary-lesions in bacon pigs (observations made at the slaughterhouse). *Annales de Zootechnie* 31, 341.
- Merlot E, Thomas F and Prunier A 2013. Comparison of immune and health markers in intact and neonatally castrated male pigs. *Veterinary Record* 173, 317.
- Odink J, Smeets JFM, Visser IJR, Sandman H and Snijders JMA 1990. Hematological and clinicochemical profiles of healthy swine and swine with inflammatory processes. *Journal of Animal Science* 68, 163–170.
- Pescovitz MD, Sakopoulos AG, Gaddy JA, Husmann RJ and Zuckermann FA 1994. Porcine peripheral-blood CD4(+) CD8(+) dual expressing T-cells. *Veterinary Immunology Immunopathology* 43, 53–62.
- Radojevic K, Arsenovic-Ranin N, Kosec D, Pesic V, Pilipovic I, Perisic M, Plecas-Solarovic B and Leposavic G 2007. Neonatal castration affects intrathymic kinetics of T-cell differentiation and the spleen T-cell level. *Journal of Endocrinology* 192, 669–682.
- Shioya N, Inomata T, Ninomiya W and Nakamura T 2000. Influence of castration on development of the thymus in neonatal male rats. *Experimental Animals* 49, 57–60.
- Sutherland JS, Goldberg GL, Hammett MV, Uldrich AP, Berzins SP, Heng TS, Blazar BR, Millar JL, Malin MA, Chidgey AP and Boyd RL 2005. Activation of thymic regeneration in mice and humans following androgen blockade. *Journal of Immunology* 175, 2741–2753.
- Tallet C, Brillouet A, Paulmier V, Meunier-Salaun M-C, Bonneau M and Prunier A 2011. Conséquences de l'enrichissement du milieu sur le développement sexuel et l'agressivité des porcs mâles entiers et castrés. *Journées de la Recherche Porcine* 43, 187–188.
- Tallet C, Brillouet A, Meunier-Salaun MC, Paulmier V, Guerin C and Prunier A 2013. Effects of neonatal castration on social behaviour, human-animal relationship and feeding activity in finishing pigs reared in a conventional or an enriched housing. *Applied Animal Behaviour Science* 145, 70–83.
- Viselli SM, Stanziale S, Shults K, Kovacs WJ and Olsen NJ 1995. Castration alters peripheral immune function in normal-male mice. *Immunology* 84, 337–342.
- Zuckermann FA and Husmann RJ 1996. Functional and phenotypic analysis of porcine peripheral blood CD4/CD8 double-positive T cells. *Immunology* 87, 500–512.