



Inter individual variations of the fish skin microbiota: Host genetics basis of mutualism?

Sébastien Boutin, Christopher Sauvage, Louis Bernatchez, Céline Audet,
Nicolas Derôme

► To cite this version:

Sébastien Boutin, Christopher Sauvage, Louis Bernatchez, Céline Audet, Nicolas Derôme. Inter individual variations of the fish skin microbiota: Host genetics basis of mutualism?. PLoS ONE, 2014, 9 (7), 17 p. 10.1371/journal.pone.0102649 . hal-02634270

HAL Id: hal-02634270

<https://hal.inrae.fr/hal-02634270>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Inter Individual Variations of the Fish Skin Microbiota: Host Genetics Basis of Mutualism?

Sébastien Boutin¹, Christopher Sauvage^{1,2}, Louis Bernatchez¹, Céline Audet³, Nicolas Derome^{1*}

1 Institut de Biologie Intégrative et des Systèmes (IBIS), Département de Biologie, Université Laval, Québec, Québec, Canada, **2** INRA, UR1052, Génétique et Amélioration des Fruits et Légumes (GAFL), Domaine St Maurice - Allée des Chênes, Montfavet, France, **3** Institut des sciences de la mer de Rimouski (ISMER), Université du Québec à Rimouski (UQAR), Rimouski, Québec, Canada

Abstract

The commensal microbiota of fish skin is suspected to provide a protection against opportunist infections. The skin of fish harbors a complex and diverse microbiota that closely interacts with the surrounding water microbial communities. Up to now there is no clear evidence as to whether the host regulates the recruitment of environmental bacteria to build a specific skin microbiota. To address this question, we detected Quantitative Trait Loci (QTL) associated with the abundance of specific skin microbiota bacterial strains in brook charr (*Salvelinus fontinalis*), combining 16S RNA tagged-amplicon 454 pyrosequencing with genetic linkage analysis. Skin microbiota analysis revealed high inter-individual variation among 86 F2 fish progeny based upon the relative abundance of bacterial operational taxonomic units (OTUs). Out of those OTUs, the pathogenic strain *Flavobacterium psychrophilum* and the non-pathogenic strain *Methylobacterium rhodesianum* explained the majority of inter-individual distances. Furthermore, a strong negative correlation was found between *Flavobacterium* and *Methylobacterium*, suggesting a mutually competitive relationship. Finally, after considering a total of 266 markers, genetic linkage analysis highlighted three major QTL associated with the abundance of *Lysobacter*, *Rheinheimera* and *Methylobacterium*. All these three genera are known for their beneficial antibacterial activity. Overall, our results provide evidence that host genotype may regulate the abundance of specific genera among their surface microbiota. They also indicate that *Lysobacter*, *Rheinheimera* and *Methylobacterium* are potentially important genera in providing protection against pathogens.

Citation: Boutin S, Sauvage C, Bernatchez L, Audet C, Derome N (2014) Inter Individual Variations of the Fish Skin Microbiota: Host Genetics Basis of Mutualism? PLoS ONE 9(7): e102649. doi:10.1371/journal.pone.0102649

Editor: Stefano Mariani, School of Environment & Life Sciences, University of Salford, United Kingdom

Received: December 13, 2013; **Accepted:** June 20, 2014; **Published:** July 28, 2014

Copyright: © 2014 Boutin et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was financially supported by a Partnership grant (Strategic program) of the "Conseil de recherches en sciences naturelles et en génie" (CRSNG) to Louis Bernatchez, Nicolas Derome, and Céline Audet and the "Collaborative Research And Training Experience" program (CREATE). This is a contribution to the research program of Resources Aquatiques Québec (RAQ). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: nicolas.derome@bio.ulaval.ca

Introduction

The study of the beneficial effects of bacteria and their influence on host health is a growing field. Namely, research that has explored host microbiota variability in space and time suggests the presence of a host genetic component into the development of mutualism communities [1,2]. Similar conclusions were found in several studies on twins and on their core gut microbiota [3,4,5,6]. Furthermore, microbiome homeostasis seems to be the key to resistance against some diseases previously considered exclusively influenced by genetic factors [7]. To determine precisely which genes are involved in the recruitment of specific bacterial strains, some studies looked at gene expression in presence of symbiotic bacteria. Recent studies specifically targeted genes already known as being involved in innate or adaptive immunity (for example IgA) [8,9,10].

In fish, the influence of host genetic background on commensal bacterial community structure is poorly known. However, advances in the field of probiotic development indicate that endogenous bacteria are able to outcompete pathogens [11,12]. Results obtained in aquaculture settings are consistent with previous work showing that non-pathogenic bacteria associated

with animal mucosa can contribute to the host health by providing protection against pathogen infections [13,14,15,16]. The genetic basis of the host immune response, especially at the major histocompatibility complex (MHC), toll-like receptor (TLR) and immunoglobulin loci, has been well studied [17,18,19,20]. However, aside from the immune response, the influence of the host on the development of bacterial symbiosis remains poorly understood.

Here, the main objective of the present work was to investigate the potential link between host genotype and skin microbiota composition. As a host model, we focused on brook charr (*Salvelinus fontinalis*), a salmonid that harbors a complex and dynamic skin microbiota [21,22]. The first specific objective sought to document the genetic variation present in the host population that might underpin the inter-individual variations in the skin microbiota. The second specific objective was to characterize the relationship (cooperation/competition) existing among the different bacterial Operational taxonomic units (OTU) within the skin mucus, which influences the overall structure of the skin microbial community. The third objective was to identify host genetic regions associated to the abundance of specific skin bacterial OTUs (i. e. quantitative trait loci (QTL) associated with

Table 1. Descriptive statistics of sequencing to determine the microbiota for each individual.

Sample ID	number of reads	coverage	npshannon index	Number of OTU total	Number of OTU >10 reads
92	229	0.955975	1.690045	76	19
93	1584	0.99131	0.436879	74	28
94	669	0.977401	1.507996	242	20
95	738	0.986166	1.416086	226	27
96	503	0.953079	1.395581	149	29
97	385	0.946203	1.774329	86	33
98	955	0.98389	0.663968	79	30
99	770	0.978852	1.306709	114	29
100	1018	0.985946	0.658787	107	30
101	1482	0.994792	0.842304	188	21
102	1667	0.992283	0.691189	96	31
103	711	0.99179	0.946539	97	23
104	1072	0.987599	0.719994	159	25
105	1662	0.991645	0.371924	73	26
106	1304	0.995734	0.658557	93	18
107	1617	0.99416	0.290016	49	21
108	1237	0.990451	0.634742	73	30
109	1063	0.987778	0.679124	137	26
110	752	0.97281	0.656197	105	29
111	731	0.986217	0.63276	79	22
112	628	0.975779	0.630137	63	23
113	892	0.983549	0.446389	49	24
114	1333	0.993127	0.713271	127	22
115	1109	0.983313	1.020009	266	35
116	532	0.967611	2.709733	269	32
117	649	0.963145	2.108945	237	41
118	404	0.923977	2.645631	222	29
119	969	0.97625	1.019139	177	40
120	581	0.967647	1.975631	205	26
121	468	0.948413	2.692236	202	34
122	744	0.980551	0.876389	117	27
123	953	0.983204	0.832745	133	32
124	530	0.967846	2.29117	186	30
125	797	0.977901	0.703269	79	30
126	411	0.97561	0.904985	116	18
127	242	0.76	3.452235	182	32
128	1230	0.989637	0.504536	69	26
129	831	0.978723	1.537117	115	41
130	1449	0.983421	0.548428	123	38
131	499	0.969697	2.118408	144	29
132	431	0.945813	2.133064	181	25
133	494	0.96114	1.408929	117	35
134	746	0.980254	0.508333	38	22
135	662	0.977887	2.41624	180	36
136	754	0.987755	0.956343	118	18
137	2008	0.993154	0.557917	74	30
138	714	0.976619	1.632525	101	36
139	798	0.967632	1.534968	129	35
140	469	0.917197	3.267019	261	35

Table 1. Cont.

Sample ID	number of reads	coverage	npshannon index	Number of OTU total	Number of OTU >10 reads
141	224	0.980892	0.662835	33	6
142	517	0.957377	2.388226	180	36
143	483	0.947195	2.446711	157	36
144	414	0.958848	3.124298	188	38
145	930	0.975443	1.391363	168	43
146	769	0.979661	2.024708	140	28
147	1219	0.995122	1.641489	118	17
148	1254	0.98244	1.127563	148	40
149	1074	0.972678	2.309049	318	55
150	1108	0.980583	2.793443	321	47
151	3082	0.998643	0.332436	58	16
152	1429	0.996561	1.820656	203	25
153	1955	0.995286	0.815557	167	27
154	1449	0.983366	1.74373	354	54
155	1074	0.973611	1.716763	321	41
156	1016	0.989316	0.588719	62	21
157	1483	0.984836	1.374063	216	46
158	2004	0.998944	0.791157	44	10
159	2043	0.997359	0.812388	92	18
160	2468	0.995741	0.305525	70	23
161	1215	0.979566	1.928985	346	48
162	1904	0.995286	0.821609	164	37
163	1046	0.985799	1.569251	188	40
164	1145	0.993814	0.744587	98	19
165	995	0.981481	1.421104	243	37
166	725	0.97921	2.459141	211	36
167	727	0.988743	2.142562	131	24
168	841	0.98908	2.297402	151	31
169	552	0.968116	2.76388	178	35
170	1204	0.99177	1.912832	162	32
172	2010	0.993946	0.610072	155	36
173	1454	0.987296	1.893132	316	51
174	1056	0.979651	2.949199	335	53
175	824	0.95098	2.845177	368	46
176	851	0.96729	2.039242	202	47
177	778	0.985507	2.23077	221	25
178	2142	0.998093	0.359619	52	13

The coverage was estimated by a Good's estimator index. npshannon: non parametric index of Shannon. OTU classification was done with a threshold of 97% identity. doi:10.1371/journal.pone.0102649.t001

the abundance of each constituent bacterial genus). Accordingly, we analyzed the taxonomic structure of skin bacterial community using 16S tagged-amplicon pyrosequencing in 86 F2 fish progenies obtained in the previous work of Sauvage *et al.* (2012). Inter-individual variation in abundance of each bacterial strain detected in host microbiota was afterwards projected on the results of the genetic linkage map in order to identify quantitative trait loci (QTL) of specific skin bacterial OTUs.

Materials and Methods

Ethics statement

All fish were reared and the experiment conducted strictly following guidelines required by the "Comité de Protection des Animaux de l'Université Laval (CPAUL, <http://www.vrr.ulaval.ca/deontologie/cpa/index.html?accueil>). The CPAUL reviewed and approved all experimental procedures used in this study.




Figure 1. Taxonomic structure of the bacterial community of fish skin mucus at three different taxonomic levels: Phylum, Class and OTU. a) Proteobacteria; b) Bacteroidetes, c) Alphaproteobacteria, d) Flavobacteria, e) Gammaproteobacteria, f) OTU 50 (*Methylobacterium rhodesianum*), g) OTU 36 (*Flavobacterium psychrophilum*).

doi:10.1371/journal.pone.0102649.g001

Fish sampling

The study population was generated using two different populations of brook charr. The first one (D) is a domestic population used in aquaculture for more than 100 years

(Pisciculture de la Jacques Cartier - Cap-Santé, Québec, Canada). The other population (L), is an anadromous population from the Laval River near Forestville (North of the St. Lawrence River, Québec, Canada). Breeders from the L population were kept in captivity at the Station aquicole de l'ISMER (Québec, Canada, 48°31'N, 68°28'W) under natural photoperiod and temperature. Crosses between 10 sires of each population (L and D; F₀ generation) with 10 dams (L and D) were performed to generate 10 full-sib outbred hybrid families (L×D - F₁ generation). Subsequently, six F₁ individuals were bi-parentally crossed to obtain three F₂ families. The F₂ family selected for the present study demonstrated the lowest post-hatch mortality rate (<2%) and the greatest abundance. Fish were raised in the same tank and had fasted for 12 hours before sampling. Fish were measured (mean = 28.8±1.77 cm) and weighed (mean = 276.7±59.48 g) to calculate Fulton index [23].

Mucus was sampled using a sterile swab on the same area for all the fish [24]. We choose to sample the skin between the adipose fin and the caudal fin because this area was undisturbed by fish handling. Samples were stored in a sterile micro-centrifuge tube containing lysis buffer (Tris 50 mM, EDTA 40 mM, sucrose 0.75 g) and immediately stored in -80°C until DNA extraction.

DNA extraction

DNA was extracted using a modified protocol of salt-extraction from Aljanabi & Martinez (1997). During the first lysis step, 22.6 µL of lysozyme (40 mg/mL) was added to the sample and incubated 45 minutes at 37°C. After this step, 22.6 µL of proteinase K (20 mg/µL) and 90 µL of SDS (10%) were added to the lysate and incubated at 55°C over night with agitation. The aqueous phase was transferred into a clean Eppendorf tube containing 600 µL of NaCl 6M, mixed and centrifuged 20 min at 14,800 rpm. The supernatant was transferred again into a clean Eppendorf tube containing 1 volume of cold isopropanol, mixed and stored 30 minutes at -20°C. The mixture was centrifuged 20 minutes at 14,800 rpm and the supernatant was thrown away. The pellet was washed with cold ethanol 70%, air-dried and finally resuspended in 25 µL of sterile MilliQ H₂O. Subsequently, DNA integrity and quantity were controlled using a Nanodrop instrument (ND-1000, Nanodrop).

Microbial 16S pyrosequencing

Each DNA sample, the 16S gene was PCR amplified using Takara Ex taq premix (Fisher). All PCR reactions were performed in a reaction volume of 50 µL containing 25 µL of premix Taq, 1 µM of each primer and sterile MilliQ H₂O to up to 50 µL. A general reverse primer (R519) combined with B primer (Roche) was used for amplifications in combination with one of 86 uniquely tagged forward primers (F63-targeted) combined with A primer (Roche) (for primer sequences see [25,26]). The mean length of the amplified fragment was 450 bp. This procedure facilitates the parallel sequencing of thousands of samples on the same run and to reassign each reads to their respective samples. PCR conditions were applied as follows: after a denaturing step of 30 s at 98°C, samples were processed through 30 cycles consisting of 10 sec at 98°C, 30 sec at 55°C, and 30 sec at 72°C. The final extension step was done at 72°C for 4 min 30 sec. Following the amplification step, samples were purified using AMPure Beads (Beckman Coulter Genomics). Samples were adjusted to 100 µL with EB (Qiagen), 63 µL of beads were added. Samples were mixed and incubated for 5 min at RT. Using a Magnetic Particle Concentrator (MPC), the beads were pelleted against the wall of the tube and supernatant was removed. The beads were washed twice with 500 µL of 70% ethanol and incubated for 30 sec each

Table 2. Taxonomic variation observed between male and female.

Condition	N° OTU	Best Hit on Blast	Identity	Relative abundance
More abundant in female	82	<i>Amaricoccus kaplicensis</i>	98%	0.00037647
	115	<i>Loktanella salsilacus</i>	99%	0.00015338
	6	<i>Methylovirgula ligni</i>	91%	0.00026492
	163	<i>Micrococcus antarcticus</i>	99%	0.00061351
	88	<i>Paracoccus yeei</i>	94%	0.00019521
	75	<i>Pseudochrobactrum kiredjianiae</i>	99%	0.00050196
	128	<i>Psychrobacter faecalis</i>	99%	0.00100392
	131	<i>Rheinheimera pacifica</i>	99%	0.00057168
	7, 11	<i>Singularimonas variicoloris</i>	89%	0.00037647
	104	<i>Sphingopyxis witflariensis</i>	99%	0.00015338
	113	<i>Stella humosa</i>	100%	0.00018126
	142	<i>Stenotrophomonas nitritireducens</i>	99%	0.00022309
More abundant in male	186	<i>Flavobacterium aquatile</i>	97%	0.00029281
	190	<i>Hyphomicrobium sulfonivorans</i>	95%	0.00039041
	4	<i>Methylosinus sporium</i>	92%	0.00027887
	176	<i>Polynucleobacter necessarius</i> subsp. <i>Asymbioticus</i>	99%	0.00015338
	129	<i>Pseudomonas hibiscicola</i>	99%	0.00075294
	12	<i>Pseudorhodofex soli</i>	85%	0.00018126
	140	<i>Psychrobacter alimentarius</i>	99%	0.00015338
	116	<i>Rhodobacter blasticus</i>	95%	0.00016732
	175	<i>Simplicispira limi</i>	99%	0.00034858

This table summarize the OTUs differentially abundant between skin mucus communities between male and female (Metastat using a FDR correction, p-value<0.01). doi:10.1371/journal.pone.0102649.t002

time. Supernatant was removed and beads were air dried for 5 min. Tubes were removed from the MPC and 24 µL of EB were added. Samples were vortexed in order to suspend the beads. Finally, using the MPC, the beads were pelleted against the wall once more and supernatant was transferred to a new clean tube. Samples were quantified with Nanodrop before the amplification step. Amplicons were then quantified with Quant-iT PicoGreen dsDNA Assay Kit and mixed equally before being sent to the Plateforme d'Analyses Biomoléculaires (Institut de Biologie Intégrative et des Systèmes, Université Laval) for pyrosequencing on a 454 GS-FLX DNA Sequencer with the Titanium Chemistry (Roche), according to manufacturer's procedure.

16S sequence analysis

All sequences are available on MG-RAST server (MG-RAST IDs: 4539915.3). The data were analyzed in two steps. First, CLC Genomics Workbench 3.1 (CLC Bio, Aarhus, Denmark CLC workbench BIO) was used to trim sequences for quality, recover and remove the primers' sequences and tags (minimum average quality score: 35 for a window of 50, number of differences to the primer sequence = 0, maximum number of differences to the barcode sequence = 0, number of ambiguous base calls = 0, maximum homopolymer length = 8). In a second step, pre-processing and analysis were performed using the MOTHUR software [27]. All datasets were checked for chimeras with the chimera slayer algorithm implemented in MOTHUR. Standardization of the different samples was done by using the zscore which calculates the normalized abundance as follows: normalized abundance = (abundance - mean) / standard deviation

[3,28,29]. An alternative method for normalization was also tested, which consisted of subsampling equal number of reads from each sample [30]. This method greatly reduced the numbers of sequences (224 sequences per samples), but gave the same results as zscore (Figure S2, Figure S3, Table S1). Therefore we preferred to keep zscore normalization to base our conclusions on a larger dataset. We used the Operational Taxonomic Unit-based method described by Costello *et al.* (2009) because it is not biased towards a predefined taxonomy. One index was retained to assess the quality of pyrosequencing: the sequence coverage index (Good's estimator). The sequence coverage index is a metric used to estimate the quality of the depth sequencing. All sequences were clustered into Operational Taxonomic Unit (OTU) using a 97% identity threshold and OTU were classified from phylum to genus using the program MOTHUR with the default settings. For all interesting OTUs (explaining the inter-individual variation or linked to a genetic region), we extracted all the sequences classified in that OTU and used the single best BLAST hit to identify the different species. Statistical differences in the abundance of each OTU were calculated with the software for metagenomic analysis (Metastats).

To visualize potential differences across host progenies in terms of mucus bacterial community structure, distances between those communities were computed using the Yue & Clayton measure of dissimilarity (Thetayc). This index developed by Yue and Clayton (2005) is a measure of dissimilarity between the structures of two communities, meaning that this calculator takes in account the abundances of each OTU. Then, Distances were represented using a dendrograms based on this index and statistical robustness of the dendrogram was determined by a Unifrac weighted test.




Figure 2. Dendrogram analysis based upon ThetaYC index of bacteria found on the skin of the 86 brook charr individuals.

Groups are defined with the Weighted Unifrac distance. The first and second groups are composed of closely related bacterial communities. The third group is an assemblage of dissimilar communities and is considered as an outgroup. The most differentiated groups are groups 1 and 3 (Unifrac Score: 0.710811, $p < 0.0010$) followed by the distance between groups 2 and 3 (Unifrac Score: 0.685361, $p < 0.0010$), and the distance between groups 1 and 2 (Unifrac Score: 0.401674, $p < 0.0010$). doi:10.1371/journal.pone.0102649.g002

This test allows determining whether any of the samples have a significantly different structure than the other samples. A random (Monte Carlo) permutation test was performed to test whether or not the distance between two communities was greater than expected by chance alone.

To visualize the distances between communities, we also applied a Principal Coordinate Analysis (PCoA) based on the Yue & Clayton measure of dissimilarity (Thetayc) using an eigenvector-based approach to represent multidimensional data in as few dimensions as possible. This method allows determining which OTUs are responsible for shifting the samples along the PCoA axes by measuring the correlation of the relative abundance of each OTU clustered in genera with the axes of the PCoA dataset. Statistical differences in the abundance of each OTU was calculated to highlight which genera were the causative agents of the differentiation between groups obtained in the dendrogram based upon the ThetaYC index [31]. All statistical analyses and graphics were carried out in the R environment (<http://www.r-project.org>).

QTL detection

QTL analysis was carried out using the [R] package J/ql [v. 1.3.1, August 2012, <http://churchill.jax.org/software/jql.shtml/>]. QTL were projected on the consensus brook charr linkage map (see Sauvage *et al.*, 2012) according to the following steps. (1) A single QTL analysis was performed using the Haley-Knott (HK) regression method (10,000 permutations) to reveal which LGs were carrying QTL. The most probable position of the QTL was defined at the position giving the largest logarithm of odds (LOD) score indicating the QTL was fixed. (2) A two QTL model based upon the Haley-Knott regression was used to refine the QTL detection across the genome with a resolution of 5 cM and eventually to detect two QTL on a single LG linked to a particular trait. (3) The best model fitting our data was used to compute the percent variance explained (PVE expressed in %) by the QTL. The chromosome-wide and the genome-wide thresholds were calculated for each LG using 10,000 permutations. The 1.5 LOD confidence intervals were determined for all analyses following the Bayesian method implemented in the “bayesint” function in R/ql. The bayesint function calculates an approximate interval (end points around the maximum LOD) for a given chromosome using the genome scan output.

Results

Mucus samples were obtained from 86 F2 fish progenies (44 males and 42 females) issued from the same family. A total of 87,940 high-quality, classifiable sequences were obtained with an average number of 1022 ± 540 per sample, which were subsequently classified in OTUs [1]. All of these sequences were successfully clustered into 9520 OTUs with 97% identity. A final trimming step was performed to focus on the most abundant OTU (which are represented by at least 10 sequences for the whole project) and result in a dataset containing 71,719 sequences (81% of the initial data set) clustered in 192 OTUs. The depth of

Bacterial diversity




Figure 3. Alpha diversity of each group. Each group was defined by the dendrogram analysis based upon ThetaYC index (see figure 2). Alpha-diversity was calculated by the non-parametric index of Shannon.
doi:10.1371/journal.pone.0102649.g003

sequencing and the coverage were estimated using the Good's estimator index; the coverage found was always higher than 90% except for one sample (#127), which exhibited 76% of coverage (Table 1, Figure S1). The number of OTUs per sample was not equally distributed and the mean number was 156.5 ± 81.7 (ranging from 33 to 368). The alpha-diversity was estimated by calculating the non-parametric index of Shannon (npShannon) because of the non-equally distributed abundance of each OTU in the samples. The npShannon index ranged from 0.29 to 3.45, with a mean index of 1.43 ± 0.83 .

Phylum, class, and OTU composition of all samples were represented in the Figure 1. Most of the OTUs belong to the *Proteobacteria* phylum (88.7%) and the *Bacteroidetes* phyla (9.7%).

At the class level, most of the OTUs were classified as *Alpha-proteobacteria* (78.9%), *Gamma-proteobacteria* (6.5%) and *Flavobacteria* (13%). At the OTU level, the most abundant sequence was identified via BLAST as *M. rhodesianum* (69.8%) followed by *F. psychrophilum* (8%) (Figure 1 and Table S1).

Male and female brook charr harbored slightly different microbial communities. The Unifrac weighted distance between samples from males and females was statistically significant indicating that structures of bacterial communities were differentiated according to the sex of the host (Unifrac Score: 0.241695; p-value < 0.001). However, there was not a sex difference in the two most abundant species, as this difference was based on 22 OTUs




Figure 4. Descriptive analysis of the Metastats. Numbers indicates the numbers of OTUs differentially present in the different groups of individuals. For details on the identities of the OTUs, please refer to the table 3; 4 and 5.
doi:10.1371/journal.pone.0102649.g004

classified on 20 genera and 21 species, all of them being less abundant than 0.1% (Table 2).

In terms of community structure, clustering based on ThetaYC index clearly defined two groups of samples, whilst a third group was less well defined and composed of highly variable communities (Figure 2). There was little variation among samples in the first and second group (group 1 and group 2). The third group (group 3) was an assemblage of dissimilar communities and is considered as an external group. The variation of the alpha-diversity visualized by the npShannon index (Figure 3) indicated that alpha-diversity increases from group 1 to group 3. Furthermore, the bacterial communities within the third group were more diverse than groups 1 and 2. The same clustering was found with the normalization by subsampling (Figure S2, Figure S3, table S1). The difference between each group was greater than the difference between males and females (Figure 2). The most differentiated groups were groups 1 and 3 (Unifrac Score: 0.710811, p-value < 0.001) followed by the distance between groups 2 and 3 (Unifrac Score: 0.685361, p-value < 0.001), and the distance between groups 1 and 2 (Unifrac Score: 0.401674, p-value < 0.001). This

was correlated with the numbers of OTU that were differentially abundant between those 3 groups (Figure 4). The divergence between groups 1 and 2 was based upon 54 OTU classified in 26 genera and 41 species (Table 3). The microbiota of the first group was dominated by *M. rhodesianum*, and host individuals from this group were tightly clustered. The second group was more diverse, dominated by *Flavobacterium* and had a lower abundance of *M. rhodesianum*. For groups 1 and 3, the difference was based upon 109 OTU classified in 45 genera and 70 species (Table 4). Finally, 26 OTU were differently abundant between group 2 and 3 and classified in 17 genera and 22 species (Table 5).

PCoA analysis displayed the same patterns of divergence between the three groups (Figure 5). Each group found in the Unifrac analysis was highlighted in the PCoA. To understand which OTUs were responsible for the differentiation of the cluster on the PCoA axes 1 and 2, individual correlation coefficients were calculated. Three OTUs were highly correlated with the two axes ($\sigma > 0.6$); OTU 36, 50 and 149. Those OTU belonged to two species: *Flavobacterium psychrophilum* (OTU 36) and *M. Rhodesianum* (OTU 50 and 149). The negative correlation

Table 3. Detailed information related to the OTU differentially abundant between individuals belonging to group 1 and group 2.

N° OTU	mean abundance in group 1	mean abundance in group 2	Best hit on Blast
36	0.00484	0.037926	<i>Flavobacterium psychrophilum</i>
122	0.000778	0.010066	<i>Pseudomonas cedrina subsp. Fulgida</i>
37	0.000471	0.006593	<i>Flavobacterium terrigena</i>
68	0.0014	0.006629	<i>Methylocella palustris</i>
128	0	0.004734	<i>Psychrobacter faecalis</i>
69	0.000531	0.004942	<i>Pseudorhodobacter ferrugineus</i>
190	0	0.004394	<i>Hyphomicrobium sulfonivorans</i>
139	0.000153	0.003833	<i>Lysobacter capsici</i>
70	0.000831	0.0043	<i>Methylobacterium salsuginis</i>
29	0.000028	0.003344	<i>Methylobacterium adhaesivum</i>
67	0.000518	0.003772	<i>Rhodobacter blasticus</i>
127	0.000218	0.003382	<i>Acinetobacter haemolyticus</i>
21	0.000117	0.002977	<i>Flavobacterium aquatile</i>
74	0	0.002743	<i>Paracoccus yeei</i>
52	0.000331	0.002911	<i>Sphingopyxis alaskensis</i>
126	0.000969	0.003524	<i>Pseudomonas xanthomarina</i>
89	0	0.002528	<i>Sphingomonas dokdonensis</i>
27	0.000113	0.001814	<i>Sphingopyxis chilensis</i>
63	0.00071	0.002374	<i>Paracoccus haeundaensis</i>
141	0.000156	0.001642	<i>Acinetobacter johnsonii</i>
84	0.000079	0.001484	<i>Pseudorhodobacter ferrugineus</i>
38	0.000063	0.001334	<i>Sejongia jeonii</i>
138	0.000014	0.001205	<i>Acinetobacter junii</i>
192	0	0.001099	<i>Methylovirgula ligni</i>
81	0.000248	0.00132	<i>Ruegeria atlantica</i>
8	0	0.001062	<i>Methylosinus sporium</i>
6	0	0.001033	<i>Methylovirgula ligni</i>
12	0	0.001029	<i>Pseudorhodoferrax soli</i>
86	0.000282	0.00131	<i>Methylobacterium rhodesianum</i>
32	0.000125	0.001139	<i>Methylobacterium zatmanii</i>
22	0.000011	0.000871	<i>Flavobacterium psychrophilum</i>
107	0.000082	0.000932	<i>Rhodobacter capsulatus</i>
43	0.000258	0.001098	<i>Flavobacterium aquatile</i>
94	0.000073	0.000886	<i>Rhodobacter sphaeroides</i>
189	0.000043	0.000809	<i>Methylobacterium album</i>
78	0.000047	0.000761	<i>Rhodobacter capsulatus</i>
91	0.000226	0.000929	<i>Caulobacter leidyia</i>
46	0.00002	0.000712	<i>Flavobacterium psychrolimnae</i>
101	0.00042	0.001111	<i>Brevundimonas variabilis</i>
154	0.00005	0.000728	<i>Methylobacterium adhaesivum</i>
82	0.000156	0.000815	<i>Amaricoccus kaplicensis</i>
54	0.000176	0.000782	<i>Methylopila capsulata</i>
19	0.000154	0.000756	<i>Brevundimonas variabilis</i>
71	0	0.00054	<i>Sphingomonas faeni</i>
182	0.000046	0.000579	<i>Flavobacterium aquatile</i>
90	0.000282	0.000812	<i>Methylobacterium adhaesivum</i>
44	0.000094	0.000609	<i>Flavobacterium psychrophilum</i>
143	0	0.000513	<i>Dokdonella koreensis</i>
112	0	0.000511	<i>Sphingomonas sanguinis</i>

Table 3. Cont.

N° OTU	mean abundance in group 1	mean abundance in group 2	Best hit on Blast
97	0	0.00049	<i>Hyphomicrobium facile subsp. Tolerans</i>
105	0	0.000359	<i>Paracoccus haeundaensis</i>
137	0	0.000339	<i>Acinetobacter johnsonii</i>
75	0.000558	0	<i>Pseudochrobactrum kiredjianiae</i>
50	0.889343	0.677999	<i>Methylobacterium rhodesianum</i>

Differentially abundance were calculated by using Metastats (using a FDR correction, $p < 0.01$).
doi:10.1371/journal.pone.0102649.t003

between *Methylobacterium* and *Flavobacterium* was further confirmed by the co-occurrence analysis, which identified potential taxonomic interactions between genera. Two significant correlations between genera were detected: a positive correlation between *Maritimibacter* and *Micrococcus* ($\sigma = 0.69$, p -value < 0.01) and a negative correlation between *Methylobacterium* and *Flavobacterium* ($\sigma = -0.63$, p -value < 0.01). Furthermore, variance analysis of OTU 36' abundance showed that it was influenced by the Fulton index (Shapiro test for Normality: p -value = 0.07, Linear model: p -value = 0.02741, $F = 5.0388$, $Rsq = -0.1496$). A negative relationship between Fulton index and the OTU 36 was observed (Figure 6).

In the F2 fish progeny, three significant QTLs (at the genome-wide level) were identified on two linkage groups (LG 11 and LG 16). One major QTL per strain was detected respectively for *Lysobacter*, *Rheinheimera* and *Methylobacterium* counts (LOD score = 9.89, 7.46 and 3.48 respectively). For each of these traits, the total PVE (percent variance explained) of the QTL were estimated to 17.01%, 31.05% and 41.1% for *Methylobacterium*, *Rheinheimera* and *Lysobacter* respectively. The most probable positions of these QTL, their respective 95% CIs, the closest linked molecular markers (one per QTL) as well as additive and dominance effects are presented in Table 6.

Discussion

Inter-individual variations in host microbiota has been well documented (e.g. [1]). Such variations occur even in hosts with identical genetic background, as observed in monozygotic twins [3] indicating that both genetic and environmental conditions play a role in the modulation of host microbiota. The gold standard forward genetics technique to identify areas of the genome that relate to certain phenotypes is to make a cross between genetically divergent individuals [32]. In this study, we focused on the structural variation of the skin microbiota (i.e abundance of each bacterial genus) of an F2 intercross originally generated from two genetically contrasted strains of brook charr. To our knowledge, this is the first study that identified host genomic regions associated with the abundance of specific microbiota strains in a non-model vertebrate.

The deep taxonomic analysis of the fish skin microbiota indicates that two phyla dominate: *Proteobacteria* (88.7%), followed by *Bacteroidetes* (9.7%). Those two phyla are mainly represented by a single OTU each; 50 and 36 respectively. OTU 50 is classified as *M. rhodesianum*. OTU 36 is classified as *F.*

psychrophilum, the causative agent of the cold water disease, a pervasive infectious disease in farmed fish [33,34]. This disease especially affects salmonids at early life-stages, and it is well documented that both stressful conditions, and injuries facilitate infection triggering [33,35]. According to the dendrogram (Figure 2), each individual's bacterial communities clustered into three groups. Similar clustering was also found with the second normalization method (subsampling of the same number of sequences for each samples). The same result was obtained with PCoA analysis and furthermore the Pearson correlation indicates that the genus *Flavobacterium* and *Methylobacterium* are negatively correlated. In the PCoA, two species (*F. psychrophilum* and *M. rhodesianum*) are correlated in opposite ways with the axis 1 which discriminate the three groups. We assume that two or more species which seem to be mutually exclusive are involved in a negative relationship [36]. All together those results indicate that the antagonistic relationship between those two dominant species is very likely playing a key role in shaping the structure of the individual's microbiota.

Because all fish progenies were reared in the same tank, and under the same environmental conditions since birth, environment had likely little influence on the phenotypic variation observed in this study. Yet, the individual's microbiota either clustered in one of two closely related groups, or formed divergent outliers. Sex-specific variations had little influence on individual clustering since both males and females were present in each defined group and there was no correlation between sex and groups. The difference between male and female is based on 22 OTUs which are less abundant than 0.1% of the community and are then classified as part of the rare biosphere. Previous studies observed similar individual variations of the microbiota in human or mouse [1,2,32].

Three OTUs classified into two species were responsible for the inter-individual differentiation; *M. rhodesianum* and *F. psychrophilum*. Interestingly, *Flavobacterium* and *Methylobacterium*, which are the most abundant genera, negatively co-occurred in the samples, indicating that the relationship between those two genera is based on competition. Furthermore, a strong negative correlation was found between the OTU 36 and 50, thus supporting a strong antagonistic relationship between the two species, *F. psychrophilum* and *M. rhodesianum*.

M. rhodesianum produces poly- β -hydroxybutyrate, a polymer of short-chain fatty acid, known to inhibit the growth of pathogens like enterobacteria and *Vibrio sp.* [37,38,39,40,41,42]. Taken together, this suggests that a reduction of *M. rhodesianum*

Table 4. Detailed information related to the OTU differentially abundant between individuals belonging to group 1 and group 3.

OTU	mean abundance in group 1	mean abundance in group 3	Best hit on Blast
36	0.00484	0.15627	<i>Flavobacterium psychrophilum</i>
52	0.000331	0.052599	<i>Sphingopyxis alaskensis</i>
51	0.002582	0.045797	<i>Sphingomonas paucimobilis</i>
119	0.001237	0.034255	<i>Pseudomonas peli</i>
125	0.001567	0.030681	<i>Acinetobacter lwoffii</i>
120	0.002302	0.030821	<i>Rheinheimera texasensis</i>
63	0.00071	0.027857	<i>Paracoccus haeundaensis</i>
122	0.000778	0.018741	<i>Pseudomonas cedrina subsp. Fulgida</i>
61	0.003372	0.02093	<i>Porphyrobacter dokdonensis</i>
21	0.000117	0.011826	<i>Flavobacterium aquatile</i>
127	0.000218	0.00999	<i>Acinetobacter haemolyticus</i>
55	0	0.008105	<i>Bradyrhizobium pachyrhizi</i>
54	0.000176	0.007992	<i>Methylopila capsulata</i>
162	0.002112	0.009611	<i>Knoellia sinensis</i>
126	0.000969	0.008421	<i>Pseudomonas xanthomarina</i>
67	0.000518	0.007951	<i>Rhodobacter blasticus</i>
69	0.000531	0.007849	<i>Pseudorhodobacter ferrugineus</i>
81	0.000248	0.006693	<i>Ruegeria atlantica</i>
134	0.000245	0.00635	<i>Acinetobacter lwoffii</i>
143	0	0.005914	<i>Dokdonella koreensis</i>
129	0	0.005644	<i>Pseudomonas hibiscicola</i>
124	0	0.004908	<i>Nevskia soli</i>
40	0.000014	0.004576	<i>Flavobacterium aquatile</i>
38	0.000063	0.004547	<i>Sejongia jeonii</i>
68	0.0014	0.005784	<i>Methylocella palustris</i>
71	0	0.004232	<i>Sphingomonas faeni</i>
79	0.000259	0.004408	<i>Rhodobacter capsulatus</i>
78	0.000047	0.003657	<i>Rhodobacter capsulatus</i>
37	0.000471	0.004035	<i>Flavobacterium terrigena</i>
141	0.000156	0.003673	<i>Acinetobacter johnsonii</i>
156	0.000033	0.003487	<i>Sphingopyxis alaskensis</i>
39	0	0.00318	<i>Flavobacterium psychrophilum</i>
24	0.000095	0.00295	<i>Pseudomonas peli</i>
106	0.000106	0.002945	<i>Bosea vestrisii</i>
137	0	0.002615	<i>Acinetobacter johnsonii</i>
101	0.00042	0.002997	<i>Brevundimonas variabilis</i>
97	0	0.002498	<i>Hyphomicrobium facile subsp. Tolerans</i>
139	0.000153	0.002639	<i>Lysobacter capsici</i>
157	0.00006	0.002488	<i>Sphingomonas sanguinis</i>
116	0	0.002339	<i>Rhodobacter blasticus</i>
177	0.000289	0.00255	<i>Aquabacterium commune</i>
25	0.000069	0.002236	<i>Acinetobacter lwoffii</i>
182	0.000046	0.002172	<i>Flavobacterium aquatile</i>
47	0	0.002082	<i>Flavobacterium psychrophilum</i>
84	0.000079	0.00214	<i>Pseudorhodobacter ferrugineus</i>
164	0.000219	0.002164	<i>Rhodococcus fascians</i>
105	0	0.001788	<i>Paracoccus haeundaensis</i>
43	0.000258	0.002022	<i>Flavobacterium aquatile</i>
180	0.000017	0.001709	<i>Acidovorax defluvii</i>
92	0.000147	0.001809	<i>Sphingopyxis witflariensis</i>

Table 4. Cont.

OTU	mean abundance in group 1	mean abundance in group 3	Best hit on Blast
4	0	0.00161	<i>Methylosinus sporium</i>
145	0	0.001417	<i>Acinetobacter johnsonii</i>
5	0	0.001415	<i>Magnetospirillum magnetotacticum</i>
88	0.000036	0.00143	<i>Amaricoccus veronensis</i>
3	0	0.001358	<i>Rhodobacter changlensis</i>
138	0.000014	0.001365	<i>Acinetobacter junii</i>
70	0.000831	0.002052	<i>Methylobacterium salsuginis</i>
184	0	0.001156	<i>Flavobacterium aquatile</i>
22	0.000011	0.001144	<i>Flavobacterium psychrophilum</i>
112	0	0.001106	<i>Sphingomonas sanguinis</i>
118	0.000062	0.001143	<i>Haematobacter missouriensis</i>
93	0.000029	0.001093	<i>Roseomonas mucosa</i>
152	0.000028	0.001009	<i>Rhodobacter blasticus</i>
35	0.00005	0.000936	<i>Sphingomonas sanguinis</i>
148	0.000033	0.000915	<i>Psychrobacter arcticus</i>
172	0	0.000864	<i>Sphingomonas sanguinis</i>
146	0	0.000832	<i>Pseudomonas mohnii</i>
94	0.000073	0.000865	<i>Rhodobacter sphaeroides</i>
140	0	0.000775	<i>Psychrobacter alimentarius</i>
160	0.000061	0.000823	<i>Sphingomonas faeni</i>
83	0.000128	0.000877	<i>Paracoccus aminovorans</i>
42	0.000323	0.001071	<i>Flavobacterium psychrophilum</i>
186	0	0.000711	<i>Flavobacterium aquatile</i>
44	0.000094	0.000796	<i>Flavobacterium psychrophilum</i>
117	0.000029	0.000731	<i>Sphingopyxis taejonensis</i>
107	0.000082	0.000766	<i>Rhodobacter capsulatus</i>
72	0.000432	0.001098	<i>Beijerinckia mobilis</i>
104	0.000028	0.000693	<i>Sphingopyxis wittflariensis</i>
113	0	0.000663	<i>Stella humosa</i>
115	0	0.000622	<i>Loktanella salsilacus</i>
108	0.000051	0.000647	<i>Novosphingobium panipatense</i>
26	0	0.000559	<i>Acinetobacter johnsonii</i>
48	0	0.00052	<i>Flavobacterium psychrophilum</i>
166	0	0.000508	<i>Rhodococcus fascians</i>
65	0.00064	0.001148	<i>Methylobacterium organophilum</i>
103	0.000215	0.000655	<i>Sphingomonas ursincola</i>
144	0.00007	0.000462	<i>Acinetobacter lwoffii</i>
191	0.00006	0.000406	<i>Beijerinckia derxii</i> subsp. <i>Venezuelae</i>
19	0.000154	0.000498	<i>Brevundimonas variabilis</i>
178	0.000043	0.000378	<i>Pseudorhodoferax soli</i>
23	0.000017	0.000348	<i>Flavobacterium psychrophilum</i>
192	0	0.000328	<i>Methylovirgula ligni</i>
49	0.000058	0.000354	<i>Flavobacterium aquatile</i>
8	0	0.00029	<i>Methylosinus sporium</i>
9	0.000062	0.000245	<i>Sphingomonas wittichii</i>
169	0.000207	0.000287	<i>Corynebacterium tuberculostearicum</i>
111	0.000416	0.000202	<i>Mycoplana bullata</i>
7	0.000343	0	<i>Singularimonas variicoloris</i>
175	0.000348	0	<i>Simplicispira limi</i>

Table 5. Detailed information related to the OTU differentially abundant between individuals belonging to group 2 and group 3.

N° OTU	mean abundance in group 2	mean abundance in group 3	Best hit on Blast
63	0.002374	0.027857	<i>Paracoccus haeundaensis</i>
55	0	0.008105	<i>Bradyrhizobium pachyrhizi</i>
54	0.000782	0.007992	<i>Methylophilus capsulata</i>
134	0.000155	0.00635	<i>Acinetobacter lwoffii</i>
129	0	0.005644	<i>Pseudomonas hibiscicola</i>
143	0.000513	0.005914	<i>Dokdonella koreensis</i>
124	0.000109	0.004908	<i>Nevskia soli</i>
40	0.000306	0.004576	<i>Flavobacterium aquatile</i>
123	0.000266	0.004124	<i>Acinetobacter haemolyticus</i>
71	0.00054	0.004232	<i>Sphingomonas faeni</i>
38	0.001334	0.004547	<i>Sejongia jeonii</i>
156	0.000306	0.003487	<i>Sphingopyxis alaskensis</i>
39	0.000144	0.00318	<i>Flavobacterium psychrophilum</i>
78	0.000761	0.003657	<i>Rhodobacter capsulatus</i>
137	0.000339	0.002615	<i>Acinetobacter johnsonii</i>
47	0.000087	0.002082	<i>Flavobacterium psychrophilum</i>
164	0.000236	0.002164	<i>Rhodococcus fascians</i>
4	0	0.00161	<i>Methylosinus sporium</i>
182	0.000579	0.002172	<i>Flavobacterium aquatile</i>
3	0	0.001358	<i>Rhodobacter changlensis</i>
184	0	0.001156	<i>Flavobacterium aquatile</i>
42	0.000252	0.001071	<i>Flavobacterium psychrophilum</i>
12	0.001029	0	<i>Pseudorhodoferrax soli</i>
6	0.001033	0	<i>Methylovirgula ligni</i>
27	0.001814	0.000082	<i>Sphingopyxis chilensis</i>
50	0.677999	0.227483	<i>Methylobacterium rhodesianum</i>

Differentially abundance were calculated by using Metastats (using a FDR correction, $p < 0.01$).
doi:10.1371/journal.pone.0102649.t005

abundance allows colonization or over-growth by other bacteria, some of which are pathogenic.

Evidently, a change in skin microbiota taxonomic structure favors opportunist bacteria and especially opportunistic pathogens like *F. psychrophilum*, *Acinetobacter haemolyticus*, *Acinetobacter johnsonii* and *Acinetobacter junii*, which have a higher abundance in group 2 compared with group 1. Such a disturbance in the microbiome homeostasis is called dysbiosis, and its occurrence enlightens the importance of the function of *M. rhodesianum* in controlling the balance between both other commensal bacteria and opportunistic pathogens. Furthermore, *Flavobacterium psychrophilum* was negatively correlated to both *M. rhodesianum* abundance and Fulton index. The Fulton index is a condition factor used as a proxy for fish health status. Therefore, it suggests that skin microbiota from the weakest host individuals were unable to prevent colonization by the opportunistic pathogen *F. psychrophilum*. Furthermore, skin microbiota taxonomic structure varies significantly across the F2 progeny. This result added to the fact that *i* the genetics of the host is the only variable in the experiment, and *ii* the finding of three QTLs associated to the abundance of bacterial genera with high PVE, suggest that host

genotype influences the abundance of commensal strains e.g. *Methylobacterium*, which will regulate the abundance of *F. psychrophilum*.

We found three QTL associated with the abundance of three genera: *Lysobacter*, *Rheinheimera* and *Methylobacterium*, all of which were observed to provide antimicrobial compounds [39,40,41,42,43,44,45,46,47]. These findings strongly suggest that host genotype influences abundance of specific commensal strains, and possibly targets their recruitment. The most compelling evidence concerns the QTL associated with *Methylobacterium* (PVE = 17.01%): as previously described above, *Methylobacterium* is influential on the structure and the homeostasis of the microbiota. Its abundance is inversely correlated with those of the pathogen *F. psychrophilum*. The targeted recruitment of *M. rhodesianum* mediated by the host genotype is therefore a strategy to prevent pathogen growth *via* harnessing antagonistic relationship towards resources use [48]. To our knowledge, this is only the second study that identified QTL associated with microbial variation among individuals. A study on murine gut microbiota showed similar results, in which McKnite *et al.* (2012) found 5 QTL located on four chromosomes influencing the variation of




Figure 5. PCoA analysis of the microbiome for all 86 F₂ individuals based on the Yue & Clayton measure of dissimilarity (Thetayc).

Black circles represent individuals belonging to the group 1, red circles represent individuals belonging to the group 2 and green circles represent individuals belonging to the group 3. Arrows represent genus, which are significantly correlated with the axis. The first and second axes represented 24.5% and 7.7% of the variation respectively. The R-squared between the original distance matrix and the distance between the points in 2D PCoA space was 0.87.

doi:10.1371/journal.pone.0102649.g005

different taxa. The variance explained by their QTL is in the same range of our (20% to 27% of the variance explained for McKnight *et al.* (2012)). We also found QTLs with a major effect on the variance of genus abundances (PVE ranging for 17.02 to 41.1%). Linkage analysis on genus abundance data strongly evidenced the influence of host genetics on the modulation and/or recruitment for those genera. Therefore, brook charr immunity involves both specific commensal strains recruitment and nuclear gene expression, as previously evidenced in [49], those are under the control of the host genotype.

The study of the genetic architecture underlying the regulation of bacterial abundance further highlights the coevolutionary

pattern between host, commensals, and pathogens. However, identifying genes located in the genomic regions (QTL) linked to the abundance of microbial partners will be far more challenging as the genome of brook charr is currently not fully sequenced and poorly annotated compared to the mouse genome. The markers surrounding the QTL regions may define regions to be deeply sequenced in future work to identify potential underlying genes and their associated functions. Evidently, those markers will be invaluable to conduct highly innovative genetic breeding programs targeting disease resistant host strains via the recruitment of highly resilient microbiota.




Figure 6. Relationship between the Fulton Index and the abundance of the OTU 36. A linear regression is observed based on a linear model (Shapiro test for Normality: p -value = 0.07, Linear model: p -value = 0.02741, F = 5.0388, R^2 = 0.04536).
doi:10.1371/journal.pone.0102649.g006

Table 6. Descriptive statistics, including the LOD score, the position, 95% CI, PVE (%), the associated P-value of each QTL linked to every phenotype related to bacteria counts trait (LOD, \log_{10} of the odd ratio; 95% CI, 95% confidence interval; PVE, percent variance explained) [23].

Phenotype	Linkage Group	Pos (cM)	95% CI (cM)	LOD	p value (F)	PVE (in %)	Nearest marker
<i>Lysobacter</i>	11	62	56.2–68.2	9.89	0.000	41.1	sf003455
<i>Rheinheimera</i>	16	42.8	39.3–45.5	7.46	0.000	31.05	SFO-D91
<i>Methylobacterium</i>	16	39	34.5–43.5	3.48	0.001	17.02	SFO-D91

doi:10.1371/journal.pone.0102649.t006

Supporting Information

Figure S1 Rarefaction curves for each group defined by the dendrogram analysis based upon ThetaYC index (see figure 2). Each group curve reaches a plateau thus indicating the depth of sequencing is sufficient. (EPS)

Figure S2 Alpha diversity of each group normalized by subsampling of the same number of reads per sample. Each group was defined by the dendrogram analysis based upon ThetaYC index performed on the normalized dataset. Alpha-diversity was calculated by the non-parametric index of Shannon. Statistical differences (represented by the asterisk) were calculated with a wilcoxon test with a correction for multiple testing (Holm, p -values $< 1.10^{-4}$). (EPS)

Figure S3 PCoA analysis of the microbiome for all 86 F₂ individuals. This PCoA was performed on a normalized dataset with the second method of normalization (subsampling of the same number of reads per sample). Black circles represent individuals belonging to the group 1, red circles represent individuals belonging to the group 2 and green circles represent individuals belonging to the group 3. Arrows represent genera, which are

significantly correlated with the axis. The first and second axes represented 24.5% and 7.8% of the variation respectively. The R-squared between the original distance matrix and the distance between the points in 2D PCoA space was 0.88. (EPS)

Table S1 Differentiation between the three groups of OTU calculated with an analysis of similarity (ANOSIM, statistic R). The q -values represent p -values corrected with Bonferroni and considered as significant when $p < 0.05$. Note: The analysis was performed on a dataset normalized by two methods: z -score and subsampling. Both analyses gave the same results, each group being significantly different from the two others. (DOC)

Acknowledgments

We thank Martin Llewellyn, Scott A. Pavey, the associate editor of Plos One, and two anonymous reviewers for critical reading of the manuscript.

Author Contributions

Conceived and designed the experiments: ND CA LB SB CS. Performed the experiments: SB CS. Analyzed the data: SB CS. Contributed reagents/materials/analysis tools: ND CA LB. Wrote the paper: ND CA LB SB CS.

References

- Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, et al. (2009) Bacterial community variation in human body habitats across space and time. *Science* 326: 1694–1697.
- Fierer N, Lauber CL, Zhou N, McDonald D, Costello EK, et al. (2010) From the cover: Forensic identification using skin bacterial communities. *Proceedings of the National Academy of Sciences* 107: 6477–6481.
- Turnbaugh PJ, Quince C, Faith JJ, McHardy AC, Yatsunenko T, et al. (2010) Organismal, genetic, and transcriptional variation in the deeply sequenced gut microbiomes of identical twins. *Proceedings of the National Academy of Sciences* 107: 7503–7508.
- Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, et al. (2007) The human microbiome project. *Nature* 449: 804–810.
- Turnbaugh PJ, Henriksen B, Gordon JI (2010) Viewing the human microbiome through three-dimensional glasses: integrating structural and functional studies to better define the properties of myriad carbohydrate-active enzymes. *Acta Crystallographica Section F* 66: 1261–1264.
- Turnbaugh PJ, Gordon JI (2009) The core gut microbiome, energy balance and obesity. *The Journal of Physiology* 587: 4153–4158.
- Giongo A, Gano KA, Crabb DB, Mukherjee N, Novelo LL, et al. (2011) Toward defining the autoimmune microbiome for type 1 diabetes. *ISME J* 5: 82–91.
- Peterson DA, McNulty NP, Guruge JL, Gordon JI (2007) IgA Response to Symbiotic Bacteria as a Mediator of Gut Homeostasis. *Cell Host & Microbe* 2: 328–339.
- Bouskra D, Brézillon C, Bérard M, Werts C, Varona R, et al. (2008) Lymphoid tissue genesis induced by commensals through NOD1 regulates intestinal homeostasis. *Nature* 456: 507–510.
- Macpherson AJ, Geuking MB, McCoy KD (2011) Immunoglobulin A: a bridge between innate and adaptive immunity. *Current Opinion in Gastroenterology* 27: 529–533. 10.1097/MOG.1090b1013e32834bb32805
- Boutin S, Bernatchez L, Audet C, Derôme N (2012) Antagonistic effect of indigenous skin bacteria of brook charr (*Salvelinus fontinalis*) against *Flavobacterium columnare* and *F. psychrophilum*. *Veterinary Microbiology* 155: 355–361.
- Balcázar J, de Blas I, Ruiz-Zaruela I, Vendrell D, Muzquiz J (2004) Probiotics: a tool for the future of fish and shellfish health management. *Journal of Aquaculture in the Tropics* 19: 239–242.
- Backhed F (2005) Host-bacterial mutualism in the human intestine. *Science* 307: 1915–1920.
- Hooper LV, Gordon JI (2001) Commensal host-bacterial relationships in the gut. *Science* 292: 1115–1118.
- O'Hara AM, Shanahan F (2006) The gut flora as a forgotten organ. *EMBO Reports* 7: 688–693.
- Stecher B, Hardt W-D (2008) The role of microbiota in infectious disease. *Trends in Microbiology* 16: 107–114.
- Loiseau C, Richard M, Garnier S, Chastel O, Julliard R, et al. (2009) Diversifying selection on MHC class I in the house sparrow (*Passer domesticus*). *Molecular Ecology* 18: 1331–1340.
- Krawczyk M, Reith W (2006) Regulation of MHC class II expression, a unique regulatory system identified by the study of a primary immunodeficiency disease. *Tissue Antigens* 67: 183–197.
- Dionne M, Miller KM, Dodson JJ, Bernatchez L (2009) MHC standing genetic variation and pathogen resistance in wild Atlantic salmon. *Philosophical Transactions of the Royal Society B-Biological Sciences* 364: 1555–1565.
- Alexander C, Rietschel ET (2001) Invited review: Bacterial lipopolysaccharides and innate immunity. *Journal of Endotoxin Research* 7: 167–202.
- Boutin S, Audet C, Derôme N (2013) Probiotic treatment by indigenous bacteria decreases mortality without disturbing the natural microbiota of *Salvelinus fontinalis*. *Canadian Journal of Microbiology* In press.
- Boutin S, Bernatchez L, Audet C, Derôme N (2013) Network Analysis Highlights Complex Interactions between Pathogen, Host and Commensal Microbiota. *PLoS ONE* 8: e84772.
- Sauvage C, Vagner M, Derôme N, Audet C, Bernatchez L (2012) Coding gene single nucleotide polymorphism mapping and quantitative trait loci detection for physiological reproductive traits in brook charr, *Salvelinus fontinalis*. *G3: Genes|Genomes|Genetics* 2: 379–392.
- Livia L, Antonella P, Hovirag L, Mauro N, Panara F (2006) A nondestructive, rapid, reliable and inexpensive method to sample, store and extract high-quality DNA from fish body mucus and buccal cells. *Molecular Ecology Notes* 6: 257–260.
- Turner S, Pryer KM, Miao VPW, Palmer JD (1999) Investigating deep phylogenetic relationships among *Cyanobacteria* and plastids by small subunit rRNA sequence analysis. *Journal of Eukaryotic Microbiology* 46: 327–338.
- Marchesi JR, Sato T, Weightman AJ, Martin TA, Fry JC, et al. (1998) Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Applied and Environmental Microbiology* 64: 795–799.
- Schloss P, Westcott S, Ryabin T, Hall J, Hartmann M, et al. (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* 75: 7537–7541.
- Siddharth J, Holway N, Parkinson SJ (2013) A Western Diet Ecological Module Identified from the 'Humanized' Mouse Microbiota Predicts Diet in Adults and Formula Feeding in Children. *PLoS ONE* 8: e83689.
- Takeshita T, Suzuki N, Nakano Y, Yasui M, Yoneda M, et al. (2012) Discrimination of the oral microbiota associated with high hydrogen sulfide and methyl mercaptan production. *Sci Rep* 2.
- Horner-Devine MC, Lage M, Hughes JB, Bohannon BJM (2004) A taxa-area relationship for bacteria. *Nature* 432: 750–753.
- White JR, Nagarajan N, Pop M (2009) Statistical methods for detecting differentially abundant features in clinical metagenomic samples. *PLoS Computational Biology* 5: e1000352.
- McKite AM, Perez-Munoz ME, Lu L, Williams EG, Brewer S, et al. (2012) Murine gut microbiota is defined by host genetics and modulates variation of metabolic traits. *PLoS ONE* 7: e39191.
- Starliper CE (2011) Bacterial coldwater disease of fishes caused by *Flavobacterium psychrophilum*. *Journal of Advanced Research* 2: 97–108.

34. Wiklund T, Madsen L, Bruun MS, Dalsgaard I (2000) Detection of *Flavobacterium psychrophilum* from fish tissue and water samples by PCR amplification. *Journal of Applied Microbiology* 88: 299–307.
35. Madetoja J, Nyman P, Wiklund T (2000) *Flavobacterium psychrophilum*, invasion into and shedding by rainbow trout *Oncorhynchus mykiss*. *Diseases of Aquatic Organisms* 43: 27–38.
36. Faust K, Raes J (2012) Microbial interactions: from networks to models. *Nat Rev Micro* 10: 538–550.
37. Defoirdt T, Halet D, Vervaeren H, Boon N, Van de Wiele T, et al. (2007) The bacterial storage compound poly- β -hydroxybutyrate protects *Artemia franciscana* from pathogenic *Vibrio campbellii*. *Environmental Microbiology* 9: 445–452.
38. Halet D, Defoirdt T, Van Damme P, Vervaeren H, Forrez I, et al. (2007) Poly- β -hydroxybutyrate-accumulating bacteria protect gnotobiotic *Artemia franciscana* from pathogenic *Vibrio campbellii*. *FEMS Microbiology Ecology* 60: 363–369.
39. Yellore Desai (1998) Production of poly-3-hydroxybutyrate from lactose and whey by *Methylobacterium* sp. ZP24. *Letters in Applied Microbiology* 26: 391–394.
40. Mothes G, Ackermann J-U, Babel W (1998) Regulation of poly(β -hydroxybutyrate) synthesis in *Methylobacterium rhodesianum* MB 126 growing on methanol or fructose. *Archives of Microbiology* 169: 360–363.
41. Bormann EJ, Roth M (1999) The production of polyhydroxybutyrate by *Methylobacterium rhodesianum* and *Ralstonia eutropha* in media containing glycerol and casein hydrolysates. *Biotechnology Letters* 21: 1059–1063.
42. Bormann EJ, Roth M, Linß W, Leißner M (1999) Selection of capsule deficient strains of *Methylobacterium rhodesianum* producing polyhydroxybutyrate. *Biotechnology Techniques* 13: 539–544.
43. Balachandran C, Duraipandian V, Ignacimuthu S (2012) Cytotoxic (A549) and antimicrobial effects of *Methylobacterium* sp. isolate (ERI-135) from Nilgiris forest soil, India. *Asian Pacific Journal of Tropical Biomedicine* 2: 712–716.
44. Korotkova N, Chistoserdova L, Lidstrom ME (2002) Poly- β -Hydroxybutyrate Biosynthesis in the Facultative Methylophilic *Methylobacterium extorquens* AM1: Identification and Mutation of gap11, gap20, and phaR. *Journal of Bacteriology* 184: 6174–6181.
45. Chen W-M, Lin C-Y, Sheu S-Y (2010) Investigating antimicrobial activity in *Rheinheimera* sp. due to hydrogen peroxide generated by l-lysine oxidase activity. *Enzyme and Microbial Technology* 46: 487–493.
46. Chen W-M, Lin C-Y, Young C-C, Sheu S-Y (2010) *Rheinheimera aquatica* sp. nov., antimicrobial activity-producing bacterium isolated from freshwater culture pond. Seoul, COREE, REPUBLIQUE DE: Korean Society for Applied Microbiology. 7 p.
47. Reichenbach H (2006) The genus *Lysobacter*. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E, editors. *The Prokaryotes*: Springer New York. pp. 939–957.
48. Dillon R, Charnley K (2002) Mutualism between the desert locust *Schistocerca gregaria* and its gut microbiota. *Research in Microbiology* 153: 503–509.
49. Langevin C, Blanco M, Martin SAM, Jounneau L, Bernardet J-F, et al. (2012) Transcriptional responses of resistant and susceptible fish clones to the bacterial pathogen *Flavobacterium psychrophilum*. *PLoS ONE* 7: e39126.