



HAL
open science

Bacillus phytases: Current status and future prospects

Mohamed Ali Borgi, Samira S. Boudebouze, Hela H. Mkaouar, Emmanuelle Maguin, Moez M. Rhimi

► **To cite this version:**

Mohamed Ali Borgi, Samira S. Boudebouze, Hela H. Mkaouar, Emmanuelle Maguin, Moez M. Rhimi. Bacillus phytases: Current status and future prospects. *Bioengineered*, 2015, 6 (4), pp.233-236. 10.1080/21655979.2015.1048050 . hal-02634782

HAL Id: hal-02634782

<https://hal.inrae.fr/hal-02634782>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



Bacillus phytases: Current status and future prospects

Mohamed Ali Borgi, Samira Boudebouze, H la Mkaouar, Emmanuelle Maguin & Moez Rhimi

To cite this article: Mohamed Ali Borgi, Samira Boudebouze, H la Mkaouar, Emmanuelle Maguin & Moez Rhimi (2015) Bacillus phytases: Current status and future prospects, Bioengineered, 6:4, 233-236, DOI: [10.1080/21655979.2015.1048050](https://doi.org/10.1080/21655979.2015.1048050)

To link to this article: <http://dx.doi.org/10.1080/21655979.2015.1048050>



Accepted author version posted online: 06 May 2015.
Published online: 06 May 2015.



Submit your article to this journal [↗](#)



Article views: 195



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 2 View citing articles [↗](#)

Bacillus phytases: Current status and future prospects

Mohamed Ali Borgi¹, Samira Boudebouze², H la Mkaouer², Emmanuelle Maguin², and Moez Rhimi^{2,*}

¹Faculty of Sciences of Gafsa - Unit of Macromolecular Biochemistry and Genetic; Department of Life Sciences; Zarroug, Gafsa, Tunisia; ²INRA, UMR 1319 Micalis; Jouy-en-Josas, France

Phytases catalyze the hydrolysis of phytic acid in a stepwise manner to lower inositol phosphates, myo-inositol (having important role in metabolism and signal transduction pathways), and inorganic phosphate. These enzymes have been widely used in animal feed in order to improve phosphorus nutrition and to decrease pollution in animal waste. Compared to previously described phytases, the phytase (PhyL) from *Bacillus licheniformis* ATCC 14580 has attractive biochemical properties which can increase the profitability of several biotechnological procedures (animal nutrition, human health...etc). Due to its amino acid sequence with critical substitutions, the PhyL could be a model to enhance other phytases features, in terms of thermal stability and high activity. Otherwise, an engineered PhyL, with low pH optimum, will represent a challenge within the class of β -propeller phytases.

Phytate, the principle storage form of phosphate and inositol in cereals, legumes, oil seeds and nuts, strongly chelates charged proteins, minerals and amino acids within digestive tract.¹ Phytate phosphorus is largely unavailable to monogastric animals due to the lack or insufficient amount of phytate degrading enzymes in their gastrointestinal tract.² Degradation of phytate is catalysed by phytase (myo-inositol hexakisphosphate phosphohydrolase; EC 3.1.3.8 and EC 3.1.3.26) which releases a series of lower isomers of myo-inositol phosphates.³ This hydrolytic reaction plays an important role in energy metabolism, metabolic regulation and signal transduction pathways in biological system.⁴ Hence, phytases have been

studied intensively due to its potential application as feed additives, processing and manufacturing of human food to improve mineral nutrition.⁵ In fact, β -Propeller phytases, a class to which belong *Bacillus* phytases were shown to entirely abrogate the ability of phytate to chelate metal ions.⁶ In addition to that, phytase use protects the environment against phosphorus pollution.⁷ This phosphohydrolase has different possible sources: plant phytases, microbial phytases (fungal and bacterial phytases), Mucosal phytases derived from small intestine and gut microfloral phytases.⁸

The phytate-hydrolysing enzyme has many applications in food industries. It has a potential for producing low phytin bread.⁹ Addition of phytase improves also the nutritional value of bread through the reduction of phytate content and enhances the activation of endogenous alpha-amylase by making more calcium available.¹⁰ Phytase can also be added for the production of phytate-free soymilk.¹¹ Phytase plays a crucial role for various inositol phosphate preparations, especially in immobilized forms.^{12,13} The myo-inositol phosphates have various beneficial effects on health, as enzyme stabilizers,¹⁴ inhibitors of enzymes and thus as potential drug blockers.¹⁵ In animal nutrition, phytases are used in aquaculture feed and additive to ensure proper degradation of the phytate present in animal diets during digestion in the stomach.¹⁶ For improved phosphorus utilization in animal agriculture, several transgenic plants overexpressing bacterial phytases were generated, including alfalfa, soybean, potato, rice and wheat.¹⁷⁻²¹

Given that phytases deliver economic benefits through their ability to replace added inorganic phosphorus, many works

Keywords: *Bacillus licheniformis*, biotechnological applications, enzyme engineering, phytase, protein biochemistry

*Correspondence to: Moez Rhimi; Email: moez.rhimi@jouy.inra.fr

Submitted: 04/06/2015

Revised: 04/24/2015

Accepted: 04/29/2015

<http://dx.doi.org/10.1080/21655979.2015.1048050>

Addendum to: Borgi MA, Khila M, Boudebouze S, Aghajari N, Szukala F, Pons N, Maguin E, Rhimi M. The attractive recombinant phytase from *Bacillus licheniformis*: biochemical and molecular characterization. *Appl Microbiol Biotechnol*. 2014 Jul;98(13):5937-47. doi: 10.1007/s00253-013-5421-9. Epub 2013 Dec 13.

were undertaken with the aim to improve phytase features and make it more suitable for industrial uses. Several studies were focused on the enhancement of thermal stability to preserve the enzyme activity during the heat step of feed pelleting under high temperature.²² In this field, the PhyL is well suited due to its remarkable thermal stability. According to Farhat-Khemakhem et al. (2013)²³, it seems to be interesting to substitute the residue Ala 257 into Pro inside PhyL. This mutation should enhance more and more the thermal stability of PhyL.

The phytase of *Bacillus licheniformis* ATCC 14580 gathers the best features to be involved in animal feed formulation.²⁴ Its high specific activity toward phytic acid is a major parameter to be used for myo-inositol phosphates production.

Unlike other phytases from *Bacillus* strains, the low Ca²⁺ requirement of PhyL for its optimal activity seems to be explained by the fact that Ca²⁺ ions are not involved alone ovoid the on the maintain of the enzyme leading to an active state.²⁵

Due to their atypical features compared to other phytases, the phyL from *Bacillus licheniformis* ATCC14580 could be promising to overcome the inhibitory effect of phytic acid and polyphenols as they chelate minerals in feeds. Such fact was demonstrated to be useful to limit zinc

deficiency and ovoid the fortification process of cereals staples with zinc.²⁶ Supplementation of phyL in feeds should improve growth performance and nutrient digestibility as well as the increase of gene expression encoding for the peptide transporter.²⁷ In the same context, Zeng et al. (2014)²⁸ described the use of higher phytase amount produced from *E. coli* (having lesser interests than PhyL) to further improved mineral use, protein use and performance of young pigs.

Otherwise, the high thermal stability of PhyL compared to phytases from Bacteria, Fungi and Yeast (Table 1) is especially of interest for pelleting purposes. In fact, Park et al. (2003)²⁹ demonstrated the suitability of the phytase from *Bacillus amyloliquefaciens* (which is less thermostable compared to PhyL) in this field. Thereby, we believe that using PhyL in pelleting process is an attractive strategy.

On the basis on the works of Sanz-Penella et al. (2009)³⁰ and Haros et al. (2007)³¹, it will be interesting to explore the PhyL genes within Bifidobacteria for bread fermentation technology. Taking into account the high specific activity of PhyL, such feature constitutes a promising way to reduce the content of InsP(6) in rich fiber products for human consumption, in favor of InsP(3) production. Dephosphorylation of other

phosphorylated molecules could also be performed by using PhyL.

Engineered PhyL with low pH optimum can constitute a remarkable perspective since the obvious drawback of phytase from *Bacillus* is their inability to act at acidic conditions. Even so, no truly reliable methods for modifying the pH activity profile of an enzyme are yet available and the decrease of the pH optimum of phytases from *Bacillus* became a challenge. In this context, different strategies could be applied in order to modify the enzyme pH feature. The first is the mutation of ionizable groups that are implicated in substrate binding or catalysis by nonionizable ones or by amino acids having different charge or pK values.³² The second is the replacement of residues interacting with Alanine residues by forming hydrogen bonds and/or salt bridges. Substitution of such residues may disturb the hydrogen-bonding network in the active site or alter the electronic environment of Alanine residues.^{33,34} The third is the modification of the enzyme surface charge, which can be achieved by chemical modification of residues located at the protein surface.^{35,36} In fact, making the surface more positively charged lowers the pKa values of ionizable groups and, thus the pH optimum. Such fact is favoured at low ionic strength.³⁶

Table 1. The PhyL properties compared to those from other previously described phytases

Phytase source	Optimal temperature (°C)/ Activity at low temperature ^a	pH optimum	Specific activity (U.mg ⁻¹)	Molecular weight (kDa)	Ca ²⁺ demand (mM)	Reference
Bacillus						
<i>B. licheniformis</i> ATCC 14580	75/40% at 4°C	6.5–7.0	316	42	0.6	24
<i>B. subtilis</i> US417	55/50% at 37°C	7.5	25	41	1.0	40
<i>B. subtilis</i> 168	55/>5% at 25°C	7.0	36.9	44	5.0	37
<i>B. licheniformis</i>	65/>10% at 25°C	7.0	23,6	47	5.0	37
<i>B. subtilis</i> VTT E-68013	55/>20% at 37°C	7.0	88	43	1.0	41
<i>B. sp</i> MD2	67–73/-	6–7	39	47.5	2.0–5.0	38
<i>B. laevolacticus</i>	70/30% at 30°C	7.0–8.0	12.69	46	5.0	42
<i>B. sp</i> KHU-10	60/20% at 20°C	6.5–8.5	36	44	10.0	43
Fungi						
<i>Buttiauxella</i> sp. GC21	55/40% at 30°C	4.5	1180	45	No effect	44
<i>Aspergillus ficuum</i> NTG-23	67/40% at 30°C	1.3	150.1	65.5	No effect	45
Yeasts						
<i>Hansenula fabianii</i> J640	50/>20% at 20°C	4.5	25.67	49	No effect	46
<i>Kodamaea ohmeri</i> BG3	65/>20% at 30°C	5.0	16.5	51	No effect	47
<i>S. cerevisiae</i> CY	40/>20% at 20°C	3.6	71.06	55	inhibited	48
<i>Debaryomyces castellii</i> CBS 2923	60	4.0–4.5	182	51.2	—	49

^arelative activity is indicated.

The inspection of the PhyL amino acid sequence in comparison with previously reported phytases from *Bacillus* genus showed some original substitutions. It was found that more than 40 substitutions were encountered inside the PhyL, compared to the most related phytase from *B. licheniformis* previously characterized by Tye et al. (2002).³⁷ In spite of their high sequence homology, the two phytases have significant differences in their specific activity, thermostability and efficiency at low temperature and requirement of Ca²⁺ ions. Among the 40 substitutions the N86/K, N139/S, N239/D, G251/D, D302/E could impact the performance of PhyL. Site-directed mutagenesis, crystallization and enzyme modeling procedures should certainly shed light on the role of these substitutions. These observations increasingly confirmed by the works of Tran et al. (2010)³⁸, which introduced the E229V and S283R mutations in phytase from *Bacillus* sp. MD2 and the recent work of Xu et al. (2015)³⁹ who concluded that the mutations D148E and S197E increased activity and thermostability of the phytase of *Bacillus amyloliquefaciens* DSM 1061. It is worthy to note that all newly introduced residues already existed or had their homologous ones inside PhyL amino acid sequence. Finally, amino acid sequence originality of PhyL gave it better physicochemical and kinetic properties, compared to phytases derived from bacterial, fungal and yeast species.⁴⁰⁻⁴⁹

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

- Cheryan M, Rackis JJ. Phytic acid interactions in food systems. *Crit Rev Food Sci Nutr* 1980; 13:297-335; PMID:7002470; <http://dx.doi.org/10.1080/10408398009527293>
- Boling SD, Douglas MW, Johnson ML, Wang X, Parsons CM, Koelkebeck KW, et al. The effects of dietary available phosphorus levels and phytase performance of young and older laying hens. *Poultry Science* 2000; 79:224-30; PMID:10735751; <http://dx.doi.org/10.1093/ps/79.2.224>
- Mitchell DB, Vogel K, Weimann BJ, Pasamontes L, van Loon APMG. The phytase subfamily of acid histidine phosphatases: isolation of genes for two novel phytases from the fungi *Aspergillus terreus* and *Myceliophthora thermophila*. *Microbiology* 1997; 143:245-52; PMID:9025298
- Vats P, Banerjee UC. Production studies and catalytic properties of phytases (myo-inositol hexakisphosphate phosphohydrolases): an overview. *Enz Microb Tech* 2004; 35:3-14; <http://dx.doi.org/10.1016/j.enzmictec.2004.03.010>
- Haefner S, Knietzsch A, Scholten E, Braun J, Lohscheidt M, Zelder O. Biotechnological production and applications of phytases. *Appl Microbiol Biotechnol* 2005; 68:588-97; PMID:16041577; <http://dx.doi.org/10.1007/s00253-005-0005-y>
- Kim OH, Kim YO, Shim JH, Jung YS, Jung WJ, Choi WC, Lee H, Lee SJ, Kim KK, Auh JH, Kim H, Kim JW, Oh TK, Oh BC. β -propeller phytase hydrolyzes insoluble Ca(2+)-phytate salts and completely abrogates the ability of phytate to chelate metal ions. *Biochemistry* 2010; 49:10216-27; PMID:20964370; <http://dx.doi.org/10.1021/bi1010249>
- Baruah K, Sahu NP, Pal AK, Deb Nath D, Yengkokpam S, Mukherjee SC. Interactions of dietary microbial phytase, citric acid and crude protein level on Mineral utilization by *Robu, Labeo rohita* (Hamilton), Juveniles. *J World Aquacul Soc* 2007; 38:238-49; <http://dx.doi.org/10.1111/j.1749-7345.2007.00092.x>
- Kumar V, Sinha AK, Makkar HPS, Becker K. Dietary roles of phytate and phytase in human nutrition: A review. *Food Chem* 2010; 120:945-59; <http://dx.doi.org/10.1016/j.foodchem.2009.11.052>
- Simell M, Turunen M, Pironen J, Vaara T. 1989. Feed and food applications of phytase. Lecture-3rd Meet. Industrial Applications of Enzymes, Barcelona (Spain).
- Haros M, Rosell CM, Benedetto C. Fungal phytase as a potential breadmaking additive. *Eur Food Res Tech* 2001; 213:317-22; <http://dx.doi.org/10.1007/s002170100396>
- Khare SK, Jha K, Gupta MN. Entrapment of wheat phytase in polyacrylamide gel and its application in soy milk phytate hydrolysis. *Biotechnol Appl Biochem* 1994; 19:193-98.
- Billington DC, 1993. The inositol phosphates: Chemical synthesis and biological significance. Weinham: Verlag Chemie.
- Quan CS, Fan SD, Ohta Y. Immobilization of *Candida krusei* cells producing phytase in alginate gel beads: an application of the preparation of myo-inositol phosphates. *Appl Microbiol Biotechnol* 2003; 62:41-7; PMID:12709834; <http://dx.doi.org/10.1007/s00253-003-1247-1>
- Siren M, 1986b. New myo-inositol triphosphoric acid isomer. *Pat SW* 052950.
- Laumen K, Ghisalba O. Preparative scale chemo enzymatic synthesis of optically pure D-myo-inositol 1-phosphate. *Biosci Biotech Biochem* 1994; 58:2046-49; <http://dx.doi.org/10.1271/bbb.58.2046>
- Kumar V, Sangwan P, Verma AK, Agrawal S. Molecular and biochemical characteristics of recombinant β -propeller phytase from *Bacillus licheniformis* strain PB-13 with potential application in aquafeed. *Appl Biochem Biotechnol* 2014; 173:646-59; PMID:24687556; <http://dx.doi.org/10.1007/s12010-014-0871-9>
- Ullah AHJ, Sethumadhavan K, Mullaney EJ, Zieglerhoffer T, Austin-Phillips S. Cloned and expressed fungal *phyA* gene in alfalfa produces a stable phytase. *Biochem Biophys Res Comm* 2002; 290:1343-48; <http://dx.doi.org/10.1006/bbrc.2002.6361>
- Li J, Hegemann CE, Hanlon RW, Lacy GH, Denbow DM, Grabau EA. Secretion of active recombinant phytase from soybean cell-suspension cultures. *Plant Physiol* 1997; 114:1103-11; PMID:9232886; <http://dx.doi.org/10.1104/pp.114.3.1103>
- Ullah AHJ, Sethumadhavan K, Mullaney EJ, Zieglerhoffer T, Austin-Phillips S. Fungal *phyA* gene expressed in potato leaves produces active and stable phytase. *Biochem Biophys Res Comm* 2003; 306:603-9; PMID:12804608; [http://dx.doi.org/10.1016/S0006-291X\(03\)01002-7](http://dx.doi.org/10.1016/S0006-291X(03)01002-7)
- Lucca P, Hurrell R, Potrykus I. Approaches to improve the bioavailability and level of iron in rice seeds. *J Sci Food Agric* 2001; 81:828-34.
- Brinch-Pedersen H, Olesen A, Rasmussen SK, Holm PB. Generation of transgenic wheat (*Triticum aestivum* L.) for constitutive accumulation of an *Aspergillus* phytase. *Mol Breed* 2000; 6:195-206; <http://dx.doi.org/10.1023/A:1009690730620>
- Mullaney EJ, Daly CB, Ullah AHJ. Advances in phytase research. *Adv Appl Microbiol* 2000; 47:157-99; PMID:12876797; [http://dx.doi.org/10.1016/S0065-2164\(00\)47004-8](http://dx.doi.org/10.1016/S0065-2164(00)47004-8)
- Farhat-Khemakhem A, Ali MB, Boukhris I, Khemakhem B, Maguin E, Bejar S, Chouayekh H. Crucial role of Pro 257 in the thermostability of *Bacillus* phytases: biochemical and structural investigation. *Int J Biol Macromol* 2013; 54:9-15; PMID:23178368; <http://dx.doi.org/10.1016/j.ijbiomac.2012.11.020>
- Borgi MA, Boudebouze S, Aghajari N, Szukala F, Pons N, Maguin E, Rhimi M. The attractive recombinant phytase from *Bacillus licheniformis*: biochemical and molecular characterization. *Appl Microbiol Biotechnol* 2014; 98:5937-47; PMID:24337251; <http://dx.doi.org/10.1007/s00253-013-5421-9>
- Keruvuo J, Tykkynen S. Expression of *Bacillus subtilis* phytase in *Lactobacillus plantarum* 755. *Lett Appl Microbiol* 2000; 30:325-9; PMID:10792656; <http://dx.doi.org/10.1046/j.1472-765x.2000.00660.x>
- Brnić M, Wegmüller R, Zeder C, Senti G, Hurrell RF. Influence of phytase, EDTA, and Polyphenols on Zinc Absorption in Adults from Porridges Fortified with Zinc Sulfate or Zinc Oxide. *J Nutr* 2014; 144:1467-73; PMID:24966411
- Vigors S, Sweeney T, O'Shea CJ, Browne JA, O'Doherty JV. Improvements in growth performance, bone mineral status and nutrient digestibility in pigs following the dietary inclusion of phytase accompanied by modifications in intestinal nutrient transporter gene expression. *Br J Nutr* 2014; 7:1-10; PMID:24998244
- Zeng ZK, Wang D, Piao XS, Li PF, Zhang HY, Shi CX, Yu SK. Effects of adding super DosePhytaseto the phosphorus-deficient diets of young pigs on growth performance, bone quality, minerals and amino acids digestibilities. *Asian-Australas J Anim Sci* 2014; 27:237-46; PMID:25049948; <http://dx.doi.org/10.5713/ajas.2013.13370>
- Park SC, Oh BC, Rhee MH, Jeong KS, Lee KW, Song JC, Oh TK. The enzyme activity of a novel phytase from *Bacillus amyloliquefaciens* DS11 and its potential use as a feed pellet. *J Gen Appl Microbiol* 2003; 49:129-33; PMID:12833216
- Sanz-Penella JM, Tamayo-Ramos JA, Sanz Y, Haros M. Phytate reduction in bran-enriched bread by phytase-producing bifidobacteria. *J Agric Food Chem* 2009; 57:10239-44; PMID:19817458; <http://dx.doi.org/10.1021/jf9023678>
- Haros M, Bielecka M, Honke J, Sanz Y. Myo-inositol-hexakisphosphate degradation by *Bifidobacterium infantis* ATCC 15697. *Int J Food Microbiol* 2007; 117:76-84; PMID:17462768; <http://dx.doi.org/10.1016/j.ijfoodmicro.2007.02.021>
- Myers M, Ahealy MJ, Oakeshort JG. Effects of the residue adjacent to the reactive serine on the substrate interactions of *Drosophila* esterase 6. *Biochem Genet* 1993; 31:259-78; PMID:8274134; <http://dx.doi.org/10.1007/BF00553170>
- Mantafounis D, Pitts J. Protein engineering of chymosin: modification of the optimum pH of enzyme catalysis. *Prot Eng* 3:605-9; PMID:2217134; <http://dx.doi.org/10.1093/protein/3.7.605>
- Sakoda H, Imanaka T. Cloning and sequencing of the gene coding for alcohol dehydrogenase of *Bacillus stearothermophilus* and rational shift of the optimum pH. *J Bacteriol* 1974; 139:7-402; PMID:1735726
- Rashid MH, Siddiqui KS. Carboxy-group modification: high temperature activation of charge-neutralized and charge-reversed-glucosidases from *Aspergillus niger*. *Biotechnol Appl Biochem* 1998; 27:231-7; PMID:9664679
- Siddiqui KS, Loviny-Anderton T, Rangarajan M, Hartley BS. Arthrobaacter D-xylose isomerase: chemical modification of carboxy groups and protein engineering

- of pH optimum. *Biochem J* 1993; 296:685-91; PMID:7904154
37. Tye AJ, Siu FKY, Leung TYC, Lim BL. Molecular cloning and the biochemical characterization of two novel phytases from *B. subtilis* 168 and *B. licheniformis*. *Appl Microbiol Biotechnol* 2002; 59:190-7; PMID:12111145; <http://dx.doi.org/10.1007/s00253-002-1033-5>
 38. Tran TT, Mamo G, Mattiasson B, Hatt-Kaul R. A thermostable phytase from *Bacillus* sp. MD2: cloning, expression and high-level production in *Escherichia coli*. *J Indust Microbiol Biotechnol* 2010; 37:279-87; PMID:19997958; <http://dx.doi.org/10.1007/s10295-009-0671-3>
 39. Xu W, Shao R, Wang Z, Yan X. Improving the Neutral Phytase Activity from *Bacillus amyloliquefaciens* DSM 1061 by Site-Directed Mutagenesis. *Appl Biochem Biotechnol* 2015; 175:3184-94; PMID:25613522; <http://dx.doi.org/10.1007/s12010-015-1495-4>
 40. Farhat A, Chouayekh H, Ben Farhat M, Bouchaala K, Bejar S. Gene Cloning and Characterization of a Thermostable Phytase from *Bacillus subtilis* US417 and Assessment of its Potential as a Feed Additive in Comparison with a Commercial Enzyme. *Mol Biotechnol* 2008; 40:127-35; PMID:18543132; <http://dx.doi.org/10.1007/s12033-008-9068-1>
 41. Kerovuo J, Lauraus M, Nurminen P, Kalkkinen N, Apajalahti J. Isolation, characterization, molecular gene cloning, and sequencing of a novel phytase from *Bacillus subtilis*. *Appl Environ Microbiol* 1998; 64:2079-85; PMID:9603817
 42. Gulati HK, Chadha BS, Saini HS. Production and characterization of thermostable alkaline phytase from *Bacillus laevolacticus* isolated from rhizosphere soil. *J Indust Microbiol Biotechnol* 2007; 34:91-8; PMID:16967265; <http://dx.doi.org/10.1007/s10295-006-0171-7>
 43. Choi YM, Suh HJ, Kim JM. Purification and Properties of Extracellular Phytase from *Bacillus* sp. KHU-10. *J Prot Chem* 2001; 20:287-92; PMID:11594462; <http://dx.doi.org/10.1023/A:1010945416862>
 44. Shi P, Huang H, Wang Y, Luo H, Wu B, Meng K, Yang P, Yao B. A novel phytase gene appA from *Buttiauxella* sp. GC21 isolated from grass carp intestine. *Aquaculture* 2008; 275:70-5; <http://dx.doi.org/10.1016/j.aquaculture.2008.01.021>
 45. Zang GQ, Dong XF, Wang ZH, Zhang Q, Wang HX, Tong JM. Purification, characterization, and cloning of a novel phytase with low pH optimum and strong proteolysis resistance from *Aspergillus ficuum* NTG-23. *Biores Tech* 2010; 101:4125-31; PMID:20144543; <http://dx.doi.org/10.1016/j.biortech.2010.01.001>
 46. Watanabe T, Ikeda H, Masaki K, Fujii T, Iefuji H. Cloning and characterization of a novel phytase from wastewater treatment yeast *Hansenula fabianii* J640 and expression in *Pichia pastoris*. *J Biosci Bioeng* 2009; 108:225-30; PMID:19664557; <http://dx.doi.org/10.1016/j.jbiosc.2009.03.021>
 47. Li X, Liu Z, Chi Z, Li J, Wang X. Molecular cloning, characterization, and expression of the phytase gene from marine yeast *Kodamaea ohmeri* BG3. *Mycol Res* 2009; 113:24-32; PMID:18672057; <http://dx.doi.org/10.1016/j.mycres.2008.07.003>
 48. In MJ, Seo SW, Oh NS. Fermentative production and application of acid phytase by *Saccaromyces cerevisiae* CY strain. *Afri J Biotechnol* 2008; 17:3115-20.
 49. Ragon M, Neugnot-Roux V, Chemardin P, Moulin G, Boze H. Molecular gene cloning and overexpression of the phytase from *Debaryomyces castellii* CBS 2923. *Prot Exp Purificat* 2008; 58:275-83; PMID:18242101; <http://dx.doi.org/10.1016/j.pep.2007.12.003>