



HAL
open science

A DEMETER-like DNA demethylase governs tomato fruit ripening

Ruie Liu, Alexandre How-Kit, Linda Stammitti, Emeline Teyssier, Dominique D. Rolin, Anne Mortain-Bertrand, Stephanie Halle, Mingchun Liu, Junhua Kong, Chaoqun Wu, et al.

► **To cite this version:**

Ruie Liu, Alexandre How-Kit, Linda Stammitti, Emeline Teyssier, Dominique D. Rolin, et al.. A DEMETER-like DNA demethylase governs tomato fruit ripening. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112 (34), pp.10804-9. 10.1073/pnas.1503362112 . hal-02635649

HAL Id: hal-02635649

<https://hal.inrae.fr/hal-02635649>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

A DEMETER-like DNA demethylase governs tomato fruit ripening

Ruie Liu^{a,1}, Alexandre How-Kit^{b,1}, Linda Stammitt^{a,1}, Emeline Teyssier^a, Dominique Rolin^a, Anne Mortain-Bertrand^a, Stefanie Halle^a, Mingchun Liu^{c,d}, Junhua Kong^e, Chaoqun Wu^e, Charlotte Degraeve-Guibault^a, Natalie H. Chapman^f, Mickael Maucourt^a, T. Charlie Hodgman^f, Jörg Tost^{b,g}, Mondher Bouzayen^{c,d}, Yiguo Hong^{e,h}, Graham B. Seymour^f, James J. Giovannoniⁱ, and Philippe Gallusci^{a,2}

^aLaboratory of Fruit Biology and Pathology, University of Bordeaux, Institut National de la Recherche Agronomique, CS20032 Villenave d'Ornon, France; ^bLaboratory for Functional Genomics, Fondation Jean Dausset-Centre d'Études du Polymorphisme Humain, F-75010 Paris, France; ^cLaboratoire de Génétique et Biotechnologie des Fruits, Université de Toulouse, Institut National Polytechnique-École Nationale Supérieure d'Agronomie de Toulouse, F-31326 Castanet-Tolosan, France; ^dLaboratoire de Génétique et Biotechnologie des Fruits, Institut National de la Recherche Agronomique, F-31326 Castanet-Tolosan, France; ^eResearch Center for Plant RNA Signaling, College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou 310036, China; ^fSchool of Biosciences, University of Nottingham, Loughborough, Leics LE12 5RD, United Kingdom; ^gLaboratory for Epigenetics and Environment, Centre National de Génotypage, Commissariat à l'Énergie Atomique-Institut de Génétique, 91000 Evry, France; ^hSchool of Life Science, University of Warwick, Coventry CV4 7AL, United Kingdom; and ⁱUnited States Department of Agriculture, Robert W. Holley Center and Boyce Thompson Institute for Plant Research, Cornell University, Ithaca, NY 14853

Edited by Robert L. Fischer, University of California, Berkeley, CA, and approved July 8, 2015 (received for review March 3, 2015)

In plants, genomic DNA methylation which contributes to development and stress responses can be actively removed by DEMETER-like DNA demethylases (DMLs). Indeed, in *Arabidopsis* DMLs are important for maternal imprinting and endosperm demethylation, but only a few studies demonstrate the developmental roles of active DNA demethylation conclusively in this plant. Here, we show a direct cause and effect relationship between active DNA demethylation mainly mediated by the tomato DML, SIDML2, and fruit ripening—an important developmental process unique to plants. RNAi *SIDML2* knockdown results in ripening inhibition via hypermethylation and repression of the expression of genes encoding ripening transcription factors and rate-limiting enzymes of key biochemical processes such as carotenoid synthesis. Our data demonstrate that active DNA demethylation is central to the control of ripening in tomato.

active DNA demethylation | DNA glycosylase lyase | epigenetic | tomato | fruit ripening

Genomic DNA methylation is a major epigenetic mark that is instrumental to many aspects of chromatin function, including gene expression, transposon silencing, or DNA recombination (1–4). In plants, DNA methylation can occur at cytosine both in symmetrical (CG or CHG) and nonsymmetrical (CHH) contexts and is controlled by three classes of DNA methyltransferases, namely, the DNA Methyltransferase 1, Chromomethylases, and the Domain Rearranged Methyltransferases (5–7). Indeed, in all organisms, cytosine methylation can be passively lost after DNA replication in the absence of methyltransferase activity (1). However, plants can also actively demethylate DNA via the action of DNA Glycosylase-Lyases, the so-called DEMETER-Like DNA demethylases (DMLs), that remove methylated cytosine, which is then replaced by a non-methylated cytosine (8–11). Initially identified as enzymes necessary for maternal imprinting in *Arabidopsis thaliana* (12), the role of DMLs has since been established in various processes such as limiting extensive DNA methylation at gene promoters (13), determining the global demethylation of seed endosperm (8, 14) and promoting plant responses to pathogens (15). Of note, *Arabidopsis ros1*, *dml2*, and *dml3* single, double, or triple mutants showed little or no developmental alterations (9, 16, 17), suggesting that active DNA demethylation is not critical for development in this species. However, as mentioned above, genomic DNA methylation is an important mechanism that influences gene expression, and methylation at promoters is known to inhibit gene transcription (5, 18). Hence, it is likely that the active removal of methylation marks is an important mechanism during plant development and plant cell fate reprogramming, leading to the hypomethylation of sites important for

DNA–protein interaction and gene expression, as already observed in human cells (19).

Indeed, accumulating evidence suggests that active DNA demethylation might play a greater role in controlling gene expression in tomato. In support of this idea, recent work describing the methylome dynamics in tomato fruit pericarp revealed substantial changes in the distribution of DNA methylation over the tomato genome during fruit development, and demethylation during ripening at specific promoters such as the *NON RIPENING (NOR)* and *COLORLESS NON RIPENING (CNR)* promoters (20, 21). This observation is consistent with previous studies indicating that genome cytosine methylation levels decrease by 30% in pericarp of fruits during ripening, although DNA replication is very limited at this stage (22).

Significance

This work shows that active DNA demethylation governs ripening, an important plant developmental process. Our work defines a molecular mechanism, which has until now been missing, to explain the correlation between genomic DNA demethylation and fruit ripening. It demonstrates a direct cause-and-effect relationship between active DNA demethylation and induction of gene expression in fruits. The importance of these findings goes far beyond understanding the developmental biology of ripening and provides an innovative strategy for its fine control through fine modulation of epimarks in the promoters of ripening related genes. Our results have significant application for plant breeding especially in species with limited available genetic variation.

Author contributions: T.C.H., Y.H., and P.G. designed research; R.L., A.H.-K., L.S., E.T., D.R., A.M.-B., S.H., M.L., J.K., C.W., C.D.-G., N.H.C., M.M., T.C.H., and P.G. performed research; A.H.-K. targeted bisulfite pyrosequencing; R.L., A.H.-K., E.T., D.R., A.M.-B., S.H., T.C.H., J.T., M.B., Y.H., G.B.S., J.J.G., and P.G. analyzed data; and E.T., G.B.S., J.J.G., and P.G. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Data deposition: Full array datasets of WT (Ailsa Craig) and of the *rin*, *nor*, and *Cnr* mutants have been deposited with the Sol Genomics Network and are accessible under the following link: <ftp://ftp.solgenomics.net/microarray/>, file name *rinorcnr* microarrays Liuetal PNAS2015. These array datasets have been used to analyze the expression of the four DML genes in the tomato *rin*, *nor*, and *Cnr* mutants and in WT fruits. The results are shown in Fig. 6B and Dataset S1.

¹R.L., A.H.K., and L.S. contributed equally to this work.

²To whom correspondence should be addressed. Email: philippe.gallusci@bordeaux.inra.fr.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1503362112/-DCSupplemental.

Here, we investigated active DNA demethylation as a possible mechanism governing the reprogramming of gene expression in fruit pericarp cells at the onset of fruit ripening.

Results

The Tomato Genome Contains Four DNA Glycosylase Genes with Specific Expression Patterns. The tomato genome contains four putative *DML* genes encoding proteins with characteristic domains of functional DNA glycosylase-lyases (23) (*SI Appendix, Fig. S1 A and C and Table S1*). *SIDML1* and -2 are orthologous to the *Arabidopsis AtROS1* (Repressor of Silencing 1) gene and *SIDML3* to *AtDME* (DEMETER), whereas *SIDML4* has no closely related *Arabidopsis* ortholog (*SI Appendix, Fig. S1B*). All four *SIDML* genes are ubiquitously expressed in tomato plants, although *SIDML4* is expressed at a very low level in all organs analyzed. In leaves, flowers, and young developing fruits, the four genes present coordinated expression patterns characterized by high expression levels in young organs that decrease when organs develop. However, unlike *SIDML1*, *SIDML3*, and *SIDML4*, which are barely expressed during fruit ripening, *SIDML2* mRNA abundance increases dramatically in ripening fruits, suggesting an important function at this developmental phase (Fig. 1).

Transgenic Plants with Reduced *DML* Gene Expression Present Various Fruit and Plant Phenotypes. The physiological significance of tomato *DMLs* was addressed through RNAi-mediated gene repression using the highly conserved Helix-hairpin-Helix-Gly/Pro rich domain (HhH-GPD) specific to *DML* proteins as a target sequence (*SI Appendix, Fig. S2A*). Our goal was to repress simultaneously all tomato *SIDML* genes, anticipating potential functional redundancy among these four genes; 23 independent T0 transgenic lines were generated and 22 showed alterations of fruit development, including delayed ripening, modified fruit shape, altered color, shiny appearance, parthenocarp, or combinations of these phenotypes (Fig. 2A).

Lines 2 and 8, which showed delayed and inhibited ripening phenotypes, were chosen to investigate the possible link between ripening and DNA demethylation. In both cases, 10–25 T1 and T2 plants were grown that showed maintenance and strengthening of the nonripening phenotypes in subsequent generations coincident with the presence of the transgene. The loss of the RNAi transgene in segregating lines led to reversion to a wild-




Fig. 1. Differential expression of *SIDML* genes in tomato organs. Absolute quantification of *SIDML1*, *SIDML2*, *SIDML3*, and *SIDML4* mRNA; *SIDML4* gene expression is presented in a separate diagram because of its very low expression level. Fruit pericarp is at 5, 10, 20 dpa and at Breaker (BR, 39 dpa), orange (O), and red ripe (RR). Asterisks indicate significant difference [Student's *t* test ($n = 3$)] between *SIDML2* and all other *SIDML* genes: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. Error bars indicate means \pm SD. Ap, stem apex; CF, closed flowers; L, leaves at positions 3, 4, 5, 8, 10, 16, and 20 from apex; OP, open flowers 5, 10, and 20; R, roots; S, stem from whole seedlings.




Fig. 2. Phenotypes of tomato *DML* RNAi fruits. (A) Fruits (70 dpa) (upper lane) or fruit sections (lower lane) from eight independent representative T0 RNAi plants. (B) Fruits (85 dpa) from T2 plants (left to right); WT plants, line 2 plants (DML2A and DML2B), line 8 plants (DML8A and DML8B), and an azygous plant (AZ). (C) Ripening kinetics of WT (Top), DML8A (Middle), and DML2A (Bottom). (D) WT bicarpel (Upper) DML2B multicarpel fruits (Lower). (E) VIGS experiment on 47-dpa (Br + 5) fruits injected with PVX/SIDML2 [fruits (1) and (3)] or PVX [fruits (2) and (4)] at 12 dpa [fruits (3) and (4)] inside of fruits (1) and (2), respectively. (Scale bars: 1 cm.)

type (WT) phenotype, indicating a lack of memory effect across generations when fruit ripening is considered (Fig. 2A and B and *SI Appendix, Fig. S3A*). In plants of both RNAi lines, analysis of *SIDML* gene residual expression in 20 days postanthesis (dpa) fruits indicates that only *SIDML1* and *SIDML2* are repressed to 40–60% of the WT level, whereas *SIDML3* and *SIDML4* are either unaffected or induced compared with WT (Fig. 3A). This is most likely attributable to the lower homology level of these two genes, with *SIDML1* in the part of the gene used for the RNAi construct (*SI Appendix, Fig. S2A*). During ripening, *SIDML2* expression is reduced to 10% of WT at the Breaker (Br) stage and remains low at 55 dpa (Br + 16) but increases slightly at 70 dpa (Br + 31) (Fig. 3B and *SI Appendix, Fig. S2B*), coincident with the partial ripening observed in transgenic RNAi fruits (Fig. 2C and *SI Appendix, Fig. S3B*). Whether the increase in *SIDML2* expression at late ripening stages is attributable to a weaker effect of the RNAi remains unclear. None of the three remaining genes, *SIDML1*, *SIDML3*, and *SIDML4*, which are weakly expressed during ripening, displayed significantly reduced




Fig. 3. Residual expression of *SIDML* genes in fruits of transgenic *DML* RNAi plants. Normalized expression of the *SIDML* genes (A) in 20-dpa transgenic fruits of plants from line 2 (DML2A and -2B), line 8 (DML8A and -8B), an azygous plant (AZ), and the respective WT1 and WT2 controls (B) in WT2 and DML8A fruits at seven developmental stages. Expression of the *SIDML* genes was normalized to EF1 α and to the corresponding WT fruits at 20 dpa. For each *SIDML* gene, asterisks indicate significant difference [Student's *t* test ($n = 3$)] between transgenic plants and WT controls, respectively, at 20 dpa (A) or at the same age during fruit development (B). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. Error bars indicate mean \pm SD.

expression compared with WT fruits of the same age, indicating that observed ripening phenotypes are likely attributable to *SIDML2* gene repression. This hypothesis was further confirmed using virus induced gene silencing (VIGS) to specifically repress the *SIDML2* gene; 17.5% of the fruits injected with a PVX/*SIDML2* vector presented non ripening sectors contrary to those injected with a control PVX virus that all ripened normally (Fig. 2E and *SI Appendix*, Fig. S4B). Indeed, *SIDML2* was down-regulated in nonripening sectors of fruits injected with the PVX/*SIDML2* vector, whereas none of the three other *SIDML* genes was repressed (*SI Appendix*, Fig. S4C), demonstrating that the specific knock down of *SIDML2* is sufficient to inhibit ripening.

It was noteworthy that some plants from line 2 developed additional phenotypes affecting plant growth, leaf shape, flower development, and fruit carpal number that were not observed in T0 and T1 generations (Fig. 2D and *SI Appendix*, Fig. S3B and C). The screening of additional lines revealed other independent transgenic lines that presented flower, fruit, and plant phenotypes similar to line 2 (*SI Appendix*, Fig. S3D). These observations indicate that the severity of the phenotypes increases over generations and suggest that DMLs may also be involved in other aspects of tomato plant development beyond fruit ripening.

All Aspects of Fruit Ripening Are Delayed and Limited in RNAi Transgenic Lines. Fruits of transgenic lines 2 and 8 were further analyzed to investigate the consequences of DNA demethylation on the ripening process. Indeed, in fruits of both transgenic lines, the onset of fruit ripening was delayed from 10 to 20 d compared with WT or Azygous revertant fruits, and ripening of transgenic fruits was never completed even after 45 d or longer maturation times (Fig. 2B and C and *SI Appendix*, Fig. S3B). The ripening defect is further demonstrated by the late and extremely reduced total carotenoids and lycopene accumulation and the delayed

chlorophyll degradation (Fig. 4A). Primary metabolite composition was also modified, as visualized by principal component analysis (PCA) using the absolute concentration of 31 primary metabolites issued from $^1\text{H-NMR}$ analysis (Fig. 4B and *SI Appendix*, Fig. S5A). The first two principal components (PCs), explain more than 54% of total variability. During early development (20, 35, and 39 dpa), WT and transgenic samples follow parallel trajectories as highlighted by the PCA in which the




Fig. 4. Metabolic profiling of carotenoids and primary metabolites in transgenic *DML* RNAi fruits. (A) Chlorophylls (Top), total carotenoids (Middle), and lycopene (Bottom) content. Asterisks indicate significant difference [Student's *t* test ($n = 3$)] between DML2A and -2B, DML8A and -8B, and WT1 and WT2, respectively, at the same age: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. Error bars indicate means \pm SD. (B) PCA using primary metabolites in WT2 (Δ) and DML8A (\circ) fruits at seven developmental stages. Color indicates the fruit developmental stages: white is 20 dpa and from light gray to black are 35, 39 (Br), 55, 70, 85, and 110 dpa.

second PC (PC2) explains 21% of the total variability. However, at 55-dpa and later ripening stages, PC1, which accounts for 33.67% of the global variability, separates WT fruits from all other samples. Hence, WT fruit samples harvested at 55-dpa and older stages are clearly distinct from transgenic fruit samples of the same age. Metabolic differences between ripening WT and transgenic fruits are mainly attributable to overaccumulation of malate and reduction or delayed accumulation of compounds typical of ripening fruits, including glucose, fructose, glutamate, rhamnose, and galactose (SI Appendix, Fig. S5 B–D). Climacteric rise of ethylene production was also dramatically reduced in fruits of both *DML* RNAi lines, although low ethylene accumulation occurred to a degree and timing consistent with the late and limited ripening process of RNAi fruits (SI Appendix, Fig. S6).

Fruit-Ripening Defects Are Correlated with the Repression and Hypermethylation of Genes Necessary for This Developmental Process.

To demonstrate a causal relationship between fruit ripening defects of transgenic lines and the impairment of active DNA demethylation, the expression of *CNR* (21), *RIPENING INHIBITOR (RIN)* (24), *NOR* (25), and *PHYTOENE SYNTHASE 1 (PSY1)* (26, 27) genes was assessed in RNAi transgenic plants. These genes were

selected among others because they are necessary for the overall ripening process (*CNR*, *RIN*, *NOR*), or specifically govern carotenoid accumulation (*PSY1*), an important quality trait of mature tomato fruit. Moreover, the promoter regions of these genes showed reduced methylation levels during fruit ripening in WT tomato (20, 21). It is noteworthy that *CNR* gene induction was delayed 15 d in transgenic fruits, and all three other genes showed a dramatic reduction in expression level consistent with the ripening defect of the transgenic lines (Fig. 5A and SI Appendix, Fig. S7). To assess whether repression of *CNR*, *RIN*, *NOR*, and *PSY1* gene expression in ripening fruits results from the maintenance of a high cytosine methylation status of their promoter upon down-regulation of *SIDML2*, methylsensitive-PCR (McrBC-PCR) analysis of the corresponding promoters was performed. This approach revealed a ripening-associated demethylation of the *RIN*, *NOR*, and *PSY1* promoters in WT and Azygous revertant fruits but not in *SIDML* RNAi fruits (Fig. 5B). No detectable variations of methylation in the *CNR* promoter during ripening of WT fruits were revealed with this method. The putative differentially methylated regions (DMRs) in the *NOR* and *PSY1* promoter regions were subsequently analyzed by gene specific bisulfite pyrosequencing (28). Methylation analysis of the *CNR* promoter was targeted to a region known to be methylated at all stages (*CNR1*) (SI Appendix, Fig. S9C), used here as a control for methylation and to a previously identified DMR (*CNR2*) (SI Appendix, Fig. S9C) (20, 21). For all three promoters, cytosines that became demethylated in ripening WT fruits but not in transgenic fruits of the same age were identified (Fig. 6A and SI Appendix, Fig. S9). Two distinct situations were observed: (i) sequences corresponding to putative RIN binding sites (RIN BS) in the *CNR* and *NOR* promoters (20), where methylation is high at 20 and 35 dpa in all plants analyzed and drops to very low levels during ripening of WT fruits but is maintained to high levels in RNAi fruits of the same age; and (ii) sequences that are hypermethylated in transgenic fruits at all stages analyzed compared with WT fruits. These latter sequences include a newly identified DMR in the *PSY1* promoter and cytosines upstream and downstream to the RIN BS in the *NOR* and *CNR* promoters. These data demonstrate the absolute requirement of promoter demethylation in critical genes for ripening to occur. The data also suggest multiple patterns of cytosine demethylation occurring either specifically during ripening or at earlier stages.




Fig. 5. Expression and demethylation at key genes controlling ripening are inhibited in DML RNAi plants. (A) Expression of the *RIN*, *NOR*, *CNR*, and *PSY1* genes in transgenic DML8A and WT fruits normalized to EF1 α and to WT fruits at 20 dpa. Asterisks indicate significant difference [Student's *t* test ($n = 3$)] between WT and DML8A samples at a given stage: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. Error bars indicate means \pm SD. (B) McrBC-PCR analysis of selected promoter fragments in fruits of WT, azygous (Azy), and DML8A plants; 1 μ g of genomic DNA was digested with McrBC (NEB) during 5h (+); (-) indicates negative control for the digestion reaction that was performed without GTP. In the WT and azygous plants, the part of *NOR*, *RIN*, and *PSY1* promoter regions analyzed are methylated at 35 dpa (no amplification) but are demethylated at 55 dpa (amplification). In DML8A plants, the three promoter regions behave similarly to WT at 35 dpa but remained methylated at 55 dpa (no amplification in both cases). The pectin-methyl esterase (*PME*) promoter is used as an unmethylated control, and the *CNR* promoter fragment used here was found to be sufficiently methylated at all stages for complete digestion by McrBC.

Discussion

Previously reported analysis of DNA cytosine methylation and RIN binding during fruit development in WT and in the *rin* and *Cnr* tomato-ripening mutants suggested a significant role for DNA methylation during ripening and a feedback loop between methylation and ripening transcription factors (20, 21, 29). Here, we demonstrate for the first time to our knowledge that active DNA demethylation is an absolute requirement for fruit ripening to occur and show a direct cause and effect relationship between hypermethylation at specific promoters and repression of gene expression. In this context, *SIDML2* appears to be the main regulator of the ripening associated DNA demethylation process. (i) *SIDML2* is the only *SIDML* gene induced concomitantly to the demethylation and induction of genes that control fruit ripening; (ii) the specific knockdown of *SIDML2* in VIGS-treated fruits leads to inhibition of fruit ripening similar to DML-RNAi fruits; and (iii) the hypermethylated phenotype described in the *Cnr* and *rin* mutants (20) is associated with the specific repression of *SIDML2*, with none of the other *SIDML* genes being down-regulated (Fig. 6B and Dataset S1).

Indeed, we cannot formally rule out that *SIDML1*, which is repressed in the transgenic RNAi lines, also participates in the genomic DNA demethylation in fruits. However, *SIDML1* is mainly expressed at early stages of fruit development and only at very low levels during fruit ripening. Hence, this protein may also




Fig. 6. Bisulfite-sequencing analysis at the *NOR*, *CNR*, and *PSY1* promoter fragments in WT and transgenic DML RNAi plants. (A) Heat-map representation of DNA methylation at selected *NOR*, *CNR*, and *PSY1* promoter regions (*SI Appendix*, Fig. S8) in fruits of control (WT1 and WT2) and transgenic (DML2A, -2B, -8A, and -8B) plants at five (WT and line 8) or four (line 2) developmental stages. For each promoter, two fragments were analyzed (fragment 1, gray box; fragment 2, black box), the positions of which are shown in *SI Appendix*, Fig. S8 and Fig. S9. The position of the Cs within each promoter fragment is also shown (number in the columns on the right side), as defined in *SI Appendix*, Fig. S8. For each promoter, Cs have been clustered considering the two PCR fragments analyzed together. (B) Changes in expression of *SIDML* genes in fruits of *Ailsa Craig* (WT) and near-isogenic mutant lines *rin*, *Cnr*, and *nor*, as determined by microarrays analysis. For fruit development, days postanthesis are shown. Mature green is 40 dpa in *Ailsa Craig* and then Br is 49 dpa. For nonripening mutants, Br onward are 49 dpa + 1–7 d. Asterisks indicate significant difference (variance ratio, *F* tests) between WT and mutant lines for the *SIDML2* gene only to avoid overloading the figure: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). Details of expression results and statistical analyses for all four genes are provided in *Dataset S1*. Error bars indicate means \pm SD. (C) Proposed function of DNA demethylation in the control of fruit ripening; *SIDML2* is necessary for the active demethylation of the *NOR*, *CNR*, *RIN*, and *PSY1* promoter region, thereby allowing these gene expressions. *SIDML2* gene expression is reduced in the *rin*, *nor*, and *Cnr* background, suggesting a regulatory loop. There is at this time no evidence of direct regulation of the *SIDML2* gene by the *RIN*, *NOR*, or *CNR* protein. *SIDML2* may control the expression of additional ripening induced gene, as shown in this study for the *PSY1* gene and suggested by the demethylation of several promoters during fruit ripening (20). Arrows indicate activation. Lines indicate repression: black, direct effects; gray, direct or indirect effects.

be involved in demethylation events but mainly those occurring at the early stages of fruit development.

In addition to genes encoding major fruit ripening regulators, those encoding enzymes involved in various aspects of fruit ripening are also likely to be demethylated, as suggested by the observation that *PSY1* gene expression also requires demethylation. Combined transcriptomic, methylomic, and metabolomic analysis of the transgenic lines described here will now be required to determine the network of genes and metabolic processes primarily targeted by demethylation in tomato fruit.

SIDML2 is the likely focal point of a feedback regulation on ripening-associated DNA demethylation, because this gene is

clearly down-regulated in fruits of the *rin*, *nor*, and *Cnr* mutants, contrary to the other *SIDML* genes that are normally expressed (Fig. 6 B and C and *Dataset S1*). It is plausible that timing and extent of demethylation may represent an important source of variation in the diversity of kinetics and intensity of ripening found among tomato varieties, thus presenting a frontier for further investigation. Controlling the timing and kinetics of active DNA demethylation in fruits may therefore provide new strategies to enhance fruit shelf life. In addition, engineering DNA demethylation in tomato fruits would be an innovative and novel strategy for the improvement of traits of agronomical relevance in a species with little genetic diversity (30). Finally,

the recent demonstration that hypermethylation of a *Myb* promoter blocks anthocyanin accumulation during pear and apple ripening (31, 32) supports the notion of a more general role for demethylation in fruits. However, whether this mechanism occurs similarly during the ripening of all fleshy fruit species now requires further investigation.

Materials Methods

Plant Material and Experimental Plan. All experiments were performed using a cherry tomato variety (*Solanum lycopersicum*, cv *WVA106*) that was grown in greenhouse conditions, except for VIGS experiments, which were performed on *Solanum lycopersicum*, cv *Ailsa Craig* grown in growth chambers as described (21). For the array experiments, fruit pericarp of *Ailsa Craig* and near-isogenic mutants *rin*, *nor*, and *Cnr* were collected at 13 stages of fruit development and ripening with three independent biological replicates per line and immediately frozen in liquid nitrogen for RNA extraction and array analysis. Details of tomato transformation, selection of line 2 and 8 used in this study, and of VIGS experiments are provided in *SI Appendix, Materials and Methods*.

For all analysis, two independent transgenic T2 plants (DML2A and -B and DML8A and -B for lines 2 and 8, respectively) and an azygous plant obtained from line 8 were used. Additional T2 plants were eventually used as controls for the phenotypes of these four plants. T2 plants from line 2 presented dramatic alterations of flower development, not visible in previous generations, and were backcrossed to allow fruit development. This resulted in a limited number of fruits (see below). For this reason, not all developmental stages could be analyzed for this line.

The experimental plan was designed to span tomato fruit development and ripening in cv *West Virginia 106* (WVA106) and transgenic DML RNAi plants over a period of 85 d from fruit set to account for the strongly delayed ripening phenotype of the transgenic fruits. At stages following mature green, the DML RNAi fruits diverge from the WT, because they are significantly delayed in ripening induction and almost completely ripening inhibited. Because it was not possible to select stages equivalent to the Br (39 dpa) or red ripe stages in the transgenic lines, we have chosen to analyze fruits identically staged, which allows comparing changes in the context of a developmental parameter (days postanthesis) that can be precisely mea-

sured. Two independent cultures were performed. (i) Plants from line 2 and the relevant WT control (WT1), fruits were harvested at 20, 35, 55 (Br + 16), 70 (Br + 31), and 85 (Br + 46) dpa. Because the fruit yield was reduced in line 2, a sufficient number of fruits at the Br stage could not be harvested and older fruits were preferentially selected to allow the analysis of late effects of demethylation inhibition. (ii) Line 8 was grown together with its own WT control (WT2) and an azygous plant. Because there were more fruits available for this line, the Br stage (39 dpa) was harvested in addition of the stages used for line 2.

For all fruit samples, two individual T2 plants were used, and for each sample, a minimum of six fruits separated in three biological replicates were processed and stored at -80°C until used.

Molecular and Metabolite Analysis. Details of molecular (gene expression, microarrays, MCR-PCR analysis of gene DNA methylation, and gene-targeted bisulfite sequencing) and metabolite (Carotenoid, ethylene, and $^1\text{H-NMR}$) analysis are provided in *SI Appendix, Materials and Methods*.

ACKNOWLEDGMENTS. We thank Marie Mirouze for critical reading of the manuscript and Antoine Daunay and Nicolas Mazaleyrat for technical support in bisulfite pyrosequencing analysis. We acknowledge Syngenta and specifically Dr. Charles Baxter for help with the tomato GeneChip studies and Alex Marshall for help with the array analysis, as well as Cécile Cabasson and Jim Craigon for help in statistical analysis. Metabolomic profiling was performed on the Metabolome Facility of Bordeaux Functional Genomics Center and supported by the French National Infrastructure for Metabolomics and Fluxomics (MetaboHUB) funded by the Agence Nationale de Recherche (ANR, Project ANR-11-INBS-0010). R.L. was the recipient of a grant from the Chinese Scholarship Council and P.G. of a Fulbright grant. G.B.S., T.C.H., and N.H.C. acknowledge financial support from the Biotechnology and Biological Sciences Research Council, UK (Grants BB/F005458/1 and BB/J015598/1), and G.B.S. acknowledges support from the European Cooperation in Science and Technology (COST) Action FA1106. Y.H., J.K., and C.W. were supported by Hangzhou Normal University and the National Natural Science Foundation of China (NFSC, Grant 31370180). M.B. and M.L. received support from the Laboratoire d'Excellence (LABEX) entitled "Towards a Unified theory of biotic Interactions; the role of environmental Perturbations" (TULIP, ANR-10-LABX-41) and from the networking activities within the European COST Action FA1106. J.J.G. was supported by National Science Foundation Grant IOS-1322714.

- Law JA, Jacobsen SE (2010) Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nat Rev Genet* 11(3):204–220.
- Saze H, Tsugane K, Kanno T, Nishimura T (2012) DNA methylation in plants: Relationship to small RNAs and histone modifications, and functions in transposon inactivation. *Plant Cell Physiol* 53(5):766–784.
- Chan SW, Henderson IR, Jacobsen SE (2005) Gardening the genome: DNA methylation in *Arabidopsis thaliana*. *Nat Rev Genet* 6(5):351–360.
- Mirouze M, et al. (2012) Loss of DNA methylation affects the recombination landscape in *Arabidopsis*. *Proc Natl Acad Sci USA* 109(15):5880–5885.
- Zhang M, Kimatu JN, Xu K, Liu B (2010) DNA cytosine methylation in plant development. *J Genet Genomics* 37(1):1–12.
- Bender J (2004) DNA methylation and epigenetics. *Annu Rev Plant Biol* 55(1):41–68.
- Finnegan EJ, Kovac KA (2000) Plant DNA methyltransferases. *Plant Mol Biol* 43(2–3):189–201.
- Hsieh T-F, et al. (2009) Genome-wide demethylation of *Arabidopsis* endosperm. *Science* 324(5933):1451–1454.
- Gong Z, et al. (2002) ROS1, a repressor of transcriptional gene silencing in *Arabidopsis*, encodes a DNA glycosylase/lyase. *Cell* 111(6):803–814.
- Zhu J-K (2009) Active DNA demethylation mediated by DNA glycosylases. *Annu Rev Genet* 43(1):143–166.
- Wu SC, Zhang Y (2010) Active DNA demethylation: Many roads lead to Rome. *Nat Rev Mol Cell Biol* 11(9):607–620.
- Choi Y, et al. (2002) DEMETER, a DNA glycosylase domain protein, is required for endosperm gene imprinting and seed viability in *Arabidopsis*. *Cell* 110(1):33–42.
- Zhu J, Kapoor A, Sridhar VV, Agius F, Zhu J-K (2007) The DNA glycosylase/lyase ROS1 functions in pruning DNA methylation patterns in *Arabidopsis*. *Curr Biol* 17(1):54–59.
- Gehring M, Bubb KL, Henikoff S (2009) Extensive demethylation of repetitive elements during seed development underlies gene imprinting. *Science* 324(5933):1447–1451.
- Yu A, et al. (2013) Dynamics and biological relevance of DNA demethylation in *Arabidopsis* antibacterial defense. *Proc Natl Acad Sci USA* 110(6):2389–2394.
- Yamamoto C, et al. (2014) Overproduction of stomatal lineage cells in *Arabidopsis* mutants defective in active DNA demethylation. *Nat Commun* 5:4062.
- Penterman J, et al. (2007) DNA demethylation in the *Arabidopsis* genome. *Proc Natl Acad Sci USA* 104(16):6752–6757.
- Gehring M, Henikoff S (2007) DNA methylation dynamics in plant genomes. *Biochim Biophys Acta* 1769(5–6):276–286.
- Bhutani N, Burns DM, Blau HM (2011) DNA Demethylation Dynamics. *Cell* 146(6):866–872.
- Zhong S, et al. (2013) Single-base resolution methylomes of tomato fruit development reveal epigenome modifications associated with ripening. *Nat Biotechnol* 31(2):154–159.
- Manning K, et al. (2006) A naturally occurring epigenetic mutation in a gene encoding an SBP-box transcription factor inhibits tomato fruit ripening. *Nat Genet* 38(8):948–952.
- Teyssier E, et al. (2008) Tissue dependent variations of DNA methylation and endoreplication levels during tomato fruit development and ripening. *Planta* 228(3):391–399.
- Mok YG, et al. (2010) Domain structure of the DEMETER 5-methylcytosine DNA glycosylase. *Proc Natl Acad Sci USA* 107(45):19225–19230.
- Vrebalov J, et al. (2002) A MAD5-box gene necessary for fruit ripening at the tomato ripening-inhibitor (*rin*) locus. *Science* 296(5566):343–346.
- Giovannoni JJ (2004) Genetic regulation of fruit development and ripening. *Plant Cell* 16(Suppl):S170–S180.
- Bartley GE, Viitanen PV, Bacot KO, Scolnik PA (1992) A tomato gene expressed during fruit ripening encodes an enzyme of the carotenoid biosynthesis pathway. *J Biol Chem* 267(8):5036–5039.
- Ray J, et al. (1992) Cloning and characterization of a gene involved in phytoene synthesis from tomato. *Plant Mol Biol* 19(3):401–404.
- How-Kit A, et al. (2015) Accurate CpG and non-CpG cytosine methylation analysis by high-throughput locus-specific pyrosequencing in plants. *Plant Mol Biol* 88(4–5):471–485.
- Chen W, et al. (2015) Requirement of *CHROMOMETHYLASE3* for somatic inheritance of the spontaneous tomato epimutation *Colourless non-ripening*. *Sci Rep* 5:9192.
- Lin T, et al. (2014) Genomic analyses provide insights into the history of tomato breeding. *Nat Genet* 46(11):1220–1226.
- Telias A, et al. (2011) Apple skin patterning is associated with differential expression of MYB10. *BMC Plant Biol* 11(1):93.
- Wang Z, et al. (2013) The methylation of the PcMYB10 promoter is associated with green-skinned sport in Max Red Bartlett pear. *Plant Physiol* 162(2):885–896.