



**HAL**  
open science

## Plant hormones: a fungal point of view

Emilie Chanclud, Jean-Benoit J.-B. Morel

► **To cite this version:**

Emilie Chanclud, Jean-Benoit J.-B. Morel. Plant hormones: a fungal point of view. *Molecular Plant Pathology*, 2016, 17 (8), pp.1289-1297. 10.1111/mpp.12393 . hal-02636076

**HAL Id: hal-02636076**

**<https://hal.inrae.fr/hal-02636076>**

Submitted on 27 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Plant hormones: a fungal point of view

Emilie CHANCLUD<sup>1</sup> and Jean-Benoit MOREL<sup>2</sup>

<sup>1</sup> Université Montpellier, UMR BGPI INRA/CIRAD/SupAgro, 34398 Montpellier, France

<sup>2</sup> INRA, UMR BGPI INRA/CIRAD/SupAgro, 34398 Montpellier, France

Corresponding author:

Jean-Benoit MOREL, [jbmorel@cirad.fr](mailto:jbmorel@cirad.fr)

INRA, UMR BGPI INRA/CIRAD/SupAgro, 34398 Montpellier, France

Phone: +33 499 624 837; Fax: +33 499 624 822

Running title: Hormones from fungi

Keywords: Plant hormones, fungi, virulence

Abstract 147

Introduction 383

Main text 420

Conclusions 404

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: [10.1111/mpp.12393](https://doi.org/10.1111/mpp.12393).

## Abstract

Most classical plant hormones are also produced by pathogenic and symbiotic fungi. The way these molecules favor the invasion of plant tissues and the development of fungi inside plant tissues is still largely unknown. In this review, we examine the different roles of such hormone production by pathogenic fungi. Converging evidence suggest that these fungal-derived molecules have potentially two modes of action: (i) they may perturb plant processes, either positively or negatively, to favor invasion and nutrient uptake and (ii) they may also act as signals for the fungi themselves to engage appropriate developmental and physiological processes adapted to their environment. Indirect evidences suggest that abscisic acid, gibberellic acid and ethylene produced by fungi participate to pathogenicity. There is now evidence that auxin and cytokinins could be positive regulators required for virulence. Further research should establish whether or not fungal-derived hormones act like other fungal effectors.

## Introduction

Many fungi interact with plants in a beneficial manner as in mycorrhizal symbiosis (Sanders, 2011) or in an harmful manner in the case of fungal diseases (Dean et al., 2012). In order to obtain nutrients, both symbiotic and most pathogenic fungi penetrate their host without breaking the plant cell plasma membrane. The fungal membrane is protected by a cell-wall composed of chitin that can be recognized by plants through membrane receptors which then activate basal immunity. Chitin perception modulates responses during both mutualistic and pathogenic fungi-plant interactions (Gust et al., 2012). Fungi have evolved a repertoire of tools such as protein effectors and metabolites to impede such plant immunity and/or to establish favorable conditions for their establishment in plant tissues (Kamoun, 2007).

Besides the production of canonical effectors, fungi also produce compounds that are similar to plant hormones like auxins, cytokinins (CKs), gibberellic acids (GAs), ethylene (ET), abscisic acid (ABA), jasmonates (JA) and salicylates (SA). These hormones are well described to control plant development and to trigger important plant signaling events during biotic and abiotic stresses (reviewed in (Spence and Bais, 2015; Pozo et al., 2015; De Vleeschauwer et al., 2013; Robert-Seilaniantz et al., 2011; Peleg and Blumwald, 2011)).

There are now many examples showing that some pathogen protein effectors trigger hormone regulation to favor infection (Robert-Seilaniantz et al., 2007). By contrast, the involvement of hormonal compounds derived from microorganisms in plant-fungus interactions is poorly documented. Fungal-derived hormones were first suspected to be involved in the virulence of gall-forming pathogens (Robert-Seilaniantz et al., 2007; Denancé et al., 2013). For symbiotic fungi, such production of hormones is consistent with root modifications often required in these interactions (Hirsch et al., 1997). However, many pathogens that do not induce organ deformations can also produce and secrete plant hormones, suggesting a role of these molecules in other biological processes than organ deformation.

The role of plant-derived hormones in plant disease resistance has been extensively reviewed (Robert-Seilaniantz et al., 2011; De Vleeschauwer et al., 2014). In this review, we summarize the current knowledge on the role of fungal-derived plant hormones in plant-pathogen interactions with a focus on their putative role in virulence. When relevant, some information on plant-mycorrhiza interactions is also provided as it often sheds some light on the role of these molecules in plant-fungus interactions.

### **Auxins from fungi play a positive role in plant-fungus interactions**

Auxins are indole-derived hormones involved in plant development processes such as cell division differentiation and organ formation (Oka et al., 1999; Vanneste, 2005; Benjamins and Scheres, 2008) and senescence (Kim et al., 2011). Auxins also control biotic and abiotic stress responses in plants (Peleg and Blumwald, 2011). In plants, auxins is synthesized from tryptophan which is converted into indole-3-acetamide by tryptophan-2-monooxygenase enzymes (Zhao, 2010). Indole-3-acetamide is hydrolyzed to form indole-3-acetic acid (IAA) which is the major auxin active form in plants. These genes were also identified in fungi, e.g. *Fusarium* sp., and were confirmed for being involved in fungal auxin production (Tsavkelova et al., 2012). However, several auxin synthesis pathways were described in fungi. In some of them, like *Fusarium* sp. and *Colletotrichum gloeosporioides* (Tsavkelova et al., 2012; Gruen, 1959; Robinson et al., 1998), auxins is synthesized from the same precursor than in plants (indole-3-acetamide) but as observed in other fungal genus, for instance *Ustilago* (Reineke et al., 2008) and *Rhizoctonia* (Furukawa et al., 1996), auxins can also be produced from indole-3-pyruvate. Auxins could also be produced in a tryptophan-independent manner but the corresponding pathways are still not well described.

A lot of fungal species, and not only plant-interacting fungi, produce and secrete auxins, suggesting that these hormones could have an endogenous role in these organisms (Gruen,

1959; Ulrich, 1960). A negative correlation between the speed of fungal growth and auxin production was shown in several species (Gruen, 1959). By contrast, auxin treatment promotes cellular elongation and sporulation in the yeasts *Saccharomyces cerevisiae* and *S. ellipsoideus* (Yanagishima, 1965; Kamisaka et al., 1967). Similarly, an aberrant production of auxins leads to morphological transition in *S. cerevisiae* as well as in the human pathogen *Candida albicans*, in which auxin triggers the transition into hyphal growth, a known virulence trait (Rao et al., 2010). The auxin IAA also promotes spore germination of the filamentous fungus *Neurospora crassa* (Nakamura et al., 1978, 1982). When tested on the tomato pathogen *F. oxysporum lycopersici*, auxin reduced spore germination (Sharaf and Farrag, 2004). The effect of auxin on growth could be concentration-dependent: in the case of *F. delphinoides*, a chick pea pathogen, low concentration of exogenous auxin increased fungal growth whereas high concentration decreased it (Kulkarni et al., 2013). Thus the effects of auxins on fungal physiology can strongly differ from one species to another and depends on the dose tested.

Auxins are involved in the symbiotic interactions between plant and bacteria or fungi. They are required for the initiation of nodule formation in the nitrogen-fixative bacterial symbiosis (Hirsch and Fang, 1994) and for the invasion of mycorrhizal fungi (Hanlon and Coenen, 2011; Etemadi et al., 2014). For instance, mutants of the ectomycorrhizal species *Hebeloma cylindrosporum* overproducing auxin showed an increased ability to invade root tissues of *Pinus pinaster* (Gay et al., 1994; Laurans et al., 2001). However, there was no difference between the growth of plants colonized with the mutant and wild-type strains, suggesting that fungal auxin is involved in host invasion but not in the beneficial effects of symbiosis on host development. In most cases, plants interacting with mycorrhizal fungus contain a higher content in auxins than non-mycorrhized ones (Barker and Tagu, 2000; Meixner et al., 2005).

However, the origin of these auxins, whether from the host or the fungal symbiont, is still unclear.

Auxin involvement in plant-pathogen interactions were early suspected and studied when symptoms, like organ deformation, were reminiscent to responses to high auxin level. For instance, *Agrobacterium tumefaciens* and *Pseudomonas savastanoi* are well-known plant pathogenic bacteria that induce tumor formation in their hosts and auxins actively contribute to the virulence of these bacteria (Glass and Kosuge, 1988). Like *Agrobacterium*, some fungi are able to induce tumors, such as the corn smut causal agent, *U. maydis*. However, fungal mutants affected in auxin production were still able to induce tumors in a similar way than wild-type strain even if tumors contained a lower level of auxins (Reineke et al., 2008). This suggests that auxin production by *U. maydis* is not required for the virulence of this pathogen.

In the case of fungal pathogens not triggering organ deformations, functional evidences suggest a role for auxins. By measuring fungal biomass and auxins in plant tissues, it was suggested that *C. gloeosporioides* f. sp. *aeschynomene* produces auxins during early, biotrophic stages of plant colonization (Maor et al., 2004). In *F. oxysporum*, an enhanced expression of auxin biosynthetic genes (tryptophan-2-monooxygenase and indole-3-acetamide hydrolase) triggers over-accumulation of IAA and an hypervirulent phenotype on Orobanche (Cohen et al., 2002). Consistent with these observations, the transient silencing in *Puccinia graminis* f. sp. *tritici* of a gene required for auxin biosynthesis was obtained in wheat infected leaves. The two-fold reduction of the transcript led to a decrease of pustule formation. Although auxins were not measured after silencing, these results suggest that auxins were required for full fungal pathogenicity (Yin et al., 2014). Thus auxins seem to play a role in pathogenicity and more functional studies with fungal mutants should help better understand how they participate to virulence.

## Cytokinins from fungi: a now clear-cut positive function in virulence

Cytokinins (CKs) are diversified plant hormones derived from ATP/ADP/AMP or from the tRNA degradation pathway. CKs are well described for their role in plant development processes such as root and shoot formation through the regulation of cell cycle and cell differentiation (Barciszewski et al., 1999; Riou-Khamlichi et al., 1999; Carimi et al., 2003; Fosket and Torrey, 1969). CKs are also involved in delaying senescence and in source/sink nutrient distribution (Wingler et al., 1998; Peleg et al., 2011). The first step of CK biosynthesis in plants involves Isopentenyl Transferase enzymes (IPT or tRNA-IPT) which perform the transfer of the isopentenyl chain from the methylerythritol phosphate (MEP) on the adenosine phosphate substrate leading to the formation of the ribosylated phosphorylated CK forms (Sakakibara, 2006). Then, these CKs are activated, in part by the LONELY GUY (LOG) enzymes, into CK free active forms like trans-zeatin and isopentenyladenine (Kurakawa et al., 2007; Frébort et al., 2011). The putative *IPT* and *LOG* genes are present in several fungal genomes and some of them have been recently characterized (see below (Hinsch et al., 2015; Morrison et al., 2015b; Chanclud et al., 2016)).

A large diversity of fungal species, whether saprophytic, pathogenic or symbiotic, were shown to produce CKs (Murphy et al., 1997; Cooper and Ashby, 1998) and several studies suggest that they could play a role in several physiological processes in fungi themselves especially in hyphal development and nutrient uptake (LeJohn and Stevenson, 1973). For instance, CKs promote *in vitro* branching of ectomycorrhizal mycelia (Barker and Tagu, 2000) and affect in a dose-dependent manner hyphal membrane viscosity and therefore influence ion and water transport (Gogala, 1991; LeJohn and Stevenson, 1973). Pohleven et al., 1986, demonstrated that some CKs modify the content of K, Ca, P and Na in the mycelia of the basidiomycete *Suillus variegatus* (Gogala, 1991). The effect of CKs on hyphal growth seems to depend on the concentration and on the kind of CK molecule tested (Gryndler et al.,



1998). CKs could also be involved in growth optimization under adverse conditions. For instance, the inhibition of the mycelial growth of *Amanita muscaria* caused by aluminium is significantly correlated with a decrease in CKs amount (Kovač and Žel, 1995). In a recent report, we have shown that endogenous and exogenous CKs are required for oxidative stress tolerance in the rice blast fungus *Magnaporthe oryzae* (Chanclud et al., 2016). In the 60's Lee et al reported that CKs also affect sexual reproduction in the ascomycete *N. crassa* suggesting a role in communication within fungi (Lee, 1961; Elliott, 1967).

During mycorrhizal symbiosis, CKs promote growth of the host and of the symbiont (Allen et al., 1980; Drüge and Schonbeck, 1993; Barker and Tagu, 2000). CK accumulation in the host, root and shoot, were shown in many fungal symbiotic interactions (Allen et al., 1980). A model has emerged since the early 90's about the role of CKs in plant symbiotic interactions, proposing that plants secrete CKs that (i) promote growth of symbiotic microbes that are thus able to detect them, then (ii) this contributes to a better absorption of nutrients through the symbiont and (iii) leads to increase the photosynthesis process in the host leaves (Wullschleger and Reid, 1990; Drüge and Schonbeck, 1993). It is possible that CKs produced by mycorrhizas may initiate this whole process but this awaits the study of CK-deficient symbiotic fungal mutants to confirm this hypothesis.

During interaction with fungal pathogens, CKs content is often affected (Jiang et al., 2013; Devos et al., 2006). Since most of the necrotrophic fungi analyzed do not seem to secrete CKs, in contrast to (hemi)biotrophic ones, it was suggested that fungal CK production and secretion could depend on pathogen lifestyle. CKs are involved in many diseases caused by pathogens that induce tumor formation in their host: protists (e.g. *Plasmodiophora brassicae* (Siemens et al., 2006)), nematodes (e.g. *Heterodera schachtii* (Siddique et al., 2015)), bacteria (e.g. *P. savastanoi* (Barciszewski et al., 2000), *Agrobacterium* sp.(Barciszewski et al., 2000), *Rhodococcus fascians* (Pertry et al., 2009)) and fungi (*U. maydis* (Mills and Van Staden,

1978), *Claviceps purpurea* (Hinsch et al., 2015)). In the tumor-inducing pathogen *C. purpurea*, deletion of two genes partially abolished CK *de novo* synthesis but did not affect virulence of the fungus. By contrast the mutants exhibited a hyper-sporulating phenotype, implying that CKs are environmental factors influencing fungal development (Hinsch et al., 2015). Recently it was shown that CK accumulation in *U. maydis* infected tissues is correlated to the virulence of this pathogen but there was no direct genetic evidence that fungal-derived CKs are required for full virulence of this pathogen (Morrison et al., 2015a).

Fungal pathogens that do not induce tumors also produce CK compounds and their role in virulence is still poorly understood (see for instance (Murphy et al., 1997; Jiang et al., 2013)). CKs are probably involved in “green island” formation, a photosynthetically active zone often found around lesions caused by biotrophic fungi (Angra and Mandahar, 1991; Choi et al., 2011). In plants CK production is thought to occur in the roots (Rani Debi et al., 2005) and experiments with detached leaves could indirectly address the question of the origin of the CKs in green islands. Using this assay in wheat and maize leaves respectively infected with *Pyrenophora teres* and *Dreschslera maydis*, the increase of CKs content was attributed to the pathogen. The increase in CKs levels in susceptible hosts was also correlated with increased metabolite contents around infection sites (Angra-Sharma and Sharma, 1999). CK secretion was shown by immuno-detection in plant tissues in the case of *P. recondita* f.sp. *tritici* during wheat infection (Hu and Rijkenberg, 1998) but as for most hormones found during infection, it is not possible to unambiguously assign this accumulation to the plant or to the pathogen without characterization of mutants impaired in CK production or perception. A recent study mentioned that CK production by fungi, especially the *cis*- zeatin forms (which seems to be the main one produced by filamentous fungi), could involve tRNA-IPT enzymes that perform modification on tRNA which will then release free CKs after degradation (Morrison et al., 2015b). Among non-tumor inducing fungal pathogens, *M. oryzae* produces and secretes CKs

(Jiang et al., 2013). Knock-out mutants impaired in the only *tRNA-IPT* gene identified in *M. oryzae* were also impaired in CK production, thus confirming the hypothesis of Morrison et al., 2015b that tRNA-IPT are involved in fungal CK production (Chanclud et al., 2016). The interaction between rice and the CK-deficient strain of *M. oryzae* was characterized. This analysis demonstrated that *M. oryzae* derived CKs are required for full virulence by affecting rice defenses, nutrient distribution and fungal oxidative stress tolerance (Chanclud et al., 2016). Since the tRNA-IPT gene identified in *M. oryzae* is well conserved, this mutation could be studied in other fungi as a nice tool for distinguishing fungal CKs to plant CKs in other plant-fungus interactions. Recently, the deletion of a *tRNA-IPT* gene has also been performed in the nematode *H. schachtii* confirming the conservation of the role of this enzyme in CK production among different organisms (Siddique et al., 2015).

#### **Ancient but limited direct evidence for a role of Gibberellic acids from fungal origin**

GAs are terpenoid hormonal compounds identified for the first time as being produced by *Gibberella fujikuroi*. This fungus is the causal agent of the “bakanae” or “foolish seedlings” disease of rice in which infected plants are abnormally tall. Following this discovery, the role of GAs on plant physiology started to be studied. GAs have been involved in the control of germination, flowering, cell division and internode elongation (Brian and Elson, 1954; Pimenta Lange and Lange, 2006; Swain and Singh, 2005). The first steps of GA biosynthesis pathways identified in fungi are almost identical with those known in plants. The complex GA biosynthesis pathways were well described by Tudzynski, 2005. GA production was found in several fungal species but their effects on fungal biology are not well described.

In liquid culture, GAs were shown to increase conidial germination and to improve growth of young hyphae of the ascomycete fungus *N. crassa* (Nakamura et al., 1978; Tomita et al.,



responses in plants (Peleg and Blumwald, 2011). ABA is also involved in seed dormancy by acting antagonistically with the GAs pathway (Debeaujon and Koornneef, 2000). In plants, ABA is synthesized from both the MEP and the mevalonate pathway (Nambara and Marion-Poll, 2005).

In fungi, it is thought that the mevalonate pathway is mostly involved and that different ABA precursors could be used (Morrison et al., 2015b; Oritani and Kiyota, 2003). Fungal production of ABA was first shown in *Cercospora risicola* (Norman et al., 1983). Since then a lot of fungi with different lifestyles (saprophytic, symbiotic and pathogenic) were described as producing ABA (Crocoli et al., 1991; Esch et al., 1994; Jiang et al., 2010; Morrison et al., 2015b). There are only two reports that ABA affects mycelium growth. In *Ceratocystis fimbriata*, exogenous application of ABA slightly promoted fungal growth. In *M. oryzae*, ABA increased germination and the formation of appressorium, a specialized infection structure differentiated for breaking down the plant cell wall and allowing invasion (Spence and Bais, 2015 and references inside).

The arbuscular-mycorrhizal fungus *Glomus sp.* produces ABA and ABA concentration in the xylem sap is different between mycorrhized plants compared to non-mycorrhized ones. However the origin of this increase of ABA is not established yet (Esch et al., 1994).

In several plant-pathogen interactions, ABA was described to affect plant disease resistance in a positive or a negative manner, depending on the host-pathogen interaction studied (Jiang et al., 2010; Xu et al., 2013; De Vleeschauwer et al., 2010). Kettner and Dorffling, 1995 have inoculated tomato plants with two strains of *Botrytis cinerea* presenting differences in ABA production and showed that ABA increase is higher in leaves inoculated with the ABA-producing strain than with the less-producing ones. This suggests that ABA accumulation in the host during infection could result from or be initiated by this pathogenic fungus. Similarly, ABA was accumulated during the early stages of infection by *U. maydis* and this

accumulation could be correlated with the virulence of the fungus (Morrison et al., 2015a). Even though exogenous ABA triggered a faster development of the necrotic lesion, the role of fungal ABA in virulence was not described until recently. Knocking-out one gene homologous to the *B. cinerea* *ABA4* gene responsible for ABA biosynthesis, reduced by two-fold ABA levels in *M. oryzae* (Spence et al., 2015; Siewers et al., 2006). Appressorium formation *in vitro* was severely reduced in the *M. oryzae*  $\Delta$ *aba4* mutant, a phenotype that could be reverted by exogenous application of ABA. The virulence of the  $\Delta$ *aba4* mutant was also strongly compromised suggesting that ABA contributes to the virulence of this fungus. One may then speculate that this production of ABA by *M. oryzae* inhibits the SA-dependent defense response (Jiang et al., 2010), as observed in many other biological situations (Ton et al., 2009). However, since the  $\Delta$ *aba4* mutant did not form appressoria and infect plants at all, it is difficult to conclude on a role of fungal-produced ABA on the plant itself.

### **Ethylene: a gaseous hormone involved in plant physiology and defenses which also affects fungal development**

Ethylene (ET) is a gaseous compound first discovered for its role in fruit maturation (Bleecker and Kende, 2000; Payton et al., 1996). ET was later shown to be involved in senescence, germination, flowering as well as in the inhibition of root and shoot growth (Bleecker and Kende, 2000; Grbic and Bleecker, 1995). In *Arabidopsis*, ET has first been described to contribute, with JA, to the induction of defenses against necrotrophic pathogens. However, this dichotomy of responses, to biotrophic and necrotrophic pathogens, is not always clear in the other plants, like in rice, in which hormonal regulation of defenses is slightly different (De Vleeschauwer et al., 2013, 2014). In plants and fungi, ET biosynthesis occurs from methionine that is transformed in ACC (1-aminocyclopropane-1- carboxylic acid) via ACC-synthase enzymes (Esser et al., 2002). Moreover, fungi also produce ET from the 2-keto-4-

methylthiobutyric acid, deriving from methionine, and/or from the 2-oxoglutarate then requiring Ethylene-Forming enzymes (Bockhaven et al., 2015; Hottiger and Boiler, 1991 and references inside). Altogether, studies about fungal ET production show that it is strongly dependent on growth media and confirmed that several pathways exist among fungi (Esser et al., 2002; Strzelczyk et al., 1994). Since the first report of ET production by *Penicillium digdatum* in 1940, ET production has been measured in a lot of fungal species, in both hyphae and spores (Dasilva et al., 1974). These fungi belong to different phylum, have different lifestyles and range from pathogenic like *B. cinerea* to symbiotic ones like *F. oxysporum* f. sp. *pini* (Graham and Linderman, 1980; Dasilva et al., 1974; Arshad and Frankenberger, 1991). Several *in vitro* experiments demonstrated that ET or some precursors (ethephon and ACC) affect spore germination and hyphal growth of the pathogenic filamentous fungi, *Alternaria alternata* and *B. cinerea*, and the symbiotic ones, *Gigaspora ramisporophora* and *G. mosseae* (Kępczyńska, 1994; Chagué et al., 2006). The effects of ET on fungal development seem to be dose-dependent with a promoting effect observed at concentrations below 1mM and a negative effect at or higher than 1mM (Ishii et al., 1996).

In the case of mycorrhiza, the role of ET depends on the type of symbiotic interaction. A low content of ET was measured in mycorrhized roots (McArthur and Knowles, 1992) and an exogenous supply of ET suppressed AM development (Geil et al., 2001; Zsögön et al., 2008). Therefore, it was suggested that a repression of the ET pathway by AM fungi is required to allow the establishment of symbiosis. Indeed, the AM fungus *G. intraradices* secretes a protein (SP7: secreted protein 7) which interacts with an ethylene response factor to suppress ethylene signaling (Kloppholz et al., 2011). In contrast, ET seems to promote ectomycorrhizal symbiosis. Two species of truffles (*Tuber melanosporum* and *Tuber borchii*) were showed to produce ET (and auxin) for manipulating these hormonal pathways in the host and inducing root morphological modifications, a plant developmental process in which these hormones are



involved (Splivallo et al., 2009). Given the roles of ET in plant defense, this fungal ET production by symbionts could also be required for counteracting the establishment of host immunity.

During plant fungal pathogen interactions, ET content often increases at the beginning of the interaction (Broekaert et al., 2006). However, its origin, from plants or fungi, is still unclear.

In the case of the *Colletotrichum* sp. pathogens, ET is required for the formation of appressorium (Flaishman and Kolattukudy, 1994). Indeed appressoria formed on ripening tomato whereas none formed in plant mutants affected in ET production. An exogenous supply of ET restored appressorium formation on these plant mutants suggesting that ET produced by fruits during the ripening is perceived by the pathogen and is beneficial to initiate the development of specialized structures required for penetration and thus for full virulence. Furthermore ET production by fungi could be required to disturb host defense induction by affecting the plant hormonal homeostasis, essential for plant immunity establishment (Broekaert et al., 2006). This hypothesis was recently investigated in the interaction between rice and the necrotrophic pathogen, *Cochliobolus miyabeanus*, in which ET increases rice susceptibility (Bockhaven et al., 2015). In this study, the authors have used a specific inhibitor of fungal ET biosynthesis (2,2-bipyridyl), that abolishes fungal ET production and leads to a higher resistance of the host (or a lack of virulence of the fungus). Combined with other results showing that *C. miyabeanus* affects the 2-oxoglutarate (an ET precursor for microbes) pool in rice, the authors suggested that ET accumulation is mainly initiated and caused by the fungus and then contributes to the symptom development (Bockhaven et al., 2015). However, exogenous supplies of hormonal production inhibitors or signaling inhibitors could have side effects on the host and on the fungus, therefore the study of fungal mutants affected in ET perception or production is still missing for understanding the different roles of ET in plant-fungus interactions.



### Pathogenic fungi also produce defense-related hormones SA and JA

In most plants, SA and JA trigger defenses against fungal biotrophic and necrotrophic pathogens respectively, in an antagonist manner (Bari and Jones, 2009; Robert-Seilaniantz et al., 2011). Some fungal pathogens may produce one hormone in order to inhibit the defense pathway which is the most detrimental to their growth. In plants, SA is synthesized from chorismate and the corresponding pathway is for instance targeted by the fungal pathogen *U. maydis* which secretes a chorismate mutase that channels chorismate into the phenylpropanoid pathway, preventing SA accumulation during infection and then contributing to its virulence (Djamei et al., 2011). However, the chorismate pathways identified in fungi do not lead to SA biosynthesis. Thus, even if SA (or SA-derivatives) production was measured in different species, to date this pathway is still unknown in fungi (Packter and Steward, 1967).

Some pathogens produce both SA and JA, like for instance *Moniliophthora perniciosa* which causes the witches' broom disease of cocoa. In this case, the production of these hormones could (i) contribute to manipulate the hormonal pathways involved in the host defense responses throughout its invasion i.e. causing abnormal shoot development and necrosis and (ii) could have a direct effect on this fungus since both SA and JA promote *in vitro* growth (Kilaru et al., 2007; Chaves and Gianfagna, 2006). A few other studies mentioned the effects of SA or JA on fungal physiological processes. SA had a moderate suppressive effect on spore germination and colonial growth rate of *Harpophora maydis* (Degani et al., 2015). In *Aspergillus flavus*, the results obtained from *in vitro* experiments showed that SA significantly reduced hyphal growth at all concentrations tested (Panahirad et al., 2014).

Several studies mentioned the production of JA by pathogenic fungi like *G. fujikuroi* and *Botryodiplodia theobromae* (Miersch et al., 1992, 1991). JAs are derived from lipid

peroxidation, and thus belong to the oxylipins. Some fungal oxylipin biosynthesis pathways were identified and characterized. JA and the other oxylipins could affect both host and fungal physiological processes (Tsitsigiannis and Keller, 2007). The *in vitro* application on *F. oxysporum* f.sp. *lycopersici* of the methyl-JA reduced spore germination and mycelium growth (Król et al., 2015). Recently, *M. oryzae* was shown to secrete a monooxygenase that converts rice endogenous JA into hydroxylated JA (12OH-JA). This 12OH-JA may then inhibit JA signaling and thus impairs JA-dependent host defenses and resistance (Patkar et al., 2015).

Although the number of pathogenic fungi characterized for producing SA and/or JA increases, there is no direct evidence that fungal SA or JA are required for their virulence.

## Conclusions

Thus far, most fungi have been shown to produce almost all plant-like hormones *in vitro*. It is noteworthy that some growth medium used in many studies are made with potatoes dextrose agar and/or yeast extract, two compounds that already contain some plant derived hormones including auxins, CKs, ABA and the others, in unknown concentrations. Thus, some confusion exists between the ability of *de novo* production of plant hormones by fungi and their ability to metabolize them from the growth medium. This should be carefully addressed in future studies conducted *in vitro*. The sequencing of many genomes may also help to shed light on the presence of the hormonal biosynthesis pathways already described in some fungal species (Esser et al., 2002).

This overview shows that most known plant-hormonal compounds are produced and perceived by fungi. To date the involvement of fungal hormonal compounds, and the way that they are secreted and act in the plant cell are still poorly understood. Although most of the biosynthesis pathways of hormones in fungi are well described (reviewed in Esser et al., 2002), studies on fungal mutants affected in hormonal production are strikingly missing to confirm the involvement of fungal derived plant hormones in such interactions. In particular, the origin of hormones in colonized tissues is unclear and needs to be established to understand the complex relationships between fungal-derived and plant-derived hormones.

Some plant hormones have been shown to affect fungal development, nutrition and reproduction processes suggesting that these molecules trigger some signals in fungi (Nakamura et al., 1982; Gryndler et al., 1998; Esch et al., 1994; Elliott, 1967; Nakamura et al., 1978; LeJohn and Stevenson, 1973; Kępczyńska, 1989). **The Figure 1 non-exhaustively summarizes the main effects of hormones known to date on fungal biology.** However, the perception systems of hormones by fungi, as well as the signaling pathways triggered and the physiological responses induced, still remain to be discovered.

These plant hormone compounds are also produced, and very probably perceived, by other microbes including bacteria and nematodes (Denancé et al., 2013; Siddique et al., 2015; Kisiala et al., 2013). Moreover, some of these compounds have some effects on animal cells (Jiang et al., 2002; Slauchhaupt et al., 2004; Ishii et al., 2003), suggesting that “plant” hormones not only participate to plant-microbe dialogue but might also contribute to communication in other host-microbe interactions involving widely different organisms (animals, plants and all kind of pathogenic, saprophytic and symbiotic microbes). However, to date there is no report on the involvement of such compounds in these interactions.

## Acknowledgements

We thank Alexandre Robert-Seilaniantz, Monica Höfte and Florian Frugier for critical reading of this manuscript.

Accepted Article

Version postprint

**Figure 1. Schematic representation of the effects of hormones on fungal biology.**

A schematic view of yeast-like (light grey background), infecting and sporulating filamentous fungus is given. This model summarizes the effects of hormones on fungal biological processes reported to date; the effects due to the dose, the molecule tested, other environmental factors as well as the specie studied are provided in the text and could differ among fungi. Arrows and « T » bars respectively represent positive and negative effects. AUX : Auxins, ABA : Absciscic Acid, ET : Ethylene, CKs : Cytokinins, GAs : Gibberellic Acids, JA : Jasmonic acid, SA : Salicylic Acid.

## References

- Allen, M.F., Moore Jr., T.S., and Christensen, M.** (1980). infected by vesicular–arbuscular mycorrhizae: I. Cytokinin increases in the host plant. *Can. J. Bot.* **58**: 371–374.
- Angra, R. and Mandahar, C.L.** (1991). Pathogenesis of barley leaves by *Helminthosporium teres* I: Green island formation and the possible involvement of cytokinins. *Mycopathologia* **114**: 21–27.
- Angra-Sharma, R. and Sharma, D.K.** (1999). Cytokinins in pathogenesis and disease resistance of *Pyrenophora teres*-barley and *Drechslera maydis*-maize interactions during early stages of infection. *Mycopathologia* **148**: 87–95.
- Arshad, M. and Frankenberger, W.T.** (1991). Microbial production of plant hormones. *Plant Soil* **133**: 1–8.
- Barciszewski, J., Rattan, S.I.S., Siboska, G., and Clark, B.F.C.** (1999). Kinetin — 45 years on. *Plant Sci.* **148**: 37–45.
- Barciszewski, J., Siboska, G., Rattan, S.I.S., and Clark, B.F.C.** (2000). Occurrence, biosynthesis and properties of kinetin (N<sup>6</sup>-furfuryladenine). *Plant Growth Regul.* **32**: 257–265.
- Bari, R. and Jones, J.D.G.** (2009). Role of plant hormones in plant defence responses. *Plant Mol. Biol.* **69**: 473–88.
- Barker, S. and Tagu, D.** (2000). The Roles of Auxins and Cytokinins in Mycorrhizal Symbioses. *J. Plant Growth Regul.* **19**: 144–154.
- Beardsell, M.F. and Cohen, D.** (1975). Relationships between Leaf Water Status, Abscisic Acid Levels, and Stomatal Resistance in Maize and Sorghum. *Plant Physiol.* **56**: 207–212.

**Benjamins, R. and Scheres, B.** (2008). Auxin: the looping star in plant development. *Annu. Rev. Plant Biol.* **59**: 443–465.

**Blee, K. a. and Anderson, A.J.** (1998). Regulation of arbuscule formation by carbon in the plant. *Plant J.* **16**: 523–530.

**Bleecker, a B. and Kende, H.** (2000). Ethylene: a gaseous signal molecule in plants. *Annu. Rev. Cell Dev. Biol.* **16**: 1–18.

**Bockhaven, J. Van, Strnad, M., Asano, T., Kikuchi, S., Monica, H., and Vleesschauwer, D. De** (2015). Silicon induces resistance to the brown spot fungus *Cochliobolus miyabeanus* by preventing the pathogen from hijacking the rice ethylene pathway. *New Phytol.*: 761–773.

**Brian, P. and Elson, G.** (1954). The plant-growth-promoting properties of gibberellic acid, a metabolic product of the fungus *Gibberella fujikuroi*. *J. Sci. Food Agric.* **5**: 602–612.

**Broekaert, W.F., Delauré, S.L., De Bolle, M.F.C., and Cammue, B.P. a** (2006). The role of ethylene in host-pathogen interactions. *Annu. Rev. Phytopathol.* **44**: 393–416.

**Carimi, F., Zottini, M., Formentin, E., Terzi, M., and Lo Schiavo, F.** (2003). Cytokinins: new apoptotic inducers in plants. *Planta* **216**: 413–421.

**Chagué, V., Danit, L.-V., Siewers, V., Schulze-Gronover, C., Tudzynski, P., Tudzynski, B., and Sharon, A.** (2006). Ethylene sensing and gene activation in *Botrytis cinerea*: a missing link in ethylene regulation of fungus-plant interactions? *Mol. Plant. Microbe Interact.* **19**: 33–42.

**Chanclud, E., Kisiala, A., Emery, N.R.J., Chalvon, V., Ducasse, A., Romiti-Michel, C., Gravot, A., Kroj, T., and Morel, J.-B.** (2016). Cytokinin Production by the Rice Blast Fungus Is a Pivotal Requirement for Full Virulence. *PLOS Pathog.* **12**: e1005457.



**Chaves, F.C. and Gianfagna, T.J.** (2006). Necrotrophic phase of *Moniliophthora perniciosa* causes salicylic acid accumulation in infected stems of cacao. *Physiol. Mol. Plant Pathol.* **69**: 104–108.

**Choi, J., Choi, D., Lee, S., Ryu, C.-M., and Hwang, I.** (2011). Cytokinins and plant immunity: old foes or new friends? *Trends Plant Sci.* **16**: 388–94.

**Cohen, B. a, Amsellem, Z., Maor, R., Sharon, A., and Gressel, J.** (2002). Transgenically enhanced expression of indole-3-acetic Acid confers hypervirulence to plant pathogens. *Phytopathology* **92**: 590–596.

**Cooper, S.J. and Ashby, A.M.** (1998). Comparison of cytokinin and cytokinin-O-glucoside cleaving beta-glucosidase production in vitro by *Venturia inaequalis* and other phytopathogenic fungi with differing modes of nutrition in planta. *Physiol. Mol. Plant Pathol.* **53**: 61–72.

**Crocoli, C., Kettner, J., and Dorffling, K.** (1991). Abscisic acid in saprophytic and parasitic species of fungi. *Phytochemistry* **30**: 1059–1060.

**Dasilva, E.J., Henriksson, E., and Henriksson, L.E.** (1974). Ethylene production by fungi. *Plant Sci. Lett.* **2**: 63–66.

**Dean, R., Van Kan, J.A.L., Pretorius, Z.A., Hammond-Kosack, K.E., Di Pietro, A., Spanu, P.D., Rudd, J.J., Dickman, M., Kahmann, R., Ellis, J., and Foster, G.D.** (2012). The Top 10 fungal pathogens in molecular plant pathology. *Mol. Plant Pathol.* **13**: 414–430.

**Debeaujon, I. and Koornneef, M.** (2000). Gibberellin requirement for *Arabidopsis* seed germination is determined both by testa characteristics and embryonic abscisic acid. *Plant Physiol.* **122**: 415–424.

**Degani, O., Drori, R., and Goldblat, Y.** (2015). Plant growth hormones suppress the development of *Harpophora maydis*, the cause of late wilt in maize. *Physiol. Mol. Biol. Plants* **21**: 137–149.

**Denancé, N., Sánchez-Vallet, A., Goffner, D., and Molina, A.** (2013). Disease resistance or growth: the role of plant hormones in balancing immune responses and fitness costs. *Front. Plant Sci.* **4**: 1–12.

**Desjardins, A.E., Manandhar, H.K., Plattner, R.D., Manandhar, G.G., Poling, S.M., and Maragos, C.M.** (2000). *Fusarium* Species from Nepalese Rice and Production of Mycotoxins and Gibberellic Acid by Selected Species. *Appl. aenvironmentaenvironmenta* **66**: 1020–1025.

**Devos, S., Laukens, K., Deckers, P., Van Der Straeten, D., Beeckman, T., Inzé, D., Van Onckelen, H., Witters, E., and Prinsen, E.** (2006). A hormone and proteome approach to picturing the initial metabolic events during *Plasmodiophora brassicae* infection on *Arabidopsis*. *Mol. Plant. Microbe. Interact.* **19**: 1431–1443.

**Djamei, A. et al.** (2011). Metabolic priming by a secreted fungal effector. *Nature* **478**: 395–8.

**Drüge, U. and Schonbeck, F.** (1993). Effect of vesicular-arbuscular mycorrhizal infection on transpiration, photosynthesis and growth of flax (*Linum usitatissimum* L.) in relation to cytokinin levels. *J. Plant Physiol.* **141**: 40–48.

**Elliott, R.** (1967). Effects of Kinetin and Related Compounds on Growth and Sexual Reproduction of *Saprolegnia australis*. *Planta* **77**: 164–175.

**Esch, H., Hundeshagen, B., Schneider-Poetsch, H., and Bothe, H.** (1994). Demonstration of abscisic acid in spores and hyphae of the arbuscular-mycorrhizal fungus *Glomus* and in the N<sub>2</sub>-fixing cyanobacterium *Anabaena variabilis*. *Plant Sci.* **99**: 9–16.

**Esser, K., Bennett, J.W., and Osiewacz, H.D.** (2002). *The Mycota : Industrial Applications* H.D. Osiewacz, ed (Springer Berlin Heidelberg: Berlin, Heidelberg).

**Etemadi, M., Gutjahr, C., Couzigou, J.-M., Zouine, M., Laouressergues, D., Timmers, A., Audran, C., Bouzayen, M., Bécard, G., and Combier, J.-P.** (2014). Auxin Perception Is Required for Arbuscule Development in Arbuscular Mycorrhizal Symbiosis. *Plant Physiol.* **166**: 281–292.

**Flaishman, M. a and Kolattukudy, P.E.** (1994). Timing of fungal invasion using host's ripening hormone as a signal. *Proc. Natl. Acad. Sci. U. S. A.* **91**: 6579–6583.

**Fosket, D.E. and Torrey, J.G.** (1969). Hormonal control of cell proliferation and xylem differentiation in cultured tissues of *Glycine max* var. *Biloxi*. *Plant Physiol.* **44**: 871–880.

**Frébort, I., Kowalska, M., Hluska, T., Frébortová, J., and Galuszka, P.** (2011). Evolution of cytokinin biosynthesis and degradation. *J. Exp. Bot.* **62**: 2431–52.

**Furukawa, T., Koga, J., Adachi, T., Kishi, K., and Syono, K.** (1996). Efficient Conversion of L-Tryptophan to Indole-3-Acetic Acid and/or Tryptophol by Some Species of *Rhizoctonia*. **37**: 899–905.

**Gay, G., Normand, L., Marmeisse, R., Sotta, B., and Debaud, J. c.** (1994). Auxin overproducer mutants of *Hebeloma cylindrosporum* Romagnesi have increased mycorrhizal activity. *New Phytol.* **128**: 645–657.

**Geil, R.D., Peterson, R.L., and Guinel, F.C.** (2001). Morphological alterations of pea (*Pisum sativum* cv. *Sparkle*) arbuscular mycorrhizas as a result of exogenous ethylene treatment. *Mycorrhiza* **11**: 137–143.

**Glass, N.L. and Kosuge, T.** (1988). Role of Indoleacetic Acid Lysine Synthetase in Regulation of Indoleacetic-Acid Pool Size and Virulence of *Pseudomonas-Syringae*

Subsp Savastanoi. *J. Bacteriol.* **170**: 2367–2373.

**Gogala, N.** (1991). Regulation of mycorrhizal infection by hormonal factors produced by hosts and fungi. *Experientia* **47**: 331–340.

**Graham, J.H. and Linderman, R.G.** (1980). Ethylene production by ectomycorrhizal fungi, *Fusarium oxysporum* f. sp. *pini*, and by aseptically synthesized ectomycorrhizae and *Fusarium*-infected Douglas-fir roots. *Can. J. Microbiol.* **26**: 1340–1347.

**Grbic, V. and Bleecker, A.B.** (1995). Ethylene Regulates the Timing of Leaf Senescence in *Arabidopsis*. *Plant J.* **8**: 595–602.

**Gruen, H.E.** (1959). Auxins and Fungi. *Annu. Rev. Plant Physiol.* **10**: 405–440.

**Gryndler, M., Hrselova, H., Chvatalova, I., and Jansa, J.** (1998). The effect of selected plant hormones on in vitro proliferation of hyphae of *Glomus fistulosum*. *Biol. Plant.* **41**: 255–263.

**Gust, A. a., Willmann, R., Desaki, Y., Grabherr, H.M., and Nürnberger, T.** (2012). Plant LysM proteins: modules mediating symbiosis and immunity. *Trends Plant Sci.* **17**: 495–502.

**Hanlon, M.T. and Coenen, C.** (2011). Genetic evidence for auxin involvement in arbuscular mycorrhiza initiation. *New Phytol.* **189**: 701–709.

**Hinsch, J., Vrabka, J., Oeser, B., Novák, O., Galuszka, P., and Tudzynski, P.** (2015). De novo biosynthesis of cytokinins in the biotrophic fungus *Claviceps purpurea*. *Environ. Microbiol.* **17**: 2935–2951.

**Hirsch, A.M. and Fang, Y.** (1994). Plant hormones and nodulation: what's the connection? *Plant Mol. Biol.* **26**: 5–9.

**Hirsch, A.M., Fang, Y., Asad, S., and Kapulnik, Y.** (1997). The role of phytohormones in

plant-microbe symbioses. *Plant Soil* **194**: 171–184.

**Hottiger, T. and Boiler, T.** (1991). Ethylene biosynthesis in *Fusarium oxysporum* f. sp. *tulipae* proceeds from glutamate/2-oxoglutarate and requires oxygen and ferrous ions in vivo. *Arch Microbiol.* **157**: 18–22.

**Hu, G.G. and Rijkenberg, E.H.J.** (1998). Ultrastructural localization of cytokinins in *Puccinia recondita* f.sp. *tritici*-infected wheat leaves. *Physiol. Mol. Plant Pathol.* **52**: 79–94.

**Ishii, T., Shrestha, Y., Matsumoto, I., and Kadoya, K.** (1996). Effect of Ethylene on the Growth of Vesicular-Arbuscular Mycorrhizal Fungi and on the Mycorrhizal Formation of Trifoliolate Orange Roots. *J. Japanese Soc. Hortic. Sci.* **65**: 525–529.

**Ishii, Y., Sakai, S., and Honma, Y.** (2003). Cytokinin-induced differentiation of human myeloid leukemia HL-60 cells is associated with the formation of nucleotides, but not with incorporation into DNA or RNA. *Biochim. Biophys. Acta - Mol. Cell Res.* **1643**: 11–24.

**Jiang, C.-J., Shimono, M., Sugano, S., Kojima, M., Liu, X., Inoue, H., Sakakibara, H., and Takatsuji, H.** (2013). Cytokinins act synergistically with salicylic acid to activate defense gene expression in rice. *Mol. Plant. Microbe. Interact.* **26**: 287–96.

**Jiang, C.-J., Shimono, M., Sugano, S., Kojima, M., Yazawa, K., Yoshida, R., Inoue, H., Hayashi, N., Sakakibara, H., and Takatsuji, H.** (2010). Abscisic acid interacts antagonistically with salicylic acid signaling pathway in rice-*Magnaporthe grisea* interaction. *Mol. Plant. Microbe. Interact.* **23**: 791–8.

**Jiang, G., Dallas-Yang, Q., Liu, F., Moller, D.E., and Zhang, B.B.** (2002). Salicylic Acid Reverses Phorbol 12-Myristate-13-Acetate (PMA)- and Tumor Necrosis Factor alpha (TNFalpha)-induced Insulin Receptor Substrate 1 (IRS1) Serine 307 Phosphorylation

and Insulin Resistance in Human Embryonic Kidney 293 (HEK293) Cells. *J. Biol. Chem.* **278**: 180–186.

**Kamisaka, S., Yanagishima, N., and Masuda, Y.** (1967). Effect of Auxin and Gibberellin on Sporulation in Yeast. *Physiol. Plant.* **20**: 90–97.

**Kamoun, S.** (2007). Groovy times: filamentous pathogen effectors revealed. *Curr. Opin. Plant Biol.* **10**: 358–365.

**Kępczyńska, E.** (1994). Involvement of ethylene in spore germination and mycelial growth of *Alternaria alternata*. *Mycol. Res.* **98**: 118–120.

**Kępczyńska, E.** (1989). Ethylene requirement during germination of *Botrytis cinerea* spores. *Physiol. Plant.* **77**: 369–372.

**Kettner, J. and Dorffling, K.** (1995). Biosynthesis and Metabolism of Abscisic-Acid in Tomato Leaves Infected with *Botrytis cinerea*. *Planta* **196**: 627–634.

**Kilaru, A., Bailey, B. a., and Hasenstein, K.H.** (2007). *Moniliophthora perniciosa* produces hormones and alters endogenous auxin and salicylic acid in infected cocoa leaves. *FEMS Microbiol. Lett.* **274**: 238–244.

**Kim, J.I., Murphy, A.S., Baek, D., Lee, S.W., Yun, D.J., Bressan, R. a., and Narasimhan, M.L.** (2011). YUCCA6 over-expression demonstrates auxin function in delaying leaf senescence in *Arabidopsis thaliana*. *J. Exp. Bot.* **62**: 3981–3992.

**Kisiala, A., Laffont, C., Emery, R.J.N., and Frugier, F.** (2013). Bioactive cytokinins are selectively secreted by *Sinorhizobium meliloti* nodulating and nonnodulating strains. *Mol. Plant. Microbe. Interact.* **26**: 1225–31.

**Kloppholz, S., Kuhn, H., and Requena, N.** (2011). A secreted fungal effector of glomus intraradices promotes symbiotic biotrophy. *Curr. Biol.* **21**: 1204–1209.

**Kovač, M. and Žel, J.** (1995). The effect of aluminum on cytokinins in the mycelia of *Amanita muscaria*. *J. Plant Growth Regul.* **14**: 117–120.

**Król, P., Igielski, R., Pollmann, S., and Kępczyńska, E.** (2015). Priming of seeds with methyl jasmonate induced resistance to hemi-biotroph *Fusarium oxysporum* f.sp. *lycopersici* in tomato via 12-oxo-phytodienoic acid, salicylic acid, and flavonol accumulation. *J. Plant Physiol.* **179**: 122–132.

**Kulkarni, G.B., Sanjeevkumar, S., Kirankumar, B., Santoshkumar, M., and Karegoudar, T.B.** (2013). Indole-3-Acetic Acid Biosynthesis in *Fusarium delphinoides* Strain GPK, a Causal Agent of Wilt in Chickpea. *Appl. Biochem. Biotechnol.* **169**: 1292–1305.

**Kurakawa, T., Ueda, N., Maekawa, M., Kobayashi, K., Kojima, M., Nagato, Y., Sakakibara, H., and Kyojuka, J.** (2007). Direct control of shoot meristem activity by a cytokinin-activating enzyme. *Nature* **445**: 652–655.

**Laurans, F., Pepin, R., and Gay, G.** (2001). Fungal auxin overproduction affects the anatomy of *Hebeloma cylindrosporum*-*Pinus pinaster* ectomycorrhizas. *Tree Physiol.* **21**: 533–540.

**Lee, B.O.** (1961). Effect of Kinetin on the Fertility of some Strains of *Neurospora crassa*. *Nature* **192**: 288–288.

**LeJohn, H.B. and Stevenson, R.M.** (1973). Cytokinins and magnesium ions may control the flow of metabolites and calcium ions through fungal cell membranes. *Biochim. Biophys. Res. Commun.* **54**: 1061–1066.

**Mańka, M.** (1980). Auxin and gibberellin -like substances synthesis by *Fusarium* isolates. *Acta Microbiol Pol* **29**: 365–374.

**Maor, R., Haskin, S., Levi-kedmi, H., and Sharon, A.** (2004). In Planta Production of Indole-3-Acetic Acid by *Colletotrichum gloeosporioides* f. sp. *aeschynomene*. *Society* **70**: 3–6.

**McArthur, D. a and Knowles, N.R.** (1992). Resistance Responses of Potato to Vesicular-Arbuscular Mycorrhizal Fungi under Varying Abiotic Phosphorus Levels. *Plant Physiol.* **100**: 341–351.

**Meixner, C., Ludwig-Müller, J., Miersch, O., Gresshoff, P., Staehelin, C., and Vierheilig, H.** (2005). Lack of mycorrhizal autoregulation and phytohormonal changes in the supernodulating soybean mutant nts1007. *Planta* **222**: 709–715.

**Miersch, O., Brückner, B., Schmidt, J., and Sembdner, G.** (1992). Cyclopentane fatty acids from *Gibberella fujikuroi*. *Phytochemistry* **31**: 3835–3837.

**Miersch, O., Schneider, G., and Sembdner, G.** (1991). Hydroxylated jasmonic acid and related compounds from *Botryodiplodia theobromae*. *Phytochemistry* **30**: 4049–4051.

**Mills, L.J. and Van Staden, J.** (1978). Extraction of cytokinins from maize , smut tumors of maize and *Ustilago maydis* cultures. *Physiol. Plant Pathol.* **13**: 73–80.

**Morrison, E.N., Emery, R.J.N., and Saville, B.J.** (2015a). Phytohormone Involvement in the *Ustilago maydis*– *Zea mays* Pathosystem: Relationships between Abscisic Acid and Cytokinin Levels and Strain Virulence in Infected Cob Tissue. *PLoS One* **10**: e0130945.

**Morrison, E.N., Knowles, S., Hayward, a., Thorn, R.G., Saville, B.J., and Emery, R.J.N.** (2015b). Detection of phytohormones in temperate forest fungi predicts consistent abscisic acid production and a common pathway for cytokinin biosynthesis. *Mycologia* **107**: 245–257.

**Murphy, A.M., Pryce-Jones, E., Johnstone, K., and Ashby, A.M.** (1997). Comparison of



cytokinin production in vitro by *Pyrenopeziza brassicae* with other plant pathogens.

Physiol. Mol. Plant Pathol. **50**: 53–65.

**Nakamura, T., Kawanabe, Y., Takiyama, E., Takahashi, N., and Murayama, T.** (1978).

Effects of auxin and gibberellin on conidial germination in *Neurospora crassa*. Plant Cell

Physiol. **19**: 705–709.

**Nakamura, T., Tomita, K., Kawanabe, Y., and Murayama, T.** (1982). Effects of auxin and

gibberellin on conidial germination in *Neurospora crassa* II. “conidial density effect” and

auxin. Plant Cell Physiol. **23**: 1363–1369.

**Nambara, E. and Marion-Poll, A.** (2005). Abscisic Acid Biosynthesis and Catabolism.

Annu. Rev. Plant Biol. **56**: 165–185.

**Norman, S.M., Bennett, R.D., Maier, V.P., and Poling, S.M.** (1983). Cytokinins Inhibit

Abscisic Acid Biosynthesis in *Cercospora Rosicola*. Plant Sci. Lett. **28**: 255–263.

**Oka, M., Miyamoto, K., Okada, K., and Ueda, J.** (1999). Auxin polar transport and flower

formation in *Arabidopsis thaliana* transformed with indoleacetamide hydrolase (*iaaH*)

gene. Plant Cell Physiol. **40**: 231–237.

**Oritani, T. and Kiyota, H.** (2003). Biosynthesis and metabolism of abscisic acid and related

compounds. Nat. Prod. Rep. **20**: 414–425.

**Packer, N.M. and Steward, M.W.** (1967). Studies On the biosynthesis of Phenols by Fungi.

Biochemistry: 122–132.

**Panahirad, S., Zaare-Nahandi, F., Mohammadi, N., Alizadeh-Salteh, S., and Safaie, N.**

(2014). Effects of salicylic acid on *Aspergillus flavus* infection and aflatoxin B<sub>1</sub>

accumulation in pistachio (*Pistacia vera* L.) fruit. J. Sci. Food Agric. **94**: 1758–1763.

**Patkar, R.N., Benke, P.I., Qu, Z., Constance Chen, Y.Y., Yang, F., Swarup, S., and**

**Naqvi, N.I.** (2015). A fungal monooxygenase-derived jasmonate attenuates host innate immunity. *Nat. Chem. Biol.*: 1–10.

**Payton, S., Fray, R.G., Brown, S., and Grierson, D.** (1996). Ethylene receptor expression is regulated during fruit ripening, flower senescence and abscission. *Plant Mol. Biol.* **31**: 1227–1231.

**Peleg, Z. and Blumwald, E.** (2011). Hormone balance and abiotic stress tolerance in crop plants. *Curr. Opin. Plant Biol.* **14**: 290–5.

**Peleg, Z., Reguera, M., Tumimbang, E., Walia, H., and Blumwald, E.** (2011). Cytokinin-mediated source/sink modifications improve drought tolerance and increase grain yield in rice under water-stress. *Plant Biotechnol. J.* **9**: 747–758.

**Pertry, I. et al.** (2009). Identification of *Rhodococcus fascians* cytokinins and their modus operandi to reshape the plant. *Proc. Natl. Acad. Sci. U. S. A.* **106**: 929–34.

**Pimenta Lange, M.J. and Lange, T.** (2006). Gibberellin biosynthesis and the regulation of plant development. *Plant Biol.* **8**: 281–290.

**Pozo, M.J., López-Ráez, J.A., Azcón-Aguilar, C., and García-Garrido, J.M.** (2015). Phytohormones as integrators of environmental signals in the regulation of mycorrhizal symbioses. *New Phytol.* **205**: 1431–1436.

**Rani Debi, B., Taketa, S., and Ichii, M.** (2005). Cytokinin inhibits lateral root initiation but stimulates lateral root elongation in rice (*Oryza sativa*). *J. Plant Physiol.* **162**: 507–15.

**Rao, R.P., Hunter, A., Kashpur, O., and Normanly, J.** (2010). Aberrant synthesis of indole-3-acetic acid in *Saccharomyces cerevisiae* triggers morphogenic transition, a virulence trait of pathogenic fungi. *Genetics* **185**: 211–220.

**Reineke, G., Heinze, B., Schirawski, J., Buettner, H., Kahmann, R., and Basse, C.W.**

(2008). Indole-3-acetic acid (IAA) biosynthesis in the smut fungus *Ustilago maydis* and its relevance for increased IAA levels in infected tissue and host tumour formation. *Mol. Plant Pathol.* **9**: 339–355.

**Riou-Khamlichi, C., Huntley, R., Jacquard, A., and Murray, J.A.H.** (1999). Cytokinin Activation of Arabidopsis Cell Division Through a D-Type Cyclin. *Science* (80-. ). **283**: 1541–1545.

**Robert-Seilaniantz, A., Grant, M., and Jones, J.D.G.** (2011). Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annu. Rev. Phytopathol.* **49**: 317–43.

**Robert-Seilaniantz, A., Navarro, L., Bari, R., and Jones, J.D.G.** (2007). Pathological hormone imbalances. *Curr. Opin. Plant Biol.* **10**: 372–9.

**Robinson, M., Riou, J., and Sharon, A.** (1998). Indole-3-acetic acid biosynthesis in *Colletotrichum gloeosporioides* f. sp. *aeschynomene*. *Appl. Environ. Microbiol.* **64**: 0–5030.

**Sakakibara, H.** (2006). Cytokinins: activity, biosynthesis, and translocation. *Annu. Rev. Plant Biol.* **57**: 431–49.

**Sanders, I.R.** (2011). Mycorrhizal symbioses: How to be seen as a good fungus. *Curr. Biol.* **21**: R550–R552.

**Sharaf, E.F. and Farrag, A.A.** (2004). Induced Resistance in Tomato Plants by IAA against *Fusarium oxysporum lycopersici*. *Pol. J. Microbiol.* **53**: 111–116.

**Siddique, S. et al.** (2015). A parasitic nematode releases cytokinin that controls cell division and orchestrates feeding site formation in host plants. *Proc. Natl. Acad. Sci.*: 201503657.

**Siemens, J., Keller, I., Sarx, J., Kunz, S., Schuller, A., Nagel, W., Schmülling, T., Parniske, M., and Ludwig-Müller, J.** (2006). Transcriptome analysis of Arabidopsis

clubroots indicate a key role for cytokinins in disease development. *Mol. Plant. Microbe Interact.* **19**: 480–94.

**Siewers, V., Kokkelink, L., Smedsgaard, J., and Tudzynski, P.** (2006). Identification of an abscisic acid gene cluster in the grey mold *Botrytis cinerea*. *Appl. Environ. Microbiol.* **72**: 4619–4626.

**Slaugenhaupt, S. a, Mull, J., Leyne, M., Cuajungco, M.P., Gill, S.P., Hims, M.M., Quintero, F., Axelrod, F.B., and Gusella, J.F.** (2004). Rescue of a human mRNA splicing defect by the plant cytokinin kinetin. *Hum. Mol. Genet.* **13**: 429–36.

**Spence, C. and Bais, H.** (2015). Role of plant growth regulators as chemical signals in plant–microbe interactions: a double edged sword. *Curr. Opin. Plant Biol.* **27**: 52–58.

**Spence, C.A., Lakshmanan, V., Donofrio, N., and Bais, H.P.** (2015). Crucial Roles of Abscisic Acid Biogenesis in Virulence of Rice Blast Fungus *Magnaporthe oryzae*. *Front. Plant Sci.* **6**: 1–13.

**Splivallo, R., Fischer, U., Göbel, C., Feussner, I., and Karlovsky, P.** (2009). Truffles regulate plant root morphogenesis via the production of auxin and ethylene. *Plant Physiol.* **150**: 2018–2029.

**Strzelczyk, E., Kampert, M., and Pachlewski, R.** (1994). The influence of pH and temperature on ethylene production by mycorrhizal fungi of pine. *Mycorrhiza* **4**: 193–196.

**Studt, L., Schmidt, F.J., Jahn, L., Sieber, C.M.K., Connolly, L.R., Niehaus, E.-M., Freitag, M., Humpf, H.-U., and Tudzynski, B.** (2013). Two Histone Deacetylases, FfHda1 and FfHda2, Are Important for *Fusarium fujikuroi* Secondary Metabolism and Virulence. *Appl. Environ. Microbiol.* **79**: 7719–7734.

- Swain, S.M. and Singh, D.P.** (2005). Tall tales from sly dwarves: Novel functions of gibberellins in plant development. *Trends Plant Sci.* **10**: 123–129.
- Tomita, K., Murayama, T., and Nakamura, T.** (1984). Effects of auxin and gibberellin on elongation of young hyphae in *Neurospora crassa*. *Plant Cell Physiol.* **25**: 355–358.
- Ton, J., Flors, V., and Mauch-Mani, B.** (2009). The multifaceted role of ABA in disease resistance. *Trends Plant Sci.* **14**: 310–317.
- Tsavkelova, E. a., Klimova, S.Y., Cherdyntseva, T. a., and Netrusov, a. I.** (2006). Microbial producers of plant growth stimulators and their practical use: A review. *Appl. Biochem. Microbiol.* **42**: 117–126.
- Tsavkelova, E., Oeser, B., Oren-Young, L., Israeli, M., Sasson, Y., Tudzynski, B., and Sharon, A.** (2012). Identification and functional characterization of indole-3-acetamide-mediated IAA biosynthesis in plant-associated *Fusarium* species. *Fungal Genet. Biol.* **49**: 48–57.
- Tsitsigiannis, D.I. and Keller, N.P.** (2007). Oxylipins as developmental and host–fungal communication signals. *Trends Microbiol.* **15**: 109–118.
- Tudzynski, B.** (2005). Gibberellin biosynthesis in fungi: genes, enzymes, evolution, and impact on biotechnology. *Appl. Microbiol. Biotechnol.* **66**: 597–611.
- Ulrich, J.M.** (1960). Auxin production by mycorrhizal fungi. *Physiol. Plant.* **13**: 429–443.
- Vanneste, S.** (2005). Auxin coordinates cell division and cell fate specification during lateral root initiation. *Physiol. Plant.* **123**: 139–146.
- De Vleeschauwer, D., Gheysen, G., and Höfte, M.** (2013). Hormone defense networking in rice: tales from a different world. *Trends Plant Sci.* **18**: 555–65.
- De Vleeschauwer, D., Xu, J., and Höfte, M.** (2014). Making sense of hormone-mediated

defense networking: from rice to Arabidopsis. *Front. Plant Sci.* **5**: 1–15.

**De Vleesschauwer, D., Yang, Y., Cruz, C.V., and Höfte, M.** (2010). Abscisic acid-induced resistance against the brown spot pathogen *Cochliobolus miyabeanus* in rice involves MAP kinase-mediated repression of ethylene signaling. *Plant Physiol.* **152**: 2036–52.

**Wingler, A., Leegood, R.C., Lea, P.J., and Quick, W.P.** (1998). Regulation of Leaf Senescence by Cytokinin, Sugars, and Light. *Plant Physiol.* **116**: 329–335.

**Wullschleger, S.D. and Reid, C.P.P.** (1990). Implication of Ectomycorrhizal Fungi in the Cytokinin Relations of Loblolly Pine (*Pinus-Taeda* L). *New Phytol* **116**: 681–688. *New Phytol.*: 681–688.

**Xu, J., Audenaert, K., Hofte, M., and de Vleesschauwer, D.** (2013). Abscisic Acid Promotes Susceptibility to the Rice Leaf Blight Pathogen *Xanthomonas oryzae* pv *oryzae* by Suppressing Salicylic Acid-Mediated Defenses. *PLoS One* **8**.

**Yanagishima, N.** (1965). Role of Gibberellic Acid in the Growth Response of Yeast Cells to Auxin. *Physiol. Plant.* **18**: 306–312.

**Yin, C., Park, J.-J., Gang, D.R., and Hulbert, S.H.** (2014). Characterization of a tryptophan 2-monooxygenase gene from *Puccinia graminis* f. sp. *tritici* involved in auxin biosynthesis and rust pathogenicity. *Mol. Plant. Microbe. Interact.* **27**: 227–35.

**Zhao, Y.** (2010). Auxin Biosynthesis and Its Role in Plant Development. *Annu. Rev. Plant Biol.* **61**: 49–64.

**Zsögön, A., Lambais, M.R., Benedito, V.A., Figueira, A.V.D.O., and Peres, L.E.P.**

(2008). Reduced arbuscular mycorrhizal colonization in tomato ethylene mutants. *Sci. Agric.* **65**: 259–267.




Figure 1. Schematic representation of the effects of hormones on fungal biology.

A schematic view of yeast-like (light grey background), infecting and sporulating filamentous fungus is given. This model summarizes the effects of hormones on fungal biological processes reported to date; the effects due to the dose, the molecule tested, other environmental factors as well as the specie studied are provided in the text and could differ among fungi. Arrows and « T » bars respectively represent positive and negative effects. AUX : Auxins, ABA : Abscisic Acid, ET : Ethylene, CKs : Cytokinins, GAs : Gibberellic Acids, JA : Jasmonic acid, SA : Salicylic Acid.

254x190mm (96 x 96 DPI)

Acce]
   
 Accepted Article