



**HAL**  
open science

# Identification of the Key Weather Factors Affecting Overwintering Success of *Apolygus lucorum* Eggs in Dead Host Tree Branches

Hongsheng Pan, Bing Liu, Yanhui Lu, Nicolas Desneux

► **To cite this version:**

Hongsheng Pan, Bing Liu, Yanhui Lu, Nicolas Desneux. Identification of the Key Weather Factors Affecting Overwintering Success of *Apolygus lucorum* Eggs in Dead Host Tree Branches. PLoS ONE, 2014, 9 (4), 10.1371/journal.pone.0094190 . hal-02636580

**HAL Id: hal-02636580**

**<https://hal.inrae.fr/hal-02636580>**

Submitted on 27 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Identification of the Key Weather Factors Affecting Overwintering Success of *Apolygus lucorum* Eggs in Dead Host Tree Branches

Hongsheng Pan<sup>1</sup>, Bing Liu<sup>1</sup>, Yanhui Lu<sup>1\*</sup>, Nicolas Desneux<sup>2</sup>

**1** State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China, **2** French National Institute for Agricultural Research (INRA), Sophia-Antipolis, France

## Abstract

Understanding the effects of weather on insect population dynamics is crucial to simulate and forecast pest outbreaks, which is becoming increasingly important with the effects of climate change. The mirid bug *Apolygus lucorum* is an important pest on cotton, fruit trees and other crops in China, and primarily lays its eggs on dead parts of tree branches in the fall for subsequent overwintering. As such, the eggs that hatch the following spring are most strongly affected by ambient weather factors, rather than by host plant biology. In this study, we investigated the effects of three major weather factors: temperature, relative humidity and rainfall, on the hatching rate of *A. lucorum* eggs overwintering on dead branches of Chinese date tree (*Ziziphus jujuba*). Under laboratory conditions, rainfall (simulated via soaking) was necessary for the hatching of overwintering *A. lucorum* eggs. In the absence of rainfall (unsoaked branches), very few nymphs successfully emerged under any of the tested combinations of temperature and relative humidity. In contrast, following simulated rainfall, the hatching rate of the overwintering eggs increased dramatically. Hatching rate and developmental rate were positively correlated with relative humidity and temperature, respectively. Under field conditions, the abundance of nymphs derived from overwintering eggs was positively correlated with rainfall amount during the spring seasons of 2009–2013, while the same was not true for temperature and relative humidity. Overall, our findings indicate that rainfall is the most important factor affecting the hatching rate of overwintering *A. lucorum* eggs on dead plant parts and nymph population levels during the spring season. It provides the basic information for precisely forecasting the emergence of *A. lucorum* and subsequently timely managing its population in spring, which will make it possible to regional control of this insect pest widely occurring in multiple crops in summer.

**Citation:** Pan H, Liu B, Lu Y, Desneux N (2014) Identification of the Key Weather Factors Affecting Overwintering Success of *Apolygus lucorum* Eggs in Dead Host Tree Branches. PLoS ONE 9(4): e94190. doi:10.1371/journal.pone.0094190

**Editor:** Guy Smagghe, Ghent University, Belgium

**Received:** January 23, 2014; **Accepted:** March 11, 2014; **Published:** April 4, 2014

**Copyright:** © 2014 Pan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This research was supported by the Special Fund for Agro-scientific Research in the Public Interest (201103012) and the National Natural Science Funds (No. 31222046, 31321004). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: yhlu@ippcaas.cn

## Introduction

Understanding insect population dynamics can be very challenging due to the large number of biotic and abiotic factors that must be taken into account. Of these factors, weather is generally considered to be one of the most important. For instance, temperature, rainfall, relative humidity and other weather-related factors can strongly affect the survival, development, fecundity and behavior of individual insects [1,2,3,4,5], and the population dynamics of insects [6,7,8,9]. In addition to such direct effects, weather can indirectly impact insect populations by affecting life-history traits of their host plants, competitors, and natural enemies [10,11,12].

Global climate change is influencing the geographical distributions and phenology of many insect species [13,14,15,16], some of which are major crop pests. Such changes will inevitably lead to the appearance of new pests in some areas and changes to the biology of well established pests in others [17,18,19,20], Guo et al. [21] found that higher temperatures can lead to earlier egg hatching and eclosion in three Inner Mongolian grasshopper species [*Dasyhippus barbipes* (Fischer de Waldheim), *Oedaleus asiaticus*

B. Bienko, and *Chorthippus fallax* (Zubovskiy)], which will likely extend their distribution northward as climate change continues. Another good example is the diamondback moth. Indeed, it has been estimated that *Plutella xylostella* (L.) should be able to complete two additional generations for every 2°C of environmental warming [22].

*Apolygus lucorum* (Meyer-Dür) (Hemiptera: Miridae) is the dominant mirid bug species found on many crop species in Northern China, including cotton, Chinese dates, apples, pears, and grapes [23,24]. Both adults and nymphs feed on plant terminal meristems, young squares and fruits (bolls for cotton), and various other tissues, and results in bushy plants and abscission of squares and fruits (bolls). Eggs are mainly laid inside plant tissues. In each October, *A. lucorum* adults relocate to specific host plants for ovipositing and overwintering eggs [25]. Although *A. lucorum* eggs can successfully overwinter on at least 86 plant species, adults prefer to deposit eggs on fruit trees, such as the Chinese date *Ziziphus jujuba* Mill. and the grape *Vitis vinifera* L. [26]. In April, the overwintering eggs begin to hatch and the newly emerged nymphs feed on tender leaves, buds and flowers. It can seriously damage fruit trees; in particular, the highly mobile nymphs can induce

significant cryptic damage which is difficult to be detected, during the initial period of feeding and infestation [27]. Therefore, it is important to determine the factors that affect the hatching rate of overwintering *A. lucorum* eggs in order to develop more precise forecasting techniques.

*A. lucorum* adults generally lay eggs for overwintering in the dead parts of tree branches [26] where the eggs are less likely to be affected by plant growth, which makes weather conditions the most important factor evoking egg hatching. Previous studies showed that temperature and relative humidity strongly affect survival, development, fecundity, and life-history traits of laboratory populations of this insect [28,29,30,31]. In the field, *A. lucorum* as well as other mirid bugs, prefer shady and moist environments [31], and it has been reported that periods of high rainfall result in rapid increases of their populations during the summer [23,31,32,33,34,35,36]. However, the precise effects of factors such as temperature, relative humidity and rainfall on the hatching rate of overwintering *A. lucorum* eggs have not been characterized. The aim of this study was to test the hypothesis that rainfall or other weather factors played a pivotal role in the hatching of *A. lucorum* eggs overwintered in dead host plant (tree) tissues, and then select one or two key weather factors for future forecasting *A. lucorum* population in spring season. We quantified the importance of weather variables on *A. lucorum* hatching by combining with laboratory trials and field surveys.

## Materials and Methods

### Ethics Statement

No specific permits were required for the described field studies.

### Collection of overwintering eggs

Field sampling was carried out in a 1-hectare Chinese date orchard (116.6 °E, 39.5 °N) at the Langfang Experiment Station, Chinese Academy of Agricultural Sciences (CAAS), in Hebei Province, China (116.4 °E, 39.3 °N). Overwintering *A. lucorum* eggs are primarily laid in the summer-pruning wounds of Chinese date trees [25,35,37]. During early April, dead branches containing summer-pruning wounds and overwintering eggs were collected from the Chinese date trees and placed in one of four baskets (0.2 m length \* 0.2 m width \* 0.2 m height) made of bamboo splints. Each basket was affixed with wire to a main branch of a Chinese date tree keeping order to maintain the overwintering eggs under the actual environmental conditions found in the orchard. To eliminate the effects of rain, the baskets were covered

with plastic film (0.15 mm thickness, Shandong Jinan Ninglu Economic Plastic Industry Co., Ltd.) prior to each period of rainfall over the course of the experiment; the baskets were then uncovered after the rain had ended.

### Hatching dynamics of overwintering eggs in laboratory conditions

Overwintering *A. lucorum* eggs laid on Chinese date trees primarily hatch between late April and early May in the region of northern China [35,37]. Therefore, we assessed the effects of various environmental factors on the hatching of overwintering eggs over three time periods: mid April (11<sup>th</sup> April), late April (21<sup>st</sup> April), and early May (1<sup>st</sup> May). At each of these time points, the Chinese date branches containing the overwintering eggs were sampled from the baskets in the orchard, and the numbers of eggs laid on each branch were determined by microscopic examination (SZX16, Olympus, Japan) in the laboratory. To simulate the effects of rain, half of the branches were fully submerged in distilled water for 4 h; the other half of the branches were left dry to act as controls. The treated branches were then placed separately into glass vials (5 cm height × 1.5 cm diameter) and sealed with nylon mesh. Next, the glass vials were placed into environmental growth chambers (Ningbo Jiangnan Instrument Factory, Ningbo, China) adjusted to one of nine combinations of temperature (15, 20, and 25 °C) and relative humidity (40, 55, and 70%) according to a completely randomized design with 4 replicates. All chambers were subjected to a photoperiod of 14:10 (L:D) h, and a total of 100 eggs were tested per replicate. Nymph emergence was recorded daily for 60 d, and newly emerged nymphs were removed from the glass vials. *A. lucorum* nymphs were identified following the keys reported by Lu and Wu [25].

### Hatching dynamics of overwintering eggs in field conditions


Field experiments were conducted at the Chinese date orchard at the CAAS Langfang Experiment Station over five consecutive spring seasons (April 1<sup>st</sup> through May 31<sup>st</sup> from 2009 to 2013). No insecticides were applied over the course of the study period. The hatching dynamics of the overwintering *A. lucorum* eggs were monitored every 1–3 days of the observation period. During each sampling, 100–200 branches (30 cm long) with tender leaves were randomly selected and surveyed for the presence of *A. lucorum* nymphs at different instar stages. The emergence of 1<sup>st</sup>-instar

**Table 1.** *F*- and *P*-values from the Analyses of Variance tests of the effects of temperature, relative humidity and rainfall (simulated via soaking) on the hatching rate of overwintering *A. lucorum* eggs.

Source of variation	df	<i>F</i> value ( <i>P</i> value)		
		Mid April	Late April	Early May
Temperature	2	18.59 (< 0.0001)	15.67 (< 0.0001)	3.36 (0.0335)
Relative humidity	2	35.39 (< 0.0001)	35.59 (< 0.0001)	12.74 (< 0.0001)
Rainfall	1	407.26 (< 0.0001)	319.49 (< 0.0001)	215.07 (< 0.0001)
Temperature * Relative humidity	4	1.32 (0.2764)	2.21 (0.0810)	0.09 (0.9852)
Temperature * Rainfall	2	16.83 (< 0.0001)	14.24 (< 0.0001)	4.95 (0.0109)
Relative humidity * Rainfall	2	29.42 (< 0.0001)	17.10 (< 0.0001)	10.96 (0.0001)
Temperature * Relative humidity * Rainfall	4	1.28 (0.2886)	0.88 (0.4835)	0.19 (0.9445)

df = degrees of freedom.

doi:10.1371/journal.pone.0094190.t001



**Figure 1. Hatching rates of overwintering *A. lucorum* eggs under different laboratory conditions.** (a) Mid April. (b) Late April. (c) Early May.

doi:10.1371/journal.pone.0094190.g001

nymphs was used to calculate the hatching dynamics of the overwintering eggs.

*Artemisia annua* L., *Artemisia argyi* Lévl. et Vant., *Artemisia lavandulaefolia* DC., and *Humulus scandens* (Lour.) Merr. are the four major host plants of *A. lucorum* adults of last generation prior to egg laying [38], whereas adult females prefer to lay eggs on tree hosts (e.g., Chinese date trees) for overwintering [26]. Therefore, adult population levels on these four fall plants hosts could indirectly influence the abundance of overwintering eggs in nearby

tree hosts [26]. To account for this, adult *A. lucorum* population levels on *A. annua*, *A. argyi*, *A. lavandulaefolia*, and *H. scandens* plants near the Chinese date orchard were surveyed using knock-down techniques during early and mid September 2008–2012 [38]. Knock-down techniques basically consisted of pulling parts of the plants over a rectangular 40 cm × 26 cm × 11-cm white-colored pan, after which plant material was struck four times, and the number of dislodged individuals was counted [39].

**Table 2.** *F*- and *P*-values from the Analyses of Variance tests of the effects of temperature, relative humidity and sampling period on the hatching rate and developmental duration of overwintering *A. lucorum* eggs with simulated rainfall (via soaking).

Source of variation	df	<i>F</i> value ( <i>P</i> value)	
		Hatching rate	Developmental duration
Temperature	2	21.25 (< 0.0001)	315.81 (< 0.0001)
Relative humidity	2	45.98 (< 0.0001)	4.70 (0.0119)
Sampling period	2	48.11 (< 0.0001)	61.47 (< 0.0001)
Temperature * Relative humidity	4	0.78 (0.5436)	5.65 (0.0005)
Temperature * Sampling period	4	0.55 (0.7024)	1.67 (0.1667)
Relative humidity * Sampling period	4	2.15 (0.0821)	3.44 (0.0121)
Temperature * Relative humidity * Sampling period	8	0.25 (0.9788)	0.72 (0.6692)

df = degrees of freedom

doi:10.1371/journal.pone.0094190.t002

### Data analysis

A three-way ANOVA test was used to analyze the effects of temperature, relative humidity, rainfall and their interactions, on *A. lucorum* egg hatching. We also analyzed the effects and interactions of temperature, relative humidity, and sampling period on egg hatching rate and developmental duration. One-way ANOVA was used to analyze the effects of temperature or relative humidity on developmental duration. Prior to analysis, egg hatching rates were transformed using the arcsine square root, and developmental duration data were transformed using  $\log_{10}(n+1)$  in order to satisfy the assumption of normality.

Data on temperature, relative humidity and rainfall at the Langfang location between 2009 and 2013 were collected by the Chinese Meteorological Data Sharing Service (<http://cdc.cma.gov.cn/>). A Simple linear model ( $y = a + bx$ ) was used for analyzing the relationship between 1<sup>st</sup>-instar *A. lucorum* nymph abundance and (1) all nymphs recorded during that spring (late April to mid May); (2) each weather factor (temperature, relative humidity and rainfall amount) during that spring (mid April to mid

May); and (3) adult population levels on different host plants during the previous fall. All the above periods selected for different parameters (inc. *A. lucorum* abundance and weather factors) were determined according to the hatching dynamics of overwintering eggs during the whole study. Prior to analysis, the population numbers of the fall adults and spring nymphs were  $\log_{10}(n+1)$  transformed.

### Results

#### Hatching dynamics of overwintering eggs in laboratory conditions


Temperature, relative humidity, and rainfall, as well as their interaction, affected the hatching rate of overwintering *A. lucorum* eggs ( $P < 0.05$ ) (Table 1). Of these factors, rainfall was the most important for egg hatching, since the *F* values for rainfall in the ANOVA tests ranged from 215.07–407.26 during the three study periods (mid April, late April, and early May), while those for temperature and relative humidity ranged from 3.36–18.59 and

**Table 3.** The developmental duration of overwintering *A. lucorum* eggs under different combinations of temperature and relative humidity with simulated rainfall (via soaking).

Sampling period	Temperature (°C)	Relative humidity (%)		
		40	55	70
Middle April	15	40.03±3.04 aA	22.49±1.71 aC	29.07±0.60 aB
	20	17.54±1.34 bA	13.57±0.66 bA	15.12±1.03 bA
	25	10.13±0.14 cA	10.55±0.48 cA	12.11±0.85 bA
Late April	15	31.83±2.56 aA	23.47±1.16 aB	23.67±0.62 aB
	20	13.43±1.15 bA	12.26±0.69 bA	10.36±0.70 bA
	25	9.14±0.37 cB	11.61±0.91 bA	8.87±0.18 bB
Early May	15	23.24±5.85 aA	18.56±0.70 aA	17.33±1.13 aA
	20	10.41±2.03 abA	10.62±0.97 bA	10.40±0.19 bA
	25	6.20±0.46 bA	6.62±0.27 cA	6.66±0.38 cA

Note: Different lowercase letters indicate significant differences between the temperature treatments under the same relative humidity within one sampling period; different uppercase letters indicate significant differences between relative humidity treatments at the same temperature within one sampling period (LSD test;  $P < 0.05$ ).

doi:10.1371/journal.pone.0094190.t003



**Figure 2. Hatching dynamics of overwintering *A. lucorum* eggs soaked for 4 h under different laboratory conditions.** (a) Mid April. (b) Late April. (c) Early May.  
doi:10.1371/journal.pone.0094190.g002


12.74–35.59, respectively. Moreover, under conditions of no rainfall, very few nymphs emerged (Figure 1).

We found that egg-hatching rate increased following rainfall, as well as with higher temperatures and relative humidity. Egg-hatching rate also varied according to sampling time. The hatching rate of overwintering eggs sampled in early May was higher than during mid and late April ( $P < 0.05$ ), whereas no difference in hatching rate was observed between the latter two periods ( $P > 0.05$ ) (Figure 1, Table 2). Furthermore, both temperature and relative humidity significantly affected the developmental duration of the overwintering eggs ( $P < 0.05$ ), with temperature playing a stronger role ( $F$  value = 315.81) over relative humidity ( $F$  value = 4.70). In other words, high temperatures could hasten the development of overwintering eggs. The developmental duration of the overwintering eggs was different between each sampling period ( $P < 0.05$ ) (Table 2, Table 3). For example, the eggs sampled in mid April showed the longest developmental periods as well as the greatest variation in developmental duration between individuals. In contrast, the shortest developmental periods and the least

variation in developmental duration between individuals were observed in early May (Table 3, Figure 2).

#### Hatching dynamics of overwintering eggs in field conditions

Between 2009 and 2013, a variety of weather conditions (rainfall, temperature, and relative humidity) that were observed have affected the hatching dynamics of overwintering *A. lucorum* eggs and the population levels of newly emerged nymphs (Figure 3). Egg hatching began following a period of rainfall in late April and peaked in early May. The levels of newly emerged nymphs, as well as nymphs of all stages, were positively correlated with rainfall abundance during that spring ( $P < 0.05$ ). Neither temperature or relative humidity during the same spring, nor adult abundance on any of the four studied host plants during the previous fall, were correlated with nymph population levels ( $P > 0.05$ ) (Table 4, Figure 4).



**Figure 3. Hatching dynamics of overwintering *A. lucorum* eggs under field conditions during April-May.** (a) 2009. (b) 2010. (c) 2011. (d) 2012. (e) 2013. doi:10.1371/journal.pone.0094190.g003

**Discussion**

The number of newly emerged *A. lucorum* nymphs in the spring is determined by the initial number of overwintering eggs laid in the fall, the survival rate over the winter (which is dependent on weather conditions), and the hatching rate during the spring season. Very few eggs hatched in the absence of soaking or under

conditions of low relative humidity (e.g. 40% RH) following soaking (Figure 1). Consistent with the importance of rainfall and relative humidity for egg hatching and nymph survival [29], the number of 1<sup>st</sup>-instar *A. lucorum* nymphs, as well as nymphs of all stages, was positively correlated with rainfall in the field; no relationships were observed between nymph population levels and the other studied factors. For example, compared with 2011 and

**Table 4.** *P* values from simple linear regression analyses of the relationships between *A. lucorum* nymph abundance in the spring and (1) weather conditions in the spring and (N) and (2) *A. lucorum* adult abundance the previous fall (N-1).

Period	Parameters	Spring of year (N)	
		Abundance of 1 <sup>st</sup> instar nymphs	Abundance of all nymphs
Fall of year (N-1)	Adult abundance in <i>Artemisia argyi</i>	0.4787	0.6042
	Adult abundance in <i>Artemisia annua</i>	0.8060	0.8311
	Adult abundance in <i>Artemisia lavandulaefolia</i>	0.6806	0.5035
	Adult abundance in <i>Humulus scandens</i>	0.3397	0.8075
	Mean adult abundance in these four host plants	0.5011	0.6353
Spring of year (N)	Temperature	0.1404	0.3643
	Relative humidity	0.6782	0.8362
	Rainfall amount	0.0262	0.0162

Note: Fall was defined as the period between early and mid September, and spring as the period between mid April to mid May.  
doi:10.1371/journal.pone.0094190.t004


2012, the amount of rainfall was extraordinary low during April–May 2013, which resulted in very few overwintering *A. lucorum* eggs hatching. Niu et al. [35] also reported that the drought

directly decreased *A. lucorum* population levels on Chinese dates in spring season. Therefore, we conclude that rainfall is the most important factor for forecasting the population levels of *A. lucorum* during the spring.

In other species it has been noted that specific rainfall characteristics (e.g. intensity, duration and frequency of rainfalls) might exert strong effects on insect population growth and their seasonal dynamics [31,35,36]. We described year-to-year fluctuations in the numbers of 1<sup>st</sup>-instar *A. lucorum* nymphs, and it is possible that these fluctuations are due to differences in specific rainfall characteristics. For instance, the intensity, duration, and frequency of rainfall were greater during the 2012 study period than in 2011, which might explain why overwintering eggs hatched earlier in 2012 and why the population density of 1<sup>st</sup>-instar nymphs was markedly higher than in the previous year (Figure 3). Although a similar correlation between nymph population dynamics and specific rainfall characteristics was observed between 2009 and 2010, this relationship requires further study before it can be validated. As changes in rainfall are associated with global climate change [40], it is likely that *A. lucorum* nymph population levels and dynamics will be altered in years to come. The temporal and spatial variation of rainfall distribution also has been recorded in China [41], whereas their potential effects on *A. lucorum* population dynamics need further assessment.

*Apolygus lucorum* overwinters as diapause eggs [42,43,44]. Over the course of the study period, which was carried out from mid April to early May, we found that the average length of egg development as well as the variation in development length between individuals greatly decreased. In other words, we found that the rate of diapause termination of the overwintering eggs gradually increased during the study period. It is already known that high temperatures promote diapause termination in overwintering eggs and reduce developmental duration [35,44]. Consistent with this, we observed that developmental duration became shorter as temperatures increased, given constant relative humidity within the same sampling period (Table 3, Figure 2). Therefore, we conclude that temperature levels should be a useful tool for predicting the developmental status of *A. lucorum* in the field [28,30].

More than 80 different plant species have been recorded as overwintering hosts for *A. lucorum* [26], some of which are living



**Figure 4.** The linear relationships between rainfall and the abundance of *A. lucorum* 1<sup>st</sup>-instar nymphs and nymphs of all stages in a Chinese date orchard during 2009–2013. Linear model: (a) 1<sup>st</sup>-instar nymphs:  $y = 0.2848 + 0.0025 x$ ;  $F = 16.82$ ;  $df = 1,3$ ;  $P = 0.0262$ . (b) nymphs of all stages:  $y = 0.6899 + 0.0025 x$ ;  $F = 24.08$ ;  $df = 1,3$ ;  $P = 0.0162$ .

doi:10.1371/journal.pone.0094190.g004



hosts, whereas others are dead. Eggs that are embedded within live plant tissues share a complex relationship with the host plant, and processes such as egg dormancy and postembryonic development allow egg hatching to coincide with certain phases of host phenology [45]. As a result, eggs generally hatch at a time when food, water and other necessary nutrients are available on the host plant to allow for survival of the young nymphs. In contrast, eggs that hatch on dead plant material can encounter relatively unfavorable conditions for survival and development. However, as *A. lucorum* reportedly feeds upon a variety of common weeds and cultivated plants [25,33], nymphs that emerge on dead plant material can find necessary resources on surrounding plants.

In this study, we determined that rainfall was the most important weather determinant for the hatching of overwintering *A. lucorum* eggs in the dead parts of tree hosts. In addition, temperature was found to significantly hasten egg development, and relative humidity was found to greatly increase hatching rate.

## References

- Pizzol J, Pintureau B, Khoulidia O, Desneux N (2010) Temperature-dependent differences in biological traits between two strains of *Trichogramma cacoeciae* (Hymenoptera: Trichogrammatidae). *J Pest Sci* 83: 447–452.
- Willmer P (1982) Microclimate and the environmental physiology of insects. *Adv insect physiol* 16: 1–57.
- Wilstermann A, Vidal S (2013) Western corn rootworm egg hatch and larval development under constant and varying temperatures. *J Pest Sci* 86: 419–428.
- Dong YC, Clarke AR, Pereira R, Desneux N, Niu CY (2013) Pupal diapause development and termination is driven by low temperature chilling in *Bactrocera minax*. *J Pest Sci* 86: 429–436.
- Andreadis SS, Kagkellaris NK, Eliopoulos PA, Savopoulou-Soutani M (2013) Temperature-dependent development of *Sesamia nonagrioides*. *J Pest Sci* 86: 409–417.
- Day WH (2006) The effect of rainfall on the abundance of tarnished plant bug nymphs [*Lygus lineolaris* (Palisot)] in alfalfa fields. *Trans Am Entomol Soc* 132: 445–450.
- Mullens BA, Peterson NG (2005) Relationship between rainfall and stable fly (Diptera: Muscidae) abundance on California dairies. *J Med Entomol* 42: 705–708.
- Shang ZT, Wei JS, Kuai J, Shan C, Chen YW, et al. (2010) Analyses of the relations between cotton mirid dynamics and meteorological conditions. *Plant Prot* 36: 92–95.
- Shang ZT, He L, Liu SC, Lu ZG, Chen L (2012) Effect of climate change on cotton *Lygus lucorum* occurrence degree. *Agr Sci Technol* 13: 784–788, 887.
- Boggs CL, Inouye DW (2012) A single climate driver has direct and indirect effects on insect population dynamics. *Ecol Lett* 15: 502–508.
- Mattson WJ, Haack RA (1987) The role of drought in outbreaks of plant-eating insects. *Bio Sci* 37: 110–118.
- White TCR (2008) The role of food, weather and climate in limiting the abundance of animals. *Biol Rev* 83: 227–248.
- Cannon RJ (1998) The implications of predicted climate change for insect pests in the UK, with emphasis on non-indigenous species. *Glob Change Biol* 4: 785–796.
- Parmesan C, Ryrholm N, Stefanescu C, Hill JK, Thomas CD, et al. (1999) Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature* 399: 579–583.
- Parmesan C (2007) Influences of species, latitudes and methodologies on estimates of phenological response to global warming. *Glob Change Biol* 13: 1860–1872.
- Liu XF, Jin Y, Ye H (2013) Recent spread and climatic ecological niche of the invasive guava fruit fly, *Bactrocera correcta*, in mainland China. *J Pest Sci* 86: 449–458.
- Simoglou KB, Karataraki A, Roditakis NE, Roditakis E (2012) *Euzophera bigella* (Zeller) (Lepidoptera: Pyralidae) and *Dasineura oleae* (F. Low) (Diptera: Cecidomyiidae): Emerging olive crop pests in the Mediterranean? *J Pest Sci* 85: 169–177.
- Gutierrez AP, Daane KM, Ponti L, Walton VM, Ellis CK (2008) Prospective evaluation of the biological control of vine mealybug: refuge effects and climate. *J Appl Ecol* 45: 524–536.
- Jepsen JU, Hagen SB, Ims RA, Yoccoz NG (2008) Climate change and outbreaks of the geometrids *Operophtera brumata* and *Epirrita autumnata* in subarctic birch forest: evidence of a recent outbreak range expansion. *J Ani Ecol* 77: 257–264.
- Wu G, Ge F, Wan FH, Xiao NW, Li JS (2011) Responses of invasive insects to global climate change. *Chin J Appl Entomol* 48: 1170–1176.
- Guo K, Hao SG, Sun OJ, Kang L (2009) Differential responses to warming and increased precipitation among three contrasting grasshopper species. *Glob Change Biol* 15: 2539–2548.
- Morimoto N, Imura O, Kiura T (1998) Potential effects of global warming on the occurrence of Japanese pest insects. *Appl Entomol Zool* 33: 147–156.
- Lu YH, Qiu F, Feng HQ, Li HB, Yang ZC, et al. (2008) Species composition and seasonal abundance of pestiferous plant bugs (Hemiptera: Miridae) on Bt cotton in China. *Crop Prot* 27: 465–472.
- Lu YH, Wu KM, Jiang YY, Xia B, Li P, et al. (2010) Mirid bug outbreaks in multiple crops correlated with wide-scale adoption of Bt cotton in China. *Science* 328: 1151–1154.
- Lu YH, Wu KM (2008) Biology and control of cotton mirids, Golden Shield Press, China.
- Lu YH, Wu KM, Wyckhuys KAG, Guo YY (2010) Overwintering hosts of *Apolygus lucorum* (Hemiptera: Miridae) in northern China. *Crop Prot* 29: 1026–1033.
- Lu YH, Wu KM (2011) Mirid bugs in China: pest status and management strategies. *Outlooks Pest Manag* 22: 248–252.
- Lu YH, Wu KM, Wyckhuys KAG, Guo YY (2010) Temperature-dependent life history of the green plant bug, *Apolygus lucorum* (Meyer-Dür) (Hemiptera: Miridae). *Appl Entomol Zool* 45: 387–393.
- Lu YH, Wu KM (2011) Effect of relative humidity on population growth of *Apolygus lucorum* (Heteroptera: Miridae). *Appl Entomol Zool* 46: 421–427.
- Men XY, Yu Y, Zhang AS, Li LL, Zhang JT, et al. (2008) Life table of the laboratory population of *Lygus lucorum* Meyer-Dür (Hemiptera: Miridae) at different temperatures. *Acta Entomol Sin* 51: 1216–1219.
- Ting YQ (1964) Studies on the population fluctuations of cotton mirids in the cotton cultivation region of Kwanchung, Shensi, China. *Acta Entomol Sin* 13: 298–310.
- Chu HF, Meng HL (1958) Studies on three species of cotton plant bugs, *Adelphocoris taeniphorus* Reuter, *A. lineolatus* (Goeze), and *Lygus lucorum* Meyer-Dür (Hemiptera: Miridae). *Acta Entomol Sin* 8: 97–118.
- Cao CY, Wan CS (1983) Management of cotton mirids. Shanghai Science and Technology Press, Shanghai.
- Liu HM (1991) Studies on the bionomics and control strategy of black striped leaf bug in northern coastal reclamation area of Jiangsu province. *Acta Phytophyl Sin* 18: 147–153.
- Niu SG, Liu XH, Zhang SJ, Liu YS, Li JZ, et al. (2011) Study on relationship between occurrence of *Lygus lucorum* on winter jujube and climate in north Shandong province. *Acta Agr Jiangxi* 23: 97–99.
- Wu KM, Li W, Feng HQ, Guo YY (2002) Seasonal abundance of the mirids, *Lygus lucorum* and *Adelphocoris* spp. (Hemiptera: Miridae) on Bt cotton in northern China. *Crop prot* 21: 997–1002.
- Wang ZL, Han HZ, Liu MJ, Shao XH, Zhang XH (2011) Distribution and hatching of overwintering eggs of *Lygus lucorum* Meyer-Dür (Hemiptera: Miridae). *J Northwest A & F Univ* 39: 148–153.
- Pan HS, Lu YH, Wyckhuys KAG, Wu KM (2013) Preference of a polyphagous mirid bug, *Apolygus lucorum* (Meyer-Dür) for flowering host plants. *PLoS ONE* 8: e68980.
- Geng HH, Pang HS, Lu YH, Yang YZ (2012) Nymphal and adult performance of *Apolygus lucorum* (Hemiptera: Miridae) on a preferred host plant, mungbean *Vigna radiata*. *Appl Entomol Zool* 47: 191–197.
- Solomon S, Qin D, Manning M, Chen Z, Marquis M, et al. (eds). (2007) Climate Change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge: Cambridge University Press.
- Wand Y, Cao MK, Tao B, Li KR (2006) The characteristics of spatio-temporal patterns in precipitation in China under the background of global climate change. *Geographical Research* 25: 1031–1040.

Overall, spring rainfall and temperature were the key factors for predicting population level and emerging period of *A. lucorum* nymphs in this season, respectively. Hence, further forecast the emergence of *A. lucorum* based on weather data will be available for effective management of its spring population, which is the source of *A. lucorum* in different host crops in summer season.

## Acknowledgments

We thank Dr. Giovanni Benelli for valuable comments on earlier version of the manuscript.

## Author Contributions

Conceived and designed the experiments: YL HP. Performed the experiments: HP YL BL. Analyzed the data: YL HP BL. Contributed reagents/materials/analysis tools: YL. Wrote the paper: HP YL ND.

42. Chen PY, Feng HQ, Li GP, Guo XR (2010) Morphological observation of diapause and non-diapause eggs of *Apolygus lucorum* (Meyer-Dür)(Heteroptera: Miridae). *J Henan Agr Univ* 44: 83–85.
43. Feng HQ, Chen PY, Li GP, Qiu F, Guo XR (2012) Diapause Induction in *Apolygus lucorum* and *Adelphocoris suturalis* (Hemiptera: Miridae) in Northern China. *Environ Entomol* 41: 1606–1611.
44. Zhuo DG, Li ZH, Men XY, Yu Y, Zhang AS, et al. (2011) Effects of low temperature and photoperiod on diapause termination and developmental duration of the overwintering egg of *Apolygus lucorum* Meyer-Dür (Hemiptera: Miridae). *Acta Entomol Sin* 54: 136–142.
45. Wheeler AG (2001) *Biology of the plant bugs* (Hemiptera: Miridae): pests, predators, opportunists, Cornell University Press.