



**HAL**  
open science

## Use of feathers to assess polychlorinated biphenyl and organochlorine pesticide exposure in top predatory bird species of Pakistan

Naeem Akhtar Abbasi, Igor Eulaers, Veerle L. B. Jaspers, Muhammad Jamshed Iqbal Chaudhry, Adrien Frantz, Per Lennart Ambus, Adrian Covaci, Riffat Naseem Malik

### ► To cite this version:

Naeem Akhtar Abbasi, Igor Eulaers, Veerle L. B. Jaspers, Muhammad Jamshed Iqbal Chaudhry, Adrien Frantz, et al.. Use of feathers to assess polychlorinated biphenyl and organochlorine pesticide exposure in top predatory bird species of Pakistan. *Science of the Total Environment*, 2016, 569, pp.1408-1417. 10.1016/j.scitotenv.2016.06.224 . hal-02637665

**HAL Id: hal-02637665**

**<https://hal.inrae.fr/hal-02637665v1>**

Submitted on 15 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**This item is the archived peer-reviewed author-version of:**

Use of feathers to assess polychlorinated biphenyl and organochlorine pesticide exposure in top predatory bird species of Pakistan

**Reference:**

Abbasi Naeem Akhtar, Eulaers Igor, Jaspers Veerle, Chaudhry Muhammad Jamshed Iqbal, Frantz Adrien, Ambus Per Lennart, Covaci Adrian, Malik Riffat Naseem.- Use of feathers to assess polychlorinated biphenyl and organochlorine pesticide exposure in top predatory bird species of Pakistan

The science of the total environment - ISSN 0048-9697 - 569(2016), p. 1408-1417

Full text (Publisher's DOI): <http://dx.doi.org/doi:10.1016/J.SCITOTENV.2016.06.224>

To cite this reference: <http://hdl.handle.net/10067/1356960151162165141>

1 **Use of feathers to assess polychlorinated biphenyl and organochlorine**  
2 **pesticide exposure in top predatory bird species of Pakistan**

3 Naeem Akhtar Abbasi <sup>a\*</sup>, Igor Eulaers <sup>b</sup>, Veerle Leontina Bernard Jaspers <sup>c</sup>, Muhammad Jamshed  
4 Iqbal Chaudhry <sup>a, d</sup>, Adrien Frantz <sup>e</sup>, Per Lennart Ambus <sup>f</sup>, Adrian Covaci <sup>g</sup>, Riffat Naseem Malik  
5 <sup>a</sup>

6  
7 <sup>a\*</sup> Environmental Biology and Ecotoxicology Laboratory, Department of Environmental Sciences, Faculty of  
8 Biological Sciences, Quaid-i-Azam University, Islamabad 45320, Pakistan  
9

10 <sup>b</sup> Department of Bioscience, Aarhus University, Frederiksborgevej 399, P.O. Box 358, 4000 Roskilde, Denmark

11 <sup>c</sup> Environmental Toxicology, Department of Biology, Norwegian University of Science and Technology (NTNU),  
12 Trondheim, Norway

13 <sup>d</sup> WWF-Pakistan, Ferozpur Road, PO Box 5180, Lahore 54600, Pakistan

14 <sup>e</sup> Sorbonne Universités, UPMC Univ Paris 06, UPEC, Paris 7, CNRS, INRA, IRD, Institut d'Ecologie et des  
15 Sciences de l'Environnement de Paris, F-75005, Paris, France  
16

17 <sup>f</sup> Center for Permafrost, Department of Geosciences and Natural Resource Management, University of Copenhagen,  
18 Øster Voldgade 10, 1350 København K, Denmark  
19

20 <sup>g</sup> Toxicological Centre, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

21 \* Corresponding Author's e-mail address: [abbasi\\_akhtar4045@yahoo.com](mailto:abbasi_akhtar4045@yahoo.com)

22

23 **Abstract**

24 Little is known about the levels of organochlorines (OCs) in predatory bird species from Asia or  
25 the factors governing their concentration. This study is the first report on concentration of  
26 polychlorinated biphenyls (PCBs) and organochlorine pesticides (OCPs) in predatory birds of  
27 Pakistan. The concentrations of PCBs and OCPs were investigated using tail feathers of ten  
28 different species of predatory birds. In addition, concentration differences among body, tail,  
29 primary and secondary feathers were investigated for six individuals of black kite (*Milvus*  
30 *migrans*). Ranges of concentrations were highest for dichlorodiphenyldichloroethylene (*p,p'*-  
31 DDE: 0.11-2163 ng g<sup>-1</sup> dry wt.) followed by dichlorodiphenyltrichloroethane (*p,p'*-DDT: 0.36-  
32 345 ng g<sup>-1</sup> dry wt.), hexachlorobenzene (HCB: 0.02-34 ng g<sup>-1</sup> dry wt.),  $\Sigma$ PCBs (0.03-16 ng g<sup>-1</sup>  
33 dry wt.) and *trans*-nonachlor (TN; 0.01-0.13 ng g<sup>-1</sup> dry wt.). CB 118, 153, 138, and 180 along  
34 with *p,p'*-DDE were found as most prevalent compounds.  $\Sigma$ PCBs and  $\Sigma$ DDTs were significantly  
35 different among species (both  $p < 0.01$ ) and omnivorous, scavengers, carnivorous and piscivorous  
36 trophic guilds (all  $p < 0.03$ ). Whereas only  $\Sigma$ PCBs were significantly different ( $p < 0.01$ ) among  
37 different families of birds. Values of stable isotopes ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) differed significantly (all  
38  $p < 0.01$ ) among species, families, trophic guilds as well as terrestrial and aquatic habitat but not  
39 between nocturnal and diurnal predators ( $p = 0.22$  for  $\delta^{13}\text{C}$ ;  $p = 0.50$  for  $\delta^{15}\text{N}$ ). Concentrations of  
40  $\Sigma$ PCBs,  $\Sigma$ DDTs and *trans*-nonachlor, but not HCB ( $p = 0.86$ ), were significantly different among  
41 different feather types (all  $p < 0.01$ ). Trophic and taxonomic affiliation as well as dietary carbon  
42 sources ( $\delta^{13}\text{C}$ ) for species were identified as the variables best explaining the observed variation  
43 in exposure to the studied compounds. The significance of contributing factors responsible for  
44 OC contamination differences in predatory birds should be further elucidated in future studies.

45

46 **Key words:** trophic guild, feathers, habitat, POPs,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$

## 47 **1. Introduction**

48 During the past few decades, prolific discharge of legacy persistent organic pollutants (POPs)  
49 from industrial, urban and agricultural sources have remained a cause of many environmental  
50 concerns particularly related to their toxic effects in humans and wildlife (Letcher et al., 2010).  
51 These chemicals are persistent, bioaccumulative, toxic and travel large distances through long-  
52 range transport (Vorkamp and Rigét, 2014). Two major classes of POPs, i.e. polychlorinated  
53 biphenyls (PCBs) and organochlorine pesticides (OCPs) have been introduced after the  
54 industrial revolution in 1920 and are still widespread in the environment (Lohmann et al., 2007)  
55 despite being legally mitigated on a worldwide scale (UNEP, 2011). PCBs were used in a wide  
56 array of substances as a coolant or additives and escape into the environment during their usage,  
57 packaging and storage as well as through leaching from landfills (Covaci et al., 2006). On the  
58 other hand, OCPs are chlorine-containing organic pesticides which were predominately used as  
59 insecticides (Ali et al., 2014). Among a variety of compounds, dichlorodiphenyltrichloroethane  
60 (DDT) was the mostly heavily used pesticide after World War I and its production was banned  
61 from USA, Europe, China and Japan after 1972 when its toxic effects became established  
62 (Tanabe et al., 1998). Hexachlorobenzene (HCB) was found among the most prevalent OCPs  
63 because of its use as insecticide as well as an industrial by-product (Corsolini et al., 2006).  
64 Besides these, metabolites of chlordanes (CHLs) were also found as compounds of concern  
65 because of their exacerbated use as insecticides and their reported adverse effects upon wildlife  
66 (Letcher et al., 2010). These OCPs mainly get their way into the environment during their  
67 production, application and storage and are eventually dispersed through runoff and air currents  
68 (Guan et al., 2009).

69 Ever since the toxicological significance of POPs was suspected, predatory birds have been  
70 successfully used as sentinels to assess the levels of these compounds in the environment (et al.,  
71 1993; Dauwe et al., 2005; Jaspers et al., 2006). However, sampling of predatory birds often  
72 encounters various practical and ethical impediments. Use of non-destructive tissues, such as  
73 blood, feathers and preen oil, is usually recommended as a preferable choice in case of  
74 predatory birds. Among these, the use of feathers has become more and more applicable  
75 because it is less invasive, comes along with easy collection and storage, and provides a  
76 valuable assessment of internal body burdens of POPs (Jaspers et al., 2006). Some of the recent  
77 studies have also emphasized to evaluate different types of feathers i.e. body, tail, primary and

78 secondary which could best represent the level of the studied compounds (Eulaers et al.,  
79 2014b; García-Fernández et al., 2013; Jaspers et al., 2011).

80 Levels of POPs in avian tissues are influenced by a multitude of biological, spatial and  
81 ecological factors (Eulaers et al., 2013; Lavoie et al., 2010). Trophic levels/feeding guilds and  
82 taxonomic affiliation of species, locational and dietary exposure well as individual condition  
83 factors such as gender, age and reproductive status may significantly influence the concentration  
84 of POPs in birds (Eulaers et al., 2013; Behrooz et al., 2009). In general, POP concentrations at  
85 higher trophic level species mainly stem from dietary intake, which can be quantified using  
86 ratios of stable nitrogen and carbon isotopes (SIs; Eulaers et al., 2014a). The ratio of heavier  $^{15}\text{N}$   
87 to lighter  $^{14}\text{N}$  ( $\delta^{15}\text{N}$ ) provides information about the trophic level of an individual because it  
88 enriches with each trophic level (Huang et al., 2013). The ratio of carbon SIs ( $\delta^{13}\text{C}$ :  $^{13}\text{C}/^{12}\text{C}$ ) is  
89 used as an indicator for dietary origin because of the varying degree of depletion of  $^{13}\text{C}$  stable  
90 isotopes in primary producers from different habitats (Boecklen et al., 2011; Jardine et al., 2006).  
91 Although the use of SIs has shown promising to investigate trophodynamics of POPs, it has had  
92 less focus in predatory birds, particularly those from the Asian continent.

93 Predatory birds of the southern Asian region are particularly exposed to a high magnitude of  
94 legacy POPs because of their historical and current use in this region (Ali et al., 2014). Levels of  
95 POPs have been documented in biotic as well as abiotic components of the environment from  
96 South Asia (Sarkar et al., 2008; Yadav et al., 2015), but predatory birds have received less  
97 attention (Abbasi et al., 2016). Contamination of birds with POPs has only been reported in eggs  
98 of little (*Egretta garzetta*) and cattle egret (*Bubulcus ibis*) from Pakistan (Khan et al., 2014;  
99 Malik et al., 2011; Sanpera et al., 2003). Seeing this scarcity of exposure data, the present study  
100 was designed to investigate the current concentration levels of different OC compounds using  
101 feathers of multiple predatory bird species of Pakistan. Further, we evaluated the importance of  
102 various factors governing interspecific variation of OC exposure including intraspecific  
103 variations through carbon and nitrogen SI values. Lastly, the suitability of different feather types  
104 to characterize OC exposure was evaluated by comparing body, tail, primary and secondary  
105 feathers from black kites (*Milvus migrans*).

106

## 107 2. Methodology

## 108 2.1. Sample collection

109 Feather samples ( $N=76$ ) from ten different species of predatory birds were collected between  
110 June 2012 to September 2014 (Fig. 1). Species selected for this study included black kite  
111 ( $N=13$ ), Eurasian sparrowhawk (*Accipiter nisus*,  $N=10$ ), common kestrel (*Falco tinnunculus*,  
112  $N=4$ ), red-necked falcon (*Falco chicquera*,  $N=2$ ), Indian vulture (*Gyps indicus*,  $N=9$ ), white-  
113 rumped vulture (*Gyps bangalensis*,  $N=12$ ), spotted owlet (*Athene brama*,  $N=10$ ), little owl  
114 (*Athene noctua*,  $N=6$ ), great cormorant (*Phalacrocorax carbo*,  $N=4$ ) and grey heron (*Ardea*  
115 *cinerea*,  $N=6$ ). Tail feathers were obtained from all these species. In addition, tail, body,  
116 primary and secondary feathers were collected from six individuals of black kite to investigate  
117 concentration differences among feather types. Sampling details of each site are summarized in  
118 Table S1 (supplementary information). Predatory birds were sampled mainly from different  
119 towns and cities and their outskirts in Punjab province, which is considered a hub of agricultural  
120 activities of the country. Samples of black kite and spotted owlet were also collected from two  
121 metropolitan cities, i.e. Lahore and Rawalpindi, with higher expected anthropogenic input than  
122 other sites. Samples of both the vulture's species were obtained from their isolated and remotely  
123 located colonies (S1&S2) at Nagar Parker, Sindh Province. Grey heron was the only species  
124 sampled from northern regions at Lulusar Lake (S13), which is a remote waterbody. Each  
125 species was sampled from one location except black kite, Eurasian sparrowhawk and spotted  
126 owlet, which were sampled from two different locations (Table S1). Black kites and spotted  
127 owlet were sampled around the outskirts of Lahore, which is a metropolitan city with extensive  
128 agricultural activities in its suburbs, and Islamabad, which is a relatively smaller city with very  
129 small scale agricultural activities in its premises. The third species, Eurasian sparrowhawk, was  
130 sampled from Mianwali and Khaniwal, which are both small cities with extensive agricultural  
131 lands around. Further, species are discussed under various categories based on their taxonomic  
132 affiliation (families; accipitridae, ardeidae, falconidae, phalacrocoracidae, strigidae), trophic  
133 guilds (Omnivorous, scavenger, carnivorous, piscivorous, habitats (terrestrial or aquatic) and  
134 feeding regimes (diurnal or nocturnal). All the samples used in this study were taken from birds  
135 captured in the framework of other studies. A special permit from CITES authorities in Pakistan  
136 was acquired for shipping and transport of the samples of the two critically endangered vulture  
137 species. After collection, feathers were kept in zipped plastic bags and stored at  $-20^{\circ}\text{C}$  until  
138 chemical analysis.

139 2.2. Quantification of PCBs and OCPs

140 The procedure for cleanup and extraction of POPs was adapted from previous described  
141 methods (Dauwe et al., 2005; Jaspers et al., 2006). Feathers were thoroughly washed with  
142 deionized water to remove exogenous dust particles and other unwanted depositions. After  
143 washing, feathers were covered with standard laboratory paper and dried overnight at ambient  
144 temperature. Dried feathers were cut into pieces of ~1 mm, weighed and transferred to  
145 analytical glass recipients. Initially, feather samples were spiked with the internal standard CB  
146 143 (50 $\mu$ L of 200 pg  $\mu$ L<sup>-1</sup>) and incubated overnight at 45°C in HCl (4M) and  
147 hexane:dichloromethane (4:1; v:v). From the incubated mixture, analytes were liquid-liquid  
148 extracted using hexane:dichloromethane (4:1; v:v). Cleanup of the resulting extract was  
149 performed on acidified silica (800 mg; 44% H<sub>2</sub>SO<sub>4</sub>) topped with anhydrous Na<sub>2</sub>SO<sub>4</sub> (400 mg),  
150 and analytes were eluted with hexane:dichloromethane (4:1; v:v). Finally, the cleaned-up  
151 extracts were concentrated using a gentle flow of Nitrogen gas, reconstituted in 80  $\mu$ L *iso*-  
152 octane, and transferred to injection vials. The whole process of clean-up and extraction was  
153 performed at the Bird ecotoxicology laboratory, Norwegian University of Science and  
154 Technology (Trondheim, Norway), whereas the concentrations of PCBs and OCPs were  
155 quantified at the Toxicological Center, University of Antwerp (Wilrijk, Belgium).

156 The concentrations of PCBs and OCPs were quantified using a mass spectrometer (Agilent MS  
157 5973, Palo Alto, CA, USA) operated in electron-capture negative ionization mode to a gas  
158 chromatograph (Agilent GC 6890, Palo Alto, CA, USA). A total of 19 PCB congeners (CB 105,  
159 118, 146, 153, 138, 187, 183, 128, 174, 177, 171, 156, 180, 170, 199, 196/203, 194, 206, 209),  
160 HCB, *trans*-nonachlor (TN), *cis*-nonachlor (CN), oxychlordan (OXC), and DDTs, i.e. *p,p'*-  
161 DDE, *p,p'*-DDT, were measured. In all samples, only high chlorinated PCBs were measured.  
162 However, in a few samples (those with the highest concentrations of PCBs), we have attempted  
163 to measure lower-chlorinated PCB congeners. Yet, detection limits were higher and the lower  
164 chlorinated PCB congeners measured (CB28, CB52, CB95 etc.) were below the limit of  
165 quantification (<LOQ) and are thus not reported. Concentrations of analytes were expressed as  
166 ng g<sup>-1</sup> dry weight (dw). Internal standards were purchased from Accustandard (New Haven, CT,  
167 USA), while pesticide-grade solvents (Merck, Darmstadt, Germany) were used throughout the  
168 entire process. Mean recoveries of internal standards were 52% $\pm$ 13 for PCBs in all samples.  
169 The same internal standard (CB 143) was used for other OCPs. For quality assurance, in each



170 batch of 10 samples, a procedural blank was prepared and analyzed. LOQs for different analytes  
171 were set at 3\*SD of the procedural blank values. When analytes were not detected in blanks, the  
172 LOQ was calculated using a 10:1 signal to noise ratio.

### 173 2.3. Stable isotopes measurement

174 Composition of stable nitrogen and carbon isotopes was measured at the Center for Permafrost  
175 (University of Copenhagen, Denmark). We adapted the previously reported procedure by  
176 [Eulaers et al. \(2014a\)](#) for the measurement of SIs in feathers. Briefly, a representative  
177 homogenized subsample of 0.5 to 2.0 mg was wrapped into a tin combustion cup, and the ratios  
178 for stable carbon and nitrogen isotopes were measured by continuous flow using an elemental  
179 analyzer (CE 1110, Thermo Electron, Milan, Italy) coupled to a mass spectrometer (Finnigan  
180 MAT Delta PLUS, Thermo Scientific, Bremen, Germany). The ratios of SIs were expressed as

$$\delta X(\text{‰}) = \left( \frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right)$$

181 with X representing the C or N SIs and R representing their corresponding ratios ( $^{13}\text{C}/^{12}\text{C}$ ,  
182  $^{15}\text{N}/^{14}\text{N}$ ) in the sample or standard. Reference samples (Atropin) were included for the positive  
183 evaluation of analytical performance. The instrument was calibrated by employing pure gases of  
184  $\text{CO}_2$  and  $\text{N}_2$  against the certified reference material of sucrose and  $(\text{NH}_4)_2\text{SO}_4$  provided by the  
185 International Atomic Energy Agency (IAEA, Vienna, Austria). The SI ratios were calculated  
186 against the international standards Vienna PeeDee Belemnite (vPDB) and atmospheric  $\text{N}_2$  (AIR)  
187 respectively. Analytical precision was maintained at 0.1‰ SD.

### 188 2.4. Statistical analysis

189 All the statistical computations were performed using SPSS (IBM 20) and R (version 3.2.3).  
190 Firstly, screening of the data was performed as suggested by [Zuur et al. \(2010\)](#) to avoid common  
191 statistical errors. Data was  $\log_{10}$  transformed after testing for normality using Q-Q plots and  
192 Shapiro-Wilk's tests (all  $p < 0.05$ ). Only those compounds which were detected above the limit of  
193 quantification ( $>\text{LOQ}$ ) in at least 50% of the samples of a species were treated for further  
194 statistical analysis. Missing values for these compounds were substituted with the proportion of  
195 detected samples\*LOQ. The null-hypothesis was rejected at  $\alpha=0.05$ . Firstly, differences of PCB  
196 and OCP concentrations among species, families, omnivorous, scavengers, carnivorous and  
197 piscivorous trophic guilds, habitats (aquatic/terrestrial), feeding regime (diurnal or nocturnal) as

198 well their associations to dietary proxies ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) were tested through analysis of variance  
199 (ANOVA). Subsequent post-hoc Tukey's tests for honest significant differences (HSD) were  
200 used for multiple comparisons. Further, above-mentioned variables were evaluated for their  
201 capacity to explain the observed variation in levels of PCBs and OCPs using Akaike's  
202 Information Criteria (AICc) as discussed previously (Johnson and Omland, 2004). A separate  
203 AIC-based selection was run for each compound to evaluate the factors best governing the  
204 observed variation in PCB and OCPs concentrations. Separate AIC based model was run for  
205 each compound based on of the fact that they have different physicochemical properties and may  
206 be influenced differently by different factors. Associations of PCBs and OCPs with dietary  
207 proxies ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) were tested through linear regression. A separate ANOVA was performed to  
208 determine the variation in concentrations among different feather types sampled from black kite.  
209

### 210 **3. Results and Discussion**

#### 211 *3.1. Variation in OC concentrations and profiles*

212 To the best of our knowledge, so far PCBs and OCPs have never been quantified in feathers of  
213 predatory birds from Pakistan. The measured concentrations for different compounds of PCBs  
214 and OCPs are summarized in table S2. Out of the 19 PCBs congeners targeted, 14 congeners,  
215 i.e. CB 105, 118, 146, 153, 138, 187, 183, 128, 156, 180, 170, 199, 196/203, 194 were detected  
216 above LOQ in  $\geq 50\%$  samples for minimum one to maximum all the species. Among the studied  
217 compounds CB 153, HCB, *p,p'*-DDE and *p,p'*-DDT were detected in all the studied species  
218 whereas all other compounds were variably detected (Figure S1). Compounds such as CB 171,  
219 174, 177, 206, 209, as well as OXC and CN were not detected above LOQ in  $\geq 50\%$  of the  
220 samples in any species (Figure S1), hence not further discussed. In general, the trend  
221  $\sum\text{DDTs} > \text{HCB} > \text{PCBs} > \text{TN}$  was depicted in tail feathers of predatory birds from Pakistan. The  
222 concentration ranges (minimum-maximum) recorded in this study were  $0.11\text{-}2163 \text{ ng g}^{-1}$  dry wt.  
223 for DDTs,  $0.02\text{-}34 \text{ ng g}^{-1}$  dry wt. for HCB and  $0.03\text{-}16 \text{ ng g}^{-1}$  dry wt. for  $\sum\text{PCBs}$  and  $0.01\text{-}0.13$   
224  $\text{ ng g}^{-1}$  dry wt. for TN respectively. Previously, screening of OCs has only been carried out in  
225 eggs of little egret (Sanpera et al., 2003) and cattle egret (Khan et al., 2014; Malik et al., 2011)  
226 from Pakistan. In those studies, reported concentrations of OCs were higher probably due to the  
227 higher lipid content in egg. In recent global literature, PCBs and OCPs in predatory birds have  
228 been mostly reported in egg, muscle, liver, kidney and other non-keratinous tissues (Chen et al.,

229 2009; Jaspers et al., 2006; Kocagöz et al., 2014; Lavoie et al., 2010; Peng et al., 2015; Sun et  
230 al., 2014; Zhang et al., 2011) whereas only few studies are available for comparison of PCBs  
231 and OCPs in feathers. Compared to findings of the present study,  $\Sigma$ DDTs were found  
232 comparable whereas  $\Sigma$ PCBs and HCB were approximately 5 to >50 fold higher in feathers from  
233 different predatory bird species from south-west of Iran (Behrooz et al., 2009). Similarly,  
234 feather concentrations of  $\Sigma$ DDTs and HCB were found comparable to our findings, whereas  
235  $\Sigma$ PCBs levels were relatively higher in different waterbird species from the Caspian Sea coast,  
236 Northern Iran (Rajaei et al., 2011). Regarding the European scenario, compared to our study  
237 Jaspers et al., (2007) reported relatively higher  $\Sigma$ PCBs, comparable HCBs and lower  $\Sigma$ DDTs  
238 levels in tail feathers of multiple predatory species from Belgium. Further, Jaspers et al., (2009)  
239 reported a comparable level of *p,p'*-DDE (1.07-139 ng g<sup>-1</sup> dry wt.), relatively lower level of  
240 *p,p'*-DDT (0.38-11.8 ng g<sup>-1</sup> dry wt.) and fairly high range of  $\Sigma$ PCBs (2.92-236 ng g<sup>-1</sup> dry wt.) in  
241 tail feathers of common magpie (*Pica pica*) from Belgium. Similarly, concentrations of *p,p'*-  
242 DDE, HCB and TN but not  $\Sigma$ PCBs of this study were found comparable or slightly higher than  
243 reported in tail feathers of white-tailed eagle (*Haliaeetus albicilla*) from western Greenland  
244 (Jaspers et al., 2011).

245 The contribution of the detected compounds is illustrated in figure 2a, whereas profiles for PCBs  
246 and DDTs are shown in figure S2a,b. Among the detected compounds, *p,p'*-DDE was found as  
247 the predominant compound in predatory birds of the current study followed by *p,p'*-DDT and  
248 congeners of PCBs, HCB and TN respectively, which is in line with previous studies (Chen et  
249 al., 2009; Rajaei et al., 2011). Among PCBs, CB 118, 153, 138, 180, 170 and 194 were observed  
250 as more prevalent congeners in tail feathers (figure S1). In the present study, PCB congeners  
251 with six (*hexa*-CBs) and seven chlorines (*hepta*-CBs) dominated in terrestrial species whereas  
252 those containing five chlorines (*penta*-CBs) were more prevalent in aquatic species (Figure S2a),  
253 which is in agreement with previous findings (Yu et al., 2014; Jaspers et al., 2007). Previously,  
254 Abbasi et al., (2016) reported that *p,p'*-DDE has been unanimously detected as predominant  
255 metabolite of DDTs in Asian studies on birds. In contrast, *p,p'*-DDE has been found as a  
256 predominant compound in feathers of European predatory birds (Jaspers et al., 2007; Eulaers et  
257 al., 2013). In general, the elevated level of DDTs in this study corresponds to their wide scale use  
258 as pesticide in Pakistan (Ali et al., 2014). Similar to our findings, CB 153, 180, 138 have been  
259 reported as predominant congeners in feathers of predatory birds from different parts of the

260 world (Chen et al., 2009; Behrooz et al., 2009, Jaspers et al., 2007, 2011). Earlier, Dauwe et al.  
261 (2005) suggested that elevated levels of lower-chlorinated PCB congeners in feathers may be  
262 associated with differential elimination and distribution mechanisms.

### 263 3.2. *Intraspecific variation*

264 We aimed at elucidating intraspecific variation in OC exposure through linear regression of  
265 concentrations versus the dietary proxies (SI values) and also by plotting the individuals of  
266 species on a  $\delta^{13}\text{C}/\delta^{15}\text{N}$  biplot (Figure 3a). The distribution of species in the  $\delta^{13}\text{C}/\delta^{15}\text{N}$  biplot  
267 reflects within and among species variations based on the differences in values of dietary  
268 proxies (figure 3a).  $\delta^{13}\text{C}$  values reflect dietary separation of carbon sources, whereas values of  
269  $\delta^{15}\text{N}$  are used as a proxy for their position at trophic food chain (Yu et al., 2011). Aquatic  
270 species i.e. grey heron and great cormorant, in the present study were found to be feeding at a  
271 higher trophic position compared to terrestrial birds ( $p<0.01$ ), which is in line with previous  
272 studies (Hong et al., 2014; Jaspers et al., 2007). However, relatively scattered distribution  
273 (figure 3a) of the individuals of aquatic birds depicted their wide dietary flexibility. Earlier,  
274 based on stable isotope characterization, Sørmo et al., (2011) found that the diet of coastal  
275 herring gull (*Larus argentatus*) was influenced by terrestrial sources. Moreover, Morkūnė et al.,  
276 (2011) reported that great cormorant switched its diet at various stages of life whereas grey  
277 herons showed consistent dietary habits throughout their life span. We suspect that higher  
278 trophic positions on  $\delta^{13}\text{C}/\delta^{15}\text{N}$  layout and relatively scattered distribution of aquatic birds in our  
279 study were because of their more specialized dietary habits as well as varying exposure when  
280 compared to terrestrial species. In contrast, individuals of Indian vulture and white-rumped  
281 vultures were found with a relatively clustered distribution in the  $\delta^{13}\text{C}/\delta^{15}\text{N}$  biplot which might  
282 be associated with more specialized dietary habits (Yu et al., 2011). We sampled these two  
283 vultures from their isolated remote colonies where they mostly consumed the locally available  
284 carrions, which restricts their choice for diverse food sources. Interestingly, Indian vultures  
285 were observed to be feeding at a relatively higher trophic level as compared to white-rumped  
286 vultures suggesting differences in dietary habits of these two species. Similarly, a scattered  
287 distribution of black kites in the *SI* biplot reflects the availability of diverse food choices for  
288 birds dwelling in human proximity and close to urban environments. Earlier, Barón et al.,  
289 (2014) observed black kite as a versatile feeder ranging from human refusals, small insects,  
290 invertebrates, up to small mammals, frogs and snakes in urban and township areas. The results

291 confirm our assumption of exploitation of diverse feeding sources by black kites. Similarly,  
292 Eurasian sparrowhawk, red-necked falcon and common kestrel were also found with relatively  
293 wide ranging distributions in the  $\delta^{13}\text{C} / \delta^{15}\text{N}$  biplot indicating flexibility in food choices for these  
294 species as well (Chen et al., 2009; Elliot et al., 2009, Luzardo et al., 2014). Among two owl  
295 species, spotted owl residing in urban and suburban localities depicted a relatively more  
296 scattered distribution in the  $\delta^{13}\text{C} / \delta^{15}\text{N}$  biplot suggesting its dietary flexibility compared to tight  
297 clustering of little owl. Certain overlap among terrestrial species but not aquatic species is  
298 obvious from the  $\delta^{13}\text{C} / \delta^{15}\text{N}$  biplot (figure 3a) suggesting their potential sympatric distribution  
299 (Zhang et al., 2011) as well as shared feeding sources (Elliott et al., 2009). Distribution of  
300 individuals birds of different predatory species on  $\delta^{13}\text{C} / \delta^{15}\text{N}$  layout suggest that OC  
301 bioaccumulation is considerably influenced by habitat and dietary exposure in addition to  
302 different other factors. Regression analysis revealed a weak and non-significant (except few) but  
303 positive association between dietary proxies and concentration of compounds analyzed ( $R^2$   
304 ranged between 0.01 to 0.99). For  $\delta^{15}\text{N}$ , regression was significant in black kite (for PCBs;  
305  $R^2=0.44$  and HCB;  $R^2=0.42$ , both  $p<0.01$ ) and great cormorant (for DDTs;  $R^2=0.42$ ,  $p=0.02$ ).  
306 Conversely, for  $\delta^{13}\text{C}$  values regression was only significant in Eurasian sparrowhawk (for  
307 PCBs,  $p<0.03$ ; and DDTs  $p<0.04$ ). The regressions between dietary proxies ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) and  
308 OCs concentrations were found non-significant ( $p>0.05$ ) for other studied species.

309 Locational differences of POPs accumulation were also done for three species for which  
310 samples were available for comparison between sites. We have only three species, i.e. black  
311 kite, Eurasian sparrowhawk and spotted owl, for comparison between two different locations  
312 because all other species were sampled from only one site (table S1). For black kite, significant  
313 differences for  $\delta^{13}\text{C}$  ( $p=0.03$ ),  $\delta^{15}\text{N}$  ( $p=0.02$ ) and HCB ( $p=0.02$ ) were observed between sites.  
314  $\sum\text{DDTs}$  at Lahore and  $\sum\text{PCBs}$  at Rawalpindi were found slightly higher although not  
315 significantly different ( $p>0.05$ ) between sites which is possibly associated with significant  
316 differences for the values of dietary proxies. None of the compounds nor stable isotope values  
317 differed significantly ( $p>0.05$ ) between sites for spotted owl or Eurasian sparrowhawk  
318 indicating similar exposure to pollutants at both sites. The above results for stable isotopes  
319 suggest that black kite, being a more urban dwelling species, may switch its feeding choices  
320 (Barón et al., 2014) based on availability at different sites hence reflect differential exposure to  
321 OCs. Conversely, spotted owl and Eurasian sparrowhawk, which remain consistent between

322 sites due to lower availability of choices at suburban to forested sites hence reflect similar  
323 exposure to OCs.

### 324 3.3. *Interspecific differences*

325 In the present study, highest median concentrations (minimum-maximum) of  $\sum$ PCBs at  $7.9 \text{ ng g}^{-1}$   
326  $\text{dry wt.}$  ( $0.4\text{-}15.4 \text{ ng g}^{-1} \text{ dry wt.}$ ) in red-necked falcon,  $\sum$ DDTs at  $195.5 \text{ ng g}^{-1} \text{ dry wt.}$  ( $7.1\text{-}$   
327  $1022.2 \text{ ng g}^{-1} \text{ dry wt.}$ ) in common kestrel and HCB in Eurasian sparrowhawk at  $0.7 \text{ ng g}^{-1} \text{ dry}$   
328  $\text{wt.}$  ( $0.1\text{-}34.4 \text{ ng g}^{-1} \text{ dry wt.}$ ) were recorded. All three species of the current study are terrestrial  
329 predators that mainly feed upon small birds, rats, mouse, frogs, snakes and invertebrates with  
330 some flexibility in their dietary choices (Behrooz et al., 2009). Further, these species mainly  
331 reside in suburbs and agricultural lands around cities where they can get their prey easily.  
332 Relatively higher concentrations of OCPs in common kestrel and Eurasian sparrowhawk in  
333 particular is possibly attributed to their higher dietary exposure to agricultural used pesticides  
334 (Behrooz et al., 2009). This was further corroborated through the significant regression between  
335 origin of dietary carbon ( $\delta^{13}\text{C}$ ) and PCBs ( $R^2=0.45, p=0.03$ ) as well as with DDTs ( $R^2=0.41;$   
336  $p=0.04$ ) in Eurasian sparrowhawk. Regression was not significant for common kestrel possibly  
337 because of limited movement and exposure of this species (Jaspers et al., 2007). However,  
338 relatively higher concentrations of PCBs in red-necked falcon must be considered with caution  
339 because of the low sample size. Moreover, it is observed that in winter Eurasian sparrowhawk  
340 and red-necked falcon move towards towns and cities from the agricultural lands to overcome  
341 winter harshness, which increases their exposure to the urban sources of OCs (Chen et al.,  
342 2009). Based on their utility and disposal, PCBs in particular and HCB up to some extent are  
343 originating from urban sources and hence bioaccumulate in the tissues of top predators during  
344 their winter feeding exposure at temporary stopover sites. To the best of our knowledge, the  
345 exposure to PCBs and OCPs in vultures has never been studied. In the present study, we  
346 sampled both vulture species from their remotely located colonies from Sindh province (figure  
347 1, table S1), where the exposure to urban as well as agricultural chemicals is expected to be  
348 minimal, corresponding to the lower concentrations of PCBs and OCPs in vultures of the  
349 current study. In contrast, the concentrations of PCBs and OCPs were found relatively higher in  
350 black kite and spotted owlet because of higher exposure of these urban/suburban dwelling  
351 species to both agriculture and urban sources of OCs (Barón et al., 2014). The concentrations of  
352  $\sum$ DDTs but not  $\sum$ PCBs were found relatively higher in aquatic birds of our study suggesting

353 that the surplus use of OCPs and environmental leaching of PCBs is more bioavailable in water  
354 reservoirs than terrestrial food chain (Rajaei et al., 2011). Unexpectedly, we have observed  
355 relatively high concentrations of  $\sum$ DDTs in grey herons, which were sampled from a relatively  
356 high altitude pristine location (figure 1, table S1), suggesting that these compounds may also  
357 move towards high laying areas through long range transport from their origin (Wania and  
358 Mackay 1996).

359 Previously, interspecific variation in contaminant levels was attributed to the combined  
360 influence of several biological, ecological and spatial factors (Eulaers et al., 2013, 2014; Peng et  
361 al., 2015). In the present study, we have evaluated the importance of various factors, such as  
362 trophic level, taxonomic affiliation, habitat and feeding regime, as drivers of OC exposure, and  
363 have investigated how values of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  may specifically serve as dietary tracers that  
364 govern difference in OC concentrations among species. Concentrations of compounds such as  
365  $\sum$ PCBs and  $\sum$ DDTs, but not HCB ( $p=0.08$ ), differed significantly among species ( $p<0.01$  for  
366 both) as well as trophic guilds (all  $p<0.03$ ). Multiple comparison (Tukey HSD) test revealed that  
367 more specialized predators such as Eurasian sparrowhawk and red-necked falcon differed  
368 significantly from omnivorous and piscivorous birds (Table 1). Among families, only  $\sum$ PCBs  
369 ( $p<0.01$ ), but not  $\sum$ DDTs ( $p<0.88$ ) nor HCB ( $p<0.82$ ), were significantly different. The  
370 bioaccumulation trend of PCBs in falconidae family was found significantly different from all  
371 other families. Conversely, the corresponding differences between habitats and feeding regimes  
372 were non-significant for  $\sum$ PCBs ( $p<0.27$ ;  $p<0.91$ ),  $\sum$ DDTs ( $p<0.45$ ;  $p<0.62$ ) and HCB ( $p<0.45$ ;  
373  $p<0.62$ ) respectively (Table 1). Values for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  are found significantly different  
374 ( $p<0.01$ ) for all the above mentioned variables, except for feeding regime (both  $p=0.22$ ).  
375 Further, associations between dietary habits of species and contaminants concentrations were  
376 obtained by regressing the  $\log_{10}$  normalized concentrations for  $\sum$ PCBs,  $\sum$ DDTs and HCB with  
377  $\delta^{15}\text{N}$  values (figure 3b, c, d). Regression slopes for species depicted that association of log  
378 normalized  $\sum$ PCBs and  $\sum$ DDTs with  $\delta^{15}\text{N}$  values were stronger when compared to HCBs. This  
379 suggest that different OC compounds vary in their potential of bioaccumulation in birds. This  
380 bioaccumulation differences could possibly be attributed to differences of exposure and  
381 physicochemical properties of compounds (Behrooz et al., 2009). However, species feeding at  
382 different trophic level ( $\delta^{15}\text{N}$  values) depicted similar trends for each  $\sum$ PCBs,  $\sum$ DDTs and HCBs  
383 accumulation except for few species. In case of common kestrel, regression slope was positive

384 for  $\sum$ PCBs while it is relatively straight or negative for  $\sum$ DDTs and HCB. Alternatively,  
385 regression slope for  $\sum$ PCBs was negative for Indian vulture followed by positive and straight  
386 lines for  $\sum$ DDTs and HCB respectively. Regression slopes for all other species somehow  
387 depicted similar bioaccumulation trend for  $\sum$ PCBs,  $\sum$ DDTs and HCB with varying degree of  
388 positive or negative trends. Although there is no clear differences, as depicted by slopes that  
389 bioaccumulation of PCBs and OCPs vary between terrestrial and aquatic species, however the  
390 bioaccumulation trend (slopes) for grey heron but not greater cormorant were somehow similar  
391 to most of the terrestrial species. This differences between two aquatic species could be  
392 attributed to sufficient terrestrial exposure (Ito et al., 2013) and potential influence of terrestrial  
393 feeding sources in grey heron (Sørmo et al., 2011) compared to great cormorant which strictly  
394 rely on aquatic food sources. We have also evaluated the importance of different variables in  
395 explaining the magnitude of exposure of PCBs and OCPs in feathers of predatory birds through  
396 their respective AIC values (table 3). In different models which are separately run for each of  
397 the compounds, variables with the lowest AIC value (shown bold in table 3) best explain the  
398 observed variation in concentrations of the different compounds. The models suggested that the  
399 concentrations of  $\sum$ PCBs, *p,p'*-DDE, *p,p'*-DDT and  $\sum$ DDTs are best explained by the variable  
400 species. But we have run separate models for each compounds assuming that physicochemical  
401 differences of these compounds may varyingly influenced by factors governing their  
402 bioaccumulation. Interestingly, most of the PCB congeners are best explained either by trophic  
403 guild,  $\delta^{13}\text{C}$  values or taxonomic affiliation of species. Concentration differences of lower-  
404 chlorinated compounds were found to be more governed by trophic guilds. Earlier, Behrooz et  
405 al., (2009) suggested that the interspecific variations in OC concentrations are mainly due to  
406 differences in feeding habits of the species. Very few congeners, i.e. CB 128 and 187, were best  
407 predicted by habitat or feeding regime differences. For future studies, we recommend to further  
408 elucidate the factors best predicting the bioaccumulation of OCs by investigating large sample  
409 size for each species.

#### 410 3.4. Variation based on feather type

411 To present date, very few studies have reported the differential accumulation pattern of OCs in  
412 different feather types of predatory birds. Earlier, Jaspers et al. (2011) found significant  
413 differences in levels of different organic contaminants among different feathers types of white-



414 tailed eagles from Greenland. Similarly, [Eulaers et al. \(2014b\)](#) also detected varying  
415 accumulation trends of OCs in different feather types of barn owl (*Tyto alba*) from Belgium.  
416 Based on the assumption that OCs bioaccumulate differently among feathers types, we tested  
417 the concentrations of PCBs and OCPs as well as values of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  among body, tail,  
418 primary and secondary feathers of six individuals of black kite. The concentration pattern of  
419 detected compounds in different types of feathers of black kite is shown in figure S5 (a-d)  
420 whereas detection frequencies are shown in figure S1b. Detection frequencies for body and tail  
421 feathers were similar, but higher than wing (primary and secondary) feathers. The concentration  
422 trend of OCs in different feathers types was also found as  $\sum\text{DDTs} > \sum\text{PCBs} > \text{HCB} > \text{TN}$ . Test  
423 results for significant differences of OCs among different feather types are presented in table 2.

424 Analysis of variance revealed that concentrations of  $\sum\text{PCBs}$ ,  $\sum\text{DDTs}$ , TN differed significantly  
425 ( $p < 0.01$  for all three compounds) among body, tail, primary and secondary feathers. Whereas  
426 differences were non-significant for HCB ( $p = 0.86$ ),  $\delta^{13}\text{C}$  ( $p = 0.65$ ) and  $\delta^{15}\text{N}$  ( $p = 0.64$ ) among  
427 feathers types. Among the different feather types, body feathers were found with highest mean  
428 concentrations of  $\sum\text{PCBs}$ ,  $\sum\text{DDTs}$ , TN and  $\delta^{13}\text{C}$  whereas HCB was highest in secondary  
429 feathers (table 2). Based on the higher detected concentrations and detection frequencies, we  
430 believe that body feathers could be used as a most useful tool for future biomonitoring studies in  
431 predatory birds as suggested earlier by [Jaspers et al., \(2011\)](#). However, we urge for a more  
432 elaborate investigation in the future with larger sample sizes to confirm this. [Eulaers et al.,](#)  
433 [\(2014b\)](#) suggested that relatively elevated concentrations of OCs in tail and body compared to  
434 primary feathers are mainly associated with preening activity and moult pattern of the barn owl.  
435 Further, external contamination through preening has also been suggested to alter the level of  
436 OCs in feathers ([Jaspers et al., 2008; 2013](#)). The specific moulting pattern of different feathers  
437 types in black kite is currently unknown. However, we predict that a higher influence of  
438 preening activity on body and tail feathers of black kites because of their proximity to preen  
439 gland or beak ([Eulaers et al., 2014b](#)) may be associated with higher concentrations of OCs in  
440 these feather types. It has been suggested that moulting pattern and age ([Jaspers et al., 2011](#)) as  
441 well as length and growth of feathers ([Bortolotti et al., 2010](#)) are key factors to describe the  
442 variations in OCs levels among different types of feathers. Besides these, various confounding  
443 factors such as differences in moulting strategy ([Jaeger et al., 2013](#)) as well as feeding and  
444 migratory habits ([García-Fernández et al., 2013](#)) of species can influence the levels of OCs in

445 feathers. Although we have not evaluated any of the above mentioned factors in this study  
446 because of lack of information during data collection, we suspect that a combined effect of just  
447 mentioned factors may be responsible for differences in bioaccumulation patterns of OCs. In  
448 future studies, we suggest to further elucidate the factors responsible for differential  
449 accumulation of OCs among various feather types.

450

#### 451 **4. Conclusions**

452 The present study is the first to report the levels of PCBs and OCPs in predatory birds of  
453 Pakistan. Various contributing factors explaining the intra- and interspecific differences in PCBs  
454 and OCPs levels in predatory birds were also evaluated. We concluded that PCBs and OCPs  
455 could easily be quantified in predatory birds using feathers, particularly body feathers because of  
456 their high detectability. Significant differences in concentrations of PCBs and some OCPs among  
457 different feathers types emphasize the need for appropriate feather choice in future toxicological  
458 studies. Compared to PCBs, concentrations of OCPs, particularly DDTs, were found higher in  
459 predatory birds reflecting the large scale historical and current application of pesticides in  
460 Pakistan. In general, PCB levels reported in predatory birds of Pakistan are found lower than  
461 those of European studies whereas OCPs are relatively comparable. To get a broader picture, we  
462 urge future research to investigate the significance of contributing factors influencing the levels  
463 of OCs using multiple species of predatory birds from a wide geographical range.

#### 464 **Conflict of interest**

465 The authors declare no conflicts with any person or organization.

#### 466 **Supplementary information**

467 All the supporting material cited in this manuscript is available in the online supplementary files.

#### 468 **Acknowledgements**

469 The authors are highly thankful to Higher Education Commission (HEC) Pakistan for providing  
470 the necessary funding for the project. This work was completed in collaboration with Quaid-i-  
471 Azam University, Islamabad, Pakistan, Norwegian university of Science and Technology  
472 (NTNU), and the University of Antwerp, Belgium. Sakhawat Ali and Shahid Iqbal are highly  
473 acknowledged for their support during sampling of this project.

474

475 **References**

- 476 Abbasi, N. A., Malik, R. N., Frantz, A., Jaspers, V.L.B., 2016. Science of the Total Environment  
 477 A review on current knowledge and future prospects of organohalogen contaminants  
 478 (OHCs) in Asian birds. *Sci. Total Environ.* 542, 411–426.  
 479 doi:10.1016/j.scitotenv.2015.10.088
- 480 Ali, U., Syed, J.H., Malik, R.N., Katsoyiannis, A., Li, J., Zhang, G., Jones, K.C., 2014.  
 481 Organochlorine pesticides (OCPs) in South Asian region: A review. *Sci. Total Environ.* 476-  
 482 477, 705–717. doi:10.1016/j.scitotenv.2013.12.107
- 483 Barón, E., Máñez, M., Andreu, A.C., Sergio, F., Hiraldo, F., Eljarrat, E., Barceló, D., 2014.  
 484 Bioaccumulation and biomagnification of emerging and classical flame retardants in bird  
 485 eggs of 14 species from Doñana Natural Space and surrounding areas (South-western  
 486 Spain). *Environ. Int.* 68, 118–126. doi:10.1016/j.envint.2014.03.013
- 487 Behrooz, R.D., Esmaili-Sari, A., Ghasempouri, S.M., Bahramifar, N., Covaci, A., 2009.  
 488 Organochlorine pesticide and polychlorinated biphenyl residues in feathers of birds from  
 489 different trophic levels of South-West Iran. *Environ. Int.* 35, 285–290.  
 490 doi:10.1016/j.envint.2008.07.001
- 491 Boecklen, W.J., Yarnes, C.T., Cook, B.A., James, A.C., 2011. On the use of stable isotopes in  
 492 trophic ecology. *Annu. Rev. Ecol. Evol. Syst.* 42, 411–40.
- 493 Bortolotti, G.R., 2010. Flaws and pitfalls in the chemical analysis of feathers: bad news–good  
 494 news for avian chemoecology and toxicology. *Ecol. Appl.* 20, 1766–74.
- 495 Chen, D., Zhang, X., Mai, B., Sun, Q., Song, J., Luo, X., Zeng, E.Y., Hale, R.C., 2009.  
 496 Polychlorinated biphenyls and organochlorine pesticides in various bird species from  
 497 northern China. *Environ. Pollut.* 157, 2023–2029. doi:10.1016/j.envpol.2009.02.027
- 498 Corsolini, S., Covaci, A., Ademollo, N., Focardi, S., Schepens, P., 2006. Occurrence of  
 499 organochlorine pesticides (OCPs) and their enantiomeric signatures, and concentrations of  
 500 polybrominated diphenyl ethers (PBDEs) in the Adélie penguin food web, Antarctica.  
 501 *Environ. Pollut.* 140, 371–382. doi:10.1016/j.envpol.2005.04.039
- 502 Covaci, A., Gheorghe, A., Hulea, O., Schepens, P., 2006. Levels and distribution of  
 503 organochlorine pesticides, polychlorinated biphenyls and polybrominated diphenyl ethers in  
 504 sediments and biota from the Danube Delta, Romania. *Environ. Pollut.* 140, 136–149.  
 505 doi:10.1016/j.envpol.2005.06.008
- 506 Dauwe, T., Jaspers, V.L.B., Covaci, A., Schepens, P., Eens, M., 2005. Feathers As a  
 507 Nondestructive Biomonitor for Persistent Organic Pollutants. *Environ. Toxicol. Chem.* 24,  
 508 442-449. doi:10.1897/03-596.1
- 509 Elliott, K.H., Cesh, L.S., Dooley, J. a., Letcher, R.J., Elliott, J.E., 2009. PCBs and DDE, but not  
 510 PBDEs, increase with trophic level and marine input in nestling bald eagles. *Sci. Total*  
 511 *Environ.* 407, 3867–3875. doi:10.1016/j.scitotenv.2009.02.027
- 512 Eulaers, I., Jaspers, V.L.B., Bustnes, J.O., Covaci, A., Johnsen, T. V., Halley, D.J., Moum, T.,  
 513 Ims, R. a., Hanssen, S. a., Erikstad, K.E., Herzke, D., Sonne, C., Ballesteros, M., Pinxten,  
 514 R., Eens, M., 2013. Ecological and spatial factors drive intra- and interspecific variation in  
 515 exposure of subarctic predatory bird nestlings to persistent organic pollutants. *Environ. Int.*  
 516 57-58, 25–33. doi:10.1016/j.envint.2013.03.009
- 517 Eulaers, I., Jaspers, V.L.B., Halley, D.J., Lepoint, G., Nygård, T., Pinxten, R., Covaci, A., Eens,  
 518 M., 2014a. Brominated and phosphorus flame retardants in White-tailed Eagle *Haliaeetus*

519 *albicilla* nestlings: Bioaccumulation and associations with dietary proxies ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and  
520  $\delta^{34}\text{S}$ ). *Sci. Total Environ.* 478, 48–57. doi:10.1016/j.scitotenv.2014.01.051

521 Eulaers, I., Jaspers, V.L.B., Pinxten, R., Covaci, A., Eens, M., 2014b. Legacy and current-use  
522 brominated flame retardants in the Barn Owl. *Sci. Total Environ.* 472, 454–462.  
523 doi:10.1016/j.scitotenv.2013.11.054

524 Furness, R. W. 1993. In birds as monitor of environmental change (ed. R. W., Furness and J. J.  
525 D., Greenwood) pp. 86-143. London, UK: Chapman and Hall.

526 García-Fernández, A.J., Espín, S., Martínez-López., E., 2013. Feathers as a biomonitoring tool of  
527 polyhalogenated compounds: A review. *Environ. Sci. Technol.* 47, 3028–3043.  
528 doi:10.1021/es302758x

529 Guan, Y.F., Wang, J.Z., Ni, H.G., Zeng, E.Y., 2009. Organochlorine pesticides and  
530 polychlorinated biphenyls in riverine runoff of the Pearl River Delta, China: Assessment of  
531 mass loading, input source and environmental fate. *Environ. Pollut.* 157, 618–624.  
532 doi:10.1016/j.envpol.2008.08.011

533 Hong, S.H., Shim, W.J., Han, G.M., Ha, S.Y., Jang, M., Rani, M., Hong, S., Yeo, G.Y., 2014.  
534 Levels and profiles of persistent organic pollutants in resident and migratory birds from an  
535 urbanized coastal region of South Korea. *Sci. Total Environ.* 470-471, 1463–1470.  
536 doi:10.1016/j.scitotenv.2013.07.089

537 Huang, K., Lin, K., Guo, J., Zhou, X., Wang, J., Zhao, J., Zhou, P., Xu, F., Liu, L., Zhang, W.,  
538 2013. Polybrominated diphenyl ethers in birds from Chongming Island, Yangtze estuary,  
539 China: Insight into migratory behavior. *Chemosphere* 91, 1416–1425.  
540 doi:10.1016/j.chemosphere.2013.01.042

541 Ito, A., Yamashita, R., Takada, H., Yamamoto, T., Shiomi, K., Zavalaga, C., Abe, T., Watanabe,  
542 S., Yamamoto, M., Sato, K., Kohno, H., Yoda, K., Iida, T., Watanuki, Y., 2013.  
543 Contaminants in tracked seabirds showing regional patterns of marine pollution. *Environ.*  
544 *Sci. Technol.* 47, 7862–7867. doi:10.1021/es4014773

545 Jaeger, A., Jaquemet, S., Phillips, R., Wanless, R., Richard, P., Cherel, Y., 2013. Stable isotopes  
546 document inter- and intra-specific variation in feeding ecology of nine large southern  
547 Procellariiformes. *Mar. Ecol. Prog. Ser.* 490, 255–266. doi:10.3354/meps10436

548 Jardine, T.D., Kidd, K.A., Fisk, A.T., 2006. Critical Review Applications, Considerations, and  
549 Sources of Uncertainty When Using Stable Isotope Analysis in Ecotoxicology 40, 7501–  
550 7511.

551 Jaspers, V.L.B., Voorspoels, S., Covaci, A., Eens, M., 2006. Can predatory bird feathers be used  
552 as a non-destructive biomonitoring tool of organic pollutants? *Biol. Lett.* 2, 283–285.  
553 doi:10.1098/rsbl.2006.0450

554 Jaspers, V.L.B., Voorspoels, S., Covaci, A., Lepoint, G., Eens, M., 2007. Evaluation of the  
555 usefulness of bird feathers as a non-destructive biomonitoring tool for organic pollutants : A  
556 comparative and meta-analytical approach 33, 328–337. doi:10.1016/j.envint.2006.11.011

557 Jaspers, V.L.B., Covaci, A., Deleu, P., Neels, H., Eens, M., 2008. Preen oil as the main source of  
558 external contamination with organic pollutants onto feathers of the common magpie (*Pica*  
559 *pica*). *Environ. Int.* 34, 741–748. doi:10.1016/j.envint.2007.12.002

560 Jaspers, V.L.B., Covaci, A., Deleu, P., Eens, M., 2009. Concentrations in bird feathers reflect  
561 regional contamination with organic pollutants Local contamination. *Sci. Total Environ.*  
562 407, 1447–1451. doi:10.1016/j.scitotenv.2008.10.030

563 Jaspers, V.L.B., Rodriguez, F.S., Boertmann, D., Sonne, C., Dietz, R., Rasmussen, L.M., Eens,  
564 M., Covaci, A., 2011. Body feathers as a potential new biomonitoring tool in raptors: A

565 study on organohalogenated contaminants in different feather types and preen oil of West  
566 Greenland white-tailed eagles (*Haliaeetus albicilla*). Environ. Int. 37, 1349–1356.  
567 doi:10.1016/j.envint.2011.06.004

568 Jaspers, V.L.B., Sonne, C., Soler-Rodriguez, F., Boertmann, D., Dietz, R., Eens, M., Rasmussen,  
569 L.M., Covaci, a., 2013. Persistent organic pollutants and methoxylated polybrominated  
570 diphenyl ethers in different tissues of white-tailed eagles (*Haliaeetus albicilla*) from West  
571 Greenland. Environ. Pollut. 175, 137–146. doi:10.1016/j.envpol.2012.12.023

572 Johnson, J.B., Omland, K.S., 2004. Model selection in ecology and evolution. 19, 101-108  
573 doi:10.1016/j.tree.2003.10.013

574 Khan, M., Mohammad, A., Ahad, K., Katsoyiannis, A., Malik, S.A., Abdullaha, M., Rashid, A.,  
575 Fasola, M., Hussain, A., Bokhari, H., Eqani, S.A.M.A.S., 2013. Cattle egrets as a  
576 biosentinels of persistent organic pollutants exposure. Environ. Geochem. Health. 36, 375–  
577 384. doi:10.1007/s10653-013-9556-5

578 Kocagöz, R., Onmuş, O., Onat, İ., Çağdaş, B., Sıkı, M., Orhan, H., 2014. Environmental and  
579 biological monitoring of persistent organic pollutants in waterbirds by non-invasive  
580 versus invasive sampling. Toxicol. Lett. 230, 208–217.

581 Lavoie, R.A., Champoux, L., Rail, J., Lean, D.R.S., 2010. Organochlorines, brominated flame  
582 retardants and mercury levels in six seabird species from the Gulf of St. Lawrence (Canada):  
583 Relationships with feeding ecology, migration and molt 158, 2189–2199.  
584 doi:10.1016/j.envpol.2010.02.016

585 Letcher, R.J., Bustnes, J.O., Dietz, R., Jenssen, B.M., Jørgensen, E.H., Sonne, C., Verreault, J.,  
586 Vijayan, M.M., Gabrielsen, G.W., 2010. Exposure and effects assessment of persistent  
587 organohalogen contaminants in arctic wildlife and fish. Sci. Total Environ. 408, 2995–3043.  
588 doi:10.1016/j.scitotenv.2009.10.038

589 Lohmann, R., Breivik, K., Dachs, J., Muir, D., 2007. Global fate of POPs: Current and future  
590 research directions. Environ. Pollut. 150, 150–165. doi:10.1016/j.envpol.2007.06.051

591 Luzardo, O.P., Ruiz-Suárez, N., Henríquez-Hernández, L.A., Valerón, P.F., Camacho, M.,  
592 Zumbado, M., Boada, L.D., 2014. Assessment of the exposure to organochlorine pesticides,  
593 PCBs and PAHs in six species of predatory birds of the Canary Islands, Spain. Sci. Total  
594 Environ. 472, 146–153. doi:10.1016/j.scitotenv.2013.11.021

595 Malik, R.N., Rauf, S., Mohammad, A., Eqani, S.A.M.A.S., Ahad, K., 2011. Organochlorine  
596 residual concentrations in cattle egret from the Punjab Province, Pakistan. Environ. Monit.  
597 Assess. 173, 325–341. doi:10.1007/s10661-010-1390-y

598 Morkūnė, R., 2011. Trophic peculiarities of the Great Cormorant, Grey Heron and Long-tailed  
599 Duck on the Baltic Sea Lithuanian coast : a stable isotope approach 57, 173–178.

600 Muir, D., Braune, B., DeMarch, B., Norstrom, R., Wagemann, R., Lockhart, L., Hargrave, B.,  
601 Bright, D., Addison, R., Payne, J., Reimer, K., 1999. Spatial and temporal trends and effects  
602 of contaminants in the Canadian Arctic marine ecosystem: A review. Sci. Total Environ.  
603 230, 83–144. doi:10.1016/S0048-9697(99)00037-6

604 Peng, Y., Wu, J., Tao, L., Mo, L., Tang, B., Zhang, Q., Luo, X., Zou, F., Mai, B., 2015.  
605 Contaminants of legacy and emerging concern in terrestrial passerines from a nature reserve  
606 in South China : Residue levels and inter-species differences in the accumulation. Environ.  
607 Pollut. 203, 7–14. doi:10.1016/j.envpol.2015.03.037

608 Rajaei, F., Sari, A.E., Bahramifar, N., Savabieasfahani, M., Ghasempouri, M., 2011. Persistent  
609 organic pollutants in muscle and feather of ten avian species from Mazandaran province of

610 Iran, on the coast of the Caspian sea. *Bull. Environ. Contam. Toxicol.* 87, 678–683.  
611 doi:10.1007/s00128-011-0420-y

612 Sanpera, C., Ruiz, X., Jover, L., Llorente, G., Jabeen, R., Muhammad, A., Boncompagni, E.,  
613 Fasola, M., 2003. Persistent organic pollutants in little egret eggs from selected wetlands in  
614 Pakistan. *Arch. Environ. Contam. Toxicol.* 44, 360–368. doi:10.1007/s00244-002-2044-z

615 Sarkar, S.K., Bhattacharya, B.D., Bhattacharya, a., Chatterjee, M., Alam, a., Satpathy, K.K.,  
616 Jonathan, M.P., 2008. Occurrence, distribution and possible sources of organochlorine  
617 pesticide residues in tropical coastal environment of India: An overview. *Environ. Int.* 34,  
618 1062–1071. doi:10.1016/j.envint.2008.02.010

619 Sørmo, E.G., Lie, E., Ruus, A., Gaustad, H., Skaare, J.U., Jenssen, B.M., 2013. Ecotoxicology  
620 and Environmental Safety Trophic level determines levels of brominated flame-retardants in  
621 coastal herring gulls. *Ecotoxicol. Environ. Saf.* 74, 2091–2098.  
622 doi:10.1016/j.ecoenv.2011.06.012

623 Sun, Y.X., Hao, Q., Zheng, X.B., Luo, X.J., Zhang, Z.W., Zhang, Q., Xu, X.R., Zou, F.S., Mai,  
624 B.X., 2014. PCBs and DDTs in light-vented bulbuls from Guangdong Province, South  
625 China: Levels, geographical pattern and risk assessment. *Sci. Total Environ.* 490, 815–821.  
626 doi:10.1016/j.scitotenv.2014.05.066

627 Tanabe, S., Senthilkumar, K., Kannan, K., Subramanian, A.N., 1998. Accumulation features of  
628 polychlorinated biphenyls and organochlorine pesticides in resident and migratory birds  
629 from South India. *Arch. Environ. Contam. Toxicol.* 34, 387–397.

630 UNEP, 2011. Draft revised guidance on the global monitoring plan for persistent organic  
631 pollutants, UNEP/POPS/COP.5/INF/27. United Nations Environment Programme. UNEP  
632 Chemicals Geneva, Switzerland.

633 Vorkamp, K., Rigét, F.F., 2014. A review of new and current-use contaminants in the Arctic  
634 environment: Evidence of long-range transport and indications of bioaccumulation.  
635 *Chemosphere* 111, 379–395. doi:10.1016/j.chemosphere.2014.04.019

636 Wania, F., Mackay, D., 1996. Tracking the distribution of persistent organic pollutants. *Environ. Sci. Technol.*  
637 30, 390A–396A. doi:10.1016/S0926-3373(97)80026-4

638 Wania, F., Mackay, D., 1996. Tracking the distribution of persistent organic pollutants. *Environ.*  
639 *Sci. Technol.* 30, 390A–396A. doi:10.1016/S0926-3373(97)80026-4

640 Yadav, I.C., Devi, N.L., Syed, J.H., Cheng, Z., Li, J., Zhang, G., Jones, K.C., 2015. Current  
641 status of persistent organic pesticides residues in air, water, and soil, and their possible effect  
642 on neighboring countries: A comprehensive review of India. *Sci. Total Environ.* 511, 123–  
643 137. doi:10.1016/j.scitotenv.2014.12.041

644 Yu, L.-H., Luo, X.-J., Wu, J.-P., Liu, L.-Y., Song, J., Sun, Q.-H., Zhang, X.-L., Chen, D., Mai,  
645 B.-X., 2011. Biomagnification of Higher Brominated PBDE Congeners in an Urban  
646 Terrestrial Food Web in North China Based on Field Observation of Prey Deliveries.  
647 *Environ. Sci. Technol.* 45, 5125–5131. doi:10.1021/es200030z

648 Zhang, X.L., Luo, X.J., Liu, J., Luo, Y., Chen, S.J., Mai, B.X., 2011. Polychlorinated biphenyls  
649 and organochlorinated pesticides in birds from a contaminated region in South China:  
650 Association with trophic level, tissue distribution and risk assessment. *Environ. Sci. Pollut.*  
651 *Res.* 18, 556–565. doi:10.1007/s11356-010-0397-9

652 Zuur, A.F., Ieno, E.N., Elphick, C.S., 2010. A protocol for data exploration to avoid common  
653 statistical problems 3–14. doi:10.1111/j.2041-210X.2009.00001.x

**Table 1:** Pairwise comparisons for PCBs, DDTs, HCB (ng g<sup>-1</sup> dry wt.) and stable isotope values (‰) in tail feathers among different groups of birds based on ANOVAs. Means are shown under each variable whereas differences through multiple comparisons are illustrated through different alphabetic characters. Groups with a single different letter differ significantly from the other. Sample sizes in each group are given in parentheses.

Test variable	Compounds/SIs	Test for significance			Mean concentrations and Significance levels (alphabets) among/between groups (Tukey-HSD)									
		df	F	P	BK (10)	WRV (12)	IV (9)	ESH (10)	CK (4)	RNF (2)	GH (6)	GC (4)	SO (10)	LO (6)
Species	$\delta^{13}C$	9,60	6.14	<0.01	-20.49 AB	-17.54 B	-17.53 B	-20.89 AB	-20.34 AB	-24.73 A	-24.00 A	-21.23 AB	-17.20 B	-20.64 AB
	$\delta^{15}N$	9,60	8.28	<0.01	9.01 AB	9.57 ABC	11.51 BCD	8.58 A	8.33 A	8.51 A	12.18 CD	13.60 D	10.67 ABC	9.97 ABC
	$\Sigma PCBs$	9,66	2.55	<0.01	2.05 B	0.45 B	0.57 B	3.56 AB	5.20 AB	8.41 A	4.10 AB	2.46 AB	2.04 B	2.74 AB
	$\Sigma DDTs$	9,66	2.61	<0.01	12.97 B	2.03 B	6.15 B	647.98 A	355.03 AB	43.4 B	55.69 B	13.15 B	132.49 B	25.97 B
	HCB	9,66	1.80	0.08	0.11	0.10	0.06	5.86	0.11	0.29	0.09	0.05	0.14	0.18
Families					Accipitridae (44)		Ardeidae (6)		Falconidae (6)		Phalacrocoracidae (4)		Strigidae (16)	
	$\delta^{13}C$	4,65	5.1	<0.01	-19.17 B		-24.00 A		-21.81 AB		-21.30 AB		-18.76 B	
	$\delta^{15}N$	4,65	9.19	<0.01	9.58 C		12.18 AB		8.39 C		13.60 A		10.35 BC	
	$\Sigma PCBs$	4,71	3.24	<0.01	1.55 B		4.10 AB		6.27 A		2.46 B		2.30 B	
	HCB	4,71	0.37	0.82	152.91		55.69		251.15		13.15		92.55	
Trophic guilds					Omnivorous (13)		Scavengers (21)		Carnivorous (32)		Piscivorous (10)			
	$\delta^{13}C$	3,66	9.95	<0.01	-20.49 AB		-17.53 C		-20.23 B		-23.08 A			
	$\delta^{15}N$	3,66	12.91	<0.01	9.01 B		10.4 B		9.26 B		12.65 A			
	$\Sigma PCBs$	3,72	4.45	<0.01	2.05 AB		0.28 B		3.44 A		3.45 A			
	HCB	3,72	1.09	0.35	12.97 B		3.80 B		295.86 A		38.67 B			
Habitat					Terrestrial (66)					Aquatic (10)				
	$\delta^{13}C$	1,68	12.66	<0.01	19.36					-23.08				
	$\delta^{15}N$	1,68	27.35	<0.01	9.61					12.66				
	$\Sigma PCBs$	1,74	1.19	0.27	2.17					3.45				
	HCB	1,74	0.37	0.54	147.22					38.68				
Feeding regime					Diurnal (60)					Nocturnal (16)				
	$\delta^{13}C$	1,68	1.5	0.22	-20.04					18.77				
	$\delta^{15}N$	1,68	0.44	0.50	9.93					10.35				
	$\Sigma PCBs$	1,74	0.01	0.91	2.34					2.31				
	HCB	1,74	0.58	0.44	143.70					92.55				

**Table 2:** Tests for strength of significance for PCBs and OCPs (ng g<sup>-1</sup> dry wt.) and stable isotope residues (‰) among different feather types of 6 individuals of black kites from Pakistan. Means are shown under each feather types whereas different alphabetic characters are used to illustrate significant differences through multiple comparison (Tukey-HSD) test.

Compounds/SIs	Test for significance			Mean concentrations and Significance levels (alphabets) among feather types (Tukey-HSD)			
	df	F	P	Body	Tail	Primary	Secondary
<b>δ13C</b>	3,20	0.56	0.65	-20.24	-21.55	-20.97	-22.32
<b>δ15N</b>	3,20	0.57	0.64	9.17	10.16	8.68	9.22
<b>ΣPCBs</b>	3,20	5.13	<0.01	4.21 B	2.57 AB	1.48 A	1.92 A
<b>ΣDDTs</b>	3,20	13.57	<0.01	27.57 B	12.17 A	7.68 A	9.97 A
<b>HCB</b>	3,20	0.24	0.86	0.13	0.14	0.14	0.16
<b>TN</b>	3,20	7.37	<0.01	0.05 B	0.03 AB	0.019 A	0.016 A



Table 3. Evaluation of factors governing OCs concentration differences based on Akaike's information criteria (AIC). Variables with lowest AIC (shown in bold) best explained the concentration of the respective compound. Significant differences (ANOVA) of compounds among/between tested variables are shown with bold p values.

		Species	Families	Trophic guilds	Food chain	Feeding regime	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$
CB-105	AIC	116.41	<b>114.42</b>	116.93	115.58	115.87	118.00	119.38
	F <sub>i</sub> (p)	2.16 (0.08)	2.78 ( <b>0.04</b> )	3.99 ( <b>0.02</b> )	8.17 ( <b>0.00</b> )	3.74 (0.06)	NC*	NC
CB-118	AIC	204.85	203.05	<b>198.85</b>	202.92	213.28	210.78	212.81
	F <sub>i</sub> (p)	2.85 ( <b>0.01</b> )	4.51 ( <b>0.00</b> )	5.51 ( <b>0.00</b> )	11.53 ( <b>0.00</b> )	0.85 (0.35)	NC	NC
CB-146	AIC	132.21	128.37	127.41	125.65	124.94	<b>124.24</b>	125.88
	F <sub>i</sub> (p)	0.88 (0.51)	0.82 (0.51)	0.60 (0.55)	0.00 (0.99)	1.56 (0.21)	NC	NC
CB-153	AIC	226.48	240.69	<b>223.07</b>	245.41	251.33	249.54	250.66
	F <sub>i</sub> (p)	2.67 ( <b>0.01</b> )	2.88 ( <b>0.02</b> )	4.05 ( <b>0.01</b> )	1.32 (0.25)	0.07 (0.79)	NC	NC
CB-138	AIC	209.73	214.53	<b>202.81</b>	212.32	215.32	214.25	215.42
	F <sub>i</sub> (p)	1.56 (0.15)	1.38 (0.25)	2.66 ( <b>0.05</b> )	1.10 (0.29)	0.36 (0.55)	NC	NC
CB-187	AIC	114.26	108.47	108.06	<b>106.07</b>	107.79	107.69	107.69
	F <sub>i</sub> (p)	1.27 (0.29)	1.91 (0.14)	1.14 (0.32)	1.24 (0.27)	0.01 (0.89)	NC	NC
CB-183	AIC	84.34	<b>82.36</b>	82.79	83.89	84.09	84.07	84.05
	F <sub>i</sub> (p)	2.07 (0.10)	2.85 ( <b>0.05</b> )	1.34 (0.27)	0.72 (0.40)	0.21 (0.64)	NC	NC
CB-128	AIC	154.67	149.27	148.62	147.42	<b>145.59</b>	146.78	148.07
	F <sub>i</sub> (p)	0.56 (0.77)	0.75 (0.56)	0.24 (0.78)	0.32 (0.57)	2.03 (0.15)	NC	NC
CB-156	AIC	87.59	86.16	87.16	86.22	86.81	<b>84.05</b>	85.09
	F <sub>i</sub> (p)	3.41 ( <b>0.02</b> )	2.45 (0.08)	0.76 (0.47)	0.01 (0.90)	0.75 (0.39)	NC	NC
CB-180	AIC	218.61	215.15	<b>213.21</b>	216.7	216.9	214.6	216.82
	F <sub>i</sub> (p)	1.46 (0.18)	1.45 (0.22)	2.51 (0.06)	1.01 (0.31)	0.12 (0.72)	NC	NC
CB-170	AIC	202.26	196.43	196.79	197.19	197.07	<b>192.91</b>	196.50
	F <sub>i</sub> (p)	1.40 (0.21)	1.58 (0.18)	1.99 (0.12)	0.72 (0.39)	0.00 (0.95)	NC	NC
CB-199	AIC	66.40	64.51	<b>63.31</b>	<b>63.31</b>	66.57	66.66	66.66
	F <sub>i</sub> (p)	1.13 (0.36)	1.78 (0.19)	1.40 (0.25)	1.40 (0.25)	0.24 (0.62)	NC	NC
CB-196/203	AIC	139.60	135.90	136.15	134.23	134.57	<b>133.85</b>	134.52
	F <sub>i</sub> (p)	1.21 (0.32)	1.38 (0.26)	1.55 (0.22)	0.84 (0.36)	0.21 (0.64)	NC	NC
CB-194	AIC	149.63	<b>149.22</b>	150.14	153.12	153.52	150.73	153.69
	F <sub>i</sub> (p)	2.50 ( <b>0.03</b> )	2.90 ( <b>0.04</b> )	2.11 (0.10)	0.72 (0.40)	0.47 (0.49)	NC	NC
$\Sigma\text{PCBs}$	AIC	<b>217.74</b>	273.26	238.12	279.91	283.48	282.77	282.57
	F <sub>i</sub> (p)	2.55 ( <b>0.01</b> )	3.24 ( <b>0.01</b> )	4.45 ( <b>0.00</b> )	1.19 (0.27)	0.01 (0.91)	NC	NC
TN	AIC	71.03	71.03	<b>69.10</b>	75.06	74.02	75.90	80.23
	F <sub>i</sub> (p)	4.91 ( <b>0.00</b> )	4.91 ( <b>0.00</b> )	7.59 ( <b>0.00</b> )	4.40 ( <b>0.04</b> )	7.88 ( <b>0.00</b> )	NC	NC
HCB	AIC	<b>207.83</b>	245.68	233.55	241.42	243.80	242.46	243.95
	F <sub>i</sub> (p)	1.80 (0.08)	0.37 (0.82)	1.09 (0.35)	0.37 (0.54)	0.58 (0.44)	NC	NC
<i>p,p'</i> -DDE	AIC	<b>272.43</b>	315.55	277.44	317.85	317.24	318.66	318.61
	F <sub>i</sub> (p)	2.60 ( <b>0.01</b> )	0.21 (0.92)	2.71 ( <b>0.05</b> )	0.45 (0.50)	0.26 (0.60)	NC	NC
<i>p,p'</i> -DDT	AIC	<b>268.20</b>	292.00	273.85	294.74	295.88	298.21	293.90
	F <sub>i</sub> (p)	2.67 ( <b>0.01</b> )	1.51 (0.20)	3.15 ( <b>0.03</b> )	1.10 (0.29)	0.10 (0.75)	NC	NC
$\Sigma\text{DDTs}$	AIC	<b>263.33</b>	305.87	269.40	308.57	307.31	308.72	308.01
	F <sub>i</sub> (p)	2.61 ( <b>0.01</b> )	0.29 (0.88)	2.92 ( <b>0.03</b> )	0.56 (0.45)	0.24 (0.62)	NC	NC

\* NC (not calculated) are shown where ANOVA could not be quantified




Figure 1. Map showing different sampling sites in Pakistan. The details of birds collected at different sampling sites are given in table S1 in the supplementary information.







Figure 2: Contribution profile (percentage) of PCBs and OCPs in (a) tail feathers of predatory birds and (b) different feather types of black kite from Pakistan.







Figure 3. Scatter plots indicating interspecific differences in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values (a), and species-specific regressions of  $\log_{10}$  transformed concentrations of  $\Sigma\text{PCBs}$  (b),  $\Sigma\text{DDTs}$  (c) and HCB (d) on stable isotope values. Species abbreviations are as follows: BK=black kite, WRV=white-rumped vulture, IV=Indian vulture, ESH=Eurasian sparrowhawk, GH=grey heron, RNF=red-necked falcon, CK=common kestrel, GC=great cormorant; SO=spotted owlet, LO=little owl