



HAL
open science

Biochemical measurements of beef are a good predictor of untrained consumer sensory scores across muscles

S P F Bonny, G E Gardner, David William Pethick, Isabelle Legrand, Rod J. Polkinghorne, Jean-François J.-F. Hocquette

► **To cite this version:**

S P F Bonny, G E Gardner, David William Pethick, Isabelle Legrand, Rod J. Polkinghorne, et al.. Biochemical measurements of beef are a good predictor of untrained consumer sensory scores across muscles. *Animal*, 2015, 9 (1), pp.179-90. 10.1017/S1751731114002389 . hal-02638458

HAL Id: hal-02638458

<https://hal.inrae.fr/hal-02638458>

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biochemical measurements of beef are a good predictor of untrained consumer sensory scores across muscles

S. P. F. Bonny^{1,4,5a}, G. E. Gardner^{1†}, D. W. Pethick¹, I. Legrand², R. J. Polkinghorne³ and J. F. Hocquette^{4,5}

¹School of Veterinary and Life Sciences, Murdoch University, 90 South St, Murdoch, WA 6150, Australia; ²Institut de l'Élevage, Service Qualité des Viandes, MRAL, 87060 Limoges Cedex 2, France; ³Polkinghorne Pty Ltd, 431 Timor Road, Murrurundi, NSW 2338, Australia; ⁴INRA, UMR1213, Recherches sur les Herbivores, F-63122 Saint Genès Champanelle, France; ⁵Clermont Université, VetAgro Sup, UMR1213, Recherches sur les Herbivores, F-63122 Saint Genès Champanelle, France

(Received 18 March 2014; Accepted 4 August 2014; First published online 23 September 2014)

The ability of the biochemical measurements, haem iron, intramuscular fat (IMF%), moisture content, and total, soluble and insoluble collagen contents, to predict untrained consumer sensory scores both across different muscles and within the same muscle from different carcasses were investigated. Sensory scores from 540 untrained French consumers (tenderness, flavour liking, juiciness and overall liking) were obtained for six muscles; outside (m. biceps femoris), topside (m. semimembranosus), striploin (m. longissimus thoracis), rump (m. gluteus medius), oyster blade (m. infraspinatus) and tenderloin (m. psoas major) from each of 18 French and 18 Australian cattle. The four sensory scores were weighted and combined into a single score termed MQ4, which was also analysed. All sensory scores were highly correlated with each other and with MQ4. This in part reflects the fact that MQ4 is derived from the consumer scores for tenderness, juiciness, flavour and overall liking and also reflects an interrelationship between the sensory scores themselves and in turn validates the use of the MQ4 term to reflect the scope of the consumer eating experience. When evaluated across the six different muscles, all biochemical measurements, except soluble collagen, had a significant effect on all of the sensory scores and MQ4. The average magnitude of impact of IMF%, haem iron, moisture content, total and insoluble collagen contents across the four different sensory scores are 34.9, 5.1, 7.2, 36.3 and 41.3, respectively. When evaluated within the same muscle, only IMF% and moisture content had a significant effect on overall liking (5.9 and 6.2, respectively) and flavour liking (6.1 and 6.4, respectively). These results indicate that in a commercial eating quality prediction model including muscle type, only IMF% or moisture content has the capacity to add any precision. However, all tested biochemical measurements, particularly IMF% and insoluble collagen contents, are strong predictors of eating quality when muscle type is not known. This demonstrates their potential usefulness in extrapolating the sensory data derived from these six muscles to other muscles with no sensory data, but with similar biochemical parameters, and therefore reducing the amount of future sensory testing required.

Keywords: beef, biochemistry, muscle, untrained consumers, sensory scores

Implications

In a commercial eating quality prediction model including muscle type, our results indicate that only IMF% or H₂O%, unlike iron or collagen content, would add any precision. Alternatively, all biochemical measurements are strong predictors of eating quality when muscle type is not known. Therefore, for extrapolating sensory data derived from the six muscles tested in this study to other muscles with no sensory

data, insoluble collagen content might indeed be useful for predicting the unknown muscle and therefore reduce the amount of future testing required.

Introduction

For beef to remain competitive in the market place, the product must satisfy the demands of consumers. Variable eating quality is seen as a major factor in the decline in beef consumption (Morgan *et al.*, 1991, Polkinghorne *et al.*, 2008). A system that could guarantee beef eating quality would be well accepted by the European beef consumers

^a Present address: School of Veterinary and Life Sciences, Murdoch University, 90 South St, Murdoch, WA 6150, Australia.

[†] E-mail: g.gardner@murdoch.edu.au

(Verbeke *et al.*, 2010), and they would be likely to pay a premium if eating quality could be guaranteed (Lyford *et al.*, 2010).

A large proportion of the research investigating the variability in the tenderness of beef has used either objective measures, such as Warner Bratzler shear force, or trained taste panels. Total collagen content has been found to be moderately associated with decreasing trained panel scores for tenderness, and this relationship has been demonstrated between different muscles and within a single muscle, that is, *m. longissimus thoracis* (Torrescano *et al.*, 2003; Chriki *et al.*, 2012). This is also true for insoluble collagen (Riley *et al.*, 2005; Jurie *et al.*, 2007). Studies also indicate that age-related collagen cross-linking is an important determinant of insoluble collagen and therefore sensory tenderness (Bailey, 1985; Dransfield *et al.*, 2003; Chriki *et al.*, 2012). Chriki *et al.* (2012) reported that the ratio of soluble to total collagen content was positively correlated to tenderness when evaluated by trained French panellists. The more soluble collagen in comparison with insoluble collagen in beef the more tender the beef was scored by French trained taste panelists. This indicates that increasing the proportion of soluble collagen will have a positive association with tenderness. There is very limited information in the literature comparing biochemical analysis and untrained consumer taste panel scores, which are the basis of the Meat Standards Australia (MSA) approach. In several countries, including France, MSA has been shown to be relevant (Legrand *et al.*, 2013). The degree of correlation between untrained consumer responses and both total and insoluble collagen contents is yet to be determined in beef.

Many Studies have shown that increased marbling level, or intramuscular fat (IMF%) was associated with greater tenderness, juiciness, flavour liking and overall liking (Lorenzen *et al.*, 2003; Garmyn *et al.*, 2011; Chriki *et al.*, 2012; O'Quinn *et al.*, 2012). Furthermore Thompson (2004) in cattle and (Pannier *et al.*, 2014) in lamb found a positive relationship between IMF% and the flavour liking and juiciness scores of untrained Australian consumers. IMF% is also positively associated with the oxidative capacity (Jurie *et al.*, 2007; Kelman *et al.*, 2014) and therefore the haem iron content of muscles (Turkki and Campbell, 1967; Lengyel *et al.*, 2003; Kelman *et al.*, 2014) and has a strong negative association with the moisture content of beef and pork (Barlocco *et al.*, 2006; Pflanzner and de Felício, 2011).

We hypothesise that decreasing IMF% or decreasing soluble collagen content, and increasing insoluble collagen and/or total collagen content will relate to a decrease in untrained consumer taste panel perception of tenderness, similar to those relationships seen with trained taste panels both across different muscles and within the same muscle. IMF% will have a negative correlation with moisture content and a positive correlation with haem iron content and both will function as a proxy for IMF% with similar predictive capacities for untrained consumer taste panel scores both across different muscles and within the same muscle.

Material and methods

Animals and sample source

As previously described (Legrand *et al.*, 2013), 18 Australian cattle were slaughtered at an abattoir in Northern New South Wales, Australia and a further 18 French cattle were slaughtered in an abattoir in the western part of France following French industry practice. The cattle were chosen to reflect the different commercial production practices of the two countries. The Australian cattle were Murray Grey steers, ~18 months of age (Thompson *et al.*, 2010), while the French cattle consisted of three young French Holstein bulls (age range 19–20 months, carcass weight range 275–304 kg), three young crossbred Prim'Holstein × Hereford cows (age range 24–31 months, carcass weight range 288–307 kg), six cull French Holstein or Normande dairy cows (age range 51–98 months, carcass weight range 314–370 kg) and six cull Limousine or Charolaise beef cows (age range 68–201 months, carcass weight range 321–373 kg). The French cows were culled for age-related reasons. All cattle were hormonal growth promotant free.

Muscle samples

Six muscles from each animal were selected to represent a wide range of eating qualities: outside (*m. biceps femoris*), topside (*m. semimembranosus*), striploin (*m. longissimus thoracis*), rump (*m. gluteus medius*), oyster blade (*m. infraspinatus*) and tenderloin (*m. psoas major*). The Australian samples had two ageing times, with half of the samples aged for 5 days and the other half aged for 21 days. All French samples were aged for 10 days, except for the tenderloin which was aged for 7 days only.

Demographic profile of consumers

French consumers ($n = 540$) were involved in the sensory analysis, including 306 women (57%) and 234 men (43%). There was an even distribution between the six age classes, except for the oldest class (≥ 65 years) which consisted of only 3.3% of the consumers. Women were most represented in the younger classes (average 35 years) and there was a higher proportion of men in the oldest class (≥ 65 years). For a more detailed analysis of the consumer demographics, see Legrand *et al.* (2012).

Meat preparation and French consumer panels

Consumer assessment of eating quality was done according to the protocols for MSA testing described by Watson *et al.* (2008). Following the specific procedure also outlined in (Legrand *et al.*, 2013). Each sample (muscle) was sectioned into five steaks of 25 mm thickness. These steaks were halved after cooking making 10 portions available for tasting from each muscle (sample). Each consumer received seven portions: the first portion (a link sample) was a steak derived from either a generic striploin or rump muscle and designed to be of average quality – the sensory scores for this steak were not part of the final statistical analysis. The remaining six steaks were derived from the experimental samples collected.

Grilled steaks were cooked on a Silex clamshell grill (Silex, Hamburg, Germany) set to 220°C for 4.75 min for 'medium' and to 200°C for 3.25 min for 'rare' (Watson *et al.*, 2008). There was a high correlation between medium and rare cooking for the consumer palatability scores ($R^2 > 0.9$) (Legrand *et al.*, 2013).

In total, 360 French consumers took part in the 'medium' cooking test, with each consumer first consuming a medium quality 'link' sample then eating three Australian beef samples and three French beef samples. French consumers ($n = 180$) took part in the 'rare' cooking test, these portions were from French origin only and prepared as paired samples from the same muscles used for the 'medium' cooking test. Both medium and rare samples were 25 mm thick. Consumers scored portions for tenderness, juiciness, flavour liking and overall liking, by making a mark on a 100-mm line scale, with the left hand of the scale representing dislike and the right hand of the scale representing very liked. For a more detailed description of the testing procedures see Legrand *et al.* (2013).

Meat quality score (MQ4)

Values of consumer scores for tenderness, juiciness, flavour liking and overall liking were used to create a single MQ4 score that reflected the MSA star rating of the muscle. The weightings of the four sensory parameters (tenderness, juiciness, flavour liking and overall liking) to create the MQ4 score and hence the final rating ('unsatisfactory', 3*, 4* and 5*) were $0.31 \times$ tenderness, $0.04 \times$ juiciness, $0.30 \times$ flavour liking and $0.36 \times$ overall liking (Legrand *et al.*, 2013). The weightings were calculated using a discriminant analysis, as described by Watson *et al.* (2008). These weightings predicted the actual rating given by consumers for over 70% of the total number of samples, indicating that a high level of prediction is possible for French consumers.

Biochemical analysis

Proximate analyses were conducted to determine the chemical percentage of fat, moisture, protein, total and insoluble collagen of each muscle as previously described (Allais *et al.*, 2010). A single $2.5 \times 5 \times 5$ cm steak from each muscle was used for proximate analysis of Australian and French samples. Frozen French and Australian samples were thawed at 2°C to 4°C for 24 h before proximate analysis. All exterior fat and connective tissue were removed before proximate analysis, leaving only the lean meat.

Intramuscular lipid (IMF%) content was measured by the Soxhlet method using a Soxtherm apparatus (Gerhardt France SARL, Les Essarts Le Roi, France). Total and insoluble collagen content (%) were measured according to the INRA method (Listrat and Hocquette, 2004) on muscle samples that had undergone 2 h of heat treatment in a buffer solution at 90°C. Total and insoluble percentages were then estimated from the measurement of hydroxyproline content (collagen = $8 \times$ L-hydroxyproline). Soluble collagen content was calculated as the difference between total and insoluble collagen. Haem iron was measured using the technique described by Hornsey (1956). The sample was ground and the

pigments were extracted with HCl and acetone. The resulting liquid was stored in a light free environment for 24 h before the absorbance was measured at 510 nm with a spectrophotometer and the haem iron in mg/g of sample was calculated.

Statistical analysis

The sensory scores for tenderness, juiciness, flavour liking, overall liking and the composite score MQ4 were analysed using a linear mixed effects model (SAS v9.1). Initially, a core model was established with source country/ageing time (Aust5, Aus21, France10) and cooking 'doneness' (rare, medium) included as fixed effects. Cooking 'doneness' was fitted within source country/ageing time as only the French 10-day-aged samples were cooked rare. Animal ID was included as a random term to account for the samples being evaluated at two different degrees of cooking doneness. The same core model was then established with the same effects plus muscle type included as a fixed effect.

Biochemical measurements including total collagen, insoluble collagen, soluble collagen, iron, moisture and IMF contents were then incorporated one at a time as covariates into each of the core models (with and without muscle), as well as their interactions with all fixed effects to assess their association with the sensory scores. In all cases, non-significant terms ($P > 0.05$) were removed in a step-wise fashion. The oyster blade was not used in the analysis of the collagen parameters due to erroneous sample selection for collagen determinations.

Results

There was a large range in both IMF% and collagen content (Table 1) across the muscles and cattle sampled. Moisture content had a small variation.

Sensory scores

MQ4 is strongly correlated with all of the sensory parameters with average (across all muscles) correlations of ≥ 0.91 for tenderness, flavour liking and overall liking. Juiciness had the lowest average correlation with MQ4 (≥ 0.76 ; Table 2). Tenderness, flavour liking and overall liking had strong correlations with each other. Flavour liking and overall liking had the strongest correlation averaged across each source country, ageing period and cooking doneness (≥ 0.91). Juiciness and flavour had the weakest correlation averaged across each source country, ageing period and cooking doneness (≥ 0.65 ; Table 2). Sensory score correlations were relatively consistent for the striploin, rump and topside, with averages between 0.83 and 0.87, while the tenderloin, outside and the oyster blade had correlations slightly lower with averages ranging between 0.77 and 0.78 (data not shown).

Prediction of sensory scores using biochemical measurements

The tenderloin had the highest MQ4 of all the muscles for all of the different countries, degrees of doneness and ageing periods (Table 3). The outside had the lowest MQ4 for the

Table 1 Descriptive statistics of the biochemical parameters measured on six different muscles taken from 18 French cattle and 18 Australian steers

Biochemical parameter	<i>n</i>	Mean	s.d.	Minimum	Maximum	Coefficient of variation
IMF%	216	3.0	2.6	0.2	19.5	0.85
Moisture content (%)	216	73.2	1.8	62.1	76.6	0.02
Haem iron (µg/g)	216	17.3	4.1	8.1	30.3	0.24
Soluble collagen	180	0.2	0.2	0.0	2.0	0.92
Insoluble collagen	180	0.6	0.2	0.2	1.6	0.39
Total collagen	180	0.8	0.4	0.3	3.3	0.46

n = number of muscles; IMF% = intramuscular fat; collagen measurements are expressed in g/100 g fresh meat.

All measurements are on a fresh meat basis.

Muscles sampled: outside (*m. biceps femoris*), topside (*m. semimembranosus*), striploin (*m. longissimus thoracis*), rump (*m. gluteus medius*), oyster blade (*m. infraspinatus*) and tenderloin (*m. psoas major*).

The number of muscles samples is reduced for the collagen measurements due to the oyster blade (*m. infraspinatus*) being excluded from the analysis as a result of erroneous sample selection.

Table 2 Correlation coefficients between the sensory scores from untrained French consumers tasting six different muscles from French and Australian beef for the two countries and each ageing period and doneness by country

DF = 300	Tenderness	Juiciness	Flavour liking	Overall liking
Juiciness				
Average	0.66			
Australia 21 d medium	0.70			
Australia 5 d medium	0.65			
France 10 d medium	0.74			
France 10 d rare	0.52			
Flavour				
Average	0.76	0.65		
Australia 21 d medium	0.88	0.72		
Australia 5 d medium	0.82	0.75		
France 10 d medium	0.77	0.66		
France 10 d rare	0.71	0.58		
Overall liking				
Average	0.88	0.73	0.91	
Australia 21 d medium	0.90	0.77	0.93	
Australia 5 d medium	0.92	0.76	0.92	
France 10 d medium	0.91	0.77	0.89	
France 10 d rare	0.83	0.65	0.92	
MQ4				
Average	0.93	0.76	0.92	0.98
Australia 21 d medium	0.94	0.81	0.96	0.98
Australia 5 d medium	0.94	0.77	0.95	0.98
France 10 d medium	0.95	0.81	0.91	0.98
France 10 d rare	0.90	0.67	0.92	0.97

DF = degrees of freedom; MQ4 = a weighted combination (0.3, 0.1, 0.3, 0.3) of the other four sensory scores, tenderness, juiciness, flavour liking and overall liking.

The average is the correlation using all the different cooking doneness, ageing periods and countries.

21 d = muscle aged for 21 days post slaughter; 5 d = muscle aged for 5 days post slaughter; 10 d = muscle aged for 10 days post slaughter (except the tenderloin, which was aged for 7 days); medium = cooked to a medium doneness; rare = cooked to a rare doneness.

French cattle and the topside had the lowest MQ4 for the Australian cattle. The standard error between the different countries, degrees of doneness and ageing periods was consistently higher for tenderness, with the highest being 2.1 for the topside.

Analysis without 'muscle' as a fixed effect

The core model alone (source country/ageing time (Australian aged 5 days, Australian aged 21 days, French aged 10 days)

and degrees of doneness (rare, medium) as fixed effects, and animal identification number as a random term) explained from 0.6% to 8.6% of the variation (Table 4) for each of the sensory traits (Table 4). When insoluble collagen (g/100g fresh meat) was included in the core model, it accounted for the greatest amount of variation of all the biochemical parameters for all the sensory scores and described from 18.3% to 67.1% of the variation (Table 4). This was followed closely by total collagen content and IMF% (Table 4). The biochemical

Table 3 Quality scores predicted from the core model for each muscle from the 36 cattle for the two countries and each ageing period and doneness by country

	Outside				
	Tenderness ¹	Juiciness ¹	Flavour liking ¹	Overall liking ¹	MQ4 ¹
Australia 21 d medium	32.4 ^a	41.2 ^a	42.6 ^a	38.5 ^a	38.7 ^{ab}
Australia 5 d medium	35.8 ^a	44.2 ^a	49.4 ^a	42.8 ^a	43.0 ^b
France 10 d medium	21.7 ^b	42.8 ^a	40.3 ^a	30.0 ^b	31.8 ^a
France 10 d rare	22.1 ^b	43.4 ^a	41.9 ^a	30.2 ^b	32.7 ^a
Standard error	2.0	1.9	1.7	1.8	1.7
	Oyster blade				
	Tenderness ²	Juiciness ²	Flavour liking ²	Overall liking ²	MQ4 ²
Australia 21 d medium	82.0 ^a	72.5 ^a	75.5 ^a	77.9 ^a	77.4 ^a
Australia 5 d medium	80.3 ^{ac}	72.4 ^a	75.7 ^a	78.0 ^a	76.8 ^a
France 10 d medium	60.3 ^b	65.6 ^a	58.2 ^b	59.0 ^b	59.0 ^b
France 10 d rare	70.9 ^c	71.4 ^a	63.7 ^b	65.1 ^c	66.2 ^c
Standard error	2.0	1.9	1.7	1.8	1.7
	Rump				
	Tenderness ³	Juiciness ³	Flavour liking ³	Overall liking ³	MQ4 ³
Australia 21 d medium	57.3 ^a	48.9 ^a	59.1 ^a	55.9 ^a	56.8 ^a
Australia 5 d medium	59.3 ^a	52.1 ^a	61.0 ^a	57.6 ^a	58.2 ^a
France 10 d medium	54.7 ^a	54.0 ^a	58.5 ^a	57.6 ^a	56.7 ^a
France 10 d rare	57.2 ^a	64.5 ^b	60.4 ^a	58.5 ^a	58.7 ^a
Standard error	2.0	1.9	1.7	1.8	1.7
	Striploin				
	Tenderness ⁴	Juiciness ³	Flavour liking ³	Overall liking ³	MQ4 ⁴
Australia 21-day medium	72.1 ^a	55.5 ^{ab}	64.3 ^{ab}	65.5 ^a	66.4 ^{ab}
Australia 5 d medium	64.9 ^a	52.1 ^a	62.2 ^{ab}	60.9 ^{ab}	61.5 ^a
France 10 d medium	51.4 ^b	54.9 ^a	57.4 ^a	55.3 ^b	54.4 ^b
France 10 d rare	64.1 ^{ab}	64.0 ^b	66.1 ^b	64.3 ^a	64.1 ^a
Standard error	2.1	1.9	1.7	1.8	1.7
	Tenderloin				
	Tenderness ⁵	Juiciness ²	Flavour liking ⁴	Overall liking ⁴	MQ4 ⁵
Australia 21 d medium	86.8 ^a	68.9 ^{ab}	79.3 ^a	79.2 ^a	79.7 ^a
Australia 5 d medium	87.9 ^a	61.6 ^a	80.5 ^a	82.5 ^a	80.9 ^a
France 10 d medium	85.8 ^a	71.0 ^b	79.6 ^a	80.8 ^a	80.7 ^a
France 10 d rare	89.7 ^a	72.6 ^b	80.5 ^a	81.2 ^a	81.9 ^a
Standard error	2.0	2.0	1.7	1.8	1.7
	Topside				
	Tenderness ¹	Juiciness ⁴	Flavour liking ¹	Overall liking ¹	MQ4 ¹
Australia 21 d medium	30.6 ^a	31.8 ^a	40.9 ^a	34.1 ^a	34.3 ^a
Australia 5 d medium	30.9 ^a	34.8 ^{ab}	42.9 ^a	36.5 ^a	36.5 ^a
France 10 d medium	35.4 ^a	41.5 ^b	47.5 ^a	41.0 ^a	41.1 ^b
France 10 d rare	42.5 ^b	52.2 ^c	54.5 ^b	49.1 ^b	48.7 ^c
Standard error	2.1	1.9	1.7	1.878	1.7

Rare = cooked to a rare doneness; Medium = cooked to a medium doneness; MQ4 = a weighted combination (0.3, 0.1, 0.3, 0.3) of the other four sensory scores, tenderness, juiciness, flavour liking and overall liking.

Superscript numbers = different numbers in a column indicate the scores are significantly different per muscle ($P < 0.05$); superscript letters = for each muscle different letters in a column indicate the scores are significantly different ($P < 0.05$); 21 d = muscle aged for 21 days post slaughter; 5 d = muscle aged for five days post slaughter; 10 d = muscle aged for 10 days post slaughter (except the tenderloin which was aged for 7 days).

The core model comprised the fixed effects source country/ageing time (Australian samples aged 5 days, Australian samples aged 21 days and French samples aged 10 days) and cooking 'doneness' (rare, medium). Cooking 'doneness' was fitted within source country/ageing time as only the French 10-day-aged samples were cooked rare. Muscle type was included as a fixed effect. Animal ID was included as a random term.

Table 4 The percentage of the variation in the sensory scores of French untrained consumers evaluating six different muscles taken from 18 French cattle and 18 Australian steers prepared at medium or rare cooking doneness and aged for 5, 10 or 21 days, which is explained by the core models and biochemical parameters

	Percentage of variance in sensory scores explained by the core model				
	MQ4	Overall liking	Tenderness	Flavour liking	Juiciness
Core model	1.2	0.9	0.6	2.0	8.6
	Additional percentage of variance explained by biochemical parameters				
IMF%	36.9	35.8	36.5	29.4	35.9
Haem iron ($\mu\text{g/g}$ wet matter)	1.8	3.5	4.4	4.1	11.6
Moisture content	7.9	8.3	5.4	6.9	7.4
Total collagen	41.0	41.8	36.7	30.9	31.0
Insoluble collagen	48.6	48.4	67.1	24.0	18.3

IMF% = intramuscular fat; collagen measurements are expressed in g/100g fresh meat; MQ4 = a weighted combination (0.3, 0.1, 0.3, 0.3) of the other four sensory scores, tenderness, juiciness, flavour liking and overall liking.

All measurements are on a fresh meat basis.

Muscles sampled: outside (*m. biceps femoris*), topside (*m. semimembranosus*), striploin (*m. longissimus thoracis*), rump (*m. gluteus medius*), oyster blade (*m. infraspinatus*) and tenderloin (*m. psoas major*).

The core model comprised the fixed effects source country/ageing time (Australian samples aged 5 days, Australian samples aged 21 days and French samples aged 10 days) and cooking 'doneness' (rare, medium). Cooking 'doneness' was fitted within source country/ageing time as only the French 10-day-aged samples were cooked rare. Animal ID was included as a random term.

parameter that described the least amount of the variance for all of the sensory scores was haem iron content (Table 4).

Total collagen had the greatest *F*-values for tenderness, flavour liking, overall liking and MQ4. Insoluble collagen had the greatest *F*-values for juiciness (Table 5). There were no effects on the biochemical measurements between source country and ageing period except for haem iron, where there was a greater, positive, impact of haem iron on the eating quality of the Australian beef than the French beef (data not shown). Haem iron also had the smallest *F*-values for all the sensory scores (Table 5).

There were significant relationships between all the biochemical parameters and sensory scores except for soluble collagen (Table 5; Figure 1). Of the biochemical parameters tested, total and insoluble collagen contents demonstrated the strongest association with consumer eating quality with more than 30% of the variance in sensory scores explained by these two factors (Table 1) and with all sensory scores reduced across the increasing total and insoluble collagen range (Figure 1). The magnitude of these associations varied for the different sensory scores, being strongest for tenderness, which reduced by 35.5 units across the insoluble collagen range, and weakest for juiciness, which reduced by only 33.0 units across the same range (Figure 1). For each of the different sensory scores, the majority of this response occurred between insoluble collagen of 0.2–0.7 g/100 g fresh meat. Total collagen had a similar relationship with the sensory scores as insoluble collagen although the magnitude was greater for insoluble collagen. Hence, the greatest relationship for total collagen was with tenderness, reducing the scores by 63 units, and the lowest was with juiciness, reducing the scores by 27.7 units across the range of total collagen, which varied between 0.4 and 1.6 g/100 g fresh meat.

For IMF%, the strongest association was with tenderness which increased by 57.0 units across an IMF% range of

almost 9 units. The lowest association was with flavour liking, which only increased by 34.5 units across the IMF% range. Of all the biochemical parameters measured, juiciness was most strongly associated with IMF%, increasing by 39.0 units across the IMF% range (Figure 1). In all cases, the impact of IMF% on sensory scores was curvilinear, but no plateau was reached within the range of IMF%, 0.2% to 9%, in this study.

Haem iron had a positive relationship with all sensory traits. The impact was greatest on tenderness, which increased by 34 eating quality points across the haem iron range, respectively, with less impact on flavour liking, MQ4 and juiciness, which increased by only 23.2, 10 and 12 units across the range. This effect of haem iron was reduced in the French 10-day-aged samples, with tenderness, flavour liking and overall liking scores increasing by 14.5, 3.6 and 13.9 eating quality points across the haem iron range.

H₂O% had a negative relationship with all the sensory scores. The magnitude of change in the different sensory scores was quite consistent, ranging from –19.9 for flavour liking to –23.5 for overall liking.

Analysis with 'muscle' as a fixed effect

The percentage of variation in sensory scores explained by the core model increased markedly when muscle was included, increasing to 79.5% for MQ4, 77.8% for overall liking, 82.0% for tenderness, 70.1% for flavour liking and 66.0% for juiciness (data not shown). Of the biochemical parameters tested as covariates, the only ones retaining significance in the model corrected for muscle were IMF% and H₂O% (Table 6, Figure 2). When IMF% was added to the model, it explained an additional 0.06% of the variance in overall liking scores and an additional 6.3% of the variance in flavour liking scores over the variance explained by the core model. This is much reduced from the 35.8% and

Table 5 F-values for the core model and the impact of six biochemical measurements on untrained French consumer perceptions of tenderness, juiciness, flavour liking and overall liking, and on the combined consumer MQ4 score of six muscles taken from 18 cattle from France and 18 steers from Australia prepared at medium or rare cooking doneness and aged for 5, 10 or 21 days

Effect	Core model			IMF%	H2O%	Haem iron	Total collagen	Insoluble collagen
	NDF	DDF	F-values	F-values	F-values	F-values	F-values	F-values
Tenderness								
Doneness	1	280	3.2	5.1*	3.4	3.3	3.0	3.5
Source country and days aged	2	280	1.9	0.1	0.6	4.8**	1.9	1.2
Covariate tested (linear)	1	280	–	117.7**	17.7**	3.2	133.4**	128.9**
Covariate tested ¹ (curvilinear)	1	280	–	35.6**	–	4.9*	79.1**	64.0**
Covariate tested × source country and days aged	1	280	–	–	–	5.0**	–	–
Covariate tested ¹ × source country and days aged	1	280	–	–	–	5.5**	–	–
Flavour liking								
Doneness	1	280	3.6	5.2*	3.9	3.7	3.7	4.1*
Source country and days aged	2	280	0.6	0.2	0.0	3.3*	0.1	0.5
Covariate tested (linear)	1	280	–	90.0**	32.1**	2.7	103.2**	91.6**
Covariate tested ¹ (curvilinear)	1	280	–	24.6**	–	4.5*	61.3**	46.4**
Covariate tested × source country and days aged	1	280	–	–	–	3.8*	–	–
Covariate tested ¹ × source country and days aged	1	280	–	–	–	4.4*	–	–
Juiciness								
Doneness	1	280	8.8**	14.9**	9.6**	9.1**	11.1**	12.5**
Source country and days aged	2	280	2.3	9.0**	6.1**	1.1	2.3	3.8*
Covariate tested (linear)	1	280	–	144.8**	32.6**	11.9**	60.1**	68.2**
Covariate tested ¹ (curvilinear)	1	280	–	43.7*	–	–	36.5**	37.5**
Overall liking								
Doneness	1	280	2.2	3.5	2.4	3.9*	2.3	2.6
Source country and days aged	2	280	1.0	0.1	0.0	2.3	0.5	0.3
Covariate tested (linear)	1	280	–	109.8**	30.5**	2.2	130.6**	107.4**
Covariate tested ¹ (curvilinear)	1	280	–	30.1**	–	3.8	76.7**	50.7**
Covariate tested × source country and days aged	1	280	–	–	–	4.1*	–	–
Covariate tested ¹ × source country and days aged	1	280	–	–	–	4.5*	–	–
MQ4								
Doneness	1	280	3.2	5.1	3.5	3.3	3.3	3.8
Source country and days aged	2	280	0.9	0.1	0.1	1.9	0.6	0.5
Covariate tested (linear)	1	280	–	116.6**	28.3**	7.4**	128.5**	115.8**
Covariate tested ¹ (curvilinear)	1	280	–	33.2	–	–	77.1**	57.5**

NDF = numerator degrees of freedom; DDF = denominator degrees of freedom; IMF% = intramuscular fat; collagen measurements are expressed in g/100 g fresh meat; haem iron measurements are expressed as µg/g fresh meat; all measurements are on a fresh meat basis; * $P < 0.1$; ** $P < 0.05$; *** $P < 0.01$.

The core model comprised fixed effects source country/ageing time (Australian samples aged 5 days, Australian samples aged 21 days and French samples aged 10 days) and cooking 'doneness' (rare, medium). Cooking 'doneness' was fitted within source country/ageing time as only the French 10-day-aged samples were cooked rare. Animal ID was included as a random term. The biochemical measurements were then introduced to the core model individually, and the model was regressed in a step wise fashion until only significant terms remained.

¹The term is run in the model interacted with itself, testing for a curvilinear response; the oyster blade results were excluded for the collagen models, hence the DDF values decreased to 224.

29.4% variation explained by IMF% in overall liking and flavour liking when muscle was not included in the model (Table 4). Across the IMF% range, flavour liking and overall liking increased by 6.1 and 5.9 units. When moisture content was added to the model, it explained an additional 0.08% of the variance in overall liking scores and an additional 6.3% of the variance in flavour liking scores over the variance explained by the core model. This is reduced from the 8.3% and 6.9% variation explained by moisture content in overall liking and

flavour liking, respectively, when muscle was not included in the model (Table 4). Across the H₂O% range, the flavour liking and overall liking scores decreased by 6.4 and 6.2 units.

Discussion

Relationship between the different sensory scores

The high correlations between all the four sensory scores and MQ4 reflect the fact that the MQ4 value has been derived




Figure 1 The effect of biochemical parameters on untrained French consumer perceptions of tenderness, juiciness, flavour liking and overall liking, and on the combined consumer MQ4 score of six muscles taken from 18 cattle from France and 18 steers from Australia prepared at medium or rare cooking doneness and aged for 5, 10 or 21 days without muscle included in the model. Mag = magnitude of effect. All measurements are on a fresh meat basis; MQ4 = a weighted combination (0.3, 0.1, 0.3, 0.3) of the other four sensory scores, tenderness, juiciness, flavour liking and overall liking.

from these scores and demonstrates the ability of MQ4 to reflect the whole of the consumer eating experience. The development of the MQ4 value from untrained consumer scores was first described in detail by Watson *et al.* (2008). High correlations (range 0.5 to 0.6) between the sensory scores were reported using Australian consumers tasting grilled steaks to a medium doneness, with the only exception being flavour liking (range 0.1 to 0.3). The comparative accuracy of the MQ4 value for both French and Australian consumers was investigated by Legrand *et al.* (2013) with the conclusion that it is at least as accurate at describing the consumer eating experience for French consumers as it is for Australian consumers. The correlations for French consumers between the four sensory scores and with MQ4 in this study are similar or stronger on average than those reported by Watson *et al.* (2008) with Australian consumers.

Relationship between collagen measurements and untrained consumer scores

Across muscles. The hypothesis that increasing total and insoluble collagen or decreasing soluble collagen would reduce consumer sensory scores across muscles was partially

supported by our results, with both total and insoluble collagen having negative relationships with eating quality. Yet contrary to this hypothesis, soluble collagen showed no relationship with eating quality. Total collagen comprised both soluble and insoluble collagen and therefore reflects the impact of insoluble collagen on untrained consumer scores, with the magnitude diminished by incorporating the soluble collagen component. This aligns well with the findings of Chriki *et al.* (2012) who showed a relationship between both total and insoluble collagen and eating quality ($r = -0.15$, $r = -0.20$).

This result is further supported by the work of Schonfeldt and Strydom (2011) who found that collagen content was the main driver of the differences in tenderness between different muscles ($r = 0.986$) and by Light *et al.* (1985) who reported that increasing total collagen and collagen cross-linking (insoluble collagen) was associated with increasing toughness across six different muscles. Large variations in collagen content exist across different muscles due to their different roles in the live animal. The negative relationship and relatively high magnitude of effect between insoluble collagen and eating quality demonstrates that a large

Table 6 F-values for the core model including cut and the impact of IMF% and H2O% on untrained French consumer perceptions of tenderness, juiciness, flavour liking and overall liking, and on the combined consumer MQ4 score of six muscles taken from 18 cattle from France and 18 steers from Australia prepared at medium or rare cooking doneness and aged for 5, 10 or 21 days

Effect	Core model			IMF%			H ₂ O%
	NDF	DDF	F-values	NDF	DDF	F-values	F-values
Tenderness							
Doneness	1	265	17.6**	–	–	–	–
Muscle	5	255	173.7**	–	–	–	–
Source country and days aged	2	255	264.0	–	–	–	–
Muscle × source country and days aged	10	255	4.8**	–	–	–	–
Juiciness							
Doneness	1	265	23.1**	–	–	–	–
Muscle	5	255	68.2**	–	–	–	–
Source country and days aged	2	255	2.6	–	–	–	–
Muscle × source country and days aged	10	255	2.9**	–	–	–	–
Flavour liking							
Doneness	1	265	11.7**	1	264	11.7**	11.7**
Muscle	5	255	90.7**	5	264	61.7**	85.9**
Source country and days aged	2	255	1.00	2	264	0.7	0.6
Muscle × source country and days aged	10	255	4.8**	10	264	4.0**	3.5**
Covariate tested		–	–	1	264	4.7*	5.7*
Overall liking							
Doneness	1	265	9.9**	1	264	9.9**	9.9**
Muscle	5	255	133.8**	5	264	9.2**	126.5**
Source country and days aged	2	255	1.3	2	264	1.0	0.8
Muscle × source country and days aged	10	255	6.2**	10	264	5.2**	4.7**
Covariate tested		–	–	1	264	4.1*	4.9*
MQ4							
Doneness	1	265	15.5**	–	–	–	–
Muscle	5	255	148.6**	–	–	–	–
Source country and days aged	2	255	1.1	–	–	–	–
Muscle × source country and days aged	10	255	6.1**	–	–	–	–

NDF = numerator degrees of freedom; DDF = denominator degrees of freedom; IMF% = intramuscular fat; all measurements are on a fresh meat basis; * $P < 0.1$; ** $P < 0.05$; *** $P < 0.01$; MQ4 = a weighted combination (0.3, 0.1, 0.3, 0.3) of the other four sensory scores, tenderness, juiciness, flavour liking and overall liking. The core model comprised the fixed effects of cut and source country/ageing time (Australian samples aged 5 days, Australian samples aged 21 days and French samples aged 10 days) and cooking 'doneness' (rare, medium). Cooking 'doneness' was fitted within source country/ageing time as only the French 10-day-aged samples were cooked rare. Animal ID was included as a random term. The biochemical measurements were then introduced to the core model individually, and the model was regressed in a step-wise fashion until only significant terms remained.

proportion of the well-known differences in eating quality between different muscles is due to differences in insoluble collagen.

The lack of association between soluble collagen and eating quality conflicts with the results of Purslow (2005) and Renand *et al.* (2001) who both demonstrated that increasing collagen solubility had a positive relationship with trained taste panel scores both across muscles and within muscle. However, in contrast to our study where soluble collagen was expressed as a percentage of total muscle mass, Purslow (2005) and Renand *et al.* (2001) expressed soluble collagen as a proportion of total collagen. When soluble collagen is expressed as a proportion of total collagen, it incorporates the influence of insoluble collagen through the denominator

of this ratio, limiting the capacity to understand soluble collagen independently. Consequently, the lack of any relationship between soluble collagen and untrained consumer eating quality in this study implies that previously published relationships between collagen solubility and tenderness may be driven by the insoluble collagen component of the proportion. This is supported by Chriki *et al.* (2012) who failed to find a relationship between soluble collagen and eating quality. Knowledge of soluble collagen levels would therefore fail to add any further information in the prediction of eating quality.

Within muscle. Contrary to our hypothesis, total collagen content, insoluble collagen content and soluble collagen content did not demonstrate any relationship with untrained




Figure 2 The effect of IMF% and H2O% on untrained French consumer perceptions of tenderness, juiciness, flavour liking and overall liking, and on the combined consumer MQ4 score of six muscles taken from 18 cattle from France and 18 steers from Australia prepared at medium or rare cooking doneness and aged for 5, 10 or 21 days with muscle included in the model. Mag = magnitude of effect. All measurements are on a fresh meat basis; MQ4 = a weighted combination (0.3, 0.1, 0.3, 0.3) of the other four sensory scores, tenderness, juiciness, flavour liking and overall liking. The graphed range of intramuscular fat has been truncated at 9%, excluding three samples with higher levels.

consumer scores within any one muscle. This expectation was based on evidence that as the amount of insoluble collagen increases, more collagen remains in the final cooked product, increasing the resistance to mastication and therefore toughness (Purslow, 2005). Although both Chriki *et al.* (2012) and Schonfeldt and Strydom (2011) were able to demonstrate small effects of collagen solubility on sensory tenderness, this was done using trained consumer scores, contrasting with the present study where untrained consumer scores were used. Hence, the greater variability inherent within untrained consumer data (Watson *et al.*, 2008) was likely the reason that we were not able to detect these small differences. Furthermore, the relatively small data set and experimental factors such as the different animal breeds, different abattoirs and different production methods would also preclude the detection of subtle relationships.

Alternatively, these results align with the work of Jeremiah and Martin (1981) who found no relationships between total collagen content, insoluble collagen content and soluble collagen content and tenderness within muscle. However, unlike in our study, the carcasses were slaughtered at a similar chronological age, which would reduce the within-muscle variation in collagen solubility resulting from age-related collagen cross-linking. The effect of animal age on collagen cross-linking and subsequent increases in insoluble collagen content is well documented in the literature (Bailey *et al.*, 1998). However, the expected positive relationship between animal age and insoluble collagen was not observed in this particular data set. Given the large range in animal ages in our study, this was expected to deliver a large range in insoluble collagen content improving the possibility of identifying an association between collagen

and tenderness. We hypothesise that the lack of the expected positive relationship between animal age and insoluble collagen may be a result of the multiple different animal production methods, confounded with animal age, gender and breed, for example, older French dairy cows, older French beef cows, young Australian steers and young French bulls, combined with low animal numbers, inherent biological variability and a relatively high technical error of the biochemical test itself. The lack of this expected positive relationship may possibly explain the absence of any relationship between collagen measurements and eating quality within muscles.

Relationship between IMF% and consumer scores

Aligning with our hypothesis, IMF% had a positive relationship with eating quality both across muscles and within muscle. Lipids carry the majority of the species-specific flavour profiles of meat, which has a large, positive impact on eating quality (Hornstein and Crowe, 1960). IMF% also improves juiciness scores by the stimulation of salivation in the consumer and prohibits the absorption of moisture by meat due to its hydrophobic nature (Thompson, 2004). IMF% also affects tenderness, as shown in sheep meat by Pannier *et al.* (2014) where a 1-unit increase in IMF% increased tenderness scores by 2.2 eating quality points. This may be due to the weaker structural properties of fat than other muscle tissue components (Thompson, 2004). These relationships are supported by O'Quinn *et al.* (2012) who found that as the IMF% in the *m. longissimus thoracis* increased American untrained consumer scores for tenderness, flavour liking, juiciness and overall liking increased, though this effect was not consistent between the samples of beef derived from Australian and American cattle. This mirrors the findings of Thompson (2004), who also utilised untrained consumer panels and the *m. longissimus thoracis*. The results of our study show that French untrained consumer scores have a similar positive response to increasing IMF% compared with Australian and American consumers both across muscles and within muscle. This demonstrates that IMF% or a proxy measurement will add value to a predictive grading system based on meat quality irrespective of whether muscle is in the prediction model. This is consistent with its use within numerous quality grading systems around the world including the Australian MSA system (for a review, see Polkinghorne and Thompson, 2010).

The ability of moisture content to act as a proxy for IMF%

Our hypothesis that H₂O% would act as a proxy for IMF% and have similar eating quality predictive capacities as IMF% was partially supported. As expected, while exhibiting effects of similar magnitude, IMF% and H₂O% had contrasting relationships with eating quality because of the inverse relationship between moisture content and IMF% (Barlocco *et al.*, 2006; Pflanzler and de Felício, 2011). However, when the two covariates were run in the model simultaneously, their *F*-values reduced, though they both retained significance. This demonstrates that the correlation between IMF% and

H₂O% does not completely account for the relationship between H₂O% and eating quality. H₂O% is not simply acting as a proxy for IMF% and also explains some of the variation in eating quality completely independent of its relationship with IMF%. This is reinforced by Renand *et al.* (2001) who found that within the *m. longissimus thoracis* of Charolais young bulls the dry matter, the inverse of moisture content, was the principle muscle component related to flavour liking ($r = 0.40$) when compared with other biochemical measures including IMF%. This may be due to the flavour contribution of the water-soluble components of meat and smaller errors in the measurements of H₂O% when compared with IMF% (Renand *et al.*, 2001). These results indicate that H₂O% or its inverse, dry matter, may have a similar value in an eating quality prediction system in addition to IMF%.

The ability of haem iron to act as a proxy for IMF%

Our results only partially supported our hypothesis that haem iron would function as a proxy for IMF% when predicting eating quality. When muscle was excluded from the model and the IMF% and haem iron were both describing variation across and within muscle, both covariates had a positive relationship with eating quality. In addition, when the two biochemical measurements were included in the prediction model concurrently, only IMF% retained significance. This indicates that the correlation between the two biochemical measurements is the sole basis for the relationship that exists between haem iron and eating quality and haem iron is therefore functioning as a proxy and not describing any additional variation in eating quality. However, IMF% had an average magnitude of effect 21.3 points higher than haem iron across all the sensory scores. This is almost double the magnitude of effect of haem iron and highlights the reduced ability for haem iron to predict eating quality when compared with IMF%. This contradicts our hypothesis that haem iron had no relationship with eating quality scores within individual muscles. This result contrasts with the work of Renand *et al.* (2001) who found that, within the *m. longissimus thoracis*, haem iron content had a positive correlation with trained taste panel scores for flavour and juiciness ($r = 0.21$, $r = 0.19$, respectively). This may be explained by the different animal types used in the two studies. As with insoluble collagen, the greater variability inherent within untrained consumer data (Watson *et al.*, 2008) used in this study was likely one major reason among others that we were not able to detect these small differences. Haem iron does function as a proxy for IMF% when predicting eating quality across muscles, however, the magnitude of the effect is heavily reduced and this relationship is not present within muscle. This shows that haem iron, despite being correlated with IMF% (Turkki and Campbell, 1967) is not suitable as a proxy when predicting eating quality.

Conclusion

Biochemical measurements, particularly IMF% and insoluble collagen have the capacity to explain a large amount of the

variation in the eating quality of beef. However, this capacity of insoluble collagen is eliminated when muscle type is known. Muscle type is enough to explain the effect of connective tissue on the eating quality of beef. Our results show that a combination of muscle type and IMF% allows for a very good prediction of the eating quality of beef and that an industrial model would not be greatly improved by the estimation of collagen content. This is reflected in the Australian MSA model where muscle type has the strongest influence on the final quality grade of a piece of beef (Polkinghorne *et al.*, 2008). If a model like MSA was to be implemented in the European market other factors such as animal age, cooking method and the effects of the country of origin for the beef and the consumer would have to be investigated.

Acknowledgements

This research was funded by Meat and Livestock Australia and Murdoch University. The authors would also like to acknowledge the financial contributions of the European research project ProSafeBeef (Contract No. FOOD-CT-2006-36241), the French 'Direction Générale de l'Alimentation' and FranceAgriMer. Furthermore, this project would not have been possible without the practical support of the Association Institut du Charolais, the Syndicat de Défense et du promotion de la Viande de Boeuf de Charolles and the gourmet restaurants 'Jean Denaud'. The authors are also very grateful to the staff of the Meat Laboratory from the French Livestock Institute in Villers Bocage, which provided all muscle biochemical data. The international travel required for this project has been funded by 'Egide/Fast' funds from the French and Australian governments, respectively. The assistance and participation of the Beef CRC, Janine Lau, Alex Ball (MLA, Australia), Alan Gee (Cosign, Australia), Ray Watson (Melbourne University, Australia), Paul Allen (Teagasc, Irish Republic) and John Thompson (UNE) are also gratefully acknowledged.

References

- Allais S, Levézil H, Payet-Duprat N, Hocquette JF, Lepetit J, Rousset S, Denoyelle C, Bernard-Capel C, Journaux L, Bonnot A and Renand G 2010. The two mutations, Q204X and nt821, of the myostatin gene affect carcass and meat quality in young heterozygous bulls of French beef breeds. *Journal of Animal Science* 88, 446–454.
- Bailey AJ 1985. The role of collagen in the development of muscle and its relationship to eating quality. *Journal of Animal Science* 60, 1580–1587.
- Bailey AJ, Paul RG and Knott L 1998. Mechanisms of maturation and ageing of collagen. *Mechanisms of Ageing and Development* 106, 1–56.
- Barlocco N, Vadell A, Ballesteros F, Galietta G and Cozzolino D 2006. Predicting intramuscular fat, moisture and Warner-Bratzler shear force in pork muscle using near infrared reflectance spectroscopy. *Animal Science* 82, 111–116.
- Chriki S, Gardner GE, Jurie C, Picard B, Micol D, Brun JP, Journaux L and Hocquette JF 2012. Cluster analysis application identifies muscle characteristics of importance for beef tenderness. *BMC Biochemistry* 13, 29.
- Dransfield E, Martin JF, Bauchart D, Abouelkaram S, Lepetit J, Culioli J, Jurie C and Picard B 2003. Meat quality and composition of three muscles from French cull cows and young bulls. *Animal Science* 76, 387–399.
- Garmyn AJ, Hilton GG, Mateescu RG, Morgan JB, Reecy JM, Tait RG, Beitz DC, Duan Q, Schoonmaker JP, Mayes MS, Drewnoski ME, Liu Q and VanOverbeke DL 2011. Estimation of relationships between mineral concentration and fatty acid composition of longissimus muscle and beef palatability traits. *Journal of Animal Science* 89, 2849–2858.

- Hornsey HC 1956. The colour of cooked cured pork. I. Estimation of the nitric oxide-haem pigments. *Journal of the Science of Food and Agriculture* 7, 534–540.
- Hornstein I and Crowe PF 1960. Meat flavor chemistry, flavor studies on beef and pork. *Journal of Agricultural and Food Chemistry* 8, 494–498.
- Jeremiah LE and Martin AH 1981. Intramuscular collagen content and solubility: their relationship to tenderness and alteration by post mortem aging. *Canadian Journal of Animal Science* 61, 53–61.
- Jurie C, Picard B, Hocquette J-F, Dransfield E, Micol D and Listrat A 2007. Muscle and meat quality characteristics of Holstein and Salers cull cows. *Meat Science* 77, 459–466.
- Kelman KR, Pannier L, Pethick DW and Gardner GE 2014. Selection for lean meat yield in lambs reduces indicators of oxidative metabolism in the longissimus muscle. *Meat Science* 96, 1058–1067.
- Legrand I, Hocquette J-F, Polkinghorne RJ and Pethick DW 2012. Prediction of beef eating quality in France using the Meat Standards Australia system. *Animal* 7, 524–529.
- Legrand I, Hocquette JF, Polkinghorne RJ and Pethick DW 2013. Prediction of beef eating quality in France using the Meat Standards Australia system. *Animal* 7, 524–529.
- Lengyel Z, Husvéth F, Polgár P, Szabó F and Magyar L 2003. Fatty acid composition of intramuscular lipids in various muscles of Holstein-Friesian bulls slaughtered at different ages. *Meat Science* 65, 593–598.
- Light N, Champion AE, Voyle C and Bailey AJ 1985. The role of epimysial, perimysial and endomysial collagen in determining texture in six bovine muscles. *Meat Science* 13, 137–149.
- Listrat A and Hocquette J-F 2004. Analytical limits of total and insoluble collagen content measurements and of type I and III collagen analysis by electrophoresis in bovine muscles. *Meat Science* 68, 127–136.
- Lorenzen CL, Miller RK, Taylors JF, Neely TR, Tatum JD, Wise JW, Buyek MJ, Reagan JO and Savell JW 2003. Beef customer satisfaction: trained sensory panel ratings and Warner-Bratzler shear force values. *Journal of Animal Science* 81, 143–149.
- Lyford C, Thompson J, Polkinghorne R, Miller M, Nishimura T, Neath K, Allen P and Belasco E 2010. Is willingness to pay (WTP) for beef quality grades affected by consumer demographics and meat consumption preferences? *Australasian Agribusiness Review* 18, 1–15.
- Morgan JB, Savell JW, Hale DS, Miller RK, Griffin DB, Cross HR and Shackelford SD 1991. National beef tenderness survey. *Journal of Animal Science* 69, 3274–3283.
- O'Quinn TG, Brooks JC, Polkinghorne RJ, Garmyn AJ, Johnson BJ, Starkey JD, Rathmann RJ and Miller MF 2012. Consumer assessment of beef strip loin steaks of varying fat levels. *Journal of Animal Science* 90, 626–634.
- Pannier L, Gardner GE, Pearce KL, McDonagh M, Ball AJ, Jacob RH and Pethick DW 2014. Associations of sire estimated breeding values and objective meat quality measurements with sensory scores in Australian lamb. *Meat Science* 96, 1076–1087.
- Pflanzer SB and de Felício PE 2011. Moisture and fat content, marbling level and color of boneless rib cut from Nellore steers varying in maturity and fatness. *Meat Science* 87, 7–11.
- Polkinghorne RJ and Thompson JM 2010. Meat standards and grading: a world view. *Meat Science* 86, 227–235.
- Polkinghorne R, Thompson JM, Watson R, Gee A and Porter M 2008. Evolution of the Meat Standards Australia (MSA) beef grading system. *Australian Journal of Experimental Agriculture* 48, 1351–1359.
- Purslow PP 2005. Intramuscular connective tissue and its role in meat quality. *Meat Science* 70, 435–447.
- Renand G, Picard B, Touraille C, Berge P and Lepetit J 2001. Relationships between muscle characteristics and meat quality traits of young Charolais bulls. *Meat Science* 59, 49–60.
- Riley DG, Johnson DD, Chase CC Jr, West RL, Coleman SW, Olson TA and Hammond AC 2005. Factors influencing tenderness in steaks from Brahman cattle. *Meat Science* 70, 347–356.
- Schonfeldt HC and Strydom PE 2011. Effect of age and cut on tenderness of South African beef. *Meat Science* 87, 206–218.
- Thompson J 2004. The effects of marbling on flavour and juiciness scores of cooked beef, after adjusting to a constant tenderness. *Australian Journal of Experimental Agriculture* 44, 645–652.
- Thompson J, Polkinghorne R, Gee A, Motiang D, Strydom P, Mashau M, Ng'ambi J, deKock R and Burrow H 2010. Beef palatability in the Republic of South Africa: implications for niche-marketing strategies. In *ACIAR Technical Reports*, pp. 1–56. Australian Centre for International Agricultural Research (ACIAR), Canberra, ACT, Australia.
- Torrescano G, Sánchez-Escalante A, Giménez B, Roncalés P and Beltrán JA 2003. Shear values of raw samples of 14 bovine muscles and their relation to muscle collagen characteristics. *Meat Science* 64, 85–91.
- Turkki PR and Campbell AM 1967. Relation of phospholipids to other tissue components in two beef muscles. *Journal of Food Science* 32, 151–154.
- Verbeke W, Van Wezemael L, de Barcellos MD, Kügler JO, Hocquette J-F, Ueland Ø and Grunert KG 2010. European beef consumers' interest in a beef eating-quality guarantee: insights from a qualitative study in four EU countries. *Appetite* 54, 289–296.
- Watson R, Gee A, Polkinghorne R and Porter M 2008. Consumer assessment of eating quality – development of protocols for Meat Standards Australia (MSA) testing. *Australian Journal of Experimental Agriculture* 48, 1360–1367.