



**HAL**  
open science

## Vertebrate bacterial gut diversity: size also matters

Jean-Jacques Godon, Pugazhendi Arulazhagan, Jean-Philippe Steyer, Jérôme Hamelin

► **To cite this version:**

Jean-Jacques Godon, Pugazhendi Arulazhagan, Jean-Philippe Steyer, Jérôme Hamelin. Vertebrate bacterial gut diversity: size also matters. *BMC Ecology*, 2016, 16 (1), pp.12. 10.1186/s12898-016-0071-2 . hal-02639327

**HAL Id: hal-02639327**

**<https://hal.inrae.fr/hal-02639327v1>**

Submitted on 28 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access



# Vertebrate bacterial gut diversity: size also matters

Jean-Jacques Godon<sup>1\*</sup>, Pugazhendi Arulazhagan<sup>1,2</sup>, Jean-Philippe Steyer<sup>1</sup> and Jérôme Hamelin<sup>1</sup>

## Abstract

**Background:** One of the central issues in microbial ecology is to understand the parameters that drive diversity. Among these parameters, size has often been considered to be the main driver in many different ecosystems. Surprisingly, the influence of size on gut microbial diversity has not yet been investigated, and so far in studies reported in the literature only the influences of age, diet, phylogeny and digestive tract structures have been considered. This study explicitly challenges the underexplored relationship connecting gut volume and bacterial diversity.

**Results:** The bacterial diversity of 189 faeces produced by 71 vertebrate species covering a body mass range of 5.6 log. The animals comprised mammals, birds and reptiles. The diversity was evaluated based on the Simpson Diversity Index extracted from 16S rDNA gene fingerprinting patterns. Diversity presented an increase along with animal body mass following a power law with a slope  $z$  of  $0.338 \pm 0.027$ , whatever the age, phylogeny, diet or digestive tract structure.

**Conclusions:** The results presented here suggest that gut volume cannot be neglected as a major driver of gut microbial diversity. The characteristics of the gut microbiota follow general principles of biogeography that arise in many ecological systems.

**Keywords:** Biodiversity, Biogeography, Gut, Fingerprint, Species-area relationship

## Background

Among a number of parameters, the 'size' of an ecosystem is often assumed to have a key impact on the management of diversity. In fact, the species-area relationship is central to the ecological theory [1] and was first described for macro-organisms [2]. For bacteria, the species-area relationship is generally expressed in terms of habitat volume (i.e., volume-area relationship) and has been illustrated in liquid sump tanks of metal-cutting machines [3], membrane bioreactors [4] and tree holes (i.e., rainwater accumulated in holes at the base of large trees) [5]. However, until present, the microbial species-volume relationship has never yet been studied for gut or body size, even though vertebrate gut size covers a wide range of magnitudes. There is a  $10^6$  body mass factor between a tiny bird or a shrew and an elephant.

The vertebrate gut hosts a microbial community that fulfils many vital functions for the host: it enhances resistance to infection, stimulates mucosal immune defences, synthesizes essential vitamins and promotes caloric uptake by hydrolysing complex carbohydrates. The bacterial populations inhabiting the gut are complex, varying considerably from individual to individual and from species to species. However, gut microbial ecosystems are not a random association of microbes but are shaped by the host. A transfer occurs vertically from mothers to offspring or horizontally between individuals within a specific group. Such transfers have given rise to the long-standing co-evolution of microbiota and their hosts [6].

The benefit of bacterial diversity in the human gut has often been highlighted [7] and driving factors such as age [8], diverse lifestyles [9] and diet variations [10] have already been explored. Despite such an interest, the relationship between body mass and gut microbiota has never been explored whereas, in contrast, the positive

\*Correspondence: jean-jacques.godon@supagro.inra.fr

<sup>1</sup> UR0050, Laboratoire de Biotechnologie de l'Environnement, INRA, 102 avenue des étangs, 11100 Narbonne, France

Full list of author information is available at the end of the article

links between the abundance of parasitic organisms or protozoal faunas and animal body size have been thoroughly referenced [11] [12]. The aim of the present study is to analyse a large bacterial dataset, comprising faeces collected from 71 different vertebrate species, in order to examine the effect of the volume-microbial diversity relationship in animal digestive tracts.

## Methods

### Sampling

All the animal samples were obtained from domesticated or captive populations in France (zoo, farm, aquarium, recreational farm or individual keeper). There is non-experimental research dedicated for this study, faeces samples were collected on ground with the animal keeper or animal owner without stresses for the animals. We obtained permissions from Lunaret zoo, Montpellier; Océanopolis, Brest; Réserve Africaine, Sigean; Mini Ferme Zoo, Cessenon sur Orb and consent from the animal owners (Jean-Philippe Steyer, Anais Bonnafous, Jean-Jacques Godon). Animal were living alone or in small groups (1 to 5). Furthermore, their food (meat, seeds, fruits or hay) were more standardized in comparison to wild diets.

Human stool specimens used in the present study were from infant and adult subjects included in international multicentric studies. Samples were collected between 2001 to 2005 and used on previous published studies. Infants samples were collected in the frame of the European project INFABIO (<http://www.gla.ac.uk/departments/infabio/>), ethical permission was obtained from Yorkhill Research Ethics Committee P16/03 and parents gave written informed consent [13]. Adults samples were collected in the frame of the European project Crownalife, the studies were approved by the Ethics Committee of Versailles Hospital Centre and written informed consent was obtained from all participants [14]. Approval for Institut National de la Recherche Agronomique to manage human-derived biological samples in accordance with Articles L.1243-3, R.1243-49 of “Code de la Santé Publique” was granted by the Ministry of Research and Education under number DC-2012-1728.

Faeces from 189 individuals belonging to 71 vertebrate species (31 mammals, 37 birds and 3 reptiles) were collected (Table 1). They were sub-divided into 80 categories according to species or to body mass (i.e., age (young–adult), sex (female–male), breed size (small–big–domesticated–wild), see Table 1). Body masses were provided by the breeder for large animals or obtained from literature for small animals. Body masses, along with diversity, were displayed with a logarithmic scale in order to highlight the linear shape of the power-law relationship. Except for the distinct dimorphism of male and female turkey samples, an average value of male and female

body mass values was used. Dwarf or young individuals from the same species were also classified in specific body mass categories. For example, human samples were divided into two body mass categories: babies between 1 and 10 months old (mean of 5.8 kg) and adults between 29 and 61 years old (set at 70 kg). Composite faeces samples were avoided except for those that could not provide enough material for DNA extraction (less than 0.5 g).

### DNA extraction, PCR amplification and Capillary Electrophoresis Single Strand Conformation Polymorphism (CE-SSCP) fingerprinting

Genomic DNAs were extracted from 0.5 g of raw material using the procedure described by Godon et al. [15]. The V3 region of the 16S rRNA gene was amplified with *Bacteria*-specific primers and PCR products were analysed by CE-SSCP analysis using an ABI3130 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA) in accordance with a previously described method [16]. All raw CE-SSCP data are available on Additional file 4.

### Calculation of diversity and statistical computing

Diversity was estimated by the Simpson Diversity Index from CE-SSCP fingerprinting patterns. The Simpson Diversity Index was expressed as  $D = 1 / \sum_{i=1}^p a_i^2$  where  $a_i$  is the relative abundance of each CE-SSCP peak  $p$ . This index was directly calculated from each CE-SSCP fingerprint [17] using the R StatFingerprints library [18].

Preference was given to the Simpson Diversity Index from CE-SSCP fingerprinting rather than the Richness estimation because: (1) neither fingerprinting nor sequencing data can provide a robust estimation of richness [19]; (2) the Simpson Diversity Index can be estimated accurately with CE-SSCP fingerprinting [17, 20].

A generalized linear model was applied to fit the relationship between body mass and diversity. ANOVA followed by Tukey post hoc tests were used for determining the statistical difference between (sub-) categories and body mass or diversity, both expressed in a logarithmic scale. All statistics were performed under R software (version 3.1.2) [21]. The calculation of the slope  $z$  was based on the exponent of the power-law relationship as follows: diversity =  $c \text{ weight}^z$

## Results and discussion

The bacterial diversity of faeces from 189 vertebrates belonging to 71 species (31 mammals, 37 birds and 3 reptiles) was analysed (Table 1; Fig. 1). Analysis was only focused on diversity (Simpson Diversity Index), which can be accurately measured according to CE-SSCP fingerprinting patterns [15] (see the “Methods” section and Additional file 1). Apart from their phylogenetic position, animals can also be classified according to: (1) their

**Table 1 Animal data ranked by body mass**

Name (common name)	Phylogeny	Body mass (kg)	Feeding type	Type of digestive tract	Size of animal husbandry group	Diversity	SD	Number of samples
<i>Taeniopygia guttata</i> (zebra finch)	Aves, Passeriformes	0.012	Granivorous	Hindgut colon	Large	1.2	0.1	3
<i>Serinus canaria</i> (canary)	Aves, Passeriformes	0.024	Granivorous	Hindgut colon	Large	1.6	0.5	2
<i>Ramphocelus bresilius</i> (brazilian tanager)	Aves, Passeriformes	0.035	Frugivorous	Hindgut colon	Small	3.4	0.4	4
<i>Melospiza undulatus</i> (budgerigar)	Aves, Psittaciformes	0.04	Granivorous	Hindgut colon	Large	1.9	3.0	3
<i>Ploceus cucullatus</i> (village weaver)	Aves, Passeriformes	0.04	Granivorous	Hindgut colon	Small	2.3	0.0	2
<i>Agapornis fischeri</i> (Fischer's lovebird)	Aves, Psittaciformes	0.05	Granivorous	Hindgut colon	Large	1.5		1
<i>Agapornis roseicollis</i> (rosy-faced lovebird)	Aves, Psittaciformes	0.05	Granivorous	Hindgut colon	Large	2.1	0.0	2
<i>Amblyramphus holosericeus</i> (scarlet-headed blackbird)	Aves, Passeriformes	0.08	Carnivorous	Hindgut colon	Small	3.4	0.2	2
<i>Nymphicus hollandicus</i> (cockatiel)	Aves, Psittaciformes	0.08	Granivorous	Hindgut colon	Small	1.4	0.3	2
<i>Guira guira</i> (guira cuckoo)	Aves, Cuculiformes	0.14	Carnivorous	Hindgut colon	Small	2.9	0.5	4
<i>Poicephalus senegalus</i> (senegal parrot)	Aves, Psittaciformes	0.14	Granivorous	Hindgut colon	Small	2.6		1
<i>Streptopelia decaocto</i> (eurasian collard dove)	Aves, Columbidae	0.19	Granivorous	Hindgut colon	Large	2.4	0.5	3
<i>Corvus monedula</i> (eurasian jackdaw)	Aves, Passeriformes	0.22	Omnivorous	Hindgut colon	Small	1.8		1
<i>Psarocolius decumanus</i> (crested oropendola)	Aves, Passeriformes	0.3	Omnivorous	Hindgut colon	Small	3.8	0.3	3
<i>Columba livia</i> (pigeon)	Aves, Columbidae	0.3	Granivorous	Hindgut colon	Large	1.9		1
<i>Gallus gallus</i> (dwarf chicken) <sup>a</sup>	Aves, Galliformes	0.3	Granivorous	Hindgut caecum	Large	2.1	0.5	2
<i>Tauraco erythrolophus</i> (red-crested turaco)	Aves, Cuculiformes	0.35	Frugivorous	Hindgut colon	Small	3.8		1
<i>Agamia agami</i> (agami heron)	Aves, Ciconiiformes	0.46	Piscivorous	Hindgut colon	Small	3.6		1
<i>Coracopsis vasa</i> (vasa parrot)	Aves, Psittaciformes	0.5	Frugivorous	Hindgut colon	Small	2.4		1
<i>Chinchilla laniger</i> x <i>Chinchilla brevicaudata</i> (chinchilla)	Mammalia, Rodentia	0.6	Herbivorous	Hindgut caecum	Small	4.2	0.1	2
<i>Ramphastos tucanus</i> (white-throated toucan)	Aves, Piciformes	0.675	Frugivorous	Hindgut colon	Small	3.6	0.4	3
<i>Chrysolophus pictus</i> (golden pheasant)	Aves, Galliformes	0.700	Granivorous	Hindgut caecum	Small	3.4		1
<i>Cavia porcellus</i> (domestic guinea pig)	Mammalia, Rodentia	0.8	Herbivorous	Hindgut caecum	Large	5.0	0.2	3
<i>Anas acuta</i> (northern pintail)	Aves, Anatidae	0.9	Granivorous	Hindgut caecum	Small	3.6		1
<i>Elaphe guttata</i> (corn snake)	Sauropsida, Serpentes	0.9	Carnivorous	Hindgut colon	Small	4.1		1
<i>Lampropeltis getula</i> (common kingsnake)	Sauropsida, Serpentes	1	Carnivorous	Hindgut colon	Small	3.3		1
<i>Ara ararauna</i> (blue-and-yellow macaw)	Aves, Psittaciformes	1	Granivorous	Hindgut colon	Small	2.8	0.1	2
<i>Anas platyrhynchos</i> (wild type duck) <sup>a</sup>	Aves, Anatidae	1.1	Granivorous	Hindgut caecum	Large	3.1		1
<i>Neochen jubata</i> (orinoco goose)	Aves, Anatidae	1.25	Granivorous	Hindgut caecum	Small	3.5		1

**Table 1 continued**

Name (common name)	Phylogeny	Body mass (kg)	Feeding type	Type of digestive tract	Size of animal husbandry group	Diversity	SD	Number of samples
<i>Gallus gallus</i> (chicken) <sup>a</sup>	Aves, Galliformes	1.5	Granivorous	Hindgut caecum	Large	1.7	0.5	6
<i>Numida meleagris</i> (guinea-fowl)	Aves, Galliformes	2	Granivorous	Hindgut caecum	Large	3.0	0.7	3
<i>Branta sandvicensis</i> (nene)	Aves, Anatidae	2	Granivorous	Hindgut caecum	Small	2.6		1
<i>Oryctolagus cuniculus</i> (domestic rabbit)	Mammalia, Lagomorpha	2.2	Herbivorous	Hindgut caecum	Large	4.5	0.7	3
<i>Anas platyrhynchos</i> (domestic duck) <sup>a</sup>	Aves, Anatidae	2.3	Granivorous	Hindgut colon	Small	2.3	1.4	3
<i>Eudyptes chrysocome</i> (western rockhopper penguin)	Aves, Sphenisciformes	2.6	Piscivorous	Hindgut colon	Large	1.6	0.6	3
<i>Testudo hermanni boettgeri</i> (Hermann's tortoise)	Sauropsida, Testudines	3	Herbivorous	Hindgut colon	Large	5.6		1
<i>Meleagris gallopavo</i> (turkey female) <sup>a</sup>	Aves, Galliformes	3	Granivorous	Hindgut caecum	Small	2.2		1
<i>Thylogale sp.</i> (pademelon)	Mammalia, Marsupials	3.5	Herbivorous	Hindgut colon	Small	4.3		1
<i>Cairina moschata</i> (muscovy duck)	Aves, Anatidae	4	Granivorous	Hindgut colon	Large	3.3		2
<i>Chauna torquata</i> (southern screamer)	Aves, Anseriformes	4	Herbivorous	Hindgut caecum	Small	3.4		1
<i>Canis lupus familiaris</i> (puppy) <sup>a</sup>	Mammalia, Carnivora	4	Carnivorous	Hindgut colon	Small	2.6		1
<i>Pavo cristatus</i> (blue peafowl)	Aves, Galliformes	5	Granivorous	Hindgut caecum	Small	3.7	0.1	2
<i>Anser anser domesticus</i> (domestic goose)	Aves, Anatidae	5	Granivorous	Hindgut caecum	Small	3.5	0.1	2
<i>Homo sapiens</i> (baby human caucasian) <sup>a</sup>	Mammalia, Primates	6	Omnivorous	Hindgut colon	Small	3.2	0.7	15
<i>Meleagris gallopavo</i> (turkey male) <sup>a</sup>	Aves, Galliformes	8	Granivorous	Hindgut caecum	Small	3.8		1
<i>Wallabia bicolor</i> (black wallaby)	Mammalia, Marsupials	9	Herbivorous	Hindgut colon	Small	4.8		1
<i>Hylobates lar</i> (gibbon)	Mammalia, Primates	10	Frugivorous	Hindgut colon	Small	5.5		1
<i>Aptenodytes patagonicus</i> (king penguin)	Aves, Sphenisciformes	13	Piscivorous	Hindgut colon	Small	2.8	1.1	4
<i>Capra hircus</i> (dwarf goat)	Mammalia, Ruminantia	15	Herbivorous	Ruminants foregut	Small	5.2	1.1	2
<i>Canis lupus familiaris</i> (medium size dog) <sup>a</sup>	Mammalia, Carnivora	20	Carnivorous	Hindgut colon	Small	3.2	1.1	2
<i>Ovis aries</i> (dwarf sheep) <sup>a</sup>	Mammalia, Ruminantia	20	Herbivorous	Ruminants foregut	Small	5.7	0.7	2
<i>Hippotragus equinus</i> (roan antelope)	Mammalia, Ruminantia	20	Herbivorous	Ruminants foregut	Small	5.7		1
<i>Tragelaphus streps</i> (greater kudu)	Mammalia, Ruminantia	20	Herbivorous	Ruminants foregut	Small	5.5		1
<i>Hystrix cristata</i> (crested porcupine)	Mammalia, Rodentia	25	Herbivorous	Hindgut caecum	Small	5.7		1
<i>Rhea americana</i> (greater rhea)	Aves, Rheiformes	31	Granivorous	Hindgut caecum	Large	4.0	0.5	4
<i>Ovis aries</i> (sheep) <sup>a</sup>	Mammalia, Ruminantia	40	Herbivorous	Ruminants foregut	Small	5.0		1
<i>Canis lupus familiaris</i> (big size dog) <sup>a</sup>	Mammalia, Carnivora	40	Carnivorous	Hindgut colon	Small	3.1		1
<i>Pan troglodytes</i> (chimpanzee)	Mammalia, Primates	40	Omnivorous	Hindgut colon	Small	5.3		1

**Table 1 continued**

Name (common name)	Phylogeny	Body mass (kg)	Feeding type	Type of digestive tract	Size of animal husbandry group	Diversity	SD	Number of samples
<i>Dromaius novaehollandiae</i> (emu)	Aves, Casuariiformes	40	Granivorous	Hindgut colon	Small	3.9		1
<i>Capra hircus</i> (goat)	Mammalia, Ruminantia	50	Herbivorous	Ruminants foregut	Small	7.0		1
<i>Sus scrofa</i> (dwarf pig) <sup>a</sup>	Mammalia, Suina	55	Omnivorous	Hindgut colon	Small	5.4	0.4	2
<i>Lama glama</i> (llama)	Mammalia, Tylopoda	55	Herbivorous	Ruminants foregut	Small	5.4		1
<i>Homo sapiens</i> (adult human caucasian) <sup>a</sup>	Mammalia, Primates	70	Omnivorous	Hindgut colon	Large	4.4	0.8	34
<i>Sus scrofa</i> (pig) <sup>a</sup>	Mammalia, Suina	100	Omnivorous	Hindgut colon	Small	5.8	1.1	4
<i>Tragelaphus spekei</i> (sitatunga)	Mammalia, Ruminantia	100	Herbivorous	Ruminants foregut	Small	7.5		1
<i>Struthio camelus</i> (ostrich)	Aves, Struthioniformes	120	Herbivorous	Hindgut colon	Small	4.4	0.2	3
<i>Equus asinus</i> (donkey)	Mammalia, Equidae	150	Herbivorous	Hindgut caecum	Small	5.3	0.4	2
<i>Ammotragus lervia</i> (aoudad)	Mammalia, Ruminantia	150	Herbivorous	Ruminants foregut	Small	6.1		1
<i>Equus caballus</i> (pony)	Mammalia, Equidae	160	Herbivorous	Hindgut caecum	Small	5.6	0.1	2
<i>Panthera leo</i> (african lion)	Mammalia, Carnivora	160	Carnivorous	Hindgut colon	Small	4.4		1
<i>Equus zebra hartmannae</i> (mountain zebra)	Mammalia, Equidae	350	Herbivorous	Hindgut caecum	Small	5.4		1
<i>Syncerus caffer nanus</i> (forest buffalo)	Mammalia, Ruminantia	450	Herbivorous	Ruminants foregut	Small	2.9		1
<i>Camelus dromedarius</i> (arabian Camel)	Mammalia, Tylopoda	500	Herbivorous	Ruminants foregut	Small	3.2		1
<i>Bos grunniens</i> (yak)	Mammalia, Ruminantia	600	Herbivorous	Ruminants foregut	Small	5.3		1
<i>Tragelaphus oryx</i> (eland antelope)	Mammalia, Ruminantia	600	Herbivorous	Ruminants foregut	Small	6.2		1
<i>Bos taurus</i> (cow)	Mammalia, Ruminantia	750	Herbivorous	Ruminants foregut	Large	6.2	0.7	4
<i>Giraffa camelopardalis reticulata</i> (somali giraffe)	Mammalia, Ruminantia	1100	Herbivorous	Ruminants foregut	Small	6.4		1
<i>Giraffa camelopardalis peralta</i> (nigerian giraffe)	Mammalia, Ruminantia	1100	Herbivorous	Ruminants foregut	Small	6.6		1
<i>Ceratotherium simum</i> (white rhinoceros)	Mammalia, Rhinocerotidae	2500	Herbivorous	Hindgut colon	Small	5.6		1
<i>Elephas maximus</i> (asian elephant)	Mammalia, Proboscidea	3500	Herbivorous	Hindgut colon	Small	4.9		1


SD standard deviation

<sup>a</sup> species with different sizes (young-adult, female-male, small-big or domesticated-wild)

diet (herbivorous, granivorous, omnivorous, carnivorous, piscivorous and frugivorous); (2) their metabolic body mass (from 12 g (zebra finch) to 3500 kg (Asian elephant)); (3) the structure of their digestive tracts; (4) and the size of the animal husbandry group (small and large). The present study focused on bacterial diversity, although changes within the structure of the bacterial communities were not taken into account. This study is

also based on two assumptions: (1) the gut size should be proportional to the animal body mass, as has been demonstrated for herbivores [22] and birds [23]; and (2) the microbial diversity of faeces should be similar to that in the gut [24].

Results point to a correlation between animal body mass and microbial diversity (linear regression with a slope  $z$  of  $0.338 \pm 0.027$ ;  $p$  value  $< 2.2 \times 10^{-16}$ ),



irrespective of the diet, phylogeny or structure of the digestive tracts (Fig. 1). Consequently, the use of a greater amount of samples over a wider size range confirms previous works on unrelated bacterial communities that have suggested the existence of a link between volume and diversity in tree holes [5], membrane bioreactors [4] and metal-cutting fluid sump tanks [3]. In the present results, the Simpson Diversity Index ranges between 3.3 and 1789.5, thus corresponding to a 5.6 log body mass range (Fig. 1).

A wide variability in the diversity between individuals for a given species was observed. However the average diversity value for species that were represented by several individuals was close to the regression line (Fig. 1). For example, the average diversity value for adult human microbiota (34 samples) was 80.8 with a standard

deviation of 294.2, and  $23.7 \pm 20.3$  for the 15 baby human microbiota. As a matter of comparison, Trosvik et al. [25] observed a similar range of diversity (over 2 log-units of Shannon index) when analysing a time-series of 332 sequencings over 443 days, on a single male adult individual.

Animal gut microbiota covered a broad range of diversity ranging from 2.2 to 1808.0. This was comparable to the values found in various types of environment, like drinking water, raw milk, plant roots, activated sludge in wastewater treatment plants, compost or soil (Additional file 2). On one hand, the lowest diversity in gut microbiota varied around 2, similarly to those found in drinking water. On the other hand, the highest diversity in gut microbiota reaching about 1808 resembled the values found in soils (Additional file 2).

**Table 2 Bacterial diversity and animal weight within sub-categories, correlation between diversity and weight, and slope of the relationship of the diversity versus log-weight**

Category	Number of samples	Simpson diversity mean (SD)	Weight in kg mean (SD)	Pearson correlation between diversity and weight		Power law relationship diversity = c weight <sup>z</sup>		
				cor	p value	Slope z	Confidence interval	
Sub-categories								
Diet								
Carnivorous	13	32.0 (23.4)	19.0 (44.1)	0.277	0.359 (NS)	0.075 (NS)	-	
Frugivorous	10	55.2 (68.5)	1.3 (3.1)	0.533	0.113 (NS)	0.234 (NS)	-	
Granivorous	54	20.4 (21.7)	4.4 (9.4)	0.667	3.7e-08 (***)	0.298 (***)	0.205-0.391	
Herbivorous	44	301.6 (342.3)	354.5 (668.1)	0.338	0.025 (*)	0.137 (*)	0.018-0.256	
Omnivorous	60	111.3 (149.0)	50.5 (32.4)	0.542	7.7e-06 (***)	0.361 (***)	0.214- 0.508	
Piscivorous	8	20.4 (20.8)	7.5 (5.9)	0.030	0.944 (NS)	0.029 (NS)	-	
Phylogeny								
Bird	85	25.1 (23.6)	7.8 (23.0)	0.456	1.1e-05 (***)	0.202 (***)	0.116-0.288	
Mammal	101	194.9 (268.8)	183.3 (464.0)	0.415	1.6e-05 (***)	0.272 (***)	0.153-0.391	
Reptile	3	119.3 (131.9)	1.6 (1.2)	0.964	0.172 (NS)	1.686 (NS)	-	
Gut structure								
Caecum	46	70.4 (80.6)	26.1 (65.3)	0.528	1.7e-04 (***)	0.397 (***)	0.203-0.591	
Colon	124	78.5 (122.5)	79.0 (382.7)	0.678	<2.2e-16 (***)	0.293 (***)	0.236-0.350	
Rumen	19	484.9 (449.7)	411.8 (384.5)	-0.036	0.883 (NS)	-0.031 (NS)	-	
Group size								
Large	85	93.1 (151.0)	65.8 (156.4)	0.734	1.3e-15 (***)	0.380 (***)	0.303-0.457	
Small	104	137.2 (253.7)	130.6 (449.1)	0.632	6.6e-13 (***)	0.300 (***)	0.227-0.372	
Age								
Adult	173	125.6 (222.2)	110.3 (364.6)	0.687	<2.2e-16 (***)	0.337 (***)	0.283-0.391	
Baby	16	28.9 (20.1)	6.3 (2.0)	0.205	0.446 (NS)	0.427 (NS)	-	
All	189	117.4 (214.3)	101.5 (350.0)	0.675	<2.2e-16 (***)	0.338 (***)	0.284-0.391	

NS not significant

\* low significance, \*\*\* high significance



This vast range of variations in gut diversity is often associated with factors that are different to the body mass: diet [10], phylogeny [26], digestive tract structure [27], age [8] [28], way of life [29], ethnic origin [30], state of health (immune system, pregnancy, obesity) [31] [32], or genetic background [32]. Among these parameters, age has been well documented as the major one to explain these variations and the diversity or richness between human baby microbiota and those of adults [33] [34]. However the size of the gut also varies during infant growth. In this case, a difference in the microbial diversity between infant ( $29.9 \pm 20.3$ ) and adults ( $106.6 \pm 76.0$ ) was observed, concomitantly with changes in body mass when comparing human babies ( $6.5 \pm 1.9$ ) and adults (70 kg). The same observation was made for young and adult dog samples (Table 1). Furthermore, when comparing two penguin species that only differ in their body mass (only adult specimens, with the same diet and living in the same location), the relationship between microbial diversity and body mass still remain valid.

The correlation between body mass and diversity has been assessed for homogenous sub-categories (Table 2 and Additional file 3), thus excluding the potential effects of the different parameters. Indeed, the 189 samples could also be analysed according to phylogeny (reptile, bird, and mammal), diet (carnivorous, herbivorous, granivorous, omnivorous and piscivorous), gut structure (hindgut caecum, hindgut colon and foregut ruminant), age (baby and adult), and size of the animal husbandry group (small and large). Except for the latter category, all of them depended on the body mass (e.g. body mass was related to phylogeny, related to age or to ruminants). Significantly positive body mass/diversity correlations were observed for each sub-category, provided that a sufficient amount of data was available (over 50 samples minimum per sub-category) (Table 2; Additional file 3). The significant slopes  $z$  of the mass-diversity relationships generally ranged from  $0.202 \pm 0.043$  to  $0.380 \pm 0.039$ . As the herbivorous group only contained 44 samples, the interestingly weak body mass diversity correlation with a  $z$  value of 0.137 could not be correctly interpreted.

The observed slope  $z$  was similar to that reported for 'island' patterns of bacterial diversity such as metal-cutting fluid sump tanks ( $z = 0.245\text{--}0.295$ ) [3] and tree holes ( $z = 0.26$ ) [5] and varied within a similar range to that reported for plants and animals from discrete islands ( $z = 0.25\text{--}0.35$ ). The slope  $z$ -values reported for continuous patterns (such as marsh sediment [35] with  $z$ -values between 0.02 and 0.04) are generally much lower than those reported for discrete habitats.

According to these results, which confirm the assumption that species and volume are related, guts can

compared to an archipelago, where microbes originating from feed tend to colonise the available niches provided by the gut. This is also in line with the MacArthur and Wilson biogeography theory [1]. Size, similarly to island environments appears to reflect the heterogeneity of the environment. Hence, a large gut size should provide more space, enabling a large microbial diversity to settle in [36].

## Conclusions

The aim of this study was not to explain the genesis of bacterial diversity in vertebrate guts but was rather focused on producing evidence on the role of gut size in the maintenance of a level of microbial diversity. This work highlights the hitherto unexplored relationship between volume and diversity in the case of gut microbiota. Gut volume should henceforth be taken into account along with other parameters to explain the level of diversity. Finally, this work confirms the relevance of the microbial world when addressing ecological issues such as the relationship between species diversity and the size of the habitat [37].

## Availability of supporting data

Our data are provided in the electronic supplementary materials (Additional file 4).

## Additional files

**Additional file 1.** Examples of representative CE-SSCP fingerprinting patterns from low-diverse to high-diverse samples.

**Additional file 2.** Simpson Diversity Index calculated from 671 CE-SSCP fingerprinting patterns. Samples were grouped according to their ecosystem of origin.

**Additional file 3.** Simpson Diversity Index as a function of animal weight for (A) all the samples and for the sub-categories; (B) diet; (C) phylogeny; (D) gut structure; (E) age and (F) group size. The slope of the linear regression is indicated.

**Additional file 4.** Raw 671 CE-SSCP profiles. Columns correspond to the samples. The 7359 lines correspond to the fluorescence intensity according to time.

## Abbreviations

CE-SSCP: capillary electrophoresis single strand conformation polymorphism; PCR: polymerase chain reaction; 16S rRNA: 16S ribosomal RNA; ANOVA: analysis of variance.

## Authors' contributions

JJG designed the experiment, JJG and JH collected the data, PA performed the experiments and contributed to the analysis of the data. Analysis and interpretation were carried out by JH and JJG. JH and JJG contributed to the first draft, which was completed by JPS. All authors read and approved the final manuscript.

## Author details

<sup>1</sup> UR0050, Laboratoire de Biotechnologie de l'Environnement, INRA, 102 avenue des étangs, 11100 Narbonne, France. <sup>2</sup> Present Address: Centre of Excellence in Environmental Studies, King Abdulaziz University, Jeddah, Saudi Arabia.

### Acknowledgements

We thank the following people and organizations who generously provided samples: Joël Doré; Thierry Gidenne; 'Reserve africaine', Sigean, France; 'Mini-ferme zoo', Cessenon/Orb, France; 'Oceanopolis', Brest, France; and Lunaret Zoo, Montpellier, France. We thank Biswarup Sen, Anais Bonnafous and Valérie Bru-Adan for technical assistance. INRA funded this research.

### Competing interests

The authors declare that they have no competing interests.

Received: 18 May 2015 Accepted: 10 March 2016

Published online: 23 March 2016

### References

- MacArthur RH, Wilson EO. The theory of island biogeography. Princeton Univ Press Monogr Popul Biol. 1967;1:202.
- Arrhenius O. Species and area. *J Ecol*. 1921;9:95–9.
- van der Gast CJ, Lilley AK, Ager D, Thompson IP. Island size and bacterial diversity in an archipelago of engineering machines. *Environ Microbiol*. 2005;7:1220–6.
- Van der Gast CJ, Jefferson B, Reid E, Robinson T, Bailey MJ, Judd SJ, Thompson IP. Bacterial diversity is determined by volume in membrane bioreactors. *Environ Microbiol*. 2006;8:1048–55.
- Bell T, Ager D, Song JI, Newman JA, Thompson IP, Lilley AK, van der Gast CJ. Larger islands house more bacterial taxa. *Science*. 2005;308:1884.
- Ley RE, Peterson DA, Gordon JI. Ecological and evolutionary forces shaping microbial diversity in the human intestine. *Cell*. 2006;124:837–48.
- Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, Almeida M, Arumugam M, Batto J-M, Kennedy S, Leonard P, Li J, Burgdorf K, Grarup N, Jorgensen T, Brandslund I, Nielsen HB, Juncker AS, Bertalan M, Levenez F, Pons N, Rasmussen S, Sunagawa S, Tap J, Tims S, Zoetendal EG, Brunak S, Clement K, Dore J, Kleerebezem M, et al. Richness of human gut microbiome correlates with metabolic markers. *Nature*. 2013;500:541.
- Yatsunencko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, Magris M, Hidalgo G, Baldassano RN, Anokhin AP, Heath AC, Warner B, Reeder J, Kuczynski J, Caporaso JG, Lozupone CA, Lauber C, Clemente JC, Knights D, Knight R, Gordon JI. Human gut microbiome viewed across age and geography. *Nature*. 2012;486:222.
- Schnorr SL, Candela M, Rampelli S, Centanni M, Consolandi C, Basaglia G, Turrioni S, Biagi E, Peano C, Severgnini M, Fiori J, Gotti R, De Bellis G, Luiselli D, Brigidi P, Mabulla A, Marlowe F, Henry AG, Crittenden AN. Gut microbiome of the Hadza hunter-gatherers. *Nat Commun*. 2014;5:3654.
- Vital M, Gao J, Rizzo M, Harrison T, Tiedje JM. Diet is a major factor governing the fecal butyrate-producing community structure across mammalia aves and reptilia. *ISME J*. 2015;9:832–43.
- Kamiya T, O'Dwyer K, Nakagawa S, Poulin R. What determines species richness of parasitic organisms? A meta-analysis across animal, plant and fungal hosts. *Biol Rev*. 2014;89:123–34.
- Clauss M, Mueller K, Fickel J, Streich WJ, Hatt J-M, Suedekum K-H. Macroecology of the host determines microecology of endobionts: protozoal faunas vary with wild ruminant feeding type and body mass. *J Zool*. 2011;283:169–85.
- Fallani M, Amarri S, Uusjarvi A, Adam R, Khanna S, Aguilera M, Gil A, Vieites JM, Norin E, Young D, Scott JA, Doré J, Edwards CA, Team I. Determinants of the human infant intestinal microbiota after the introduction of first complementary foods in infant samples from five European centres. *Microbiol SGM*. 2011;157:1385–92.
- Mueller S, Saunier K, Hanisch C, Norin E, Alm L, Midtvedt T, Cresci A, Silvi S, Orpianesi C, Verdenelli MC, Clavel T, Koebnick C, Zunft HJF, Dore J, Blaut M. Differences in fecal microbiota in different European study populations in relation to age, gender, and country: a cross-sectional study. *Appl Environ Microbiol*. 2006;72:1027–33.
- Godon JJ, Zumstein E, Dabert P, Habouzit F, Moletta R. Molecular microbial diversity of an anaerobic digester as determined by small-subunit rDNA sequence analysis. *Appl Environ Microbiol*. 1997;63:2802–13.
- Wéry N, Bru-Adan V, Minervini C, Delgènes JP, Garrelly L, Godon JJ. Dynamics of *Legionella* spp. and bacterial populations during the proliferation of *L. pneumophila* in a cooling tower facility. *Appl Environ Microbiol*. 2008;74:3030–7.
- Haegeman B, Sen B, Godon JJ, Hamelin J. Only Simpson diversity can be estimated accurately from microbial community fingerprints. *Microb Ecol*. 2014;68:169–72.
- Michelland RJ, Dejean S, Combes S, Fortun-Lamothe L, Cauquil L. Stat-Fingerprints: a friendly graphical interface program for processing and analysis of microbial fingerprint profiles. *Mol Ecol Resour*. 2009;9:1359–63.
- Haegeman B, Hamelin J, Moriarty J, Neal P, Dushoff J, Weitz JS. Robust estimation of microbial diversity in theory and in practice. *ISME J*. 2013;7:1092–101.
- Lalande J, Villemur R, Deschenes L. A new framework to accurately quantify soil bacterial community diversity from DGGE. *Microb Ecol*. 2013;66:647–58.
- R-Core-Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2014.
- Clauss M, Schwarm A, Ortmann S, Streich WJ, Hummel J. A case of non-scaling in mammalian physiology? Body size, digestive capacity, food intake, and ingesta passage in mammalian herbivores. *Comp Biochem Physiol A: Mol Integr Physiol*. 2007;148:249–65.
- Lavin SR, Karasov WH, Ives AR, Middleton KM, Garland T Jr. Morphometrics of the avian small intestine compared with that of nonflying mammals: a phylogenetic approach. *Physiol Biochem Zool*. 2008;81:526–50.
- Michelland RJ, Monteils V, Zened A, Combes S, Cauquil L, Gidenne T, Hamelin J, Fortun-Lamothe L. Spatial and temporal variations of the bacterial community in the bovine digestive tract. *J Appl Microbiol*. 2009;107:1642–50.
- Trosvik P, de Muinck EJ, Stenseth NC. Biotic interactions and temporal dynamics of the human gastrointestinal microbiota. *ISME J*. 2015;9:533–41.
- Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI. Worlds within worlds: evolution of the vertebrate gut microbiota. *Nat Rev Microbiol*. 2008;6:776–88.
- Stevens CE, Hume ID. Comparative physiology of the vertebrate digestive system. Cambridge: University Press; 2004.
- Waite DW, Eason DK, Taylor MW. Influence of hand rearing and bird age on the fecal microbiota of the critically endangered kakapo. *Appl Environ Microbiol*. 2014;80:4650–8.
- Dicksved J, Floistrup H, Bergstrom A, Rosenquist M, Pershagen G, Scheynius A, Roos S, Alm JS, Engstrand L, Braun-Fahrlander C, von Mutius E, Jansson JK. Molecular fingerprinting of the fecal microbiota of children raised according to different lifestyles. *Appl Environ Microbiol*. 2007;73:2284–9.
- Kwok L, Zhang J, Guo Z, Gesudu Q, Zheng Y, Qiao J, Huo D, Zhang H. Characterization of fecal microbiota across seven Chinese ethnic groups by quantitative polymerase chain reaction. *PLoS One*. 2014;9(4):93631.
- Rescigno M. Intestinal microbiota and its effects on the immune system. *Cell Microbiol*. 2014;16:1004–13.
- Turnbaugh PJ, Hamady M, Yatsunencko T, Cantarel BL, Duncan A, Ley RE, Sogin ML, Jones WJ, Roe BA, Affourtit JP, Egholm M, Henrissat B, Heath AC, Knight R, Gordon JI. A core gut microbiome in obese and lean twins. *Nature*. 2009;457:480–7.
- Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, Knight R, Angenent LT, Ley RE. Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci*. 2011;108:4578–85.
- Henderson G, Cox F, Ganesh S, Jonker A, Young W, Collaborators GRC, Janssen PH. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci Rep*. 2015;5:14567.
- Horner-Devine MC, Lage M, Hughes JB, Bohannan BJM. A taxa-area relationship for bacteria. *Nature*. 2004;432:750–3.
- Kassen R. The experimental evolution of specialists, generalists, and the maintenance of diversity. *J Evol Biol*. 2002;15:173–90.
- Jessup CM, Kassen R, Forde SE, Kerr B, Buckling A, Rainey PB, Bohannan BJM. Big questions, small worlds: microbial model systems in ecology. *Trends Ecol Evol*. 2004;19:189–97.