



HAL
open science

A comprehensive review on flavanones, the major citrus polyphenols

Muhammad Kamran Khan, - Zill-E-Huma, Olivier O. Dangles

► **To cite this version:**

Muhammad Kamran Khan, - Zill-E-Huma, Olivier O. Dangles. A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 2014, 33, pp.85-104. 10.1016/j.jfca.2013.11.004 . hal-02640844

HAL Id: hal-02640844

<https://hal.inrae.fr/hal-02640844v1>

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A comprehensive review on flavanones, the major citrus polyphenols

Muhammad Kamran Khan ^{a,*}, Zill-E-Huma ^b, Olivier Dangles ^c

^a Department of Food Science, Government College University, Faisalabad, Pakistan
^b National Institute of Food Science & Technology, University of Agriculture, Faisalabad, Pakistan
^c University of Avignon, INRA, UMR408, 84000 F-Avignon, France

ABSTRACT

The consumption of *Citrus* fruits and juices has been widely investigated for its possible role in the prevention of cardiovascular disease and cancer. These beneficial effects are mainly attributed to flavanones, the typical polyphenols of *Citrus* species. Major flavanones in plant species include hesperetin, naringenin, eriodictyol, isosakuranetin and their respective glycosides. Hesperetin and its derivatives are characteristic flavanones of sweet orange, tangelo, lemon and lime, while naringenin and its derivatives are those of grapefruit and sour orange. Advances in analytical techniques like ultra high performance liquid chromatography (UPLC) coupled with mass spectrometry has facilitated (a) the estimation of flavanone contents in other plant species and in humans after ingestion and (b) the determination of flavanone metabolites more rapidly and with greater efficiency. The present review will summarize the current knowledge about flavanones from their occurrence in plants to the bioactivity of their metabolites in humans.

Keywords:
 Citrus fruit
 Orange
 Grapefruit
 Polyphenol
 Flavanone
 Chalcone
 Conjugate
 Glucuronide
 Metabolites
 Synthesis
 Extraction
 Analysis
 Bioavailability
 Bioactivity
 Food composition

Contents

1. Introduction	85
2. Structures and classification	85
3. Biosynthesis of flavanones in plants	86
4. Diversity and distribution of flavanones	86
4.1. Naringenin	86
4.2. Hesperetin	86
5. Extraction and analysis of flavanones	87
6. Chemical synthesis of flavanones	92
6.1. Aglycones	92
6.2. Chalcones	93
6.3. Glycosides	93
6.4. Glucuronides	93
7. Bioavailability of flavanones	93
7.1. Intestinal absorption	94
7.2. Metabolism	95
8. Interaction of flavanones with human serum albumin	96
9. Bioactivity of flavanones	97
9.1. Radical-scavenging effects	98

* Corresponding author. Tel.: +92 347 44 95 648; fax: +92 41 92 00 671.
 E-mail address: mk.khan@gcuf.edu.pk (M.K. Khan).

9.2.	Anti-inflammatory effects.....	98
9.3.	Anti-cancer effects	98
9.3.1.	Antimutagenic effects	99
9.3.2.	Inhibition of tumor development	99
9.3.3.	Anti-proliferation effects	99
9.4.	Cardiovascular effects.....	99
9.4.1.	Vasorelaxant and vasoprotective effects.....	99
9.4.2.	Effect on atherosclerosis and coronary heart disease	99
9.5.	Other biological effects.....	100
10.	Conclusion	100
	References	100

1. Introduction

It is now well accepted that a low consumption of fatty foods, a regular physical activity and a high consumption of plant-derived foods help to maintain a good health status. In particular, there is an association between an increased level of fruit and vegetables in the diet and a reduced risk of some life-threatening diseases such as cardiovascular disease and cancer (Parr and Bolwell, 2000). There is also growing acceptance that many phenolic secondary metabolites (polyphenols) present in plant-derived foods exert beneficial effects in the prevention of these degenerative diseases (Benavente-García and Castillo, 2008; Del Rio et al., 2010). Moreover, *Citrus* fruit and juices are highly consumed worldwide and this consumption has significantly increased during the past few years. Global production of *Citrus* fruit reached 82 million tons in the years 2009–2010, of which oranges – commercially the most important citrus fruit – accounts for about 50 million tons (USDA, 2010). Worldwide *Citrus* consumption has thus stimulated research on the most abundant *Citrus* phenols, i.e. flavanones. Based on the criterion of flavanone content, *Citrus* plants belonging to the *Rutaceae* family appear especially important. This review will cover the updated literature available on the chemistry and biochemistry of the main *Citrus* flavanones, hesperetin and naringenin.

2. Structures and classification

A few decades ago, flavanones were considered as only minor flavonoids (Bohm, 1994), like chalcones, dihydrochalcones, dihydroflavonols and aurones. However, during the past 15 years, the total number of known flavanones has increased to the point that they are now considered a major flavonoid class like flavones, isoflavones, flavanols, flavonols and anthocyanidins (Veitch and Grayer, 2006). Up to now about 350 flavanone aglycones and 100 flavanone glycosides have been identified in nature (Iwashina, 2000).

Generally, polyphenols are classified into two major classes: flavonoids and nonflavonoids. The latter include structurally simple molecules such as phenolic acids (hydroxybenzoic acids and hydroxycinnamic acids) and stilbenes, and highly complex molecules such as stilbene oligomers, tannins and lignins (Cheynier, 2005). The former, the most studied class of polyphenols, includes more than 9000 identified compounds (Martens and Mithöfer, 2005; Pietta, 2000). Flavonoids commonly share the same generic structure, the flavan nucleus, consisting of two aromatic rings (A and B) linked by a pyran ring (C). Differences in the location of the B-ring to C-ring linkage make it possible to distinguish between flavonoids (2-phenylbenzopyrans), isoflavonoids (3-phenylbenzopyrans), and neoflavonoids (4-phenylbenzopyrans). The by far the most abundant 2-phenylbenzopyran group may be further divided into 3-hydroxyflavonoids (flavonols, flavanols, anthocyanidins, dihydroflavonols), and flavonoids without substituent at C3 (flavanones and flavones). Flavones differ from flavanones by a C2–C3 double bond (Marais et al., 2006). The flavanone class encompasses a wide array of compounds with

O- and/or C-substitutions at the A- or B-ring, e.g., hydroxy, methoxy, methylenedioxy, O- and C-glycosyl, C-methyl, C-benzyl, C-hydroxymethyl, C-formyl, C-isoprenyl substituents (including furano or dihydrofurano rings), conjugations to stilbene, anastatin, phenolic acid, and diarylheptanoid moieties (Fig. 1) (Veitch and Grayer, 2006, 2008).

3. Biosynthesis of flavanones in plants

Due to the diverse physiological functions in plants and beneficial effects on human health, flavonoids are now attractive targets for genetic engineering strategies. In most plant species, the flavonoid biosynthetic pathway has been almost completely elucidated. In general, the biosynthesis of flavonoids is initiated by the two precursors, malonyl-CoA and *p*-coumaroyl-CoA, which are originated from carbohydrate metabolism and the phenylpropanoid pathway, respectively. After condensation of three molecules of malonyl-CoA with one molecule of *p*-coumaroyl-CoA, the yellow 2',4,4',6'-tetrahydroxychalcone is formed. This step is catalyzed by chalcone synthase (CHS). The unstable chalcone is then cyclized to the corresponding 4',5,7-trihydroxyflavanone by the enzyme chalcone isomerase (CHI). Flavanones may be regarded as the cornerstone of flavonoid biosynthesis as they are the precursors of all other flavonoid classes (Fig. 2) (Martens and Mithöfer, 2005; Schijlen et al., 2004). Moreover, in *Citrus* species, UDP-glucose flavanone-7-O-glucosyltransferase (UFGT) and UDP-rhamnose flavanone glucoside rhamnosyltransferase (UFGRT) sequentially convert the flavanone aglycones into their 7-O-β-D-glucosides and rhamnoglucosides (Lewinsohn et al., 1989) (Fig. 3).

Naturally occurring flavanones display the (S) configuration at C₂ (Tomas-Barberan and Clifford, 2000) as a consequence of the enantioselectivity of the chalcone isomerase (CHI)-catalyzed intramolecular Michael addition within the chalcone precursor (Jez and Noel, 2002a). However, mixtures of (2R)- and (2S)-hesperidin epimers in an approximate molar ratio of 1/6 was detected in orange juice (Si-Ahmed et al., 2010), possibly because of the propensity of flavanone glycosides to undergo epimerization at C₂ via the chalcone form. In the CHI-catalyzed cyclization, the enone moiety of 2',4,4',6'-tetrahydroxychalcone is locked in an *s-trans* conformation. Electrophilic activation of the C=O group involves a water molecule bound to the Tyr106 phenolic OH group and to the OH group of the Thr48 side chain. The main hydrogen bonds established between CHI and the flavanone product are represented on Fig. 4 (Jez et al., 2002b).

The concentrations of biosynthesis enzymes may play an important role in defining the distribution of flavanones in the different parts of the fruit. For instance, in the peel of *Solanum lycopersicum* (a tomato variety), the level of gene expression is lower for CHI than for CHS and flavanone 3-hydroxylase, which results in a high accumulation of naringenin chalcone (Iijima et al., 2008). This biosynthetic pathway is extensively investigated to outline the role of flavonoids in different physiological functions of plants such as insect-plant interactions (Simmonds, 2001), pigmentation (Mato et al., 2000), heavy metal tolerance (Keilig and Ludwig-Müller,

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. Journal of Food Composition and Analysis, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

JFCA-D-13-00747




Fig. 1. Chemical diversity of flavanones in nature.

2009), disease resistance and UV-screening (Cooper-driver and Bhattacharya, 1998). Recently, Fowler and Koffas (2009) have reviewed the biotechnological production of flavanones by using various microorganisms. Alternatively, some works have tried to reduce the levels of flavanones because of their bitter taste. For example, an *Agrobacterium*-mediated approach targeting the CHS and CHI genes has been used to reduce the naringin contents in grapefruit (Koca et al., 2009).

4. Diversity and distribution of flavanones

Flavanones are widely distributed in about 42 higher plant families especially in *Compositae*, *Leguminosae* and *Rutaceae* (Iwashina, 2000). Depending on the plant type, flavanones can be found in all plant parts, above and below ground, from

vegetative part to generative organs: stem, branches, bark, flowers, leaves, roots, rhizomes, seeds, fruits, peels etc... The highest concentrations of flavanones are found in peel as compared to the fleshy part of *Citrus* fruit (Nogata et al., 2006). Among flavanones, the naringenin and hesperetin aglycones and their glycosides are of particular interest because of their high prevalence in foods.

4.1. Naringenin

Naringenin (5,7,4'-trihydroxyflavanone) is found in high concentrations in *Citrus* fruit while low concentrations are also found in tomatoes and their products (Erlund, 2004). Naringenin can be found as aglycone and/or as glycosides. Among the latter, naringin and narirutin are especially abundant. Naringin (naringenin-7-neohesperidoside, Fig. 3) has a bitter taste due to its glucose moiety.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004




Fig. 2. Biosynthesis of flavonoids. R is generally OH or OMe, although other substitutions can be occurred at these positions. CHS, chalcone synthase; CHI, chalcone isomerase; FHT, flavanone 3-hydroxylase; DFR, dihydroflavonol 4'-reductase; LAR, leucoanthocyanidine 4-reductase; ANS, anthocyanidin synthase; FSI, flavone synthase; FLS, flavonol synthase; IFS, 2-hydroxyisoflavone synthase.

It is the major flavonoid in grapefruit and sour orange (Iguar et al., 2013), which present different naringin contents depending on their varieties (Table 1). Other *Citrus* species like sweet orange, tangelo, lemon and lime exhibit low quantities of naringin. Another major naringenin glycoside, narirutin (naringenin-7-rutinoside, Fig. 3) is most abundant in grapefruit although less than naringin. Significant levels of narirutin are also detected in tangor, sweet orange, tangerine and tangelo (Peterson et al., 2006a). The processing of these fruits may significantly affect the naringenin content due to release of bound flavanones and/or degradation of these heat-sensitive compounds. The effect can be clearly observed by comparing the naringenin contents of grapefruit and its jam (see Table 1) (Iguar et al., 2013). The naringenin chalcone is found in higher quantities in tomato peel, which also contains some other flavanone chalcones (Slimestad et al., 2008).

4.2. Hesperetin

As naringenin, hesperetin (4'-methoxy-5,7,3'-trihydroxyflavone) and its glycosides are also mainly present in *Citrus* fruit. The aglycone is less dominant in nature than the glycosides. The most widely distributed glycosides of hesperetin are hesperidin and

neohesperidin, which are conjugates with rutinose and neohesperidose, respectively. Hesperidin (hesperetin-7-rutinoside, Fig. 3) is present in higher extents in lemons, limes, sweet oranges, tangerine and tangor species of citrus fruits (Cano et al., 2008), while neohesperidin (hesperetin-7-neohesperidose, Fig. 3) is absent in them. Among all, sweet oranges are greatly studied for their phenolic contents and Table 1 presents the data for fruits from different regions (Cyprus, Mauritius). Although variations in the hesperetin content are detected, the ratio of the hesperetin to naringenin contents is approximately constant (Goulas and Manganaris, 2012; Ramful et al., 2011). Significant amounts of hesperetin also occur in grapefruits while tangelo and sour orange are especially rich in neohesperidin (Peterson et al., 2006a, 2006b).

5. Extraction and analysis of flavanones

Owing to the chemical complexity and the vast distribution of flavanones in plant, extraction and analysis of flavanones remain as challenging as ever, despite the recent advances in analytical instrumentation. After the detailed analysis of phenolic composition of citrus fruit and juice (edible parts), nowadays, more attention is focused on their by-products. In fact, the domestic and

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004




Fig. 3. Some common flavanone aglycones and their respective glycosides.

industrial use of large quantities of *Citrus* fruit, especially for the production of juice, results in the accumulation of huge amounts of by-products such as peels, seeds, cell and membrane residues, which account for about half of the fruit weight. These by-products can be used for the production of molasses, pectins, essential oils, limonene and cattle feed (Bocco et al., 1998; Jeong et al., 2004; Li et al., 2006a, 2006b). In addition, *Citrus* by-products are a good source of phenolic compounds, especially the characteristic flavanone glycosides, mainly naringin, hesperidin, narirutin, and neohesperidin. Currently, extraction of phenolic compounds from *Citrus* peels has attracted considerable scientific interest with aim to use them as natural antioxidants mainly in foods to prevent the oxidative alteration of lipids (Anagnostopoulou et al., 2006; Peschel et al., 2006; Zia-ur-Rehman, 2006). Indeed, cheap plant extracts rich in antioxidants could replace synthetic additives such

as butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT), which might be toxic (Moure et al., 2001), in particular damaging to the liver and carcinogenic (Ak and Gülçin, 2008).

Up to now, several conventional and innovative extraction techniques have been reported for the extraction of phenols from *Citrus* by-products (Table 2). The most common method used is maceration in solvents (Anagnostopoulou et al., 2006; Jeong et al., 2004; Li et al., 2006a; Manthey and Grohmann, 1996; Xu et al., 2007; Zia-ur-Rehman, 2006), which is also considered suitable at scaled-up level (Peschel et al., 2006). However, this extraction technique generally requires high temperatures (50–150 °C), long extraction times (up to several hours) and high polarity solvents like water and ethanol. Different types of solvent extraction techniques have been developed: hot water extraction (Xu et al., 2008), alkaline extraction (Bocco et al., 1998; Curto et al., 1992), extraction assisted by resins (Calvarano et al., 1996; Kim et al., 2007a), enzymes (Li et al., 2006b), electron beam and γ -irradiation (Kim et al., 2008; Oufedjikh et al., 2000). However, common drawbacks remain the degradation of the targeted compounds due to high temperature and long extraction periods and, more specifically, restrictive safety criteria (extraction under irradiation) and problems of enzyme denaturation. Supercritical fluid extraction (SFE) is a mild and efficient technique that has been little studied for the extraction of the phenolic compounds from *Citrus* by-products (Giannuzzo et al., 2003; Yu et al., 2007), possibly because of its higher costs in comparison to other innovative techniques.

With the increasing energy prices and environmental concern, chemical and food industries are challenged to find new technologies in order to reduce energy consumption, meet legal requirements on CO₂ emissions, product/process safety and control, and increase quality as well as functionality. Separation technology (such as extraction, distillation, and crystallization) is one of the promising fields that could contribute to sustainable growth of chemical and food industries. For example, existing extraction technologies have considerable technological and scientific bottlenecks to overcome: often requiring up to 50% of




Fig. 4. The hydrogen bond network in the (S)-4',7-dihydroxyflavanone-chalcone isomerase complex.

Adapted from Jez et al. (2002b)

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. Journal of Food Composition and Analysis, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

Table 1
 Flavanone glycosides in different plant varieties and their products.

No.	Plant type	Flavanone contents (mg aglycone/100g juice or edible fruit (without rind, pith and seeds))								References
		Narirutin	Naringin	Hesperidin	Neohesperidin	Eriocitrin	Neoeriocitrin	Didymin	Poncirin	
1	Black Sorghums (<i>Sorghum bicolor</i>) grains ^c	53 (Aglycone)	-	-	-	66 (Aglycone)	-	-	-	Dykes et al. (2013)
2	Clementine (<i>Citrus clementina</i>) juice ^b	46.4	0.8	399	-	-	-	-	-	Dhuique-Mayer et al. (2005) and Gattuso et al. (2007)
3	Defatted olive	-	-	8.01%	-	-	-	-	-	Alu'datt et al. (2013)
4	Defatted soybean	-	-	17.21%	-	-	-	-	-	Alu'datt et al. (2013)
5	Grapefruit (<i>C. paradisi</i>)	29.4	84	2.4	2.92	-	-	1.42	1.92	Peterson et al. (2006b) and Iguar et al. (2013)
6	Grapefruit (<i>C. paradisi</i>) juice ^b	76	230	9.3	12.1	4.1	3.2	3.0	12.6	Gattuso et al. (2007)
7	Grapefruit (<i>C. paradisi</i>) jam	20	84	1.72	2.64	-	-	1.16	0.6	Iguar et al. (2013)
8	Grapefruit red and pink	3.34	13.87	0.27	0.42	-	-	-	-	Peterson et al. (2006b)
9	Grapefruit white	5.36	16.90	3.95	0.25	0.16	0.05	0.09	0.20	Peterson et al. (2006b)
10	Lemon (<i>C. limon</i>)	0.80	0.18	15.78	-	9.46	-	0.17	-	Peterson et al. (2006b) and González-Molina ¹ et al. (2010)
11	Lemon (<i>C. limon</i>) juice ^b	3.8	-	205	14.5	167	-	-	-	Gattuso et al. (2007)
12	Lemon (<i>Citrus limetta</i> Risso) juice Mediterranean ^b	-	-	4.29	-	2.10	-	-	-	Barreca et al. (2011)
13	Lime (<i>Citrus latifolia</i>)	0.23	-	15.64	-	1.38	0.04	-	-	Peterson et al. (2006b)
14	Lime juice (<i>Citrus latifolia</i>) Mexican	-	-	386	-	-	-	44.14	-	Patil et al. (2009)
15	Mandarin (<i>C. reticulata</i>)	2.70	-	19.26	-	0.02	-	1.11	-	Peterson et al. (2006a)
16	Mandarin (<i>Citrus reticulata</i>) juice ^b	39.2	-	243	-	3.1	0.5	14.4	-	Dhuique-Mayer et al. (2005) and Gattuso et al. (2007)
17	Mandarin (<i>C. reticulata</i>) young ^f	43.9	-	261	-	-	-	-	-	Ye et al. (2011)
18	Shiikuwasha (<i>C. depressa</i>) peel ^a	39.18	11.88	467.31	5.16	-	-	-	-	Asikin et al. (2012)
19	Sour orange (<i>C. aurantium</i>)	0.08	18.83	0.00	11.09	0.53	14.01	2.89	-	Peterson et al. (2006a)
20	Sour orange (<i>C. aurantium</i>) juice ^b	-	19.7	-	8.7	-	7.7	-	7.3	Gattuso et al. (2007)
21	Sweet orange (<i>C. sinensis</i> Valencia) Cyprus ^d	787	13	6024	-	-	-	-	-	Goulas and Manganaris (2012)
22	Sweet Orange (<i>Citrus sinensis</i>) Mauritian ^e	16.77	-	19.93	-	-	1.07	2.70	0.12	Ramful et al. (2011)
23	Sweet orange (<i>C. sinensis</i> Valencia)	2.33	0.17	15.25	-	0.28	0.04	0.45	-	Peterson et al. (2006a)
24	Sweet orange (<i>C. sinensis</i> Valencia) juice ^b	51.4	-	257	-	-	-	18.9	3.1	Dhuique-Mayer et al. (2005) and Gattuso et al. (2007)
25	Sweet Orange (<i>Citrus sinensis</i> L. var. Navel late) fermented juice ^b	413	-	311	-	-	-	60.9	-	Escudero-López et al. (2013)
26	Tangelo	2.42	5.60	4.21	13.56	1.69	1.11	0.60	-	Peterson et al. (2006a)
27	Tangor	7.10	-	15.42	-	1.01	1.77	-	-	Peterson et al. (2006a)
28	Thymus Pulegioides ^c	24.64 (Aglycone)	-	-	-	3.01 (Aglycone)	-	-	-	Boros et al. (2010)
29	Tomato (<i>Lycopersicon esculentum</i>)	18.18 (Aglycone)	-	-	-	-	-	-	-	Slimestad et al. (2008) and Harnly et al. (2006)

^a mg/100 g fresh flavedo weight.

^b mg/L.

^c µg/g.

^d µg/g dry matter.

^e mg/g fresh weight.

^f mg/g dry weight.

investments in a new plant and more than 70% of total process energy used in food, chemical and pharmaceutical industries. These shortcomings have led to investigating new "green" extraction techniques, aimed at sparing energy and reducing costs, such as microwave- or ultrasound-assisted extraction, ultrafiltration, flash distillation and controlled pressure drop processing (Chemat et al., 2009). Among them, ultrasound-assisted extraction (UAE) has been studied in details for the preparation of flavanone-rich extracts from *Citrus* by-products.

With the development of the "Green Chemistry" concept during the past few years, environment-friendly techniques are becoming

more and more attractive. The extraction of bioactive compounds under ultrasound irradiation (20–100 kHz) is one of the upcoming extraction techniques that can offer high reproducibility in shorter times, simplified manipulation, reduced solvent consumption and temperature and lower energy input (Chemat et al., 2008). During sonication, the cavitation process causes the swelling of cells or the breakdown of cell walls, thereby permitting high diffusion rates across the cell wall in the first case or a simple washing-out of the cell contents in the second (Vinatoru, 2001). It is known that the bursting of tiny cavitation bubbles may result in local temperature and pressure as high as 5000 °C and 1000 atm, respectively. However,

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

Table 2
 Extraction and analysis of flavanone contents from various plant sources.

Plant part	Extraction	Flavanones studied	Analytical technique	Comments	References
Sweet orange (<i>Citrus sinensis</i>) peel	Soxhlet extraction	Hesperidin, TPC	2D-TLC (80:20:40, ethyl acetate:acetic acid:water and 15% acetic acid); Folin-Ciocalteu test	Methanolic fraction with ethyl acetate showed significant TPC and RSA	Anagnostopoulou et al. (2006)
Satsuma mandarin (<i>C. unshiu</i> Marc.) peel	Ultrasound-assisted extraction (UAE)	Narirutin, hesperidin	HPLC (RP C18 column (250 mm × 4.6 mm, 5 μm), 4% acetic acid/100% methanol (80:20, v/v))	UAE can significantly enhance the contents of phenolic compounds and antioxidant activities in samples without any intensive heat treatment	Ma et al. (2008b)
Clementines (<i>C. clementina</i>)	Solvent extraction of γ-Irradiated fruits	Naringin, hesperidin, didymin, eriocitrin, poncirin, and neoeriocitrine	HPLC (RP C18 column (150 mm × 4.5 mm, 5 μm), 4% acetic acid/100% acetonitrile)	γ-Irradiation stimulates PAL activity and the synthesis of phenolic compounds. Thus, higher amounts of flavanones are obtained	Oufedjikh et al. (2000)
Grapefruit (<i>C. paradisi</i> Macf.) seeds	Supercritical fluid extraction	Naringin	HPLC (Spherisorb ODS column (250 mm × 4.6 mm))	After experimental design implementation, crucial factor during extraction of naringin was pressure followed by temperature	Yu et al. (2007)
Citrus juices of sweet orange, tangerine, lemon and grapefruit	Solvent extraction of freeze-dried samples	Eriocitrin, narirutin, naringin, hesperidin, neohesperidin, and didymin	RP-HPLC, MS (one for the MS ¹ full scan mode and another for MS ² ion production scan mode)	Most important contribution of this work is the characterization of flavanones using HPLC-DAD-ESI-CID-MS/MS	Abad-Garcia et al. (2012)
Orange (<i>C. sinensis</i>) peel	Instant controlled pressure drop (ICPD) technology and UAE	Naringin, hesperidin	HPLC (RP C18 column (250 mm × 4 mm, 5 μm) 0.5% acetic acid and 100% acetonitrile)	Peel by-products, successively treated by ICPD (extraction of essential oil) and UAE gave extracts richer in flavanones	Allaf et al. (2013)
Tomato fruit	Solvent extraction	Naringenin, naringin	HPLC (Chromolith Performance RP-18e column (100 mm × 64.6 mm), 50 mM phosphate buffer at pH 2.2/acetoneitrile (75:25, v/v))	Faster analysis owing to shorter retention time of prominent flavonoids	Biesaga et al. (2009)
Spearmint (<i>Mentha spicata</i> L.) leaves	Supercritical carbon dioxide (SC-CO ₂) extraction	Naringenin	HPLC (RP C18 (25 cm × 4.6 mm, 5 μm), TFA 2.5 pH in deionized water and 100% methanol)	Higher amount of naringenin was obtained by SC-CO ₂ (60 °C, 200 bars, 60 min)	Bimakr et al. (2011)
Rio red grapefruits	Solvent extraction	Narirutin, naringin, neohesperidin, didymin and poncirin	HPLC, LC-MS (ESI-, capillary (250 °C, -15 V), N ₂)	Optimized parameters include extraction technique (MW vs ultrasounds), solvent, centrifugal speed, temperature, sample to solvent ratio, extraction cycles, extraction time and their interactions	Chebrolua et al. (2011)

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

Table 2 (Continued)

Plant part	Extraction	Flavanones studied	Analytical technique	Comments	References
<i>C. unshiu</i> peel	Subcritical water extraction	Hesperidin, narirutin	HPLC, LC-MS (ESI+, capillary (350 °C, 5.5 kV), N2)	This environment friendly technique gave higher amount of flavanones in comparison to conventional one	Cheigh et al. (2012)
<i>C. reticulata</i> peel	Pressurised liquid extraction (PLE)	Hesperidin	LC-DAD-ESI/MS (ESI+, capillary (350 °C, 4 kV))	PLE compared with UAE, Soxhlet and heat-reflux extractions	Li et al. (2012)
Citrus juices and beverages	Liquid-liquid extraction	Eriodictyol, naringenin, hesperetin, eriocitrin, narirutin, hesperidin, neoeriodictin, naringin, neohesperidin, didymin, poncirin	UHPLC-ESI-MS/MS (RP C18 column (2.1 mm × 50 mm, 2.6 μm), H ₂ O (0.1% HCOOH) and CH ₃ CN, 0.3 mL/min, Heated ESI+ spray voltage (4.7 kV, 270 °C))	A comprehensive and throughput assay of the flavanones in citrus juices using UPLC	Di Donna et al. (2013)
Cherry tomatoes, tomato sauce, and tomato juice	Liquid-liquid extraction	Naringenin	UHPLC-QqQ-MS (BEH C18 column (50 mm × 2.1 mm, 1.7 μm), 0.1% formic acid and acetonitrile, 0.4 mL/min, ESI-, capillary (-3.5 kV, 400 °C), N2)	An easy, fast, and sensitive UPLC-QqQ-MS method was described to identify and quantify the most abundant phenolic compounds.	Di Lecce et al. (2013)
Citrus mandarin pomace	Microwave-assisted extraction	Naringenin, naringin, hesperidin	HPLC (RP C18 column (4.6 mm × 150 mm, 5 μm), 0.1% formic acid, and 100% methanol)	Microwave treatment as an efficient process to release bound phenolic compounds from the plant matrix	Hayat et al. (2010)

Abbreviations: TPC=total phenolic content; RSA=radical scavenging activity; 2D-TLC=2 dimensional thin layer chromatography; PAL=phenylalanine ammonia-lyase; HPLC=high performance liquid chromatography; UPLC=ultra high performance liquid chromatography; MS=mass spectrometry; ESI=electron spray ionization; TFA=trifluoroacetic acid.

these local energy bursts cannot significantly affect the bulk conditions as they are dissipated to the medium in very short periods of time (Luque-Garcia and Luque de Castro, 2003). UAE actually depends on the destructive effects of ultrasonic waves on solid matrices including plant materials. In addition to optimizing the solvent and temperature conditions, a better recovery of cell contents can be obtained by optimizing ultrasound frequency, sonication power and period, as well as the ultrasonic wave distribution (Wang and Weller, 2006). UAE has been applied recently to the extraction of hesperidin from Penggan (*Citrus reticulata*) peel (Ma et al., 2008a), phenolic acids and flavanone glycosides from Satsuma Mandarin (*Citrus unshiu* Marc) peel (Ma et al., 2008b, 2009) and total phenolic contents from Penggan peel (Ma et al., 2008c). In these works, methanol came up as a suitable extraction solvent to reach good yields. However, environmentally benign and non-toxic food grade organic solvents like ethanol, n-butanol and isopropanol are recommended by the US Food and Drug Administration for extraction purposes (Bartnick et al., 2006). Using ethanol, UAE was found more efficient for the extraction of polyphenols from orange peel wastes than conventional solvent extraction (Khan et al., 2010). Moreover, for the extraction of natural products, UAE gave higher yields than conventional techniques, not only at the lab-scale but also at the pilot-plant scale (Boonkird et al., 2008).

Recently, extraction of phenols from *Citrus* peels was successfully carried out under microwave irradiation (Hayat et al., 2009, 2010a, 2010b). Moreover, extraction of onion flavonoids without any added solvent has been developed using the technique of microwave hydrodiffusion and gravity (MHG) (Zill-e-Huma et al., 2009). In the process, extraction is initiated by the selective heating of the plant water content. MHG could also be used for the efficient extraction of flavanones from plant sources.

Natural extracts from *Citrus* by-products are rich in phenols that display *in vitro* antioxidant, antibacterial, antimicrobial, anti-inflammatory and anticarcinogenic properties. As such, they could have a range of applications in the food, pharmaceutical and cosmetic industries. They could also appeal consumers due to their status of ingredients derived from natural sources without any chemical processing. One of the limiting factors in the market of natural extracts remains their significantly higher prices in comparison to synthetic additives. Natural extracts from food industry by-products could overcome this drawback due to the cheap starting material used. In addition, several studies have shown that natural extracts may exhibit higher biological activity than purified compounds, due to positive synergistic interactions between bioactive components (Gimenez-Bastida et al., 2009; Menichini et al., 2009; Sood et al., 2009, 2010).

For the determination of flavanone contents, spectrometric methods, which are fast and simple, are still used. However, they lack specificity for individual compounds. Generally, the ultraviolet spectra of flavanones and their glycosides show two strong absorption bands commonly referred to as band A (330 nm) and band B (280 nm). Band A is associated with the B-ring in conjugation with the keto group while band B is the typical absorption band of phenols and involves a strong contribution of A-ring. Substitutions on the A- or B-ring may produce hypsochromic or bathochromic shifts, which could be used for structural elucidation (Tsimogiannis et al., 2007).

The most efficient analytical techniques for flavanone identification and titration in extracts (see Table 2) are high or ultra high performance liquid chromatography (HPLC, UPLC) coupled with a diode array detector (DAD) and a mass spectrometer (Allaf et al., 2013; Bimkr et al., 2011; Cheigh et al., 2012). The development of

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004




Fig. 5. Organic synthesis of flavanones.

UPLC has significantly enhanced the performance of separation by providing greater efficiency and drastically reducing analytical time (Di Donna et al., 2013; Di Lecce et al., 2013).

6. Chemical synthesis of flavanones

6.1. Aglycones

Up to now, the most common pathway for the synthesis of flavanone aglycones is the aldol condensation of 2-hydroxyacetophenones with benzaldehydes (Claisen–Schmidt aldol condensation) (Fig. 5). The reaction is usually performed under heating using acidic or alkaline conditions. The chalcones initially formed undergo cyclization to their respective flavanones under the same conditions (French et al., 2010; Krbeček et al., 1968; Shi et al., 2010). The condensation is still under study to develop efficient and environment-friendly conditions. For instance, strongly alkaline sodium hydroxide and ethoxide were replaced by Mg–Al hydroxalates (Climent et al., 1995). Furthermore, different derivatives of chalcones and flavanones were also prepared by aldol condensation (Hsieh et al., 1998). Currently, the emphasis is on developing new catalysts that could be effective in aldol condensations and alternative methods (Chandrasekhar et al., 2005). Recently, the introduction of Li was shown to increase the surface basicity and catalytic activity of MgO in the synthesis of flavanone aglycones (Cortes-Concepcion et al., 2010).

6.2. Chalcones

The scarcity of flavanone chalcones in Nature is primarily due to their instability, as in neutral medium, 2'-hydroxychalcones undergo cyclization to the corresponding flavanones (intramolecular Michael addition). However, chalcones can be simply prepared by the reverse reaction (opening of the C-ring of flavanones) in strongly alkaline conditions. In such conditions, flavanone phenolate anions are first formed. Then, removal of a proton at C2 yields an enolate anion, from which the opening of the C-ring can take place with simultaneous formation of chalcone ions (Andújar et al., 2003). Fast acidification leads to the precipitation of the neutral chalcone, which can then be isolated by simple filtration. By contrast, in weakly basic conditions, chalcone phenolate anions undergo cyclization to flavanones through an enolate intermediate as described for naringin (González et al., 2002) and 2',6'-dihydroxy-4,4'-dimethoxychalcone (Miles and Main, 1985).

6.3. Glycosides

The most prevalent flavanone derivatives in nature are 7-O- β -glycosides. The selective glycosylation of the most acidic C7-OH group can be performed by protected glycosylbromide activated by silver carbonate (Oyama and Kondo, 2004). With some

modifications, this method is still in use not only for glycosylation of phenolic compounds (Esaki et al., 1994) but also for glucuronidation (Moon et al., 2001). A simple route to flavanone 7-O- β -D-glucoside is the partial hydrolysis of naringin and hesperidin using formic acid in cyclohexanol. However, the yields are low (ca. 10%) (Fox et al., 1953). The enzymatic synthesis of flavanone glycosides was also described (Kometani et al., 1996) as well as the synthesis of naringin analogues bearing amino groups for use as scaffolds in drug discovery (Hanessian and Kothakonda, 2005) and metal–naringin complexes to increase the antioxidant and anti-inflammatory activities (Pereira et al., 2007).

6.4. Glucuronides

A better knowledge of the biochemical mechanisms by which dietary flavanones exert their potential health effects requires investigations on appropriate cell models (e.g., endothelial or smooth muscle cells) with the authentic circulating metabolites (of which glucuronides make the largest contribution) instead of the commercially available glycosides and aglycones that are frequently used as a first approach despite the limited biological significance. As an alternative to the expensive, inconvenient and low yielding extraction of conjugates from biological fluids, chemical synthesis appears as the most direct strategy to obtain substantial amounts of polyphenol metabolites for bioavailability and *in vitro* cell studies. In particular, there is a growing interest for the synthesis of polyphenol glucuronides as standards for identification and titration of *in vivo* metabolites and as biologically pertinent compounds for cell studies aiming at elucidating the potential health effects of polyphenols. Several works have been published about the chemical synthesis of polyphenol glucuronides.

For instance, the popular procedure, based on the Lewis acid-activated coupling of methyl-2,3,4-tri-O-acetyl-1-O-(trichloroacetimidoyl)- α -D-glucuronate (Tomas-Barberan and Clifford, 2000) with partially protected polyphenols, was applied to the synthesis of isoflavone 7-O- β -D-glucuronides (Al-Maharik and Botting, 2006), quercetin 3-O- β -D-glucuronide (Needs and Kroon, 2006) and hydroxycinnamic acid O- β -D-glucuronides (Galland et al., 2008; Fumeaux et al., 2010).

Catechin O- β -D-glucuronides were also prepared with methyl-2,3,4-tri-O-acetyl-1-O-bromo- α -D-glucuronate as the glucuronyl donor (González-Manzano et al., 2009). Recently, the synthesis of a flavanone glucuronide (persicogenin 3'-O- β -D-glucuronide) was carried out with methyl-2,3,4-tri-O-acetyl-1-O-(trifluoroacetimidoyl)- α -D-glucuronate, followed by a final deprotection step involving pig liver esterase (PLE) for the hydrolysis of the methyl ester of the glucuronyl residue (Boumendjel et al., 2009). A synthesis of quercetin 3-O- β -D-glucuronide was also performed by regioselective oxidation of the corresponding 3-O- β -D-glucoside (phenolic OH groups protected as benzyl ethers) using TEMPO/NaOCl/NaBr under phase transfer conditions (Bouktaib et al., 2002). Recently, the synthesis of four flavanone glucuronides based on a regioselective protection of the flavanone nucleus was reported by our research group (Khan et al., 2010). In this work, glucuronides of naringenin (4'- and 7-O- β -D-glucuronides) and hesperetin (3'- and 7-O- β -D-glucuronides), the major flavanone aglycones in grapefruit and orange respectively, were chemically synthesized (Fig. 6A and B). On the one hand, the most reactive hydroxyl group 7-OH was protected by selective benzoylation to allow subsequent glucuronidation of 4'-OH (naringenin) or 3'-OH (hesperetin) (B-ring). On the other hand, the selective debenzoylation at 7-OH of the perbenzoylated flavanone aglycones allowed glucuronidation at the same position (A-ring). After careful deprotection, the target compounds were purified and characterized by nuclear magnetic resonance and mass spectrometry. So far, the synthesis of other less common flavanone

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. Journal of Food Composition and Analysis, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004




Fig. 6. (A) Glucuronidation of citrus flavanones on the B-ring. (i) BzCl (1.1–1.5 equiv.), Pyridin (ii) (1), BF₃-OEt₂ (2 equiv.), CH₂Cl₂ (iii) Na₂CO₃ (1.2 equiv. per ester group), H₂O/MeOH (2/5), then acidification to pH 6 by Dowex resin (H⁺ form). (B) Glucuronidation of citrus flavanones on the A-ring. (i) BzCl (excess), NEt₃, THF, 0 °C (ii) PhSH (1 equiv.), imidazole, NMP (iii) (1), BF₃-OEt₂ (2 equiv.), CH₂Cl₂ (iv) Na₂CO₃ (1.2 equiv. per ester group), H₂O/MeOH (2/5), then acidification to pH 6 by Dowex resin (H⁺ form).

metabolites (diglucuronides, sulfoglucuronides, sulfates) has not been reported.

7. Bioavailability of flavanones

The oral bioavailability of a given nutrient describes its fate once ingested: release from food matrix (bioaccessibility, typically assessed *in vitro*, Bouayed et al., 2012), intestinal absorption, metabolism, transport in the general circulation, delivery to tissues and excretion. In spite of the high consumption of citrus fruits and juices worldwide, the bioavailability of flavanones is still incompletely known. Their daily intake has not been estimated in different populations but could be quite high compared with the average flavonol intake (25 mg/day) in several European countries (Manach et al., 2003). For instance, the mean dietary intake in Finland has been evaluated to be 8.3 mg/day and 28.3 mg/day for naringenin and hesperetin, respectively (Erlund, 2004; Manach et al., 2003).

After oral intake, flavanone glucosides and other glycosides are hydrolysed in the small intestine and in the colon, respectively, and the released aglycones are converted into their respective glucuronides, sulfates and sulfoglucuronides during their passage across the small intestine and liver. Finally, the bioactive forms (metabolites) are distributed through plasma at various cell sites and significant quantities can also be found in urinary excretions (Matsumoto et al., 2004). The fate of flavanones after ingestion is summarized in Fig. 7.

7.1. Intestinal absorption

Intestinal absorption involves the uptake from the intestinal lumen of nutrients into the intestinal epithelial cells, blood, lymph, or interstitial fluids. The study of pharmacokinetics in humans and animals can provide knowledge about the rate of absorption across the intestine. For instance, naringenin and its glucuronides were detected in the plasma and brain of rats (conjugation ratio four-time

higher in plasma than in brain) only 10 min after naringenin administration (20 mg/kg) (Peng et al., 1998). The study was extended to determine the naringenin levels in rat brain, liver and bile using microdialysis coupled with a HPLC system. The results have shown a higher concentration of naringenin in liver and bile (Tsai, 2002). To the best of our knowledge, the first report on the pharmacokinetics of flavanones in human subjects was published by Erlund and co-authors in 2001. After ingestion of orange or grapefruit juice (8 mL/kg of body weight), the plasma concentration of hesperetin and naringenin aglycones was found in the range 0.6–6.0 μmol/L, with a peak plasma concentration (C_{max}) of 6.0 ± 5.4 μmol/L for naringenin from grapefruit juice and 2.2 ± 1.6 μmol/L for hesperetin from orange juice. Moreover, elimination half-lives (t_{1/2}) in the range of 1.3–2.2 h point to a relatively fast clearance from the general circulation. The percentage of flavanones excreted in urine (5–30% of total amount ingested) was lower than that of their absorption, which suggests a substantial distribution to tissues for these phenolic compounds (Erlund et al., 2001). In another study involving six volunteers, ingestion of hesperetin and naringenin (135 mg of each) under fasting conditions resulted in their appearance as metabolites in blood plasma 20 min later. The C_{max} of 2.7 μmol/L for hesperetin and 7.4 μmol/L for naringenin was reached 4.0 and 3.5 h after ingestion, respectively (Kanaze et al., 2007).

Intestinal absorption varies according to glycoside concentration and flavanone structure. After ingestion of 1 L of orange juice containing 444 mg of hesperidin and 96.4 mg of narirutin, blood analysis over a 24 h led to C_{max} values for hesperetin and naringenin (after deconjugation) at 1.28 ± 0.13 μmol/L and 0.20 ± 0.04 μmol/L, respectively. The levels of flavanones in urine were expressed as percentage of their intake and amounted to 7.87 ± 1.69% for naringenin and 6.41 ± 1.32% for hesperetin. The relative urinary excretion of flavanones was not significantly affected by the dose ingested (Manach et al., 2003). In another study, high naringenin concentrations of 128 ± 2 μM, 144 ± 8 μM and 139 ± 15 μM were detected after 10 h in the plasma of rats fed

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004




Fig. 7. Metabolic fate of flavanones.

with naringenin (0.25% of total diet), naringenin-7-*O*- β -glucoside (0.38%) and naringenin-7-*O*- β -rhamnoglucoside (0.50%), respectively. The urinary excretion of naringenin was two times higher in rats fed with naringenin than in the ones fed with naringenin-7-rhamnoglucoside (Felgines et al., 2000). Plasma and urine analyses pointed to naringenin being more bioavailable than hesperetin (Gardana et al., 2007; Kanaze et al., 2007). An *in vitro* hydrolysis showed a faster hydrolysis rate for hesperidin and narirutin (flavanone rutosides) than for naringin and neohesperidin (flavanone neohesperosides) (Wang et al., 2008). More recently, the same group studied the bioavailability of hesperetin and naringenin after the consumption of *Citrus aurantium* L. and *Citrus sinensis* Osbeck. These citrus varieties are used in the formulation of Zhi Zhu Wan, a traditional Chinese medicine used for the treatment of functional dyspepsia. The T_{max} (time to reach C_{max}) of hesperetin was found at 3.7 and 8.5 h after the oral administration of *C. aurantium* and *C. sinensis*, respectively, which probably reflects differences in the structure and environment of the hesperetin glycosides depending on the source (Cao et al., 2010).

The permeability of epithelial cells to flavanones is also a good determinant of their intestinal absorption. Flavanones are transported from apical side (gut lumen) to basolateral side (blood). In *in vitro* models, hesperetin (aglycone) was found to be efficiently absorbed across Caco-2 cell monolayers in comparison to hesperidin (hesperetin glycoside). The absorption mechanisms involved transcellular passive diffusion along with a newly proposed mechanism of proton-coupled active transport (Kobayashi et al., 2008). The study was further elaborated to explain the H^+ -driven polarized absorption and similar mechanisms were found for naringenin and eriodictyol aglycones (Kobayashi and Konishi, 2008). The faster absorption of flavanone aglycones compared to flavanone glycosides was confirmed for eriodictyol and eriocitrin in humans (Miyake et al., 2006). The transportation of flavonoids across the cell membranes involves ATP-binding cassette (ABC) transporters, which are present in the apical (lumen side) or basolateral (blood circulation side) membrane of enterocytes and facilitate excretion back into the intestinal lumen or uptake into the blood, respectively. Intestinal ABC transporters that have been related to flavonoid

transport include P-glycoprotein (Pgp/MDR1/ABCB1), multidrug resistance proteins (MRPs/ABCCs), and breast cancer resistance protein (BCRP/ABCG2), of which Pgp, MRP2, and BCRP are localized in the apical membrane. A study on hesperetin metabolism in Caco-2 monolayers has shown that hesperetin 7-*O*- β -D-glucuronide and 7-*O*-sulfate are predominantly transported to the apical side. By contrast, hesperetin aglycone also permeates to the basolateral side of the Caco-2 cell monolayer. The pattern of inhibition by different ABC transporter inhibitors suggests that the apical efflux of hesperetin metabolites mainly involves BCRP. Altogether, these findings elucidate a novel pathway of hesperetin metabolism and transport and show that BCRP-mediated transport could be one of the main limiting steps for hesperetin bioavailability (Brand et al., 2008). The work was further elaborated to distinguish between the (2*R*)- and (2*S*)-enantiomers of flavanones. It was found that the intestinal absorption of commercially available racemic flavanones was similar to that of the naturally predominant (2*S*)-enantiomers (Brand et al., 2010).

Still the missing part in most bioavailability studies is the availability of authentic conjugates for use as standards. Those conjugates are also very much needed for investigating the mechanisms of their bioactivity in cell models.

7.2. Metabolism

A great part of the bioavailability studies has been devoted to naringenin, hesperetin and their glycosides. Improvements in methods for analyzing flavanone metabolites in human plasma and urine have made possible to estimate flavanone bioavailability in humans. The first step in flavanone metabolism is the extensive deglycosylation of flavanone glycosides within the intestinal epithelium by human and bacterial enzymes like β -glucosidases, rhamnosidases, rutosidases etc. (Fig. 8). Investigations in rats demonstrated that the deglycosylation of naringenin-7-*O*- β -D-glucoside occurred early in the small intestine (Choudhury et al., 1999) while that of naringenin-7-*O*- β -D-rhamnoglucosides occurred in the colon (large intestine). Indeed, naringenin conjugates (glucuronides and/or sulfates) appeared within 3 h in the plasma of

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004




Fig. 8. Zoomed schematic presentation of flavanone metabolism in small and large intestine.

rats fed with naringenin or its 7-O-β-D-glucoside whereas no naringenin metabolites were still detected in rats fed with naringenin-7-O-β-D-rhamnoglucoside. However, 10 h after ingestion, similar naringenin concentrations were found regardless of the diet, which clearly showed the delayed intestinal absorption of naringenin rhamnoglucosides (Felgines et al., 2000). It was confirmed in humans that hesperidin and naringin are absorbed in the distal part of the intestine (cecum). Once deglycosylated, the aglycones are glucuronated and/or sulfated during their transfer from the luminal side of the gut to the portal vein by the action of UDP-glucuronosyltransferase and sulfotransferase enzymes (Manach et al., 2003). In cecum, the intestinal microflora not only cleaves the glycosidic bonds but also degrades the aglycones into phenolic acids such as *p*-hydroxyphenylpropionic acid, *p*-coumaric acid, and *p*-hydroxybenzoic acid (Felgines et al., 2000; Manach et al., 2003). Likewise, eriodictin (eriodictyol-7-O-β-D-rutinoside) is metabolized by intestinal microflora (*Bacteroides distasomis* or *Bacteroides uniformis*) to eriodictyol, which is then converted into 3,4-dihydroxycinnamic acid by *Clostridium butyricum* (Miyake et al., 2000). After intestinal absorption, aglycones and phenolic acids reach the liver, the main organ involved in flavanone metabolism, where further glucuronidation, sulfation, and in some cases methylation occurs, thus converting the rest of aglycones and phenolic acids into their respective metabolites. Due to lack of catechol groups in hesperetin and naringenin, no methylation by catechol-O-methyltransferase was observed, which is in contrast to catechin and quercetin (Felgines et al., 2000). By contrast, the conversion of eriodictyol to homoeriodictyol and hesperetin through methylation in liver was reported (Miyake et al., 2000).

Recently, a study was conducted to determine the effect of tumor on flavanone bioavailability. Similar naringenin concentrations were estimated in the liver and kidney of healthy and tumor-bearing rats (Silberberg et al., 2006).

The impact of full-fat yogurt on the bioavailability and metabolism of orange flavanones in humans was investigated by analyzing plasma and urine over different intervals of time. Addition of yogurt into orange juice significantly reduced the quantity of flavanone metabolites excreted up to 5 h after ingestion. However, a statistical analysis over a longer time span (0–24 h) did not show any significant effect of yogurt addition (Mullen et al., 2008). Thus, it can be concluded that dairy food matrices may delay the intestinal absorption of flavanones with no impact on the overall bioavailability. Another study have shown that the quantity of flavanone metabolites excreted in urine after 24 h of orange juice intake has reduced about 7 times when juice is

taken with yoghurt (Roowi et al., 2009). The authors proposed that this reduction was at the large intestine due to the alteration of flavanone metabolism by the microflora.

Flavanones appeared resistant to oxidation while their isomerization into chalcones (38–55%) was significantly observed for the orange juice (pH 7.5). Moreover, the processing (pasteurization for 30 s at 95 °C, freezing and Brix concentration to 60°) of orange juice showed no effect on the *in vitro* bioaccessibility of flavanones (Gill-izquierdo et al., 2003). However, chalcones were found in higher quantities in industrially (vs. manually) pressed juice.

Glucuronidation and sulfation are the major conjugation pathways of flavanone aglycones. Structural studies of the plasma and urinary metabolites showed that the major metabolites of naringenin are naringenin-7-O-β-D-glucuronide, naringenin-4'-O-β-D-glucuronide, naringenin-4'-O-β-D-glucuronide-7-O-sulfate, naringenin-4'-O-sulfate-7-O-β-D-glucuronide and naringenin-4',7-O-disulfate (Tripoli et al., 2007; Brett et al., 2009). Similarly, the main hesperetin conjugates are hesperetin-7-O-β-D-glucuronide, hesperetin-3'-O-β-D-glucuronide, hesperetin diglucuronide and sulfoglucuronide (Matsumoto et al., 2004; Mullen et al., 2008). Among all these metabolites, glucuronides largely prevail (87%) but the importance of the other metabolites should not be underestimated (Manach et al., 2003). The position at which glucuronidation occurs might influence the resulting bioactivity including the antioxidant activity and affinity for specific proteins (Tripoli et al., 2007). Up to now, no data have been reported about the antioxidant activity of flavanone glucuronides. However, since the common flavanones hesperetin and naringenin are devoid of catechol group, which is the critical structural determinant of the antioxidant (reducing) activity for polyphenols, both are weak antioxidants and their glucuronides (with one less free phenolic OH group) are expected to be even less potent. It is thus quite likely that the bioactivity expressed by flavanone glucuronides is largely unrelated to their redox properties and rather reflects their interactions with specific proteins.

8. Interaction of flavanones with human serum albumin

In the past decade, studies exploring the possible health effects of flavanones were devoted to assess either their metabolism and bioavailability or their possible therapeutic value as potential drugs. Unfortunately, the delivery of the circulating flavanones and their metabolites to specific biological sites is still poorly documented. The interaction of flavonoids with human serum albumin (HSA) could be an important factor controlling their




Fig. 9. Three dimensional structure of HSA with common ligands bound at different binding sites. Adapted from Ghuman et al. (2005)

half-life in plasma and transport to biological sites. Indeed, serum albumin is the major protein of blood plasma with a concentration as high as 0.6 mM. The literature on the structural aspects and binding locations of HSA is well described by a number of comprehensive reviews (He and Carter, 1992; Sugio et al., 1999). Beside its role in the maintenance of colloidal osmotic blood pressure and bodily detoxification, HSA transports fatty acids and a large variety of drugs (Varshney et al., 2010) and food components including polyphenols (Dufour and Dangles, 2005) (Fig. 9). Interestingly, HSA has been shown to accumulate in solid tumors and in inflamed joints in arthritic disease and could thus encourage the specific delivery of drugs at those sites (Kratz, 2008).

A significant amount of the literature available on albumin-flavonoid interactions reports not only quantitative thermodynamic data (binding constants), but also qualitative analyses aimed at locating the possible binding sites (Banerjee et al., 2008; Dufour and Dangles, 2005; Lu et al., 2007; Rawel et al., 2005). In particular, quercetin and its metabolites were studied for their affinity with HSA (Murota et al., 2007; Zsila et al., 2003). In addition to other conventional techniques, fluorescence spectroscopy is the analytical tool that is most widely used to investigate binding to HSA (Oravcová et al., 1996). Indeed, the single tryptophan residue of HSA (Trp214, site I) can be excited at 295 nm and emits fluorescence at 340 nm. From the quenching of this fluorescence by a given ligand, the binding constant can be estimated (Sulkowska, 2002).

The affinity of flavanones to HSA has been more recently investigated. In an original study, piezoelectric quartz crystal impedance (PQCI) analysis was performed to measure the affinity of hesperidin for immobilized HSA. The association constant was estimated at 3.42 mg ml⁻¹ using Scatchard analysis (Liu et al., 2004). The interaction of hesperidin with bovine serum albumin (BSA) was also investigated by fluorescence spectroscopy (Wang et al., 2007). Xie and co-authors used fluorescence spectroscopy with support of Fourier-transformed infrared (FT-IR) and UV-vis spectroscopies to determine the binding constant, binding site and binding mechanism of hesperetin to HSA. From the Stern-Volmer equation, a binding constant of about 81 × 10³ M⁻¹ was estimated at pH 7.4. The *K* value decreased with increasing the pH from 6.4 to 8.4 due to,

(a) conformational changes in HSA which affect the shape of the hydrophobic binding cavities and (b) the increased dissociation of the phenolic hydroxyl groups of hesperetin. Moreover, FT-IR spectroscopy suggested that hesperetin binds to subdomain IIA (Xie et al., 2005a). The study was further extended to investigate the association of naringenin ($K = 127 \times 10^3 \text{ M}^{-1}$) with HSA (Xie et al., 2005b). Conjugation of flavanone aglycones may affect their affinity for HSA. For instance, the binding constant of naringin, a naringenin diglycoside, was found lower ($K = 18 \times 10^3 \text{ M}^{-1}$) than that of naringenin (Zhang et al., 2008). Moreover, the presence of metal ions in the medium may also reduce the affinity of flavanones for HSA (Hu et al., 2010). The main mechanisms involved in the interaction include the hydrophobic effect (van der Waals interactions between the ligand and hydrophobic amino acid residues with concomitant desolvation), electrostatic interactions between Lys residues and the flavanone phenolate ion, formed after deprotonation of the most acidic OH group at position C7, and hydrogen bonding between the phenolic OH and keto groups of hesperetin and the polypeptide chain or other polar amino acid residues (Xie et al., 2005a). Thermodynamic calculations (enthalpy change ΔH , free energy change ΔG , positive entropy change ΔS) suggest that the electrostatic interactions are more effective in flavanone – HSA binding than other forces (Hu et al., 2010). Up to recently, the affinity for HSA of true circulating flavanone metabolites had not been investigated. Hence, our group studied the interaction of chemically synthesized flavanone glucuronides (main circulating metabolites in humans) with HSA (Khan et al., 2011). Compared with naringenin and hesperetin, it was noted that glucuronidation only weakly destabilizes the flavanone-HSA complexes, the effect being slightly stronger in the case of the B-ring glucuronidation. The investigations were further extended to flavanone chalcones, which proved better HSA ligands than their flavanone counterparts.

9. Bioactivity of flavanones

Over the past few decades, extensive research has been conducted on dietary compounds that could be protective against lethal diseases, in particular cardiovascular disease and cancers.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. Journal of Food Composition and Analysis, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

These potentially bioactive compounds include phytoestrogens, carotenoids, ascorbic acid, citrus limonoids, organosulfur compounds and a good number of polyphenols. The basic mechanisms implicated in the potential health effects of polyphenols are mainly the inhibition of lipid and DNA oxidation (antioxidant activity) and the regulation of gene expression (Kris-etherton et al., 2002; Patil et al., 2009). Like other polyphenols, flavanones are also studied for their effects on specific cells. However, the missing part remains the investigation of true flavanone metabolites. Examples of *in vitro/in vivo* studies conducted to explore the beneficial effects of flavanones and the mechanisms involved are discussed below.

9.1. Radical-scavenging effects

Reactive oxygen species (ROS)/reactive nitrogen species (RNS) in biological systems are typically unstable and produced in low concentration for physiological signaling pathways and in larger concentration (oxidative stress) to destroy viruses and bacteria in leucocytes during infection (inflammatory response). The chronic exposure to oxidative stress is considered an initiating event in the development of degenerative diseases (Brown and Borutaite, 2006; Forman et al., 2008). ROS/RNS include the superoxide anion ($O_2^{\bullet-}$), hydrogen peroxide (H_2O_2), the hydroxyl radical (HO^{\bullet}), the hypochlorite ion (ClO^-), nitrogen dioxide (NO_2) and peroxytrite ($ONOO^-$), lipid oxyl and peroxy radicals (RO^{\bullet} , ROO^{\bullet}) produced during the autoxidation of polyunsaturated fatty acids. Phenolic compounds are extensively studied for their ability to reduce ROS/RNS (antioxidant activity, AA), thereby preventing the oxidative damage they cause to the host's biomolecules.

The antioxidant activity of flavanones depends upon the number and spatial arrangement of phenolic OH groups (Cai et al., 2006; Sadeghipour et al., 2005). Up to now, *in vitro* and *in vivo* investigations have been performed to determine the antioxidant potential of flavanone aglycones, chalcones, and glycosides. No literature is available about the antioxidant capacity of flavanone glucuronides.

A comparative study on the antioxidant properties of nine different flavanones (naringin, neohesperidin, neoeriocitrin, hesperidin, narirutin, naringenin, hesperetin, eriodictyol and isosakuraterin) using the crocin bleaching inhibition assay showed that the presence of a catechol nucleus (3',4'-dihydroxy substitution on the B-ring) and its *O*-methylation have no significant effect on the antioxidant activity (AA) of aglycones, which is surprising. By contrast, an increase in AA was observed with the glycosides having a catechol nucleus while *O*-methylation of the catechol has an opposite effect (Di Majo et al., 2005). *O*-Glycosylation often reduces the AA, which points to the participation in the radical-scavenging reaction of the OH group involved in the glycosidic bond (Van Acker et al., 1996). The different glycosyl moieties may also have a small effect on AA. For instance, the glycosylation of hesperetin on the C7-OH group by neohesperidose affects the AA while the glycosylation by rutinose has no effect (Di Majo et al., 2005). While substitution on a hydroxyl group typically decreases the AA, addition of a hydroxyl group can strongly increase it. For example, 3',5'-dihydroxynaringin (pyrogallol B-ring) is ca. 70 times as potent as naringenin (Ye et al., 2009). Flavanones present a higher AA in a hydrophilic environment. In a lipophilic environment, some flavanones (neohesperidin, hesperetin, isosakuraterin) showed a reduced antioxidant potential while others (naringin, narirutin, naringenin, neoeriocitrin, eriodictyol) even become pro-oxidant (Finotti and Di Majo, 2003). Overall, common dietary flavanones being devoid of a catechol nucleus are only poor antioxidants and their metabolites are expected to be even less potent. Hence, the most significant mechanisms involved in their health effects must be unrelated to their antioxidant activity.

9.2. Anti-inflammatory effects

The phenomena of inflammation have been well described in literature through many reviews. Inflammation is the most obvious manifestation of immune defense. It is manifested by swelling, pain, heat, and redness in the affected tissue and helps eliminate the sources of damage (viruses, bacteria) and initiate healing. Inflammation is produced by immune cells within the tissue, releasing specific mediators that control local circulation and cell activities (Silverstein, 2009). Inflammation response to external stimuli may arise from the action of amines (histamine and 5-hydroxytryptamine), short peptides (bradykinin), long peptides (interleukin-1 (IL-1)), lipids (prostaglandins (PGs) and leukotrienes (LTs)), and many regulatory enzymes (protein kinase C, phosphodiesterase, lipoxigenase, and cyclooxygenase) (Vane and Botting, 1987). Many of the chronic and uncured diseases that plague human populations are due to a dysfunctioning of the immune response.

Hesperidin (hesperetin 7-rutinoside) was found to inhibit kinases and phosphodiesterases responsible for cellular signal transduction and activation during an inflammation response (Manthey et al., 2001). An inhibitory effect of hesperidin on pleurisy (chronic inflammation of lungs) induced by carrageenan was investigated in rats. The results showed a reduction in the volume of exudates and the number of migrating leucocytes by 48% and 34%, respectively, which makes hesperidin a mildly anti-inflammatory agent. Furthermore, this research group observed that hesperidin can reduced yeast-induced hyperthermia in rats (Emim et al., 1994). In another study, hesperidin showed an inhibitory effect on lipopolysaccharide (LPS)-induced overexpression of cyclooxygenase-2, inducible nitric oxide synthase (iNOS), overproduction of prostaglandin E2 and nitric oxide (NO) (Sakata et al., 2003). Similar anti-inflammatory effects were also found for poncirin in RAW 264.7 macrophage cells (Kim et al., 2007b). A study also showed the anti-inflammatory activity of hesperidin by inhibiting arachidonic acid and histamine release (Galati et al., 1994). Significant literature also reports on the anti-inflammatory activity of naringenin and its glycosides. These flavanones are effective in the inhibition of pro-inflammatory cytokines induced by LPS in macrophages and *ex vivo* human whole-blood models to prevent periodontitis (Bodet et al., 2008). They also help attenuate LPS/interferon (IFN)- γ -induced tumor necrosis factor (TNF)- α production in glial cells by inhibiting iNOS expression and nitric oxide production, p38 mitogen-activated protein kinase (MAPK) phosphorylation and downstream signal transducer and activator of transcription-1 (STAT-1) to protect neuro-inflammatory injury (Vafeiadou et al., 2009). Finally, they aid in the reduced production of nitrate and nitrite (indicators of inflammatory process) in DSS (dextran sodium sulfate)-induced ulcerative colitis mice models to control the formation of intestinal edema (Amaro et al., 2009).

9.3. Anti-cancer effects

Advances in cancer research have been spectacular during the past decade. However, it is very unfortunate that the rate of cancer incidence is increasing at an alarming rate. A recent estimation in France has given the figure of 320,000 cases diagnosed in 2005, among which 180,000 in man and 140,000 in woman (INC report). It is clear that serious research in combating cancer is more than ever essential.

Cancer is a complex family of diseases. In terms of molecular and cell biology, cancer is a disease characterized by abnormal gene expression. This altered gene expression occurs through a number of mechanisms, including direct insults to DNA (such as gene mutations, translocations, or amplifications) and abnormal gene transcription or translation. When the DNA of normal cells is exposed to damaging substances (carcinogens), the cells may

undergo genetic alterations that result in malignant transformations, a process known as carcinogenesis. Carcinogens include chemical agents (from industrial pollutants, tobacco *etc.*), viruses (human papilloma virus, hepatitis B&C virus), ionizing (X-rays) and ultraviolet radiations, particles (asbestos, wood dust) and many others. ROS (superoxide, hydrogen peroxide, hydroxyl radical) are also found major causes of not only DNA damage but also protein and lipid damages which lead to aging (Ames and Gold, 1998).

Unlike other flavonoids, flavanones have not been extensively studied for their anti-cancer activities. Moreover, the studies have remained limited to aglycones and glycosides. Recently, hesperetin 7-glucuronide (Hp7G) was demonstrated to affect osteoblast differentiation (Trzeciakiewicz *et al.*, 2010). The major citrus flavanones can be effective in fighting carcinogenesis by minimizing DNA damage, tumor development and proliferation.

9.3.1. Antimutagenic effects

Flavanones can protect DNA damage by their capacity to absorb UV light. The results from a UV irradiated model of plasmidic DNA showed a considerable protecting effect of naringenin against UV-induced damage of DNA (Kootstra, 1994). The moderate antioxidant capacity of flavanones can also be useful in protecting against mutation by free radicals generated near DNA. Furthermore, naringenin also inhibits H₂O₂-induced cytotoxicity and apoptosis, possibly *via* its effect on the H₂O₂-induced expression of an apoptosis-associated gene (Kanno *et al.*, 2003). Naringenin may exhibit anti-mutagenic changes by stimulating DNA repair, following oxidative damage in human prostate cancer cells (Gao *et al.*, 2006).

9.3.2. Inhibition of tumor development

The pharmacological importance of flavanones can also be evaluated by their action against tumor development. So *et al.* (1996) studied the effect of hesperetin and naringenin on the development of breast cancer induced by 7,12-dimethylbenz[*a*]anthracene in female rats. The results showed that tumor development was delayed in rats fed with an orange juice/naringenin-supplemented diet (So *et al.*, 1996). Later on, concerning the anti-angiogenic effect of flavanones, an enzyme-linked immunosorbent assay (ELISA) was used to measure the vascular endothelial growth factor (VEGF) release from mammary adenocarcinoma human breast cancer cells. Naringenin appeared more potent than rutin, apigenin, kaempferol and chrysin (Schindler and Mentlein, 2006). 8-Prenylnaringenin, a derivative of naringenin, inhibits angiogenesis induced by basic fibroblast growth factor (VEGF) or the synergistic effect of two cytokines in combination, in an *in vitro* and *in vivo* study (Pepper *et al.*, 2004). Eight flavanones, including flavanone, 2'-hydroxyflavanone, 4'-hydroxyflavanone, 6-hydroxyflavanone, 7-hydroxyflavanone, naringenin, naringin, and taxifolin, were investigated for their antitumor effects in colorectal carcinoma cells (HT29, COLO205, and COLO320HSR). 2'-Hydroxyflavanone came up as the most potent chemopreventive agent and thus showed a significant inhibitory effect on tumor formation. A recent study on hesperetin supplementation in male albino Wistar rats showed its inhibition of 1,2-dimethylhydrazine (DMH)-induced colon carcinogenesis. The investigation suggested that hesperetin may inhibit phase I enzymes (involved in carcinogen activation), induce phase II xenobiotic metabolizing enzymes and scavenge the electrophilic carcinogenic species (Aranganathan *et al.*, 2009).

9.3.3. Anti-proliferation effects

Naringenin was successfully investigated for its cell antiproliferation effect on an HT-29 colon cancer cell line. Cell proliferation measured by a colorimetric assay was significantly inhibited especially when HT-29 cells were exposed to naringenin at doses greater than 0.7 mM. The results suggested a potential role for citrus fruits as a source of chemoprotective agents against colon cancer

(Frydoonfar *et al.*, 2003). In a comparative study, flavanones showed a significant anti-proliferative activity against lung, colon, breast, prostate and melanoma cancerous cell lines. Moreover, glycosylation reduced the anti-proliferative activity in both flavonoid classes (Manthey and Guthrie, 2002). A C2–C3 double bond seems important for the anti-proliferative activity of flavonoids and indeed flavones are typically more potent than flavanones (Agullo *et al.*, 1996; Kawaii *et al.*, 1999; Manthey and Guthrie, 2002; Rodriguez *et al.*, 2002). Interestingly, methylation of hesperetin and eriodictyol at C7-OH increased the anti-proliferative capacity (Benavente-García and Castillo, 2008). Several mechanisms have been put forward to explain the antiproliferative activity of flavonoids. The most accepted hypothesis is the inhibition of several kinases involved in signal transduction such as protein kinases C, tyrosine kinases, PI 3-kinases or S6 kinase (Casagrande and Darbon, 2001).

The effects of 17 β -estradiol (E2) hormone cover a wide range of physiological processes in mammals such as reproduction, cardiovascular health, bone integrity, cognition and behavior. Besides its roles in human physiology, E2 is also involved in the development of many diseases, including various types of cancers. The mechanism proposed for explain the chemoprotective activity of naringenin suggests that the flavanone binds to ER α (estrogen receptor α) as an antagonist, thereby limiting the effect of E2 in promoting cellular proliferation (Bulzomi *et al.*, 2010).

9.4. Cardiovascular effects

Cardiovascular diseases (CVD) affect the heart and surrounding blood vessels and can take many forms, such as high blood pressure, coronary artery disease, heart disease and stroke. CVD is the largest cause of mortality in the EU and account for approximately 40% of deaths (2 million deaths per year). Chronic oxidative stress and inflammation are among the major factors causing CVD and its control by antioxidants including polyphenols is of great biological significance.

9.4.1. Vasorelaxant and vasoprotective effects

The vascular endothelial cells are very important in normal coronary functions. The regulation of vascular tone and blood flow to organs is controlled by endothelial cells, which synthesize and release a number of factors such as prostacyclin, nitric oxide (NO), endothelium-derived hyperpolarizing factor (EDHF) and endothelin. Among these factors, NO is critical in the preservation of normal vascular functions, and there is a clear relationship between coronary artery disease and NO dysfunctioning (Benavente-García and Castillo, 2008).

Flavonoids promote endothelial NO synthase (eNOS) and at the same time inhibit the inducible NOS (iNOS) (Olszanecki *et al.*, 2002). *In vitro*, inhibition of iNOS in LPS-activated RAW 264.7 cells is not significant with flavanones (naringenin) in comparison to flavones and flavonols (apigenin, luteolin, quercetin), which may demonstrate the significance of a C2–C3 double bond (Kim *et al.*, 1999). Recently, the vasorelaxant potential of hesperetin and hesperidin (Orallo *et al.*, 2004) as well as naringenin and naringin (Orallo *et al.*, 2005) was demonstrated in rats. These vasorelaxant effects are probably due to the inhibition of different phosphodiesterase isoenzymes.

9.4.2. Effect on atherosclerosis and coronary heart disease

There is compelling evidence that coronary heart disease (CHD) is principally related to an elevation of low-density lipoprotein (LDL) cholesterol. Cholesterol, cholesterol esters and triglycerides are transported within LDL particles (protein component ApoB) from their sites of absorption or synthesis to sites of bioactivity. In the artery wall, LDL oxidation and accumulation in macrophages (differentiated from circulating monocytes after adhesion to the

vessel wall) are early events of plaque formation (atherosclerosis). Atherosclerosis causes the hardening and narrowing of arteries, thus putting blood flow at risk. It is the usual cause of heart disease, stroke, and peripheral vascular disease.

The anti-atherosclerosis potential of citrus flavanones, hesperetin and naringenin, was attributed to their ability to regulate ApoB secretion and cellular cholesterol homeostasis in human hepatoma cells line HepG2. A decrease in ApoB accumulation was observed in the media following 24-h incubation with hesperetin and naringenin. This reduced ApoB secretion was related to a reduced cellular concentration of cholesteryl ester (Wilcox et al., 2001). The mechanism involved in naringenin anti-atherosclerosis activity was a reduced ApoB secretion primarily due to the inhibition of microsomal triglyceride transfer protein and the enhancement of LDL receptor (LDLr)-mediated ApoB-containing lipoprotein uptake (Borradaile et al., 2003).

Hesperetin was also shown to limit the rise of hepatic lipid contents and enzymes activities involved in triacylglycerol synthesis in rats fed with 1% orotic acid (Cha et al., 2001). Moreover, a hypolipidemic effect of hesperetin was also reported even during the high lipid concentrations (Kim et al., 2003).

In rats fed a high-cholesterol diet, 0.1% naringenin reduced the levels of plasma cholesterol and hepatic triacylglycerols. This effect was accompanied by a decrease in the activity of 3-hydroxy-3-methylglutaryl-coenzyme A reductase and acyl-CoA cholesterol acyltransferase (Lee et al., 1999, 2003). In another study on rabbits fed with cholesterol-rich diet, naringin caused an increase in superoxide dismutase and catalase activities, thereby contributing to fighting oxidative stress. Moreover, naringin was also shown to up-regulate gene expression of superoxide dismutase, catalase and glutathione peroxidase (Jeon et al., 2001). More recently, naringin, the major grapefruit flavonoid, was studied for its anti-atherosclerosis effect in diet-induced hypercholesterolemia in mice (Chanet et al., 2012a). The study proved that naringenin supplementation at a nutritionally relevant level can reduce atherosclerosis in mice fed with high fat-high cholesterol diet. This protective effect could be related to improved dyslipidemia and biomarkers of endothelial dysfunction, but also to changes in gene expression that may lead to preservation of the vascular wall, as shown by the aorta transcriptomic analysis. The authors have also published a review that outlines the role of citrus flavanones in the protection of cardiovascular disease (Chanet et al., 2012b).

9.5. Other biological effects

Some examples of other interesting bioactivities of citrus flavanone are listed below:

- The antiviral activity of hesperidin has been demonstrated against herpes simples, polio, parainfluenza, and syncytial viral infections, while naringin was inactive (Kaul et al., 1985). In a recent study, hesperetin showed a moderate antimicrobial activity against *Salmonella typhi* and *Salmonella typhimurium* (Kawaguchi et al., 2004).
- A study by Matsuda et al. (1991) suggested that hesperidin has an antiallergic activity via the inhibition of histamine release from pertinent mast cells in rats.
- Oral administration of hesperidin caused a decrease in disease activity index, myeloperoxidase (MPO) activity and malondialdehyde (MDA) content in the dextran sulfate sodium (DSS)-induced ulcerative colitis in mice (Xu et al., 2009).
- Knekt et al. (2002) found an association between a high intake of hesperetin and naringenin and a lower incidence of cerebrovascular disease and asthma. The findings support the clinical importance of monoamine oxidase (MAO-A and B) inhibitors in the treatment of several neurological and psychiatric disorders.

Among the several flavonoids (flavones, thioflavones, flavanones) tested, the most active were the flavanones with highest selective inhibitory activity against MAO-B (Chimenti et al., 2010).

- Beneficial effects of naringenin on cis-platin-induced nephrotoxicity (Badarya et al., 2005) were reported.

10. Conclusion

In the past two to three decades there has been a growing awareness of the role of diet in the etiology of chronic diseases, notably cardiovascular disease (CVD) and some cancers that are major contributors to morbidity and mortality in industrialized countries such as Canada, Australia, the USA and European countries. A wide range of bioactive substances, including polyphenols has already been identified in food and beverages, and it is likely that many more exist. Citrus fruits and juices have long been considered a valuable part of a healthy and nutritious diet, and it is generally assumed that some of the micronutrients in citrus promote health and provide protection against chronic diseases. These micronutrients include mainly the simple flavanones hesperetin, naringenin, eriodictyol and isosakuranetin, but also more complex compounds like calyxin. In-depth knowledge of the composition of all citrus products has yet to be achieved. However, it can be concluded that chemists and nutritionists are progressing more and more rapidly. Various innovative and environmentally friendly techniques are being employed to extract flavanones from a wide range of plant sources. Moreover, analytical methods are improving in their efficiency, sensitivity in detecting minute quantities of complex flavanones, simultaneous quantification of a larger range of compounds, and reduction of analysis time.

Extensive studies have been carried out on the bioactivity of flavanone aglycones and their glycosides. However, a crucial missing link is the validation of the proposed mechanisms with the actual circulating flavanone metabolites. Interestingly, a very recent investigation showed that, unlike A-ring conjugates (hesperetin-7-glucuronide and naringenin-7-glucuronide), B-ring conjugates hesperetin-3'-sulfate, hesperetin-3'-glucuronide and naringenin-4'-glucuronide in low concentrations (2 $\mu\text{mol L}^{-1}$) were as potent as the corresponding aglycones in attenuating monocyte adhesion to TNF- α -activated human umbilical vein endothelial cells (HUVEC) by affecting the expression of related genes (Chanet et al., 2013). This work provides a potential explanation for the vasculoprotective effects of flavanones.

In conclusion, it is now increasingly accepted that phytochemicals found in citrus and other plants consumed in the human diet can be helpful to reduce the risk of many chronic diseases. In our diet, citrus consumption has a considerable potential to expand as part of the overall recommended increase in fruit and vegetable consumption. There is a well-known saying:

"An apple a day keeps the doctor away."

After this survey of the literature, we would like to place emphasis on the recommendations of the French Nutrition & Health Program (PNNS) and claim:

"Five fruits and vegetables a day keep chronic diseases away."

References

- Abad-Garcia, B., Garmon-Lobato, S., Berrueta, L.A., Gallon, B., Vicente, F., 2012. Online characterization of 58 phenolic compounds in Citrus fruit juices from Spanish cultivars by high-performance liquid chromatography with photodiode-array detection coupled to electrospray ionization triple quadrupole mass spectrometry. *Talanta* 99, 213–224.
- Agullo, G., Garnet-Payraastre, L., Fernandez, Y., Anciaux, N., Demigné, C., Rémésy, C., 1996. Comparative effects of flavonoids on the growth, viability and metabolism of a colonic adenocarcinoma cell line (HT29 cells). *Cancer Letters* 105, 61–70.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

- Ak, T., Gülçin, I., 2008. Antioxidant and radical scavenging properties of curcumin. *Chemico-Biological Interactions* 174, 27–37.
- Allaf, T., Tomao, V., Ruiz, K., Chemat, F., 2013. Instant controlled pressure drop technology and ultrasound assisted extraction for sequential extraction of essential oil and antioxidants. *Ultrasonics Sonochemistry* 20, 239–246.
- Al-Maharik, N., Botting, N.P., 2006. A facile synthesis of isoflavone 7-O-glucuronides. *Tetrahedron Letters* 47, 8703–8706.
- Alu'datt, M.H., Rababah, T., Erefeij, K., Alli, I., 2013. Distribution, antioxidant and characterisation of phenolic compounds in soybeans, flaxseed and olives. *Food Chemistry* 139, 93–99.
- Amaro, I.M., Rocha, J., Vila-Real, H., Eduardo-Figueira, M., Mota-Filipe, H., Sepodes, B., Ribeiro, M.H., 2009. Anti-inflammatory activity of naringin and the bio-synthesised naringenin by naringinase immobilized in microstructured materials in a model of DSS-induced colitis in mice. *Food Research International* 42, 1010–1017.
- Ames, B.N., Gold, L.S., 1998. The causes and prevention of cancer: the role of environment. *Biotherapy* 11, 205–220.
- Anagnostopoulou, M.A., Kefalas, P., Papageorgiou, V.P., Assimopoulou, A.N., Boskou, D., 2006. Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). *Food Chemistry* 94, 19–25.
- Andújar, S.A., Filippa, M.A., Ferretti, F.H., Blanco, S.E., 2003. Isomerization of 4'-methoxy-flavanone in alkaline medium. Determination of the enolate formation constant. *Journal of Molecular Structure* 636, 157–166.
- Aranganathan, S., Panneer-Selvam, J., Sangeetha, N., Nalini, N., 2009. Modulatory efficacy of hesperetin (citrus flavanone) on xenobiotic-metabolizing enzymes during 1,2-dimethylhydrazine-induced colon carcinogenesis. *Chemico-Biological Interactions* 180, 254–261.
- Asikin, Y., Taira, I., Inafuku-Teramoto, S., Sumi, H., Ohta, H., Takara, K., Wada, K., 2012. The composition of volatile aroma components, flavanones, and polymethoxylated flavones in shiikuwasha (*Citrus depressa* Hayata) peels of different cultivation lines. *Journal of Agricultural and Food Chemistry* 60, 7973–7980.
- Badarya, O.A., Abdel-Maksoud, S., Ahmed, W.A., Owieda, G.H., 2005. Naringenin attenuates cisplatin nephrotoxicity in rats. *Life Science* 76, 2125–2135.
- Banerjee, A., Basu, K., Sengupta, P.K., 2008. Interaction of 7-hydroxyflavone with human serum albumin: a spectroscopic study. *Journal of Photochemistry and Photobiology* 90, 33–40.
- Barreca, D., Bellocco, E., Caristi, C., Leuzzi, U., Gattuso, G., 2011. Flavonoid profile and radical-scavenging activity of Mediterranean sweet lemon (*Citrus limetta* Risso) juice. *Food Chemistry* 129, 417–422.
- Bartnick, D.D., Mohler, C.M., Houlihan, M., 2006. Methods for the production of food grade extracts. U.S. Patent Application, 20060088627.
- Benavente-García, O., Castillo, J., 2008. Update on uses and properties of citrus flavonoids: new findings in anticancer, cardiovascular, and anti-inflammatory activity. *Journal of Agricultural and Food Chemistry* 56, 6185–6205.
- Biesaga, M., Ochnik, U., Pyrzynska, K., 2009. Fast analysis of prominent flavonoids in tomato using a monolithic column and isocratic HPLC. *Journal of Separation Science* 32, 2835–2840.
- Bimkr, M., Rahman, R.A., Taipa, F.S., Ganjloob, A., Salleha, L.M., Selamat, J., Hamid, A., Zaidul, I.S.M., 2011. Comparison of different extraction methods for the extraction of major bioactive flavonoid compounds from spearmint (*Mentha spicata* L.) leaves. *Food and Bioprocess Technology* 89, 67–72.
- Bocco, A., Cuvelier, M.E., Richard, H., Berset, C., 1998. Antioxidant activity and phenolic composition of citrus peel and seed extracts. *Journal of Agricultural and Food Chemistry* 46, 2123–2129.
- Bodet, C., La, V.D., Epifano, F., Grenier, D., 2008. Naringenin has anti-inflammatory properties in macrophage and ex vivo human whole-blood models. *Journal of Periodontal Research* 43, 400–407.
- Bohm, B.A., 1994. The minor flavonoids. In: Harborne, J.B. (Ed.), *The Flavonoids: Advances in Research Since 1986*. Chapman & Hall, London, UK, pp. 387–440.
- Boonkird, S., Phisalaphong, C., Phisalaphong, M., 2008. Ultrasound-assisted extraction of capsaicinoids from *Capsicum frutescens* on a lab- and pilot-plant scale. *Ultrasonic Sonochemistry* 15, 1075–1079.
- Boros, B., Jakabovaa, S., Dornyeia, A., Horvathd, G., Pluhare, Z., Kilar, F., Felinger, A., 2010. Determination of polyphenolic compounds by liquid chromatography-mass spectrometry in *Thymus* species. *Journal of Chromatography A* 1217, 7972–7980.
- Borradaile, N.M., De Dreu, L.E., Huff, M.W., 2003. Inhibition of net HepG2 cell apolipoprotein B secretion by the citrus flavonoid naringenin involves activation of phosphatidylinositol 3-kinase, independent of insulin receptor substrate-1 phosphorylation. *Diabetes* 52, 2554–2561.
- Bouayed, J., Deufler, H., Hoffmann, L., Bohn, T., 2012. Bioaccessible and dialysable polyphenols in selected apple varieties following *in vitro* digestion vs. their native patterns. *Food Chemistry* 131, 1466–1472.
- Bouktaib, M., Atmanib, A., Rolando, C., 2002. Regio- and stereo-selective synthesis of the major metabolite of quercetin, quercetin-3-O- β -D-glucuronide. *Tetrahedron Letters* 43, 6263–6266.
- Boumendjel, A., Blanc, M., Williamson, G., Barron, D., 2009. Efficient synthesis of flavanone glucuronides. *Journal of Agricultural and Food Chemistry* 57, 7264–7267.
- Brand, W., Van der Wel, P.A.I., Rein, M.J., Barron, D., Williamson, G., van Bladeren, P.J., Rietjens, I.M.C.M., 2008. Metabolism and transport of the citrus flavonoid hesperetin in Caco-2 cell monolayers. *Drug Metabolism and Disposition* 36, 1794–1802.
- Brand, W., Shao, J., Hoek-vanden Hil, E.F., Van Elk, K.N., Spenkelink, B., Haan, L.H.J.D., Rein, M.J., Dionisi, F., Williamson, G., Van Bladeren, P.J., Rietjens, I.M.C.M., 2010. Stereoselective conjugation, transport and bioactivity of S- and R-hesperetin enantiomers *in vitro*. *Journal of Agricultural and Food Chemistry* 58, 6119–6125.
- Brett, G.M., Hollands, W., Needs, P.W., Teucher, B., Dainty, J.R., Davis, B.D., Brodbelt, J.S., Kroon, P.A., 2009. Absorption, metabolism and excretion of flavanones from single portions of orange fruit and juice and effects of anthropometric variables and contraceptive pill use on flavanone excretion. *British Journal of Nutrition* 101, 664–675.
- Brown, G.C., Borutaite, V., 2006. Interactions between nitric oxide, oxygen, reactive oxygen species and reactive nitrogen species. *Biochemical Society Transactions* 34, 953–956.
- Bulzomi, P., Bolli, A., Galluzzo, P., Leone, S., Acconcia, F., Marino, M., 2010. Naringenin and 17 β -estradiol coadministration prevents hormone-induced human cancer cell growth. *IUBMB Life* 62, 51–60.
- Cai, Y., Mei, S., Jie, X., Luo, Q., Corke, H., 2006. Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. *Life Science* 78, 2872–2888.
- Calvarano, M., Postorino, E., Gionfriddo Calvarano, I., Bovalo, F., 1996. Naringin extraction from exhausted bergamot peels. *Perfumer Flavorist* 21, 1–4.
- Cano, A., Medina, A., Bermejo, A., 2008. Bioactive compounds in different citrus varieties. Discrimination among cultivars. *Journal of Food Composition and Analysis* 21, 377–381.
- Cao, H., Chen, X., Sun, H., Tetsuro, S., Zhou, J., Sun, W., Lv, H., Wang, X., 2010. Pharmacokinetics-based elucidation on disparity in clinical effectiveness between varieties of Zhi Zhu Wan, a Traditional Chinese Medical formula. *Journal of Ethnopharmacology* 128, 606–610.
- Casagrande, F., Darbon, J.M., 2001. Effects of structurally related flavonoids on cell cycle progression of human melanoma cells: regulation of cyclin-dependent kinases CDK2 and CDK1. *Biochemical Pharmacology* 61, 1205–1215.
- Cha, J.Y., Cho, Y.S., Kim, I., Anno, T., Rahman, S.M., Yanagita, T., 2001. Effect of hesperetin, a citrus flavonoid, on the liver triacylglycerol content and phosphatidate phosphohydrolase activity in orotic acid-fed rats. *Plant Foods for Human Nutrition* 56, 349–358.
- Chandrasekhar, S., Vijeender, K., Venkatram-Reddy, K., 2005. New synthesis of flavanones catalyzed by L-proline. *Tetrahedron Letters* 46, 6991–6993.
- Chanet, A., Milenkovic, D., Deval, C., Potier, M., Constans, J., Mazur, A., Bennetau-Pelissero, C., Morand, C., Bérard, A.M., 2012a. Naringin, the major grapefruit flavonoid, specifically affects atherosclerosis development in diet-induced hypercholesterolemia in mice. *Journal of Nutritional Biochemistry* 23, 469–477.
- Chanet, A., Milenkovic, D., Manach, C., Mazur, A., Morand, C., 2012b. Citrus flavanones: what is their role in cardiovascular protection? *Journal of Agricultural and Food Chemistry* 60, 8809–8822.
- Chanet, A., Milenkovic, D., Claude, S., Maier, J.A.M., Khan, M.K., Rakotomanomana, N., Shinkaruk, S., Berard, A.M., Bennetau-Pelissero, C., Mazur, A., Morand, C., 2013. Flavanone metabolites decrease monocyte adhesion to TNF- α -activated endothelial cells by modulating expression of atherosclerosis-related genes. *British Journal of Nutrition* 110, 587–598.
- Chebrolua, K.K., Jayaprakashaa, G.K., Jifon, J., Patil, B.S., 2011. Optimization of flavanones extraction by modulating differential solvent densities and centrifuge temperatures. *Talanta* 85, 353–362.
- Cheigh, C.I., Chung, E.Y., Chung, M.S., 2012. Enhanced extraction of flavanones hesperidin and narirutin from *Citrus unshiu* peel using subcritical water. *Journal of Food Engineering* 110, 472–477.
- Chemat, F., Tomao, V., Virot, M., 2008. Ultrasound-assisted extraction in food analysis. In: Otlés, S. (Ed.), *Handbook of Food Analysis Instruments*. CRC Press, Boca Raton, FL, USA, pp. 85–103.
- Chemat, F., Abert-Vian, M., Zill-e-Huma, 2009. Microwave assisted – separations: green chemistry in action. In: Pearlman, J.T. (Ed.), *Green Chemistry Research Trends*. Nova Science Publishers Inc., United States, pp. 1–30.
- Cheyrier, V., 2005. Polyphenols in foods are more complex than often thought. *American Journal of Clinical Nutrition* 81, 223S–229S.
- Chimenti, F., Fioravanti, R., Bolasco, A., Chimenti, P., Secci, D., Rossi, F., Yáñez, M., Orallo, F., Ortuso, F., Alcaro, S., Cirilli, R., Ferretti, R., Sanna, M.L., 2010. A new series of flavones, thioflavones, and flavanones as selective monoamine oxidase-B inhibitors. *Bioorganic and Medicinal Chemistry* 18, 1273–1279.
- Choudhury, R., Chowrimootoo, G., Srail, K., Debnam, E., Rice-Evans, C.A., 1999. Interaction of the flavonoid naringenin in the gastrointestinal tract and the influence of glycosylation. *Biochemical and Biophysical Research Communications* 265, 410–415.
- Climent, M.J., Corma, A., Iborra, S., Primo, J., 1995. Base catalysis for fine chemicals production: Claisen–Schmidt condensation on zeolites and hydrotalcites for the production of chalcones and flavanones of pharmaceutical interest. *Journal of Catalysis* 151, 60–66.
- Cooper-driver, G., Bhattacharya, M., 1998. Role of phenolics in plant evolution. *Phytochemistry* 49, 1165–1174.
- Cortes-Concepcion, J.A., Patcas, F., Amiridis, M.D., 2010. Effect of Li on the catalytic activity of MgO for the synthesis of flavanone. *Applied Catalysis A-General* 386, 1–8.
- Curto, R.L., Tripodo, M.M., Leuzzi, U., Giuffrè, D., Vaccarino, C., 1992. Flavonoids recovery and SCP production from orange peel. *Bioresource Technology* 42, 83–87.
- Del Rio, D., Costa, L., Lean, M., Crozier, A., 2010. Polyphenols and health: what compounds are involved? *Nutrition, Metabolism and Cardiovascular Diseases* 20, 1–6.
- Dhuique-Mayer, C., Caris-Veyrat, C., Ollitrault, P., Curk, F., Amiot, M.J., 2005. Varietal and interspecific influence on micronutrient contents in citrus from the Mediterranean area. *Journal of Agricultural and Food Chemistry* 53, 2140–2145.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

- Di Donna, L., Taverna, D., Mazzotti, F., Benabdelkamel, H., Attya, M., Napoli, A., Sindona, G., 2013. Comprehensive assay of flavanones in citrus juices and beverages by UHPLC–ESI-MS/MS and derivatization chemistry. *Food Chemistry* 141, 2328–2333.
- Di Lecce, G., Martínez-Huélamo, M., Tulipani, S., Vallverdú-Queralt, A., Lamuela-Raventós, R.M., 2013. Setup of a UHPLC–QqQ-MS method for the analysis of phenolic compounds in cherry tomatoes, tomato sauce, and tomato juice. *Journal of Agricultural and Food Chemistry* 61, 8373–8380.
- Di Majo, D., Giammanco, M., Guardia, M., Tripoli, E., Giammanco, S., Finotti, E., 2005. Flavanones in citrus fruits: structure–antioxidant activity relationships. *Food Research International* 38, 1161–1166.
- Dufour, C., Dangles, O., 2005. Flavonoid–serum albumin complexation: determination of binding constants and binding sites by fluorescence spectroscopy. *Biochimica Biophysica Acta – General Subjects* 1721, 164–173.
- Dykes, L., Rooney, W.L., Rooney, L.W., 2013. Evaluation of phenolics and antioxidant activity of black sorghum hybrids. *Journal of Cereal Science* 58, 278–283.
- Emim, J.A., Oliveira, A.B., Lapa, A.J., 1994. Pharmacological evaluation of the anti-inflammatory activity of a citrus bioflavonoid, hesperidin, and the isoflavonoids, dauricin and claussequinone, in rats and mice. *Journal of Pharmacy and Pharmacology* 46, 118–122.
- Erlund, I., 2004. Review of the flavonoids quercetin, hesperetin, and naringenin. Dietary sources, bioactivities, bioavailability, and epidemiology. *Nutritional Research* 24, 851–874.
- Erlund, I., Meririnne, E., Alfthan, G., Aro, A., 2001. Plasma kinetics and urinary excretion of the flavanones naringenin and hesperetin in humans after ingestion of orange juice and grapefruit juice. *Journal of Nutrition* 131, 235–241.
- Esaki, S., Nishiyama, K., Sugiyama, N., Nakajima, R., Takao, Y., Kamiya, S., 1994. Preparation and taste of certain glycosides of flavanones and of dihydrochalcones. *Bioscience, Biotechnology, and Biochemistry* 58, 1479–1485.
- Escudero-López, B., Cerrillo, I., Herrero-Martín, G., Hornero-Méndez, D., Gil-Izquierdo, A., Medina, S., Ferreres, F., Berná, G., Martín, F., Fernández-Pachón, M.S., 2013. Fermented orange juice: source of higher carotenoid and flavanone contents. *Journal of Agricultural and Food Chemistry* 61, 8773–8782.
- Felgines, C., Texier, O., Morand, C., Manach, C., Scalbert, A., Regeat, F., Remesy, C., 2000. Bioavailability of the flavanone naringenin and its glycosides in rats. *American Journal of Physiology: Gastrointestinal and Liver Physiology* 279, G1148–G1154.
- Finotti, E., Di Majo, D., 2003. Influence of solvents on the antioxidant property of flavonoids. *Nahrung* 47, 186–187.
- Forman, H., Fukuto, J., Miller, T., Zhang, H., Rinna, A., Levy, S., 2008. The chemistry of cell signaling by reactive oxygen and nitrogen species and 4-hydroxynonenal. *Archives of Biochemistry and Biophysics* 477, 183–195.
- Fowler, Z.L., Koffas, M.A.G., 2009. Biosynthesis and biotechnological production of flavanones: current state and perspectives. *Applied Microbiology and Biotechnology* 83, 799–808.
- Fox, D.W., Savage, W.L., Wender, S.H., 1953. Hydrolysis of some flavonoid rhamnoglucosides to flavonoid glucosides. *Journal of the American Chemical Society* 75, 2504–2505.
- French, D., Schifano, P., Cortés-Concepción, J., Hargrove-Leak, S., 2010. Li–Al layered double hydroxides as catalysts for the synthesis of flavanone. *Catalysis Communications* 12, 92–94.
- Frydoonfar, H.R., McGrath, D.R., Spigelman, A.D., 2003. The variable effect on proliferation of a colon cancer cell line by the citrus fruit flavonoid naringenin. *Colorectal Disease* 5, 149–152.
- Fumeaux, R., Menozzi-Smarrito, C., Stalmach, A., Munari, C., Kraehenbuehl, K., Steiling, H., Crozier, A., Williamson, G., Barron, D., 2010. First synthesis, characterization, and evidence for the presence of hydroxycinnamic acid sulfate and glucuronide conjugates in human biological fluids as a result of coffee consumption. *Organic and Biomolecular Chemistry* 8, 5199–5211.
- Galati, E.M., Monforte, M.T., Kirjavainen, S., Forestieri, A.M., Trovato, A., Tripodo, M.M., 1994. Biological effects of hesperidin, a citrus flavonoid. (Note I): anti-inflammatory and analgesic activity. *Pharmacology* 40, 709–712.
- Galland, S., Rakotomanomana, N., Dufour, C., Mora, N., Dangles, O., 2008. Synthesis of hydroxycinnamic acid glucuronides and investigation of their affinity for human serum albumin. *Organic and Biomolecular Chemistry* 6, 4253–4260.
- Gao, K., Henning, S., Niu, Y., Youssefian, A.A., Seeram, N.P., Xu, A., Heber, D., 2006. The citrus flavonoid naringenin stimulates DNA repair in prostate cancer cells. *Journal of Nutritional Biochemistry* 17, 89–95.
- Gardana, C., Guarnieri, S., Riso, P., Simonetti, P., Porrini, M., 2007. Flavonone plasma pharmacokinetics from blood orange juice in human subjects. *British Journal of Nutrition* 98, 165–172.
- Gattuso, G., Barreca, D., Gargiulli, C., Leuzzi, U., Caristi, C., 2007. Flavonoid composition of citrus juices. *Molecules* 12, 1641–1673.
- Ghuman, J., Zunsain, P.A., Petitpas, L., Bhattacharya, A.A., Otagiri, M., Curry, S., 2005. Structural basis of the drug-binding specificity of human serum albumin. *Journal of Molecular Biology* 353, 38–52.
- Giannuzzo, A.N., Boggetti, H.J., Nazareno, M.A., Mishima, H.T., 2003. Supercritical fluid extraction of naringin from the peel of *Citrus paradise*. *Phytochemical Analysis* 14, 221–223.
- Gil-Izquierdo, A., Gil, M.I., Tomas-Barberan, F.A., Ferreres, F., 2003. Influence of industrial processing on orange juice flavanone solubility and transformation to chalcones under gastrointestinal conditions. *Journal of Agricultural and Food Chemistry* 51, 3024–3028.
- Gimenez-Bastida, J., Martínez-Florensa, M., Espin, J.C., Tomas-Barberan, F.A., García-Conesa, M.T., 2009. A citrus extract containing flavanones represses plasminogen activator inhibitor-1 (PAI-1) expression and regulates multiple inflammatory, tissue repair, and fibrosis genes in human colon fibroblasts. *Journal of Agricultural and Food Chemistry* 57, 9305–9315.
- González, E.A., Nazareno, M.A., Borsarelli, C.D., 2002. Enthalpy–entropy compensation effect in the chalcone formation from naringin in water–ethanol mixtures. *Journal of Chemical Society Perkin Transactions 2* 2052–2056.
- González-Manzano, S., González-Paramás, A., Santos-Buelga, C., Duenãs, M., 2009. Preparation and characterization of catechin sulfates, glucuronides, and methyl ethers with metabolic interest. *Journal of Agricultural and Food Chemistry* 57, 1231–1238.
- González-Molina, I., Domínguez-Perles, R., Moreno, D.A., García-Viguer, C., 2010. Natural bioactive compounds of *Citrus limon* for food and health. *Journal of Pharmaceutical and Biomedical Analysis* 51, 327–345.
- Goulas, V., Manganaris, G.A., 2012. Exploring the phytochemical content and the antioxidant potential of *Citrus* fruits grown in Cyprus. *Food Chemistry* 131, 39–47.
- Hanessian, S., Kothakonda, K.K., 2005. Natural products as scaffolds for chemical diversification: solution-phase parallel synthesis of a series of naringin analogues. *Journal of Combinatorial Chemistry* 7, 837–842.
- Harnly, J.M., Doherty, R.F., Beecher, G.R., Holden, J.M., Haytowitz, D.B., Bhagwat, S., Gebhardt, S., 2006. Flavonoid content of U.S. fruits, vegetables, and nuts. *Journal of Agricultural and Food Chemistry* 54, 9966–9977.
- Hayat, K., Hussain, S., Abbas, S., Farooq, U., Ding, B., Xia, S., Jia, C., Zhang, X., Xia, W., 2009. Optimized microwave-assisted extraction of phenolic acids from citrus mandarin peels and evaluation of antioxidant activity *in vitro*. *Separation and Purification Technology* 70, 63–70.
- Hayat, K., Zhang, X., Chen, H., Xia, S., Jia, C., Zhong, F., 2010a. Liberation and separation of phenolic compounds from citrus mandarin peels by microwave heating and its effect on antioxidant activity. *Separation and Purification Technology* 73, 371–376.
- Hayat, K., Zhang, X., Farooq, U., Abbas, S., Xia, S., Jia, C., Zhong, F., Zhang, J., 2010b. Effect of microwave treatment on phenolic content and antioxidant activity of citrus mandarin pomace. *Food Chemistry* 123, 423–429.
- He, X.M., Carter, D.C., 1992. Atomic structure and chemistry of human serum albumin. *Nature* 358, 209–215.
- Hsieh, H.K., Lee, T.H., Wang, J.P., Wang, J.J., Lin, C.N., 1998. Synthesis and anti-inflammatory effect of chalcones and related compounds. *Pharmaceutical Research* 15, 39–46.
- Hu, Y.J., Wang, Y., Ou-Yang, Y., Zhou, J., Liu, Y., 2010. Characterize the interaction between naringenin and bovine serum albumin using spectroscopic approach. *Journal of Luminescence* 130, 1394–1399.
- Igual, M., García-Martínez, E., Camacho, M.M., Martínez-Navarrete, N., 2013. Jam processing and storage effects on β -carotene and flavonoids content in grapefruit. *Journal of Functional Foods* 5, 736–737, 44.
- Iijima, Y., Suda, K., Suzuki, T., Aoki, K., Shibata, D., 2008. Metabolite profiling of chalcones and flavanones in tomato fruit. *Journal of Japanese Society of Horticulture Science* 77, 94–102.
- Iwashina, T., 2000. The structure and distribution of the flavonoids in plants. *Journal of Plant Research* 113, 287–299.
- Jeon, S.M., Bok, S.H., Jang, M.K., Lee, M.K., Nam, K.T., Park, Y.B., Rhee, S.J., Choi, M.S., 2001. Antioxidative activity of naringin and lovastatin in high cholesterol-fed rabbits. *Life Science* 69, 2855–2866.
- Jeong, S.M., Kim, S.Y., Kim, D.R., Jo, S.C., Nam, K.C., Ahn, D.U., Lee, S.C., 2004. Effect of heat treatment on the antioxidant activity of extracts from citrus peels. *Journal of Agricultural and Food Chemistry* 52, 3389–3393.
- Jez, J.M., Noel, J.P., 2002a. Reaction mechanism of chalcone isomerase: pH dependence, diffusion control, and product binding differences. *Journal of Biological Chemistry* 277, 1361–1369.
- Jez, J.M., Bowman, M.E., Noel, J.P., 2002b. Role of hydrogen bonds in the reaction mechanism of chalcone isomerase. *Biochemistry* 41, 5168–5176.
- Kanaze, F.I., Bounartzki, M.I., Georgarakis, M., Niopas, I., 2007. Pharmacokinetics of the citrus flavanone aglycones hesperetin and naringenin after single oral administration in human subjects. *European Journal of Clinical Nutrition* 61, 472–477.
- Kanno, S., Shouji, A., Asou, K., Ishikawa, M., 2003. Effects of naringin on hydrogen peroxide-induced cytotoxicity and apoptosis in P388 cells. *Journal of Pharmacological Science* 92, 166–170.
- Kaul, T.N., Middleton, E., Ogra, P.L., 1985. Antiviral effect of flavonoids on human viruses. *Journal of Medical Virology* 15, 71–79.
- Kawaguchi, K., Kikuchi, S., Hasunuma, R., Maruyama, H., Yoshikawa, T., Kumazawa, Y., 2004. A citrus flavonoid hesperidin suppresses infection-induced endotoxin shock in mice. *Biological and Pharmaceutical Bulletin* 27, 679–683.
- Kawaii, S., Tomono, Y., Katase, E., Ogawa, K., Yano, M., 1999. Effect of citrus flavonoids on HL-60 cell differentiation. *Anticancer Research* 19, 1261–1269.
- Keilig, K., Ludwig-Müller, J., 2009. Effect of flavonoids on heavy metal tolerance in *Arabidopsis thaliana* seedlings. *Botanical Studies* 50, 311–318.
- Khan, M.K., Abert-Vian, M., Fabiano-Tixier, A.S., Dangles, O., Chemat, F., 2010a. Ultrasound-assisted extraction of polyphenols (flavanone glycosides) from orange (*Citrus sinensis* L.) peel. *Food Chemistry* 119, 851–858.
- Khan, M.K., Rakotomanomana, N., Loonis, M., Dangles, O., 2010b. Chemical synthesis of flavanone glucuronides. *Journal of Agricultural and Food Chemistry* 58, 8437–8443.
- Khan, M.K., Rakotomanomana, N., Dufour, C., Dangles, O., 2011. Binding of citrus flavanones and their glucuronides and chalcones to human serum albumin. *Food and Function* 2, 617–626.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

- Kim, H., Cheon, B.S., Kim, Y.H., Kim, S.Y., Kim, H.P., 1999. Effects of naturally occurring flavonoids on nitric oxide production in the macrophage cell line RAW 264.7 and their structure–activity relationships. *Biochemical Pharmacology* 58, 759–765.
- Kim, H., Jeong, T.S., Lee, M.K., Park, Y.B., Choi, M.S., 2003. Lipid-lowering efficacy of hesperetin metabolites in high-cholesterol fed rats. *Clinica Chimica Acta* 327, 129–137.
- Kim, M.R., Kim, W.C., Lee, D.Y., Kim, C.W., 2007a. Recovery of narirutin by adsorption on a non-ionic polar resin from a water-extract of *Citrus unshiu* peels. *Journal of Food Engineering* 78, 27–32.
- Kim, J.B., Han, A., Park, E., Kim, J.Y., Cho, W., Lee, J., Seo, E.K., Lee, K.T., 2007b. Inhibition of LPS-induced iNOS, COX-2 and cytokines expression by poncirin through the NF- κ B inactivation in RAW 264.7 macrophage cells. *Biological and Pharmaceutical Bulletin* 30, 2345–2351.
- Kim, J.W., Lee, B.C., Lee, J.H., Nam, K.C., Lee, S.C., 2008. Effect of electron-beam irradiation on the antioxidant activity of extracts from *Citrus unshiu* pomaces. *Radiation Physics and Chemistry* 77, 87–91.
- Knekt, P., Kumpulainen, J., Järvinen, R., Rissanen, H., Heliövaara, M., Reunanen, A., Hakulinen, T., Aromaa, A., 2002. Flavonoid intake and risk of chronic diseases. *American Journal of Clinical Nutrition* 76, 560–568.
- Kobayashi, S., Konishi, Y., 2008. Transepithelial transport of flavanone in intestinal Caco-2 cell monolayers. *Biochemical and Biophysical Research Communications* 368, 23–29.
- Kobayashi, S., Tanabe, S., Sugiyama, M., Konishi, Y., 2008. Transepithelial transport of hesperetin and hesperidin in intestinal Caco-2 cell monolayers. *Biochimica Biophysica Acta – Biomembranes* 1778, 33–41.
- Koca, U., Berhow, M.A., Febres, V.J., Champ, K.I., Carrillo-Mendoza, O., Moore, G.A., 2009. Decreasing unpalatable flavonoid components in the effect of transformation construct. *Physiologia Plantarum* 137, 101–114.
- Kometani, T., Nishimura, T., Nakae, T., Takii, H., Okada, S., 1996. Synthesis of neohesperidin glycosides and naringin glycosides by cyclodextrin glucanotransferase from an Alkalophilic *Bacillus* species. *Bioscience, Biotechnology, and Biochemistry* 60, 645–649.
- Kootstra, A., 1994. Protection from UV-B-induced DNA damage by flavonoids. *Plant Molecular Biology* 26, 771–774.
- Kratz, F., 2008. Albumin as a drug carrier: design of prodrugs, drug conjugates and nanoparticles. *Journal of Controlled Release* 132, 171–183.
- Krbeček, L., Inglett, G.E., Holik, M., Dowling, B., Wagner, R., Riter, R., 1968. Dihydrochalcones. Synthesis of potential sweetening agents. *Journal of Agricultural and Food Chemistry* 16, 108–112.
- Kris-etherton, P.M., Hecker, K.D., Bonanome, A., Coval, S.M., Binkoski, A.E., Hilpert, K.F., Griel, A.E., Etherton, T.D., 2002. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *American Journal of Medicine* 113, 71–88.
- Lee, S., Park, Y.B., Bae, K.H., Bok, S.H., Kwon, Y.K., Lee, E.S., Choi, M.S., 1999. Cholesterol-lowering activity of naringenin via inhibition of 3-hydroxy-3-methylglutaryl coenzyme A reductase and acyl coenzyme A: cholesterol acyltransferase in rats. *Annals of Nutrition and Metabolism* 43, 173–180.
- Lee, M., Moon, S.S., Lee, S.E., Bok, S.H., Jeong, T.S., Park, Y.B., Choi, M.S., 2003. Naringenin 7-O-cetyl ether as inhibitor of HMG-CoA reductase and modulator of plasma and hepatic lipids in high cholesterol-fed rats. *Bioorganic and Medicinal Chemistry* 11, 393–398.
- Lewinsohn, E., Britsch, L., Mazur, Y., Gresse, J., 1989. Flavanone glycoside biosynthesis in citrus. *Plant Physiology* 91, 1323–1328.
- Li, B.B., Smith, B., Hossain, M.M., 2006a. Extraction of phenolics from citrus peels: II. Enzyme-assisted extraction method. *Separation and Purification Technology* 48, 189–196.
- Li, B.B., Smith, B., Hossain, M.M., 2006b. Extraction of phenolics from citrus peels: I. Solvent extraction method. *Separation and Purification Technology* 48, 182–188.
- Li, W., Wang, Z., Wang, Y.P., Jiang, C., Liu, Q., Sun, Y.S., Zheng, Y.N., 2012. Pressurized liquid extraction combining LC–DAD–ESI/MS analysis as an alternative method to extract three major flavones in *Citrus reticulata* 'Chachi' (Guangchenpi). *Food Chemistry* 130, 1044–1049.
- Liu, Y., Li, Y., Liu, S., Li, J., Yao, S., 2004. Monitoring the self-assembly of chitosan/glutaraldehyde/cysteamine/Au-colloid and the binding of human serum albumin with hesperidin. *Biomaterials* 25, 5725–5733.
- Lu, Z., Zhang, Y., Liu, H., Yuan, J., Zheng, Z., Zou, G., 2007. Transport of a cancer chemopreventive polyphenol, resveratrol: interaction with serum albumin and hemoglobin. *Journal of Fluorescence* 17, 580–587.
- Luque-García, J.L., Luque de Castro, M.D., 2003. Ultrasound: a powerful tool for leaching. *Trends in Analytical Chemistry* 22, 41–47.
- Ma, Y., Ye, X., Hao, Y., Xu, G., Xu, G., Liu, D., 2008a. Ultrasound-assisted extraction of hesperidin from Penggan (*Citrus reticulata*) peel. *Ultrasonics Sonochemistry* 15, 227–232.
- Ma, Y., Ye, X., Fang, Z., Chen, J., Xu, G., Liu, D., 2008b. Phenolic compounds and antioxidant activity of extracts from ultrasonic treatment of Satsuma mandarin (*Citrus unshiu* Marc.) peels. *Journal of Agricultural and Food Chemistry* 56, 5682–5690.
- Ma, Y., Chen, J., Liu, D., Ye, X., 2008c. Effect of ultrasonic treatment on the total phenolic and antioxidant activity of extracts from citrus peel. *Journal of Food Science* 73, 115–120.
- Ma, Y., Chen, J., Liu, D., Ye, X., 2009. Simultaneous extraction of phenolic compounds of citrus peel extracts: effect of ultrasound. *Ultrasonics Sonochemistry* 16, 57–62.
- Manach, C., Morand, C., Gil-Izquierdo, A., Bouteloup-Demange, C., Rémésy, C., 2003. Bioavailability in humans of the flavanones hesperidin and narirutin after the ingestion of two doses of orange juice. *European Journal of Clinical Nutrition* 57, 235–242.
- Manthey, J.A., Grohmann, K., 1996. Concentrations of hesperidin and other orange peel flavonoids in citrus processing byproducts. *Journal of Agricultural and Food Chemistry* 44, 811–814.
- Manthey, J.A., Guthrie, N., 2002. Antiproliferative activities of citrus flavonoids against six human cancer cell lines. *Journal of Agricultural and Food Chemistry* 50, 5837–5843.
- Manthey, J.A., Grohmann, K., Guthrie, N., 2001. Biological properties of citrus flavonoids pertaining to cancer and inflammation. *Current Medicinal Chemistry* 8, 135–153.
- Marais, J.P.J., Deavours, B., Dixon, R.A., Ferreira, D., 2006. The stereochemistry of flavonoids. In: Grotewold, E. (Ed.), *The Science of Flavonoids*. Springer Science and Business Media Inc., USA, pp. 1–46.
- Martens, S., Mithöfer, A., 2005. Flavones and flavone synthases. *Phytochemistry* 66, 2399–2407.
- Mato, M., Onozaki, T., Ozeki, Y., Higeta, D., Itoh, Y., Yoshimoto, Y., Ikeda, H., Yoshida, H., Shibata, M., 2000. Flavonoid biosynthesis in white-flowered *Sim carnations* (*Dianthus caryophyllus*). *Scientia Horticulturae* 84, 333–347.
- Matsuda, H., Yano, M., Kubo, M., Iinuma, M., Oyama, M., Mizuno, M., 1991. Pharmacological study on citrus fruits. II. Anti-allergic effect of fruit of *Citrus unshiu* MARKOVICH (2). On flavonoid components. *Yakugaku Zasshi* 111, 193–198.
- Matsumoto, H., Ikoma, Y., Sugiura, M., Yano, M., Hasegawa, Y., 2004. Identification and quantification of the conjugated metabolites derived from orally administered hesperidin in rat plasma. *Journal of Agricultural and Food Chemistry* 52, 6653–6659.
- Menichini, F., Houghton, P.J., Jones, P., Liu, B., Loizzo, M., Tundis, R., De Luca, D., De Cindio, B., Menichini, F., Perseud, S., 2009. *In vitro* antidiabetic activity effect of *Citrus medica* cv. diamante peel extract. *Annals in Nutrient Metabolism* 55, 596.
- Miles, C.O., Main, L., 1985. Kinetics and mechanism of the cyclization of 2',6'-dihydroxy-4,4'-dimethoxychalcone; influence of the 6'-hydroxy group on the rate of cyclization under neutral conditions. *Journal of Chemical Society Perkin Translation 2* 1639–1642.
- Miyake, Y., Shimoi, K., Kumazawa, S., Yamamoto, K., Kinai, N., Osawa, T., 2000. Identification and antioxidant activity of flavonoid metabolites in plasma and urine of eriocitrin-treated rats. *Journal of Agricultural and Food Chemistry* 48, 3217–3224.
- Miyake, Y., Sakurai, C., Usuda, M., Fukumoto, S., Hiramitsu, M., Sakaida, K., Osawa, T., Kondo, K., 2006. Difference in plasma metabolite concentration after ingestion of lemon flavonoids and their aglycones in human. *Journal of Nutritional Science and Vitaminology* 52, 54–60.
- Moon, J.H., Tshushida, T., Nakahara, K., Terao, J., 2001. Identification of quercetin 3-O-beta-D-glucuronide as an antioxidative metabolite in rat plasma after oral administration of quercetin. *Free Radical Biology and Medicine* 30, 1274–1285.
- Moure, A., Cruz, J.M., Franco, D., Dominguez, J.M., Sineiro, J., Dominguez, H., Nunez, M.J., Parajo, J.C., 2001. Natural antioxidants from residual sources. *Food Chemistry* 72, 145–171.
- Mullen, W., Archeveque, M.A., Edwards, C.A., Matsumoto, H., Crozier, A., 2008. Bioavailability and metabolism of orange juice flavanones in humans: impact of a full-fat yogurt. *Journal of Agricultural and Food Chemistry* 56, 11157–11164.
- Murota, K., Hotta, A., Ido, H., Kawai, Y., Moon, J.H., Sekido, K., Hayashi, H., Inakuma, T., Terao, J., 2007. Antioxidant activity of albumin-bound quercetin metabolites after onion consumption in humans. *Journal of Medical Investigation* 54, 370–374.
- Needs, P.W., Kroon, P.A., 2006. Convenient syntheses of metabolically important quercetin glucuronides and sulfates. *Tetrahedron* 62, 6862–6868.
- Nogata, Y., Sakamoto, K., Shiratsuchi, H., Ishii, T., Yano, M., Ohto, H., 2006. Flavonoid composition of fruit tissues of citrus species. *Bioscience, Biotechnology, and Biochemistry* 70, 178–192.
- Olszanecki, R., Gebeska, A., Kozlovski, V.I., Gryglewski, R.J., 2002. Flavonoids and nitric oxide synthase. *Journal of Physiology and Pharmacology* 53, 571–584.
- Orallo, F., Alvarez, E., Basaran, H., Lugnier, C., 2004. Comparative study of the vasorelaxant activity, superoxide-scavenging ability and cyclic nucleotide phosphodiesterase-inhibitory effects of hesperetin and hesperidin. *Naunyn-Schmiedeberg's Archives of Pharmacology* 370, 452–463.
- Orallo, F., Camiña, M., Alvarez, E., Basaran, H., Lugnier, C., 2005. Implication of cyclic nucleotide phosphodiesterase inhibition in the vasorelaxant activity of the citrus-fruits flavonoid (+/–)-naringenin. *Planta Medica* 71, 99–107.
- Oravcová, J., Böhs, B., Lindner, W., 1996. Drug-protein binding studies new trends in analytical and experimental methodology. *Journal of Chromatography B: Biomedical Sciences and Applications* 677, 1–28.
- Oufedjikh, H., Mahrouz, M., Amiot, M.J., Lacroix, M., 2000. Effect of γ -irradiation on phenolic compounds and phenylalanine ammonia-lyase activity during storage in relation to peel injury from peel of *Citrus clementina* Hort. ex. Tanaka. *Journal of Agricultural and Food Chemistry* 48, 559–565.
- Oyama, K., Kondo, T., 2004. Total synthesis of apigenin 7,4'-di-O- β -glucopyranoside, a component of blue flower pigment of *Salvia patens*, and seven chiral analogues. *Tetrahedron* 60, 2025–2034.
- Parr, A.J., Bolwell, G.P., 2000. Phenols in the plant and in man. The potential for possible nutritional enhancement of the diet by modifying the phenols content or profile. *Journal of Science, Food and Agriculture* 80, 985–1012.
- Patil, B.S., Jayaprakasha, G.K., Chidambara-Murthy, K.N., Vikram, A., 2009a. Bioactive compounds: historical perspectives, opportunities, and challenges. *Journal of Agricultural and Food Chemistry* 57, 8142–8160.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004

- Patil, J.R., Murthy, K.N.C., Jayaprakasha, G.K., Chetti, M.B., Patil, B.S., 2009b. Bioactive compounds from Mexican lime (*Citrus aurantifolia*) juice induce apoptosis in human pancreatic cells. *Journal of Agricultural and Food Chemistry* 57, 10933–10942.
- Peng, H., Cheng, F., Huang, Y., Chen, C., Tsai, T., 1998. Determination of naringenin and its glucuronide conjugate in rat plasma and brain tissue by high-performance liquid chromatography. *Journal of Chromatography B: Biomedical Sciences and Applications* 714, 369–374.
- Pepper, M.S., Hazel, S.J., Hampel, M., Schleuning, W., 2004. 8-Prenylnaringenin, a novel phytoestrogen, inhibits angiogenesis *in vitro* and *in vivo*. *Journal of Cellular Physiology* 199, 98–107.
- Pereira, R.M.S., Andrade, N.E.D., Paulino, N., Sawaya, A.C.H.F., Eberlin, M.N., Maruccci, M.C., Favero, G.M., Novak, E.M., Bydlowski, S.P., 2007. Synthesis and characterization of a metal complex containing naringin and Cu, and its antioxidant, antimicrobial, antiinflammatory and tumor cell cytotoxicity. *Molecules* 12, 1352–1366.
- Peschel, W., Sánchez-Rabaneda, F., Diekmann, W., Plescher, A., Gartzia, I., Jiménez, D., Lamuela-Raventós, R., Buxaderas, S., Codina, C., 2006. An industrial approach in the search of natural antioxidants from vegetable and fruit wastes. *Food Chemistry* 97, 137–150.
- Peterson, J., Dwyer, J., Beecher, G., Bhagwat, S.A., Gebhardt, S.E., Haytowitz, D.B., Holden, J.M., 2006a. Flavanones in oranges, tangerines (mandarins), tangors, and tangelos: a compilation and review of the data from the analytical literature. *Journal of Food Composition and Analysis* 19, S66–S73.
- Peterson, J., Beecher, G., Bhagwat, S., Dwyer, J.T., Gebhardt, S.E., Haytowitz, D.B., Holden, J.M., 2006b. Flavanones in grapefruit, lemons, and limes: a compilation and review of the data from the analytical literature. *Journal of Food Composition and Analysis* 19, S74–S80.
- Pietta, P.G., 2000. Flavonoids as antioxidants. *Journal of Natural Products* 63, 1035–1042.
- Ramful, D., Tarnus, E., Aruoma, O.I., Bourdon, E., Bahorun, T., 2011. Polyphenol composition, vitamin C content and antioxidant capacity of Mauritian citrus fruit pulps. *Food Research International* 44, 2088–2099.
- Rawel, H.M., Meidtnier, K., Kroll, J., 2005. Binding of selected phenolic compounds to proteins. *Journal of Agricultural and Food Chemistry* 53, 4228–4235.
- Rodríguez, J., Yáñez, J., Vicente, V., Alcaraz, M., Benavente-García, O., Castillo, J., Lorente, J., Lozano, J.A., 2002. Effects of several flavonoids on the growth of B16F10 and SK-MEL-1 melanoma cell lines: relationship between structure and activity. *Melanoma Research* 12, 99–107.
- Roovij, S., Mullen, W., Edwards, C.A., Crozier, A., 2009. Yoghurt impacts on the excretion of phenolic acids derived from colonic breakdown of orange juice flavanones in humans. *Molecular Nutrition and Food Research* 53, S68–S75.
- Sadeghipour, M., Terreux, R., Phipps, J., 2005. Flavonoids and tyrosine nitration: structure–activity relationship correlation with enthalpy of formation. *Toxicology In Vitro* 19, 155–165.
- Sakata, K., Hirose, Y., Qiao, Z., Tanaka, T., Mori, H., 2003. Inhibition of inducible isoforms of cyclooxygenase and nitric oxide synthase by flavonoid hesperidin in mouse macrophage cell line. *Cancer Letters* 199, 139–145.
- Schijlen, E.G., Ric de Vos, C., van Tunen, A.J., Bovy, A.G., 2004. Modification of flavonoid biosynthesis in crop plants. *Phytochemistry* 65, 2631–2648.
- Schindler, R., Mentlein, R., 2006. Flavonoids and vitamin E reduce the release of the angiogenic peptide vascular endothelial growth factor from human tumor cells. *Journal of Nutrition* 136, 1477–1482.
- Shi, L., Feng, X.E., Cui, J.R., Fang, L.H., Du, G.H., Li, Q.S., 2010. Synthesis and biological activity of flavanone derivatives. *Bioorganic and Medicinal Chemistry Letters* 20, 5466–5468.
- Si-Ahmed, K., Tazerouti, F., Badjah-Hadj-Ahmed, A.Y., Aturki, Z., Orazio, G.D., Rocco, A., Fanali, S., 2010. Analysis of hesperetin enantiomers in human urine after ingestion of blood orange juice by using nano-liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis* 51, 225–229.
- Silberberg, M., Gilizquierdo, A., Combaret, L., Remesy, C., Scalbert, A., Morand, C., 2006. Flavanone metabolism in healthy and tumor-bearing rats. *Biomedical Pharmacotherapy* 60, 529–535.
- Silverstein, D.M., 2009. Inflammation in chronic kidney disease: role in the progression of renal and cardiovascular disease. *Pediatric Nephrology* 24, 1445–1452.
- Simmonds, M.S.J., 2001. Importance of flavonoids in insect–plant interactions: feeding and oviposition. *Phytochemistry* 56, 245–252.
- Slimestad, R., Fossen, T., Verheul, M.J., 2008. The flavonoids of tomatoes. *Journal of Agricultural and Food Chemistry* 56, 2436–2441.
- So, F., Guthrie, N., Chambers, A., Moussa, M., Carroll, K., 1996. Inhibition of human breast cancer cell proliferation and delay of mammary tumorigenesis by flavonoids and citrus juices. *Nutrition and Cancer* 26, 167–181.
- Sood, S., Arora, B., Bansal, S., Muthuraman, A., Gill, N.S., Arora, R., Bali, M., Sharma, P.D., 2009. Antioxidant, anti-inflammatory and analgesic potential of the *Citrus decumana* L. peel extract. *Inflammopharmacology* 17, 267–274.
- Sood, S., Muthuraman, A., Arora, B., Bansal, S., Bali, M., Sharma, P.D., 2010. Potential effect of *Citrus decumana* extract on stress induced peptic ulcer in rat. *Latin American Journal of Pharmacy* 29, 52–56.
- Sugio, S., Kashima, A., Mochizuki, S., Noda, M., Kobayashi, K., 1999. Crystal structure of human serum albumin at 2.5 Å resolution. *Protein Engineering* 12, 439–446.
- Sulkowska, A., 2002. Interaction of drugs with bovine and human serum albumin. *Journal of Molecular Structure* 614, 227–232.
- Tomas-Barberan, F.A., Clifford, M.N., 2000. Flavanones, chalcones and dihydrochalcones – nature, occurrence and dietary burden. *Journal of the Science of Food and Agriculture* 80, 1073–1080.
- Tripoli, E., Guardia, M.L., Giammanco, S., Majo, D.D., Giammanco, M., 2007. Citrus flavonoids: molecular structure, biological activity and nutritional properties: a review. *Food Chemistry* 104, 466–479.
- Trzeciakiewicz, A., Habauzit, V., Mercier, S., Barron, D., Urpi-Sarda, M., Manach, C., Offord, E., Horcajada, M.N., 2010. Molecular mechanism of hesperetin-7-O-glucuronide, the main circulating metabolite of hesperidin, involved in osteoblast differentiation. *Journal of Agricultural and Food Chemistry* 58, 668–675.
- Tsai, T.H., 2002. Determination of naringin in rat blood, brain, liver, and bile using microdialysis and its interaction with cyclosporin A, a P-glycoprotein modulator. *Journal of Agricultural and Food Chemistry* 50, 6669–6674.
- Tsimogiannis, D., Samiotaki, M., Panayotou, G., Oreopoulou, V., 2007. Characterization of flavonoid subgroups and hydroxyl substitution by HPLC-MS/MS. *Molecules* 12, 593–606.
- United States Department of Agriculture/Foreign Agricultural Service, 2010. Citrus: World Markets and Trade.. Available from: <http://www.fas.usda.gov> (accessed 28.09.2010).
- Vafeiadou, K., Vauzour, D., Lee, H.Y., Rodríguez-Mateos, A., Williams, R.J., Spencer, J.P.E., 2009. The citrus flavanone naringenin inhibits inflammatory signalling in glial cells and protects against neuroinflammatory injury. *Archives in Biochemistry and Biophysics* 484, 100–109.
- Van Acker, S.A.B.E., Groot, M.J.D., Berg, D.J.V.D., Tromp, M.N.J.L., Kelder, G.D.O.D., Vijgh, W.J.F.V.D., Bast, A., 1996. A quantum chemical explanation of the antioxidant activity of flavonoids – chemical research in toxicology. *Chemical Research in Toxicology* 9, 1305–1312.
- Vane, J., Botting, R., 1987. Inflammation and the mechanism of action of anti-inflammatory drugs. *FASEB Journal* 1, 89–96.
- Varshney, A., Sen, P., Ahmad, E., Rehan, M., Subbarao, N., Khan, R.H., 2010. Ligand binding strategies of human serum albumin: how can the cargo be utilized? *Chirality* 22, 77–87.
- Veitch, N.C., Grayer, R.J., 2006. Chalcones, dihydrochalcones, and aurones. In: Andersen, O.M., Markham, K.R. (Eds.), *Flavonoids: Chemistry, Biochemistry and Applications*. CRC Press, Taylor & Francis Group, Boca Raton, FL, pp. 1003–1100.
- Veitch, N.C., Grayer, R.J., 2008. Flavonoids and their glycosides, including anthocyanins. *Natural Products Representation* 25, 555.
- Vinatoru, F., 2001. An overview of the ultrasonically assisted extraction of bioactive principles from herbs. *Ultrasonics Sonochemistry* 8, 303–313.
- Wang, L., Weller, C.L., 2006. Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science and Technology* 17, 300–312.
- Wang, Y., Zhang, H., Zhang, G., Tao, W., Tang, S., 2007. Interaction of the flavonoid hesperidin with bovine serum albumin: a fluorescence quenching study. *Journal of Luminescence* 126, 211–218.
- Wang, X., Sakurai, T., Chen, X., Sun, H., Wang, Z., Sun, Q., Sun, W., Cao, H., 2008. Hydrolysis of flavanone glycosides and degradation of the corresponding aglycones from dried immature fruit by human fecal flora. *Planta Medica* 74, 1751–1755.
- Wilcox, L.J., Borradaile, N.M., Dreu, L.E.D., Huff, M.W., 2001. Secretion of hepatocyte ApoB is inhibited by the flavonoids, naringenin and hesperetin, via reduced activity and expression of ACAT2 and MTP. *Journal of Lipid Research* 42, 725–734.
- Xie, M., Xu, X., Wang, Y., 2005a. Interaction between hesperetin and human serum albumin revealed by spectroscopic methods. *Biochimica et Biophysica Acta* 1724, 215–224.
- Xie, M.X., Xu, X.Y., Wang, Y.D., Liu, Y., 2005b. Spectroscopic investigation of the interaction between 2,3-dihydro-4',5,7-trihydroxyflavone and human serum albumin. *Acta Chimica Sinica* 63, 2055–2062.
- Xu, G., Ye, X., Chen, J., Liu, D., 2007. Effect of heat treatment on the phenolic compounds and antioxidant capacity of citrus peel extract. *Journal of Agricultural and Food Chemistry* 55, 330–335.
- Xu, G.H., Chen, J.C., Liu, D.H., Zhang, Y.H., Jiang, P., Ye, X.Q., 2008. Minerals, phenolic compounds, and antioxidant capacity of citrus peel extract by hot water. *Journal of Food Science* 73, C11–C18.
- Xu, L., Yang, Z.L., Li, P., Zhou, Y.Q., 2009. Modulating effect of hesperidin on experimental murine colitis induced by dextran sulfate sodium. *Phytomedicine* 16, 989–995.
- Ye, H., Xu, H., Yu, C., Dai, Y., Liu, G., Xu, W., Yuan, S., 2009. Hydroxylation of naringin by *Trichoderma harzianum* to dramatically improve its antioxidative activity. *Enzymology, and Microbiology Technology* 45, 282–287.
- Ye, X.Q., Chen, J.C., Liu, D.H., Jiang, P., Shi, J., Xue, S., Wu, D., Xu, J.G., Kakud, Y., 2011. Identification of bioactive composition and antioxidant activity in young mandarin fruits. *Food Chemistry* 124, 1561–1566.
- Yu, J., Dandekar, D.V., Toledo, R.T., Singh, R.K., Patil, B.S., 2007. Supercritical fluid extraction of limonoids and naringin from grapefruit (*Citrus paradisi* Macf.) seeds. *Food Chemistry* 105, 1026–1031.
- Zhang, Y., Li, Y., Dong, L., Li, J., He, W., Chen, X., Hu, Z., 2008. Investigation of the interaction between naringin and human serum albumin. *Journal of Molecular Structure* 875, 1–8.
- Zia-ur-Rehman, 2006. Citrus peel extract – a natural source of antioxidant. *Food Chemistry* 99, 450–454.
- Zill-e-Huma, Abert-Vian, M., Maingonnat, J.F., Chemat, F., 2009. Clean recovery of antioxidant flavonoids from onions: optimising solvent free microwave extraction method. *Journal of Chromatography A* 1216, 7700–7707.
- Zsila, F., Bikádi, Z., Simonyi, M., 2003. Probing the binding of the flavonoid, quercetin to human serum albumin by circular dichroism, electronic absorption spectroscopy and molecular modelling methods. *Biochemical Pharmacology* 65, 447–456.

Comment citer ce document :

Khan, M. K. (Auteur de correspondance), Zill-E-Huma, Dangles, O. (2014). A comprehensive review on flavanones, the major citrus polyphenols. *Journal of Food Composition and Analysis*, 33, 85-104. DOI : 10.1016/j.jfca.2013.11.004