



**HAL**  
open science

## Draft Genome Sequence of the *Xanthomonas cassavae* Type Strain CFBP 4642.

Stéphanie S. Bolot, Alejandra Munoz Bodnar, Sébastien Cunnac, Erika Ortiz,  
Boris Szurek, Laurent Noel, Matthieu Arlat, Marie Agnes Jacques, Lionel  
Gagnevin, Perrine Portier, et al.

► **To cite this version:**

Stéphanie S. Bolot, Alejandra Munoz Bodnar, Sébastien Cunnac, Erika Ortiz, Boris Szurek, et al..  
Draft Genome Sequence of the *Xanthomonas cassavae* Type Strain CFBP 4642.. *Genome Announcements*, 2013, 1 (4), 10.1128/genomeA.00679-13 . hal-02642222

**HAL Id: hal-02642222**

**<https://hal.inrae.fr/hal-02642222v1>**

Submitted on 28 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Draft Genome Sequence of the *Xanthomonas cassavae* Type Strain CFBP 4642

Stéphanie Bolot,<sup>a,b</sup> Alejandra Munoz Bodnar,<sup>c</sup> Sébastien Cunnac,<sup>c</sup> Erika Ortiz,<sup>c</sup> Boris Szurek,<sup>c</sup> Laurent D. Noël,<sup>a,b</sup> Matthieu Arlat,<sup>a,b,d</sup> Marie-Agnès Jacques,<sup>e,f,g</sup> Lionel Gagnevin,<sup>h</sup> Perrine Portier,<sup>e,f,g,i</sup> Marion Fischer-Le Saux,<sup>e,f,g,i</sup> Sébastien Carrere,<sup>a,b</sup> Ralf Koebnik<sup>c</sup>

INRA, Laboratoire des Interactions Plantes Micro-organismes (LIPM), UMR 441, Castanet-Tolosan, France<sup>a</sup>; CNRS, LIPM, UMR 2594, Castanet-Tolosan, France<sup>b</sup>; UMR 186 IRD-Cirad-Université Montpellier 2 "Résistance des Plantes aux Bioagresseurs," Montpellier, France<sup>c</sup>; Université de Toulouse, Université Paul Sabatier, Toulouse, France<sup>d</sup>; INRA, UMR1345 Institut de Recherche en Horticulture et Semences, Angers, France<sup>e</sup>; Agrocampus Ouest, UMR1345 Institut de Recherche en Horticulture et Semences, Angers, France<sup>f</sup>; Université d'Angers, UMR1345 Institut de Recherche en Horticulture et Semences, SFR4207 QUASAV, PRES L'UNAM, Angers, France<sup>g</sup>; UMR "Peuplements Végétaux et Bioagresseurs en Milieu Tropical" (PVBMT), CIRAD, Saint-Pierre, La Réunion, France<sup>h</sup>; INRA, CIRM-CFBP Collection Française de Bactéries associées aux Plantes, Angers, France<sup>i</sup>

S.B. and A.M.B. contributed equally to this study.

**We report the draft genome sequence of the *Xanthomonas cassavae* type strain CFBP 4642, the causal agent of bacterial necrosis on cassava plants. These data will allow the comparison of this nonvascular pathogen with the vascular pathogen *Xanthomonas axonopodis* pv. *manihotis*, both infecting the same host, which will facilitate the development of diagnostic tools.**

Received 30 July 2013 Accepted 31 July 2013 Published 29 August 2013

**Citation** Bolot S, Munoz Bodnar A, Cunnac S, Ortiz E, Szurek B, Noël LD, Arlat M, Jacques M-A, Gagnevin L, Portier P, Fischer-Le Saux M, Carrere S, Koebnik R. 2013. Draft genome sequence of the *Xanthomonas cassavae* type strain CFBP 4642. *Genome Announc.* 1(4):e00679-13. doi:10.1128/genomeA.00679-13.

**Copyright** © 2013 Bolot et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Ralf Koebnik, koebnik@gmx.de.

Cassava (*Manihot esculenta* Crantz) is the third most important source of calories in the tropics, after rice and maize, and millions of people in Africa, Asia, and Latin America depend on cassava. South America, probably the Amazon region, is considered the center of origin for the cassava species. It was only in the 16th century that Portuguese navigators introduced cassava to the west coast of Africa, from where it later disseminated to East Africa. Most of the spread within the African continent, however, took place only during the 20th century due to colonial powers encouraging its cultivation (1). Nowadays, cassava is grown in all Sub-Saharan countries, and Africa produces more cassava than the rest of the world combined (1). Cassava plants are frugal with respect to environmental conditions (drought, poor soil) and hold great promise as a future staple crop in Africa, since this species might not only tolerate but even profit from climate change (2).

Cassava plants can suffer from two bacterial diseases, bacterial blight and bacterial necrosis, caused by two species of *Xanthomonas*. The causal agent of bacterial blight, *X. axonopodis* pv. *manihotis*, is a vascular pathogen which has been well studied over the last years (3). Recently, draft genome sequences of 65 strains have been elucidated (4). Much less is known about the nonvascular pathogen *X. cassavae*, which causes bacterial necrosis in Africa (5). The comparison of a vascular and a nonvascular pathogen, both infecting cassava, might give important clues about determinants of tissue specificity during colonization of the host plant. This prompted us to sequence the *X. cassavae* type strain CFBP 4642 (NCPPB 101, ICMP 204, LMG 673), which was isolated in Malawi in 1951.

Type strain CFBP 4642 was sequenced using the Illumina HiSeq2000 platform (GATC Biotech, Germany). The shotgun sequencing yielded 95,437,238 read pairs (64,851,255 100-bp

paired-end reads with an insert size of 250 bp and 30,585,983 50-bp mate-pair reads with an insert size of 3 kb). A combination of Velvet (6), SOAPdenovo, and SOAPGapCloser (7) yielded 83 contigs larger than 500 bp ( $N_{50}$ , 158,383 bp) with the largest contig of 425 kb for a total assembly size of 5,263,056 bp.

Multilocus sequence analysis of four housekeeping genes described earlier for xanthomonads (8) revealed that the four internal fragments are 99.94% identical (3,371/3,373) to those of the *X. cassavae* type strain ICMP 204 at PAMDB (9). The genome encodes a canonical type III protein secretion system (10) and several type III effectors, such as transcriptional activator-like (TAL) effectors (5). Interestingly, a CRISPR/*cas* defense system is present, which might be exploited to develop a powerful tool for Pan-African epidemiological surveillance (11).

**Nucleotide sequence accession number.** This whole-genome shotgun project has been deposited in GenBank under the accession no. [ATMC00000000](http://www.ncbi.nlm.nih.gov/nuccore/ATMC00000000).

## ACKNOWLEDGMENTS

This work was supported by grant ANR-2010-GENM-013 from the French Agence Nationale de la Recherche.

Strain CFBP 4642 is available at the CIRM-CFBP, French Collection for Plant-associated Bacteria, <http://www.angers-nantes.inra.fr/cfbp>.

## REFERENCES

- Hillocks RJ. 2002. Cassava in Africa, p 41–54. In Hillocks RJ, Thresh JM, Bellotti AC (ed), *Cassava: biology, production and utilization*. CAB International, Wallingford, Oxfordshire, United Kingdom.
- Jarvis A, Ramirez-Villegas J, Herrera Campo BV, Navarro-Racines C. 2012. Is cassava the answer to African climate change adaptation? *Trop. Plants Biol.* 5:9–29. doi:10.1007/s12042-012-9096-7.
- López CE, Bernal AJ. 2012. Cassava bacterial blight: using genomics for the elucidation and management of an old problem. *Trop. Plants Biol.* 5:117–126. doi:10.1007/s12042-011-9092-3.

4. Bart R, Cohn M, Kassen A, McCallum EJ, Shybut M, Petriello A, Krasileva K, Dahlbeck D, Medina C, Alicai T, Kumar L, Moreira LM, Rodrigues Neto J, Verdier V, Santana MA, Kositcharoenkul N, Vanderschuren H, Gruissem W, Bernal A, Staskawicz BJ. 2012. High-throughput genomic sequencing of cassava bacterial blight strains identifies conserved effectors to target for durable resistance. *Proc. Natl. Acad. Sci. U. S. A.* 109:E1972–E1979. doi:10.1073/pnas.1208003109.
5. Verdier V, Boher B, Maraite H, Geiger JP. 1994. Pathological and molecular characterization of *Xanthomonas campestris* strains causing diseases of cassava (*Manihot esculenta*). *Appl. Environ. Microbiol.* 60:4478–4486.
6. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829. doi:10.1101/gr.074492.107.
7. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu SM, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam TW, Wang J. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read *de novo* assembler. *Gigascience* 1:18. doi:10.1186/2047-217X-1-18.
8. Young JM, Park DC, Shearman HM, Fargier E. 2008. A multilocus sequence analysis of the genus *Xanthomonas*. *Syst. Appl. Microbiol.* 31:366–377. doi:10.1016/j.syapm.2008.06.004.
9. Almeida NF, Yan S, Cai R, Clarke CR, Morris CE, Schaad NW, Schuenzel EL, Lacy GH, Sun X, Jones JB, Castillo JA, Bull CT, Leman S, Guttman DS, Setubal JC, Vinatzer BA. 2010. PAMDB, a multilocus sequence typing and analysis database and website for plant-associated microbes. *Phytopathology* 100:208–215. doi:10.1094/PHYTO-100-3-0208.
10. Büttner D. 2012. Protein export according to schedule: architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. *Microbiol. Mol. Biol. Rev.* 76:262–310. doi:10.1128/MMBR.05017-11.
11. Horvath P, Barrangou R. 2010. CRISPR/Cas, the immune system of bacteria and archaea. *Science* 327:167–170. doi:10.1126/science.1179555.