



HAL
open science

Draft Genome Sequence of *Enterococcus faecalis* PC1.1, a Candidate Probiotic Strain Isolated from Human Feces.

Páraic O Cuív, Eline S. Klaassens, Wendy J. Smith, Stanislas Mondot, A. Scott Durkin, Derek M. Harkins, Les Foster, Jamison Mccorrison, Manolito Torralba, Karen E. Nelson, et al.

► To cite this version:

Páraic O Cuív, Eline S. Klaassens, Wendy J. Smith, Stanislas Mondot, A. Scott Durkin, et al.. Draft Genome Sequence of *Enterococcus faecalis* PC1.1, a Candidate Probiotic Strain Isolated from Human Feces.. *Genome Announcements*, 2013, 1 (1), 10.1128/genomeA.00160-12 . hal-02643385

HAL Id: hal-02643385

<https://hal.inrae.fr/hal-02643385>

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Draft Genome Sequence of *Enterococcus faecalis* PC1.1, a Candidate Probiotic Strain Isolated from Human Feces

Páraic Ó Cuív,^{a,b} Eline S. Klaassens,^{a,b*} Wendy J. Smith,^{a,b} Stanislas Mondot,^{a,b} A. Scott Durkin,^c Derek M. Harkins,^c Les Foster,^c Jamison McCarrison,^c Manolito Torralba,^c Karen E. Nelson,^c Mark Morrison^{a,b,d}

CSIRO Preventative Health Flagship Research Program, Queensland Biosciences Precinct, Queensland, Australia^a; CSIRO Animal, Food, and Health Sciences, Queensland Biosciences Precinct, Queensland, Australia^b; J. Craig Venter Institute, Rockville, Maryland, USA^c; The Ohio State University, Columbus, Ohio, USA^d

*Present address: Eline S. Klaassens, School of Molecular Bioscience, University of Sydney, Sydney, Australia.

***Enterococcus faecalis* is commonly isolated from the gastrointestinal tract of healthy infants and adults, where it contributes to host health and well-being. We describe here the draft genome sequence of *E. faecalis* PC1.1, a candidate probiotic strain isolated from human feces.**

Received 14 December 2012 Accepted 21 December 2012 Published 21 February 2013

Citation Ó Cuív P, Klaassens ES, Smith WJ, Mondot S, Durkin AS, Harkins DM, Foster L, McCarrison J, Torralba M, Nelson KE, Morrison M. 2013. Draft genome sequence of *Enterococcus faecalis* PC1.1, a candidate probiotic strain isolated from human feces. *Genome Announc.* 1(1):e00160-12. doi:10.1128/genomeA.00160-12.

Copyright © 2013 Ó Cuív et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Mark Morrison, markmorrison@csiro.au.

Enterococcus faecalis is a facultative anaerobic bacterium that is a primary colonizer of the infant gastrointestinal tract and is also prevalent in the adult gastrointestinal tract (1–3). *E. faecalis* is typically considered to be a beneficial gut bacterium that has been shown to be capable of modulating the activities of specific animal pathogens (4, 5) and the host immune response (6, 7); consequently, several strains have been applied as probiotics for the maintenance of a healthy gut and the treatment of gut diseases, including diarrhea, irritable bowel syndrome, and ulcerative colitis (8–10). Despite this, *E. faecalis* is now also widely recognized as a potential gut pathogen that is capable of compromising the epithelial barrier (11), inducing inflammation in mouse models of experimental colitis (12, 13) and causing chromosomal instability in colonic epithelial cells (14, 15). Key intraspecies variations with relevance to pathogenesis exist, and thus, we sought to elucidate further the genomic potential of an *E. faecalis* isolate from a healthy human gut and to assess its potential as a probiotic candidate. We describe here the draft genome sequence of *E. faecalis* PC1.1, isolated as part of the Australian Human Gut Microbiome Project.

E. faecalis PC1.1 was isolated from a pooled fecal sample collected from healthy human subjects by plating aerobically on brain heart infusion (BHI) medium. Genomic DNA was prepared, and a 454 Life Sciences Genome Sequencer (GS) FLX system was used at the J. Craig Venter Institute (JCVI) to generate 2,753,924 bp of DNA sequence at 30× coverage. The individual sequence reads were assembled using the Newbler Assembler v2.3 to generate 79 contigs. The contig N₅₀ was approximately 57.3 kb and the largest contig assembled was approximately 132.5 kb. Finally, the DNA sequences were annotated using the JCVI prokaryotic annotation pipeline.

The draft genome has a G+C content of 37.6% and contains 2,673 genes, with 2,619 protein-coding genes and 54 structural RNAs. As anticipated, the annotated genome of *E. faecalis* PC1.1 revealed the presence of factors relevant to the colonization of and

persistence in the human gut, including protein orthologs with putative roles in mediating adhesion to host structural factors, including collagen, fibronectin, and fibrinogen. In addition, we identified a putative lectin with a suspected role in mediating adhesion to host structural glycans and also several proteins with likely roles in harvesting host glycans. The *E. faecalis* PC1.1 genome encodes a putative gelatinase and serine proteinase; however, it does not appear to encode an enterococcal surface protein or enterococcal cytolysin, which have been shown previously to be important for virulence.

This preliminary study provides insight into the genomic factors that contribute to the colonization of and persistence in the gut. Nontoxicogenic *Clostridium difficile* can effectively colonize the gut of hamsters and humans and prevent colonization by toxicogenic *C. difficile* strains (reviewed in reference 16); further studies will be necessary to determine the probiotic potential of *E. faecalis* PC1.1, including its ability to similarly prevent colonization and/or modulate the activities of pathogenic microbes and to productively influence the immune response.

Nucleotide sequence accession numbers. This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [ADKN000000000](https://www.ncbi.nlm.nih.gov/nuccore/ADKN000000000). The version described here is the first version, [ADKN010000000](https://www.ncbi.nlm.nih.gov/nuccore/ADKN010000000).

ACKNOWLEDGMENTS

This research was performed as part of CSIRO's contribution to the International Human Microbiome Consortium. This project was supported by the Commonwealth of Australia under the International Science Linkages program, by CSIRO's Transformational Biology Capability Platform in support of the Preventative Health Flagship Research Program and NIH award number U54AI084844 to Karen E. Nelson. We gratefully acknowledge the support provided by CSIRO's OCE Science Leader scheme (to M.M.) and the OCE Postdoctoral Fellow program (awarded to E.K.).

We thank CSIRO Human Nutrition for collecting the fecal samples and Seungha Kang and Desma Grice for providing critical reading and suggestions to improve the manuscript.

REFERENCES

- Hopkins MJ, Macfarlane GT, Furrie E, Fite A, Macfarlane S. 2005. Characterisation of intestinal bacteria in infant stools using real-time PCR and northern hybridisation analyses. *FEMS Microbiol. Ecol.* 54:77–85.
- Kirtzalidou EI, Mitsou EK, Pramateftaki P, Kyriacou A. 2012. Screening fecal enterococci from Greek healthy infants for susceptibility to antimicrobial agents. *Microb. Drug Resist.* 18:578–85.
- Noble CJ. 1978. Carriage of group D streptococci in the human bowel. *J. Clin. Pathol.* 31:1182–1186.
- Cebrián R, Baños A, Valdivia E, Pérez-Pulido R, Martínez-Bueno M, Maqueda M. 2012. Characterization of functional, safety, and probiotic properties of *Enterococcus faecalis* UGRA10, a new AS-48-producer strain. *Food Microbiol.* 30:59–67.
- Soerjadi AS, Lloyd AB, Cumming RB. 1978. *Streptococcus faecalis*, a bacterial isolate which protects young chickens from enteric invasion by salmonellae. *Aust. Vet. J.* 54:549–550.
- Are A, Aronsson L, Wang S, Greicius G, Lee YK, Gustafsson JA, Pettersson S, Arulampalam V. 2008. *Enterococcus faecalis* from newborn babies regulate endogenous PPAR γ activity and IL-10 levels in colonic epithelial cells. *Proc. Natl. Acad. Sci. U. S. A.* 105:1943–1948.
- Wang S, Ng LH, Chow WL, Lee YK. 2008. Infant intestinal *Enterococcus faecalis* down-regulates inflammatory responses in human intestinal cell lines. *World J. Gastroenterol.* 14:1067–1076.
- Chen CC, Kong MS, Lai MW, Chao HC, Chang KW, Chen SY, Huang YC, Chiu CH, Li WC, Lin PY, Chen CJ, Li TY. 2010. Probiotics have clinical, microbiologic, and immunologic efficacy in acute infectious diarrhea. *Pediatr. Infect. Dis. J.* 29:135–138.
- Enck P, Zimmermann K, Menke G, Müller-Lissner S, Martens U, Klosterhalfen S. 2008. A mixture of *Escherichia coli* (DSM 17252) and *Enterococcus faecalis* (DSM 16440) for treatment of the irritable bowel syndrome—a randomized controlled trial with primary care physicians. *Neurogastroenterol. Motil.* 20:1103–1109.
- Tsuda Y, Yoshimatsu Y, Aoki H, Nakamura K, Irie M, Fukuda K, Hosoe N, Takada N, Shirai K, Suzuki Y. 2007. Clinical effectiveness of probiotics therapy (bio-three) in patients with ulcerative colitis refractory to conventional therapy. *Scand. J. Gastroenterol.* 42:1306–1311.
- Steck N, Hoffmann M, Sava IG, Kim SC, Hahne H, Tonkonogy SL, Mair K, Krueger D, Pruteanu M, Shanahan F, Vogelmann R, Schemann M, Kuster B, Sartor RB, Haller D. 2011. *Enterococcus faecalis* metalloprotease compromises epithelial barrier and contributes to intestinal inflammation. *Gastroenterology* 141:959–971.
- Balish E, Warner T. 2002. *Enterococcus faecalis* induces inflammatory bowel disease in interleukin-10 knockout mice. *Am. J. Pathol.* 160:2253–2257.
- Kim SC, Tonkonogy SL, Albright CA, Tsang J, Balish EJ, Braun J, Huycke MM, Sartor RB. 2005. Variable phenotypes of enterocolitis in interleukin 10-deficient mice monoassociated with two different commensal bacteria. *Gastroenterology* 128:891–906.
- Huycke MM, Abrams V, Moore DR. 2002. *Enterococcus faecalis* produces extracellular superoxide and hydrogen peroxide that damages colonic epithelial cell DNA. *Carcinogenesis* 23:529–536.
- Wang X, Huycke MM. 2007. Extracellular superoxide production by *Enterococcus faecalis* promotes chromosomal instability in mammalian cells. *Gastroenterology* 132:551–561.
- Gerding DN. 2012. *Clostridium difficile* infection prevention: biotherapeutics, immunologics, and vaccines. *Discov. Med.* 13:75–83.