



**HAL**  
open science

## Linking carbon and water cycles using stable isotopes across scales: progress and challenges

Christiane Werner, Franz Badeck, Enrico Brugnoli, Brian Cohn, Matthias Cuntz, Todd Dawson, Arthur Gessler, Jaleh Ghashghaie, Thorsten Grams, Zachary Kayler, et al.

### ► To cite this version:

Christiane Werner, Franz Badeck, Enrico Brugnoli, Brian Cohn, Matthias Cuntz, et al.. Linking carbon and water cycles using stable isotopes across scales: progress and challenges. *Biogeosciences Discussions*, 2012, 8 (8), pp.2659-2719. 10.5194/bgd-8-2659-2011 . hal-02643686

**HAL Id: hal-02643686**

**<https://hal.inrae.fr/hal-02643686>**

Submitted on 28 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

## Linking carbon and water cycles using stable isotopes across scales: progress and challenges

**C. Werner<sup>1</sup>, F. Badeck<sup>2</sup>, E. Brugnoli<sup>3</sup>, B. Cohn<sup>4</sup>, M. Cuntz<sup>5</sup>, T. Dawson<sup>6</sup>,  
A. Gessler<sup>7</sup>, J. Ghashghaie<sup>8</sup>, T. E. E. Grams<sup>9</sup>, Z. Kayler<sup>7</sup>, C. Keitel<sup>10</sup>,  
M. Lakatos<sup>11</sup>, X. Lee<sup>12</sup>, C. Máguas<sup>13</sup>, J. Ogée<sup>14</sup>, K. G. Rascher<sup>1</sup>, H. Schnyder<sup>15</sup>,  
R. Siegwolf<sup>16</sup>, S. Unger<sup>1</sup>, J. Welker<sup>4</sup>, L. Wingate<sup>17</sup>, and M. J. Zeeman<sup>18</sup>**

<sup>1</sup>Experimental and Systems Ecology, University Bielefeld, Universitätsstr. 25, 33615 Bielefeld, Germany

<sup>2</sup>Potsdam Institute for Climate Impact Research (PIK) PF 60 12 03 – 14412 Potsdam, Germany

<sup>3</sup>CNR – National Research Council of Italy – Institute of Agro-environmental and Forest Biology, via Marconi 2, 05010 Porano (TR), Italy

<sup>4</sup>Environment and Natural Resources Institute, University of Alaska Anchorage, 3211 Providence Dr. Anchorage, AK 99508-4614, USA

<sup>5</sup>UFZ – Helmholtz Centre for Environmental Research, Permoserstr. 15, 04318 Leipzig, Germany

2659

<sup>6</sup>Center for Isotope Biogeochemistry, Department of Integrative Biology, University of California, Berkeley, CA 94720, USA

<sup>7</sup>Institute for Landscape Biogeochemistry Leibniz-Zentrum für Agrarlandschaftsforschung (ZALF) e.V., Eberswalderstr. 84, 15374 Müncheberg

<sup>8</sup>Laboratoire d'Ecologie, Systématique et Evolution (ESE), CNRS AgroParisTech-UMR 8079, Bâtiment 362, Université de Paris-Sud (XI), 91405 Orsay Cedex, France

<sup>9</sup>Ecophysiology of Plants, Department of Ecology and Ecosystem Management, Technische Universität München, Von-Carlowitz-Platz 2, 85354 Freising, Germany

<sup>10</sup>University of Sydney, Faculty of Agriculture, Food and Natural Resources, 1 Central Avenue, Eveleigh, NSW 2015, Australia

<sup>11</sup>Experimental Ecology, University of Kaiserslautern, Erwin-Schroedinger Str. 13, 67663 Kaiserslautern, Germany

<sup>12</sup>School of Forestry and Environmental Studies, Yale University, New Haven, CT 06511, USA

<sup>13</sup>Centre for Environmental Biology, University of Lisbon, Campo Grande, 1749-016, Lisbon, Portugal

<sup>14</sup>UR 1263, INRA, Bordeaux, France

<sup>15</sup>Lehrstuhl für Grünlandlehre, Technische Universität München, 85350 Freising-Weißenstephan, Germany

<sup>16</sup>Lab for Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen-PSI, Switzerland

<sup>17</sup>Department of Plant Sciences, University of Cambridge, Cambridge, UK

<sup>18</sup>College of Oceanic and Atmospheric Sciences, Oregon State University, 104 COAS Admin Bldg, Corvallis (OR), USA

Received: 18 February 2011 – Accepted: 3 March 2011 – Published: 15 March 2011

Correspondence to: C. Werner (c.werner@uni-bielefeld.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.



## 2 Isotope effects across temporal and spatial scales

### 2.1 Leaf-level processes

#### 2.1.1 Recent advances on fractionation during leaf CO<sub>2</sub> and H<sub>2</sub>O exchange

Leaf CO<sub>2</sub> and H<sub>2</sub>O exchange is associated with isotopic fractionation, which has been widely used to investigate environmental effects on the physiology of photosynthesis and stomatal conductance (see Farquhar et al., 1989; Brugnoli and Farquhar 2000). Photosynthetic carbon isotope fractionation ( $\Delta^{13}\text{C}$ ) is primarily linked to variations in the ratio of leaf intercellular and ambient CO<sub>2</sub> concentration ( $c_i/c_a$ ) and the principle fractionations during diffusion and carboxylation (Farquhar et al., 1982). However, recent studies measuring diurnal variations in  $\Delta^{13}\text{C}$  on-line at high resolution have found that fractionations during photorespiration, day respiration and internal transfer of CO<sub>2</sub> cause the linear relationship between  $\Delta^{13}\text{C}$  and  $c_i/c_a$  to diverge at both high and low values of  $c_i/c_a$  (e.g. Wingate et al., 2007; Bickford et al., 2009). Much of this discrepancy can be resolved using the more comprehensive description of  $\Delta^{13}\text{C}$  as proposed by Farquhar et al. (1982) that quantitatively accounts for the isotopic fractionation during photorespiration ( $f = 11$ , Lanigan et al., 2008), during CO<sub>2</sub> diffusion within the leaf mesophyll (e.g. Wingate et al., 2007; Schäufele et al., 2010) and dark respiration (Ghashghaie et al., 2003; see also 2.1.2). Increasing evidence indicates that the internal mesophyll conductance ( $g_i$ ) may vary in response to several environmental factors (e.g. light, temperature, drought, salinity and CO<sub>2</sub>, see Flexas et al., 2008) that are also detectable at the canopy-scale (Schäufele et al., 2010). This reversible or adaptive component appears to be associated with variations in the conductance of membranes (Schäufele et al., 2010), that may be modified by the expression of cooporins (Hanba et al., 2004; Flexas et al., 2007): aquaporins capable of transporting CO<sub>2</sub> across plasma membranes. Although the dynamic role of  $g_i$  at the timescale of minutes to days remains somewhat controversial it is certain that  $g_i$  will vary markedly over the season with changes in leaf morphology and amongst and between species (Warren and Adams, 2006).

2663

Furthermore, our understanding on the contribution of day respiration to  $\Delta^{13}\text{C}$  dynamics is developing. Recent studies both in the field and laboratory, using either high resolution on-line  $\Delta^{13}\text{C}$  approaches (Wingate et al., 2007, 2010a; Bickford et al., 2009) or substrate labelling (Tcherkez et al., 2010), indicate that the isotopic composition of the day respiration flux is very unlikely the same as of recent photoassimilates (as was assumed in the original model of Farquhar et al., 1982). Instead, older carbon pools partially fuel this flux necessitating the need for improved models of  $\Delta^{13}\text{C}$  that account for this process (Wingate et al., 2007; Tcherkez et al., 2010) when modeling dynamic isofluxes in the field and importantly when estimating  $g_i$  using  $\Delta^{13}\text{C}$  approaches.

During photosynthetic CO<sub>2</sub> uptake there is also an exchange of oxygen isotopes between CO<sub>2</sub> and leaf water. This results in an oxygen isotope photosynthetic discrimination ( $\Delta^{18}\text{O}$ ) that can be extremely dynamic as it integrates variations in the bi-directional fluxes of CO<sub>2</sub> through the stomatal pores and in the  $\delta^{18}\text{O}$  composition of leaf water (Farquhar et al., 1993). The impact of terrestrial biosphere  $\Delta^{18}\text{O}$  on the atmosphere is highlighted in the strong Northern hemisphere meridional gradient and seasonal variations of  $\delta^{18}\text{O}$  in CO<sub>2</sub> observed by the global flask network (Farquhar et al., 1993; Cuntz et al., 2003a, b; Wingate et al., 2009; Francey and Tans, 1987).

Current leaf water isotopic enrichment models are still based on the work of Craig and Gordon (1965) and Dongmann et al. (1974) and describe how H<sub>2</sub><sup>18</sup>O enrichment in leaf water develops as a function of the  $\delta^{18}\text{O}$  in source water (water extracted from the soil by plants) and water vapor, and the ratio of vapor pressure between the atmosphere and the leaf intercellular spaces (i.e. relative humidity expressed relative to leaf temperature). Over the course of a sunny day, leaf water becomes enriched as the air gets dryer and leaf temperature increases leading to higher vapor pressure deficits between the leaf and the air, the driving force of water flux from the soil to the atmosphere. For leaves that have a high leaf water content such as conifer needles, a non-steady-state formulation is required because the turnover time of leaf water is too long to let the water pool readjust to the rapidly varying atmospheric conditions (Cernusak et al., 2002; Seibt et al., 2006, 2007; Cuntz et al., 2007; Wingate et al., 2010b).

2664

Also, most studies seem to indicate that the mixing of enriched water from the sites of evaporation in the leaf mesophyll is incomplete, and that some un-enriched (or only partially-enriched) water pools remain in the mesophyll (Yakir et al., 1998; Farquhar and Lloyd, 1993). However the way this incomplete mixing operates is still a matter of debate (Barbour et al., 2004; Ogée et al., 2007; Ferrio et al., 2009; Kahmen et al., 2009) and difficult to model as it requires knowledge on where precisely in the mesophyll water evaporates or condenses and how water molecules move and self-diffuse (Farquhar et al., 2003; Cuntz et al., 2007; Ogée et al., 2007). For this reason, current models of leaf water enrichment have not markedly evolved since the work of Dongmann et al. (1974) or Farris and Strain (1978). Similarly, and also because of a lack of understanding in the extent of isotopic exchange between CO<sub>2</sub> and water in the leaf especially in C<sub>4</sub> plants, models of oxygen isotope discrimination during CO<sub>2</sub> uptake by leaves are still based on the earlier theory of Farquhar and Lloyd (1993).

However, with new laser spectroscopic techniques it is now possible to capture leaf water enrichment and CO<sub>2</sub>-H<sub>2</sub>O exchange dynamically by measuring on-line either the δ<sup>18</sup>O of the transpiration flux (Welp et al., 2009) or the net CO<sub>2</sub> exchange from leaves (Barbour et al., 2007; Wingate et al., 2010b). By inverting either of these flux signals it is possible to reconstruct leaf water dynamics non-invasively, and to obtain new insights regarding the extent of equilibration between CO<sub>2</sub> and water for different species or environmental conditions. Therefore, the δ<sup>18</sup>O of the CO<sub>2</sub> flux is an extremely powerful tracer as δ<sup>18</sup>O of leaf water can be reconstructed both during the day and night. This approach has provided compelling evidence for the opening of stomata in the dark (Cernusak et al., 2004; Seibt et al., 2007; Wingate et al., 2010b) and may also indicate subtle variations in the extent of CO<sub>2</sub> hydration by the enzyme carbonic anhydrase between light and dark periods (Cernusak et al., 2004; Barbour et al., 2007; Kahmen et al., 2008). Thus, δ<sup>18</sup>O of CO<sub>2</sub> will prove a valuable tracer as we try to understand why stomata open in the dark and how rates of nocturnal stomatal conductance vary over the night and from day-to-day.

2665

### 2.1.2 Current view on post-carboxylation and respiratory fractionation

The isotope effects during leaf CO<sub>2</sub> and H<sub>2</sub>O exchange leave an imprint on the δ<sup>13</sup>C and δ<sup>18</sup>O signatures of newly produced assimilates, tissues and respired CO<sub>2</sub> (Farquhar et al., 1982). However, several post-carboxylation fractionation processes can alter the isotope signatures of the assimilated carbon compounds in down-stream metabolic processes.

Already within the Calvin cycle, isotopic fractionation occurs mainly due to metabolic branching points and the use of triose phosphates that can either be exported to the cytosol or continue to be used within the Calvin cycle. The triose phosphates that are not exported are subject to enzyme catalyzed reactions (aldolisation and transketolisation) with position specific discrimination during C-C bond making. As a result, the C-3 and C-4 positions within glucose are enriched in <sup>13</sup>C and thus a non-uniform intra-molecular distribution of <sup>13</sup>C within carbohydrates is established (Rossmann et al., 1991; Tcherkez et al., 2004; Gilbert et al., 2009). Subsequently, photorespiration and starch-sucrose partitioning modify the diurnal changes in photosynthetic discrimination (see Brüggemann et al., 2011). Analyses of sugar δ<sup>13</sup>C and its diurnal variations have the potential to trace changes in these metabolic activities.

The non-statistical intra-molecular distribution of <sup>13</sup>C in carbohydrates leaves its imprints on metabolites synthesized from these ("fragmentation fractionation" Tcherkez et al., 2004). If one of these products is decarboxylated then respired CO<sub>2</sub> will carry an isotopic signature different from the average sugar signature (see Brüggemann et al., 2011). Moreover, switches between substrates (Tcherkez et al., 2003), in particular during light-dark transition of leaves (i.e. light enhanced dark respiration, Barbour et al., 2007) and the oxidative pentosephosphate pathway can markedly change the δ<sup>13</sup>C of respired CO<sub>2</sub> (Bathellier et al., 2008, 2009; see Brüggemann et al., 2011). So far a full quantitative understanding of apparent respiratory fractionation has not yet been achieved (Tcherkez 2010; Werner 2010). However, measurements with a high temporal resolution indicated remarkable diel dynamics in respiratory δ<sup>13</sup>CO<sub>2</sub>, which

2666

differed between functional plant groups (Priault et al., 2009; Werner et al., 2009; reviewed in Werner and Gessler, 2011). Feeding experiments with positionally labeled glucose or pyruvate can trace changes in carbon partitioning in the metabolic branching points of the respiratory pathways (Tcherkez et al., 2004), which has been used to identify differences between functional groups (Priault et al., 2009; Wegener et al., 2010). For studies of changes in respiratory substrates measurements of respired CO<sub>2</sub> and its isotopic composition can be combined with measurements of oxygen exchange (Tcherkez et al., 2003).

The main post-photosynthetic changes in  $\delta^{18}\text{O}$  of carbohydrates are due to exchange with water pools in other plant compartments (e.g. xylem, phloem, sink tissues) or in the course of time. Thus, the isotopic signature of local water pools will be imprinted on the carbohydrate signature (Farquhar et al., 1998). The equilibration with these pools depends on the residence times within these compartments as well as on the stability of the chemical bonds of the oxygen atoms in individual positions from exchange by e.g. phosphorylation. Recent advances have further led to the improvement of process-based models that predict variability in the oxygen isotope composition of plant organic material (Barbour, 2007). These models are now employed to quantitatively relate stable carbon and oxygen isotopes to plant photosynthetic performance, e.g. leaf physiological responses to environmental changes (cf. Gessler et al., 2009a; Kahmen et al., 2008, 2009, 2011).

Further carbon and oxygen isotope fractionation steps occur during secondary metabolism (see Schmidt and Gleixner, 1998; Schmidt 2003; Schmidt et al., 2001). Fragmentation fractionation and diel variation in anabolic pathways and substrates used for between-organ transport are at the origin of inter-organ differences in isotopic composition (see Hobbie and Werner, 2004; Badeck et al., 2005; Cernusak et al., 2009 for reviews). Future progress in quantifying these effects will open new avenues for analyses of inter-organs differences as indicators of metabolic activities and their variation with plant phenology.

2667

### 2.1.3 Bulk leaf tissue $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ and water use efficiency

Bulk leaf carbon isotope discrimination ( $\Delta^{13}\text{C}$ , i.e. the difference of the leaf organic matter  $\delta^{13}\text{C}$  relative to the atmospheric source  $\delta^{13}\text{C}_a$ ; divided by  $\delta^{13}\text{C}_a$ ) provides an integrative record of supply and demand for CO<sub>2</sub> and can be used to determine the relationship between carbon gain versus water loss. As such, variation in  $\Delta^{13}\text{C}$  may be driven by changes in stomatal conductance (i.e. supply of CO<sub>2</sub>), in photosynthetic carboxylation rate (i.e. demand for CO<sub>2</sub>), or both. In spite of the above mentioned multiple fractionation processes a remarkably linear relationship between bulk leaf  $\Delta^{13}\text{C}$  and the intrinsic water use efficiency (WUE<sub>i</sub>), i.e. the ratio of net assimilation ( $P_N$ ) to stomatal conductance ( $g_s$ ) has been observed under controlled conditions (Farquhar et al., 1989). The use of  $\Delta^{13}\text{C}$  is therefore a convenient tool to study the variation in carbon water relations retrospectively, e.g. in different biological archives (see 2.5), and has been vastly used as an ecological tracer of water use efficiency at the leaf, plant and ecosystem scale (e.g. Bonal et al., 2000; Evans 2001; Lauteri et al., 2004; Ponton et al., 2006).

However, the carbon isotope ratio alone will not provide information, whether a change in  $\Delta^{13}\text{C}$  was caused by either a change in  $g_s$  or  $P_N$ . This can be answered by combining the carbon isotope ratio with information from the oxygen isotope ratio in plant organic matter (Scheidegger et al., 2000; see also Grams et al., 2007; Sullivan and Welker, 2007). The negative relationship between  $\Delta^{18}\text{O}$  and  $g_s$  was used to estimate stomatal aperture for interpreting physiological changes in  $\delta^{13}\text{C}$  of leaf material.  $\delta^{18}\text{O}$  of bulk organic matter has further been used to determine whether a change in WUE<sub>i</sub> is the result of an increase in the atmospheric CO<sub>2</sub> (Saurer and Siegwolf, 2007), or a change in VPD as a result of an increasing temperature at constant or decreasing precipitation, as it can be observed in the Mediterranean (Sarris et al., 2011) or along a Siberian North – South gradient (Siegwolf et al., unpubl.). These results suggest that the dual isotope approach allows the distinction between the contributions of  $g_s$  and/or photosynthetic capacity to changes in  $\delta^{13}\text{C}$  at the site level.

2668

However, under naturally varying environmental conditions interpretation of  $\Delta^{13}\text{C}$  in terms of  $\text{WUE}_i$  is complex, particularly when only carbon isotope information is available (e.g. Welker et al., 2003). Under natural conditions,  $\Delta^{13}\text{C}$  variations will not deliver straight-forward information on the “efficient” use of carbon and/or water resources but solely reflects changes in the  $P_N/g_s$  ratio. Thus, ecological interpretation must be cautioned if the actual rates of these processes (carbon gain and water loss, the latter requiring knowledge on VPD) are not known. Moreover, under natural conditions the trade-off between efficient carbon gain and reduction of water loss is particularly relevant when species are competing for limited water sources. Thus, ecological evaluation of an efficient resource use strategy requires knowledge on resource availability and plant interaction at the community level.

Moreover, different phenological phases and leaf structure (through its effect in mesophyll conductance, see 2.1.1) can constrain a direct comparison of bulk leaf  $\Delta^{13}\text{C}$  between different species and functional groups (Hanba et al., 2003; Warren and Adams, 2006; Werner and Máguas, 2010). The influence of mesophyll conductance and other post-photosynthetic fractionation factors may also be relevant when comparing the isotopic composition of different plant compartments (e.g., leaves, carbohydrates, phloem, roots and root exudates, Bathellier et al., 2009; Cernusak et al., 2009; Wegener et al., 2010).

For the Mediterranean evergreen sclerophyllous species deviations from the expected linear relationship between leaf  $\Delta^{13}\text{C}$  and  $c_i/c_a$  ratio are observed (see regression line Fig. 2). The expected reduction in the  $c_i/c_a$  ratio in response to increasing VPD, drought and temperatures was not manifested in leaf  $\Delta^{13}\text{C}$  (Fig. 2), due to different phenological phases, a cease of growth (in evergreens) and concomitant reduction in  $g_s$  and  $P_N$  (as identified by  $\delta^{18}\text{O}$ , Werner et al., 2001), structural regulation of plant water loss by leaf abscission (semi-malacophyllous species) and low carboxylation rates in the herbaceous species. Hence, environmental conditions during leaf formation as well as length of the growing period, leaf longevity, structure, and life form do modulate  $\Delta^{13}\text{C}$ . Furthermore, the remobilization of non-structural carbohydrates from older tissues and

2669

fractionation processes during the formation of new leaves might mask the  $\Delta^{13}\text{C}$  signal induced by environmental conditions under which the leaf was formed (Terwilliger et al., 2001).

Thus, while the dual isotope approach has proven a valuable concept for ecological applications, an interpretation of  $\text{WUE}_i$  based exclusively on  $\Delta^{13}\text{C}$  in bulk leaf material requires a careful consideration of the type of plant material and the potential environmental conditions. The complexity of this parameter may constrain its use in ecological field studies. Carbon pools with shorter turn-over times such as leaf soluble sugars or phloem allocated carbon (e.g. Gessler et al., 2004; Scartazza et al., 2004) are therefore better indicators for recent changes in  $c_i/c_a$  and environmental factors as outlined below (2.2).

## 2.2 $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ isotopes to trace plant integrated processes and plant-soil coupling

The transport of newly assimilated carbon within the plant and from the plant to the rhizosphere can be characterized by applying stable isotope techniques. For plant and canopy integrating information as well as for plant internal transport the assessment of the isotopic composition of phloem sugars is central.

### 2.2.1 Carbon and oxygen isotopic composition of phloem organic matter

For almost 15 years  $\delta^{13}\text{C}$  of phloem organic matter has been used to derive information on C-allocation and canopy integrated water use efficiency in plants as affected by environmental conditions (Yoneyama et al., 1997; Pate and Arthur, 1998; Keitel et al., 2003; Gessler et al., 2004; Scartazza et al., 2004; Barbour et al., 2005; Rascher et al., 2010). A dual isotope approach ( $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  see 2.1.3) can also be successfully applied to phloem sugars to distinguish whether net assimilation and/or stomatal conductance is changing as a result of environmental conditions (Keitel et al., 2003; Cernusak et al., 2003, 2005; Brandes et al., 2006; Keitel et al., 2006; Gessler et al.,

2670

2007b, 2008). However, it is now apparent that – even though the isotope composition of phloem organic matter can, in principle, integrate leaf physiology over the whole canopy – several points of uncertainty must be taken into account when interpreting phloem isotopic information.

- 5 1. Temporal integration: Short-term variations in the isotopic composition of leaf sugars – induced by either an environmental signal or internal plant processes – might or might not be reflected in the phloem organic matter. Twig phloem organic matter of trees (e.g. Gessler et al., 2007a) and the stem phloem of herbaceous species (e.g. Gessler et al., 2008) can be applied to monitor diel variation of evaporative  $^{18}\text{O}$  and  $^2\text{H}$  enrichment or carbon isotope fractionation. In the trunks of adult trees, however, the mixing of sugars of different metabolic origins dampens the short-term variations and the isotope signatures provide time-integrated information on canopy processes instead (Keitel et al., 2006; Rascher et al., 2010; Wingate et al., 2010a).
- 15 2. Change of the original signal: A further constraint in interpreting phloem isotope signatures is that the original isotope signal imprinted on sugars in the leaf may be altered during basipetal transport in the phloem of trees. The transport of sugar molecules itself does not fractionate to a measurable extent. However, carbon fixation by PEPc in the bark and oxygen atom exchange with stem water during metabolic processes in the stem tissue together with the continuous unloading and loading of sugars from and to the phloem might lead to the observed isotope patterns (Barnard et al., 2007; Gessler et al., 2009b). The change in  $\delta^{13}\text{C}$  along the transport path, however, varies strongly among species ranging from  $^{13}\text{C}$  enrichment (Brandes et al., 2006; Wingate et al., 2010a) and no change in  $\delta^{13}\text{C}$  (Pate and Arthur, 1998; Gessler et al., 2007a) to  $^{13}\text{C}$  depletion (Rascher et al., 2010). The nature of these species-specific differences remains to be clarified and might shed new light on assimilate partitioning in trees.

2671

- 5 3. Chemical composition: It is often assumed that only one major sugar, namely sucrose, is present in the phloem. However, besides sucrose, there are other transport carbohydrates such as myo-inositol and raffinose family sugars (Karner et al., 2004) and also hexoses present (van Bel and Hess, 2008). Phloem sugar composition varies with environmental conditions which could be one important factor for changes in phloem  $\delta^{13}\text{C}$  (Merchant et al., 2010), independent of the original leaf level isotope signal, since  $\delta^{13}\text{C}$  differs between different carbohydrate molecules (Schmidt, 2003; Devaux et al., 2009). Compound specific analysis, provided by state of the art LC- and GC-IRMS techniques, might help to differentiate between changes in phloem  $\delta^{13}\text{C}$  that result from changes in the chemical composition or changes in leaf level fractionation. In addition, we should apply comparable methods to characterize the compound specific oxygen isotope composition of phloem organic matter.

15 Thus, the isotopic composition of phloem organic matter is a useful tool for gaining physiological information on whole tree or canopy processes (Cernusak et al., 2009; Gessler et al., 2009b). There is, however, a strong need to develop more insights into the underlying mechanisms behind the spatio-temporal variation of  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  independent of leaf level processes. This is a necessary prerequisite not only to better interpret the physiological meaning of the phloem isotope signal but also to extend our understanding of whole plant carbon transport and the metabolic background behind variations in ecosystem carbon allocation.

### 2.2.2 Plant-Soil carbon coupling

25 Stable isotope approaches did enhance our awareness of tight plant-soil coupling of carbon and water flow in recent years (see accompanying paper by Brüggemann et al., 2011). Besides the application of  $\delta^{13}\text{C}$  (or even radioactive  $\delta^{14}\text{C}$ ) enriched  $\text{CO}_2$  in pulse labelling experiments (e.g. Högberg et al., 2008; Carbone et al., 2007), natural abundance  $\delta^{13}\text{C}$  approaches have been applied to characterize the coupling between

2672





atmospheric CO<sub>2</sub> isotope measurements are rarely met (Pataki et al., 2003; Kayler et al., 2010a), or that isotopic steady-state is not attained (Nickerson and Risk, 2009b; Kayler et al., 2010b). Recently, an increasing number of studies have shown that factors such as diffusivity of soil CO<sub>2</sub>, dissolution of CO<sub>2</sub> from respiration and bicarbonates and advection of soil gas may be responsible for strong isotopic disequilibria between the CO<sub>2</sub> efflux at the soil surface and concurrent soil respiration (Kayler et al., 2008, 2010a; Nickerson and Risk, 2009a; Ohlsson, 2009; Gamnitzer et al., 2011). To characterize subsurface gas transport and subsequent fractionation dynamics, soil CO<sub>2</sub> production and transport models are used (Hashimoto, 2002; Davidson et al., 2006; Nickerson and Risk, 2009c; Ohlsson, 2009; Moyes et al., 2010; Wingate et al., 2010b), but many of the above discussed processes are not yet included (e.g. Brüggemann et al., 2011). Similarly, the oxygen isotope composition of soil respired CO<sub>2</sub> ( $\delta^{18}\text{O}_S$ ) not only carries the isotopic signature of the soil water it interacted with, but has been shown to be influenced by the carbonic anhydrase present in soil microorganisms that enhances isotopic equilibration between CO<sub>2</sub> and soil water (Wingate et al., 2009; 2010b). Yet, such processes are rarely included in isotope-enabled ecosystem models.

Obviously, the main lesson learned from these recent studies is that the application of the stable isotope approach at the ecosystem scale to constrain carbon and water fluxes from different pools is more complex than previously thought. These studies also allowed us to improve our understanding of the physical and biological drivers of the isotopic signals for each of the component fluxes. This knowledge can now be imbedded into stable isotope-enabled models and must be combined with data assimilation techniques. The recent development of continuous, rapid in situ stable isotope measurement techniques (see Sect. 3) opens new possibilities to study the carbon and water budgets of terrestrial ecosystems. The assimilation of such high time resolution datasets into isotope-enabled ecosystem models can be used to improve models by constraining component fluxes when non-isotopic methods become limiting.

2677

For example, the eddy covariance (EC) technique is widely used for the development and validation of terrestrial ecosystem models. However, the EC method only provides a net flux while process-based models predict component fluxes. To overcome this long-standing incompatibility problem, it is crucial that the CO<sub>2</sub> or H<sub>2</sub>O fluxes can be separated into their component fluxes. Conventional partitioning methods make a priori assumptions about ecosystem functions and typically require several days or weeks of data to cover key phenological periods in order to obtain robust regression parameters, obscuring ecosystem functions at shorter time scales. For example, the response of soil respiration to strong rain events is pulse-like, occurring at time scales of minutes to hours (e.g. Xu et al., 2004; Unger et al., 2010b). Other sudden changes or “switches” of ecosystem states have also been documented (Baldocchi et al., 2006; Lee et al., 2007). It is imperative that the partitioning scheme resolves episodic responses of this kind, because it is the transient, non-equilibrium responses that provide a more rigorous test of model processes and their validity. Assimilating continuous measurements of CO<sub>2</sub> and H<sub>2</sub>O fluxes and their isotopic composition (e.g.  $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ ,  $\delta\text{D}$ ) into process-based models should provide a better constrained solution. Similarly, assimilating chamber-based flux measurements of these isotopic fluxes should help to understand and constrain our model predictions during metabolic switches, especially when photosynthetic products become limiting such as during droughts (Unger et al., 2010a; Fig. 4), rainy periods (Wingate et al., 2010a) or dawn (Barbour et al., 2011). Ideally, we advocate the use of multiple isotopic tracers ( $^{13}\text{CO}_2$ ,  $\text{CO}^{18}\text{O}$ ,  $\text{H}_2^{18}\text{O}$ ,  $\text{HDO}$ ) as they provide more distinct information on the governing processes and the resulting component fluxes.

## 2.5 Isotopic archives and relevant aspects of spatio-temporal integration

### 2.5.1 Isotopic archives in trees and herbaceous vegetation

Trees rings contain formidable isotopic archives, which enable retrospective analyses of intra- and interannual variation of carbon and oxygen isotope composition and the

2678

related ecophysiological drivers over many centuries (see Fig. 5a; Sidorova et al., 2009; Andreu-Hayles et al., 2011; Nock et al., 2011; Knorre et al., 2010). These archives are well suited for detailed reconstructions of Anthropocene climate change impacts from carbon and oxygen isotope discrimination, with comparatively high temporal resolution over long time scales (Fig. 5a). However, single tree ring chronologies provide limited spatial and community integration because they are retrieved from individuals and therefore only report local signals (integrating isotopic information from an area of  $10^{-1}$  to  $10^2$  m<sup>2</sup> depending on the size of the individual). Also, the tree ring signal is, or can be, species-specific. Thus, construction of community- or ecosystem-scale chronologies depends upon combining several tree ring records from the different species composing the community. This requires additional information, such as knowledge of past species dynamics for estimation of species-weighted community isotopic signals.

Life span of herbaceous vegetation is shorter than that of trees, meaning that isotopic records are not as long either; whereby long-lived perennial plants that exhibit annual growth increments can be valuable recorders of ecophysiological changes in plant performance and thus recorders of climate variation. For example,  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values of growth increments of *Cassiope tetragon* from the High Arctic depict fluctuations in the AO/NAO (Arctic and North Atlantic Oscillations) which control the amount of snow melt ( $\delta^{18}\text{O}$ ) water available to plants in summer and associated changes in leaf gas exchange ( $\delta^{13}\text{C}$ ).

However, the isotopic reconstructions of Anthropocene climate change in herbaceous vegetation (crops and grassland) are possible, if plants were sampled and preserved during the epoch. Such archives are relatively rare (Köhler et al., 2010), and are mainly represented by herbaria. In general herbarium specimens have been sampled at different locations, so that long-term isotopic records from these involve a spatially disperse representation (that exhibits inter-local isotopic noise) of a species changing isotopic composition.

Clearly, the elaboration of isotopic records presents very different methodological challenges for the different biomes. But, for grassland, a nice analogy to tree rings

2679

is given by the yearly rings (annuli) of horns (or hoofs) of obligate grazers (Barbosa et al., 2009). These can also yield isotopic records over many years, which reflect to some extent that of grassland vegetation (Barbosa et al., 2010; see Fig. 5b). The spatial integrations provided by tree and horn ring isotope composition are quite contrasting: local and stationary for the tree, and vast and cyclic for grasses, reflecting visits of the different parts of the year-round grazing ground of the grazer. Because of the ephemeral nature of grassland biomass, animal tissues (such as hair, bones or feces) have become an important isotopic proxy of grassland vegetation. Still, the use of grazer tissue for reconstructions of grassland isotopic chronologies rests on a number of assumptions which need to be verified for the different systems (e.g. selective grazing or non-constant relationships between isotopic composition of grazer tissues and diet due to differential digestibility of diet components) (Wittmer et al., 2010).

The carbon isotope composition of grazer tissues such as horn can provide an accurate reflection of that of the grazed vegetation, and this is – at least partially – due to its homogenous chemical composition (keratin), which reduces variation associated with metabolic isotope fractionation which can be a problem in chemically heterogeneous materials. Conversely, the oxygen isotope composition of horn does not directly reflect that of diet, because of modification by the isotope composition of drinking water, atmospheric oxygen and fractionation effects in animal metabolism.

## 2.5.2 Non-vascular plants as integrators of micro-environmental and long-term changes

A particular case of micro-scale environmental record can be derived from  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  of non-vascular plants (NVP), as they respond highly sensitive to micro-environmental changes. NPVs such as cyanobacteria, algae, lichens and bryophytes are often used in environmental assessment and are increasingly recognized for the important roles in biogeochemical cycling (DeLucia et al., 2003; Elbert et al., 2009) and pedosphere-biosphere-atmosphere exchanges (Shimoyama et al., 2004).

2680

Due to the desiccation tolerance and ability to reactivate the metabolism by equilibration with water vapor,  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  of NVP integrate changes of carbon and water on a micro-environmental scale and over a long period (Máguas and Brugnoli, 1996), due to their sensitivity to small-scale environmental conditions, low growth rates and long lifespan.  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  of NVP record isotopically distinct micro-resources which are not commonly traced by higher plants. This includes for example micro-scaled humidity gradients, soil-water- $\text{CO}_2$  exchange by carboanhydrase, soil respiration and evaporation recycled by vegetation cover, or diel courses of water vapor signature (e.g. Lakatos et al., 2007; Hartard et al., 2008). Moreover, organic  $\delta^{13}\text{C}$  of NVP can be used as tracer for carbon acquisition (Meyer et al., 2008), environmental change of respired  $\text{CO}_2$  (Flanagan et al., 1999; Lakatos et al., 2007), and global change (Máguas and Brugnoli, 1996). Even fossil bryophytes record ancient  $\text{CO}_2$  levels (Fletcher et al., 2005, 2006).

$\delta^{18}\text{O}$  of absorbed water and respired  $\text{CO}_2$  operates as a tracer for partitioning water sources (Flanagan et al., 1999; Hartard et al., 2008) and in full equilibration as vapor trap (Helliker and Griffiths, 2007; Hartard et al., 2009). Because epiphytic lichens are often in equilibrium with water vapor, it is inferred that organic  $\delta^{18}\text{O}$  might serve as an environmental integrator and recorder for atmospheric vapor on a regional and global level (Helliker and Griffiths, 2007; Lakatos et al., 2007; Hartard et al., 2009). Further, several studies have demonstrated the potential of stable isotopes ( $\delta^{18}\text{O}$ ,  $\delta^{13}\text{C}$ ) in peat mosses as proxies for retrospective studies of environmental change, such as temperature, humidity and/or precipitation shifts (Loader et al., 2007; Lamentowicz et al., 2008; Moschen et al., 2009; Loisel et al., 2010).

Thus, stable isotopes of NVP may be used as ecological indicators for environmental change, especially for variations in  $\text{CO}_2$ -concentrations and water vapor as well as long term indicators of water accessibility change (Niemi and McDonald, 2004).

2681

## 2.6 Continental isotope variation in precipitation at the landscape-scale

Continental-scale studies of the water isotope cycle are becoming increasingly central to our understanding of synoptic climates, ecosystem processes and the role of abiotic processes (i.e. temperature of condensation), moisture sources, and storm tracks on the ecohydrology of entire landscapes and continents (e.g. Welker 2000; Vachon et al., 2007; Birks and Edwards 2009; Liu et al., 2010). The ratio of heavy to light isotopes within precipitation for instance, records the history of the moisture, from conditions at the moisture source, moisture recycling during transport, to conditions during condensation (Vachon et al., 2010a, b). If the factors that contribute to stable isotopes in precipitation (SIP) for a given climate system are well understood, SIP represent a valuable means of quantifying changes within a given hydrologic system.

A process-based understanding is important to delineating today's climate trajectories (Welker, 2010) and evapotranspiration fluxes, such as those that have been quantified as air masses progress across the Amazon Basin (Noone and Simmonds, 2002; Henderson-Sellers et al., 2004) and into the Gulf Coast of North America (Liu et al., 2010). Concurrently, SIP have been paramount in determining which moisture sources provide vapor to regions of the eastern US, Tibetan Plateau and Canada (Tien et al., 2007; Birks and Edwards, 2009; Sjostrom and Welker, 2009).

Interannual climate differences in weather conditions can often be linked to major modes of climate variability, such as the Pacific Decadal Oscillation (PDO), El Niño Southern Oscillation (ENSO) or North Atlantic Oscillation (NAO). These conditions and indices are identified by changes in ocean temperatures, high or low pressure systems, or areas of deep convection. The phase of these modes of climate variability have been shown to affect ecosystem processes and a cascading set of consequences (Holmgren et al., 2001) along with the climate (Ault and St. George, 2010). Commensurate with the phaseology are modifications to air mass trajectories, moisture source regions and regional meteorological parameters (e.g. surface temperatures, relative humidity and precipitation amount) in regions of the globe, far from the source of the

2682

perturbations (e.g. Kumar et al., 2006). Such teleconnections make for highly variable weather, depending upon the locale which may have direct effects on ecosystem C balance (Holmgren et al., 2001). An example is given in Fig. 5c, showing that the Pacific Decadal Oscillation (PDO) is reflected in the SIP in the western US as part of an initial temporal trend analysis using USNIP (Cohn et al., 2009). There is a strong correspondence between the PDO index and  $\delta^{18}\text{O}$  in eastern Washington between the years 1989–2002.

Our understanding of the SIP and its potential impact on continental C cycling have benefited from recent discoveries, such as (i) quantifying the impact of seasonality in precipitation, paramount to driving C cycling, on the isotopic composition of annual precipitation (Vachon et al., 2007), (ii) defining how time series of monthly SIP values relate to simultaneously measured monthly temperatures (Vachon et al., 2010a) and (iii) quantifying the primary environmental causes for the average monthly isoscapes (i.e. spatial distribution maps of isotope records, Vachon et al., 2010b) (Fig. 6). The isoscapes shown in Fig. 6 related to average monthly temperatures in such a manner that the SIP-temperature slopes followed the predicted pattern of steep slopes during the colder months and more shallow slopes during warmer months. However, the observed slopes are constantly  $\sim 0.2 \delta^{18}\text{O}/^\circ\text{C}$  less than is predicted by a theoretical Rayleigh Model. Future studies should explore the storm track and moisture source conditions and examine how annual changes in the strength of such decadal oscillations as the PDO, ENSO and others affect SIP and their relationships with mean annual temperature. This would bring important insight for interpreting climate proxies.

Back trajectory analysis of weather and thus precipitation (HYSPLIT – Draxler and Hess, 2004; Sjoström and Welker, 2009) is a modeling tool that has been used extensively by the atmospheric chemistry community to quantify long-distance transport of pollutants, and more recently for studies of isotopic characteristics of precipitation (Burnett et al., 2004). More specifically, this methodology has also been very useful in determining the origin of air masses and precipitating moisture, which has been linked to excursions in SIP trends (Sjoström and Welker, 2009). Indeed, moisture source

2683

conditions are significant contributors to resultant SIP values. Each moisture source experiences its own interannual and interseasonal temperature fluctuations – key components in the most simplistic deviations of SIP. Developing this understanding will be important to fully account for changes in the C cycle.

## 2.7 GLOBAL-scale modeling

Globally, the terrestrial biosphere fractionates stable carbon isotopes during  $\text{CO}_2$  uptake by about 15‰, which includes contributions from both  $\text{C}_3$  plants (with ca. 18‰) and  $\text{C}_4$  plants (with ca. 4‰) (Lloyd and Farquhar, 1994). In contrast, the dissolution of  $\text{CO}_2$  in the oceans fractionates by only 2‰ (Tans et al., 1993). Large variations in the  $\delta^{13}\text{C}$  of atmospheric  $\text{CO}_2$  are therefore strong indicators of terrestrial biosphere activity and their observation provided one of the first corroborations of a large terrestrial carbon sink in the Northern hemisphere and have been used to quantify  $\text{CO}_2$  uptake by land masses versus uptake by the oceans on continental scale using atmospheric transport inversion techniques (Ciais et al., 1995; Francey et al., 1995). However, because even small errors in the assumed regional mean fractionations can lead to large uncertainties in derived  $\text{CO}_2$  flux partitions, atmospheric  $\delta^{13}\text{C}$  is mostly used in current atmospheric inversion studies as an additional constraint concurrently with other tracers (e.g. Rayner et al., 2008). Over continents, recent studies have also used this tracer to estimate the relative contributions of  $\text{C}_3$  and  $\text{C}_4$  plants in the continental net  $\text{CO}_2$  flux (Ballantyne et al., 2010). Even if  $\delta^{13}\text{C}$  is not providing conclusive evidence on the global scale, it can be used, for example, to decide between different formulations of stomatal conductance in global models (Ballantyne et al., 2010).

However, even when  $\delta^{13}\text{C}$  is not providing conclusive evidence on the global scale, it can be used, for example, to decide between different formulations of stomatal conductance in global models (Ballantyne et al., 2010). As stated above,  $\delta^{13}\text{C}$  is nowadays rather used as a further constrain on other atmospheric inversion studies (e.g. Rayner et al., 2008).

2684

$\delta^{13}\text{C}$  tags the net fluxes of terrestrial and oceanic  $\text{CO}_2$  fluxes. However, it is not possible to disentangle the large opposing gross fluxes such as assimilation and respiration with  $\delta^{13}\text{C}$ .  $\delta^{18}\text{O}$  in  $\text{CO}_2$ , on the other side, marks photosynthesis and respiration differently and has therefore the potential to quantify sinks and sources of carbon separately. It turns out that the global cycle of  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$  is rather difficult to understand (Francey et al., 1987). The spatial distribution of  $\delta^{18}\text{O}$ -sinks and sources seems to be well understood, while the temporal variation was not yet reproduced successfully (Cuntz et al., 2003b). So either our global carbon cycle models are not yet well adapted for  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$  (Cuntz et al., 2003a) or we are missing processes in the description of the  $\delta^{18}\text{O}$ . One of these missing processes could be the equilibration of oxygen from gaseous  $\text{CO}_2$  with water by the enzyme carbonic anhydrase in soils (Wingate et al., 2008). Including accelerated soil hydration in a global model of  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$  eliminated discrepancies on the spatial structure between model and observations (Wingate et al., 2009). However, the temporal change in the model was still out of phase with observations. This awaits an explanation at present and might bear a few more surprises.

### 3 New technical and methodological developments in stable isotope research

The past decade has seen tremendous progress in the development of new techniques that compliment or rival traditional Isotope Ratio Mass Spectrometry (IRMS) for the determination of stable isotope abundances. Most important for carbon and water cycle research was the development of instruments using alternative means for determining stable isotope abundances, as well as the introduction of innovative compound-specific sample extraction methods.

2685

#### 3.1 Laser spectroscopy

The development of absorption spectroscopy instrumentation provided new dimensions of measurement speed and number of quantifiable isotopologues based on analysis of absorption of light in selected wavelengths in the near and mid-infrared, or cavity ring down techniques, which can be used to determine the abundance of individual isotopologues. The capacity for  $\text{CO}_2$ , liquid and water vapor analysis is providing opportunities which have never been possible with a field-deployable instrument. Further, new multi-species instruments that are becoming available enable so-called “clumped isotope” measurements (Eiler, 2007), wherein the occurrence of two heavy isotopes in the same molecule can serve as a unique stable isotope tracer itself.

#### 3.2 Compound Specific Isotope Analysis (CSIA)

In parallel with the development of new optical methods, IRMS itself has experienced technological and methodological development allowing for a new dimension of our understanding in plant metabolic processes through Compound Specific Isotope Analysis (CSIA) (e.g. different structural and labile carbohydrates extracted from plant organs, leaf wax alkanes, phloem sap, soil fractions) by coupled gas chromatography (GC-IRMS) or liquid chromatography-combustion (LC-C-IRMS) isotope ratio mass spectrometry. These innovative techniques increased the repeatability, precision and sample turnover considerably, which makes them particularly useful for isotopic studies at natural abundance or in tracer experiments (see Brüggemann et al., 2011). This opens numerous possibilities for a deeper understanding of carbon metabolism, transport and allocation under different environmental conditions. For example, carbon can be traced down the stem into the soil and back into the atmosphere, using natural abundance, i.e. ‘labelling’ by changing environmental conditions, or  $^{13}\text{C}$  labelling techniques (pulse labelling for labile, non-structural carbon and long-term labelling for structural carbon) by measuring  $\delta^{13}\text{C}$  of labile and structural carbon in the phloem, the wood, the soil (or  $\delta^{13}\text{C}$  of single compounds, e.g. sucrose and cellulose), and respired  $\text{CO}_2$ . Such

2686



been introduced, large uncertainties appeared in liquid sample analysis extracted from biological sources due to spectral interference with organic contaminants (West et al., 2010).

5 The second challenge that researchers face is increased speed of detection at high instrument precision. Free of sample preparation and processing, new optical techniques achieve much faster detection than IRMS. In-situ measurements of CO<sub>2</sub> and H<sub>2</sub>O isotope ratios in ambient air, especially if made on a long-term basis and calibrated precisely, can provide a powerful tool for atmospheric inverse analysis of land carbon sink and the tracking of water transport in the atmosphere. However, to measure the source/sink signature properly, one should interface the isotopic analyzer with plant (Barbour et al., 2007; Barthel et al., 2010; Wingate et al., 2010a, b) and soil chambers (Wingate et al., 2010a, b) and deploy it in the gradient-diffusion mode either over the vegetation (Griffis et al., 2004) or over the soil surface inside the canopy (Santos et al., 2010), or combine it with a sonic anemometer for direct eddy covariance measurement of isotopic fluxes (Lee et al., 2005; Griffis et al., 2008, 2010), or landscape scale measurements in high elevation or airborne conditions (e.g. Tuszon et al., 2010). In all these configurations, highly suitable interfaces between the analyzer and the periphery are mandatory together with fast instrument response. The whole system must be robust; designed and tuned for minimal interference, memory effects or signal drifts. This is particularly critical for eddy covariance applications, which require an instrument response to be faster than 10 Hz. Fast detections are also desirable in chamber measurements in the case of episodic events such as short soil CO<sub>2</sub>-flux pulses after rain (Santos et al., 2010; Unger et al., 2010b). These requirements should be a high priority in future instrument development efforts and open a new dimension of information on the short-term temporal variation in natural systems (e.g. Unger et al., 2010a).

The third challenge that researchers face is instrument and infrastructure cost. High instrument and maintenance costs limit the broad adoption of new technologies in field research. It would be highly desirable that the costs be brought down to a level

2689

comparable to that of a broadband infrared gas analyzer, which are now an indispensable tool for ecosystem carbon and water flux monitoring worldwide (Baldocchi et al., 2001). We envision the development of a network with real-time observations of isotopic fluxes of CO<sub>2</sub> and H<sub>2</sub>O to help diagnose changes in biospheric processes and functions. This can become a realistic goal at lower instrument costs.

#### 4 Conclusions

We are in the midst of rapid growth in our process-based understanding of the application of stable isotopes. Moreover, new technological developments accelerated the scientific progress opening new frontiers in ecological and interdisciplinary research across scales.

10 Significant advances have been made recently in linking the carbon and water cycles via dual isotope approaches ( $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$ ) at the leaf, plant, and ecosystem scale. This approach largely forwarded the understanding of the coherence between carbon and water relations, in particular by uncovering plant responses to environmental changes, interactions and feed-backs within plant communities and their implications for ecosystem processes.

15 At the same time, we have achieved a new awareness on multiple isotope fractionation processes during assimilation and transpiration (e.g. fractionation during photorespiration, day respiration, leaf internal CO<sub>2</sub> conductance, or exchange between leaf water pools) and during dark respiration, carbon transport and within metabolic processes. This topic is currently in the focus of numerous investigations with the attempt to answer a large number of open questions regarding metabolic processes and within cell and plant carbohydrate formations and fluxes (see also Werner and Gessler, 2011).

20 However, a new mechanistic understanding of these processes is essential for studying ecosystem processes, where so far only moderate progress in applying stable isotopes in partitioning studies has been achieved. The recognition that the isotope

2690







- rock and plants: neglected major players in the global cycles of carbon and nitrogen?, *Biogeosciences Discuss.*, 6, 6983–7015, doi:10.5194/bgd-6-6983-2009, 2009.
- Evans, R. D.: Physiological mechanisms influencing plant nitrogen isotope composition, *Trends Ecol. Evol.*, 6, 121–126, 2001.
- 5 Farquhar, G. D. and Lloyd, J.: Carbon and oxygen isotope effects in the exchange of carbon dioxide between terrestrial plants and the atmosphere, in: *Stable isotopes and plant carbon-water relations*, edited by: Ehleringer, J. R., Hall, A. E., and Farquhar, G. D., 47–70, Academic Press, San Diego, 1993.
- Farquhar, G. D., O'Leary, M. H., and Berry, J. A.: On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves, *Aust. J. Plant Physiol.*, 9, 121–137, 1982.
- 10 Farquhar, G. D., Ehleringer, J. R. and Hubick, K. T.: Carbon isotope discrimination and photosynthesis, *Annu. Rev. Plant Phys.*, 40, 503–537, 1989.
- Farquhar, G. D., Lloyd, J., Taylor, J. A., Flanagan, L. B., Syvertsen, J. P., Hubick, K. T., Wong, S. C., and Ehleringer, E. R.: Vegetation effects on the isotope composition of oxygen in atmospheric CO<sub>2</sub>, *Nature*, 363, 439–443, 1993.
- 15 Farquhar, G. D., Barbour, M. M., and Henry, B. K.: Interpretation of oxygen isotope composition of leaf material, in: *Stable isotopes, integration of biological, ecological and geochemical processes*, edited by: Griffiths, H., 27–62, Bios Scientific Publishers, 1998.
- 20 Farris, F. and Strain, B. R.: The effect of water-stress on leaf H<sub>2</sub><sup>18</sup>O enrichment, *Radiat. Environ. Biophys.* 15, 167–202, 1978.
- Ferrio, J. P., Cuntz, M., Offermann, C., Siegwolf, R., Saurer, M., and Gessler, A.: Effect of water availability on leaf water isotopic enrichment in beech seedlings shows limitations of current fractionation models, *Plant Cell Environ.*, 32, 1285–1296, 2009.
- 25 Flanagan, L. B., Kubien, D. S., and Ehleringer, J. R.: Spatial and temporal variation in the carbon and oxygen stable isotope ratio of respired CO<sub>2</sub> in a boreal forest ecosystem, *Tellus B*, 51, 367–384, 1999.
- Fletcher, B. J., Beerling, D. J., Brentnall, S. J., and Royer, D. L.: Fossil bryophytes as recorders of ancient CO<sub>2</sub> levels: Experimental evidence and a cretaceous case study, *Global Biogeochem. Cycles*, 19, 1–13, GB3012, doi:10.1029/2005GB002495, 2005.
- 30 Fletcher, B. J., Brentnall, S. J., Quick, W. P., and Beerling, D. J.: Bryocarb: A process-based model of thallose liverwort carbon isotope fractionation in response to CO<sub>2</sub>, O<sub>2</sub>, light and temperature, *Geochim. Cosmochim. Ac.*, 70, 5676–5691, 2006.

2697

- Flexas, J., Dias-Espejo, A., Galmes, J., Kaldenhoff, R., Medrano, H., and Ribas-Carbó, M.: Rapid variations in mesophyll conductance in response to changes in CO<sub>2</sub> concentration around leaves, *Plant Cell Environ.*, 30, 1284–1298, 2007.
- Flexas, J., Ribas-Carbo, M., Diaz-Espejo, A., Galmes, J., and Medrano, H.: Mesophyll conductance to CO<sub>2</sub>: current knowledge and future prospects, *Plant Cell Environ.*, 31, 602–621, 2008.
- 5 Francey, R. J. and Tans, P. P.: Latitudinal variation in oxygen-18 of atmospheric CO<sub>2</sub>, *Nature*, 327, 495–497, 1987.
- Francey, R. J., Tans, P. P., Allison, C. E., Enting, I. G., White, J. W. C., and Troler, M.: Changes in oceanic and terrestrial carbon uptake since 1982, *Nature*, 373, 326–330, 1995.
- 10 Francey, R. J., Allison, C. E., Etheridge, D. M.: A 1000-year high precision record of δ<sup>13</sup>C in atmospheric CO<sub>2</sub>, *Tellus B*, 51, 170–193, 1999.
- Gamnitzer, U., Moyes, A. B., Bowling, D. R. and Schnyder, H.: Measuring and modelling the isotopic composition of soil respiration: insights from a grassland tracer experiment, *Biogeosciences Discuss.*, 8, 83–119, doi:10.5194/bgd-8-83-2011, 2011.
- 15 Gessler, A., Rennenberg, H., and Keitel, C.: Stable isotope composition of organic compounds transported in the phloem of European beech - Evaluation of different methods of phloem sap collection and assessment of gradients in carbon isotope composition during leaf-to-stem transport, *Plant Biol.*, 6, 721–729, 2004.
- 20 Gessler, A., Keitel, C., Kodama, N., Weston, C., Winters, A. J., Keith, H., Grice, K., Leuning, R., and Farquhar, G. D.: Delta C-13 of organic matter transported from the leaves to the roots in *Eucalyptus delegatensis*: short-term variations and relation to respired CO<sub>2</sub>, *Funct. Plant Biol.*, 34, 692–706, 2007a.
- Gessler, A., Peuke, A. D., Keitel, C., and Farquhar, G. D.: Oxygen isotope enrichment of organic matter in *Ricinus communis* during the diel course and as affected by assimilate transport, *New Phytol.*, 174, 600–613, 2007b.
- 25 Gessler, A., Tcherkez, G., Peuke, A. D., Ghashghaie, J., and Farquhar, G. D.: Experimental evidence for diel variations of the carbon isotope composition in leaf, stem and phloem sap organic matter in *Ricinus communis*, *Plant Cell Environ.*, 31, 941–953, 2008.
- 30 Gessler, A., Brandes, E., Buchmann, N., Helle, G., Rennenberg, H., and Barnard, R.: Tracing carbon and oxygen isotope signals from newly assimilated sugars in the leaves to the tree ring archive, *Plant Cell Environ.*, 32, 780–795, 2009a.
- Gessler, A., Löw, M., Heerd, C., Op de Beeck, M., Schumacher, J., Grams, T. E. E., Bahnweg,

2698







- Ohlsson, K. E. A.: Reduction of bias in static closed chamber measurement of  $\delta^{13}\text{C}$  in soil  $\text{CO}_2$  efflux, *Rapid Commun. Mass Sp.*, 24, 180–184, 2009.
- Pataki, D. E., Ehleringer, J. R., Flanagan, L. B., Yakir, D., Bowling, D. R., Still, C. J., Buchmann, N., Kaplan, J. O., and Berry, J. A.: The application and interpretation of Keeling plots in terrestrial carbon cycle research, *Glob. Biogeochem. Cy.*, 17, 1022, doi:10.1029/2001GB001850, 2003.
- Pate, J. and Arthur, D.: delta C-13 analysis of phloem sap carbon: novel means of evaluating seasonal water stress and interpreting carbon isotope signatures of foliage and trunk wood of *Eucalyptus globules*, *Oecologia*, 117, 301–311, 1998.
- Ponton, S., Flanagan, L. B., Alstad, K. P., Johnson, B. G., Morgenstern, K., Kljun, N., Black, T. A., and Barr, A. G.: Comparison of ecosystem water-use efficiency among Douglas-fir forest, aspen forest and grassland using eddy covariance and carbon isotope techniques, *Glob. Change Biol.*, 12, 294–310, 2006.
- Priault, P., Wegener, F., and Werner, C.: Pronounced differences in diurnal variation of carbon isotope composition of leaf respired  $\text{CO}_2$  among functional groups, *New Phytol.*, 181, 400–412, doi:10.1111/j.1469-8137.2008.02665.x, 2009.
- Ramírez, D. A., Querejeta, J. I., and Bellot, J.: Bulk leaf  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  reflect the intensity of intraspecific competition for water in a semi-arid tussock grassland, *Plant Cell Environ.*, 32, 1346–1356, 2009.
- Rascher, K. G., Máguas, C., and Werner, C.: On the use of phloem sap  $\delta^{13}\text{C}$  as an indicator of canopy carbon discrimination, *Tree Physiol.*, 13, 1499–1514, doi:10.1093/treephys/tpq092, 2010.
- Rayner, P. J., Law, R. M., Allison, C. E., Francey, R. J., Trudinger, C. M., and Pickett-Heaps, C.: Interannual variability of the global carbon cycle (1992–2005) inferred by inversion of atmospheric  $\text{CO}_2$  and  $\delta^{13}\text{CO}_2$  measurements, *Global Biogeochem. Cy.*, 22(3), GB3008, 1–12, doi:10.1029/2007GB003068, 2008.
- Richardson, D. M. and Pyšek, P.: Plant invasions: merging the concepts of species invasiveness and community invasibility, *Progr. Phys. Geogr.*, 30, 409–431, 2006.
- Rose, K. L., Graham, R. C., and Parker, D. R.: Water source utilization by *Pinus jeffreyi* and *Actostaphylos patula* on thin soils over bedrock, *Oecologia*, 134, 46–54, 2003.
- Rossmann, A., Butzenlechner, M., and Schmidt, H. L.: Evidence for a nonstatistical carbon isotope distribution in natural glucose, *Plant Physiol.*, 96, 609–614, 1991.
- Ruehr, N. K., Offermann, C. A., Gessler, A., Winkler, J. B., Ferrio, J. P., Buchmann, N., and

2705

- Barnard, R. L.: Drought effects on allocation of recent carbon: from beech leaves to soil  $\text{CO}_2$  efflux, *New Phytol.*, 184, 950–961, 2009.
- Santos, E. A., Wagner-Riddle, C., Lee, X., Warland, J. S., Brown, S. E., Staebler, R. M., Bartlett, P. A., and Kim, K.:  $^{13}\text{CO}_2$  and  $\text{C}^{18}\text{O}^{16}\text{O}$  temporal variation near the ground and above a temperate deciduous forest, AGU Fall Meeting, 2010.
- Sarris, D., Siegwolf, R. T. W., and Körner, C.: Ecophysiological response of *Pinus brutia* to increasing drought affecting water uptake and spring growth reflected in stable C and O isotopes, *Oecologia*, revised, 2011.
- Saurer, M. and Siegwolf, R. T. W.: Human impacts on tree-ring growth reconstructed from stable isotopes in *Stable Isotopes as Indicators of Ecological Change*, edited by: Dawson, T. E. and Siegwolf, R. T. W., *Terrestrial Ecology Series*, Elsevier, Amsterdam, Boston, 49–62, 2007.
- Scartazza, A., Mata, C., Matteucci, G., Yakir, D., Moscatello, S., and Brugnoli, E.: Comparisons of  $\delta^{13}\text{C}$  of photosynthetic products and ecosystem respiratory  $\text{CO}_2$  and their responses to seasonal climate variability, *Oecologia*, 140, 340–351, 2004.
- Schaeffer, S. M., Anderson, D. E., Burns, S. P., Monson, R. K., Sun, J., and Bowling, D. R.: Canopy structure and atmospheric flows in relation to the  $\delta^{13}\text{C}$  of respired  $\text{CO}_2$  in a subalpine coniferous forest, *Agric. For Meteorol.*, 148, 592–605, 2008.
- Schäufele, R., Santrucek, J., and Schnyder, H.: Dynamic changes of canopy-scale mesophyll conductance to  $\text{CO}_2$  diffusion of sunflower as affected by  $\text{CO}_2$  concentration and ABA, *Plant Cell Environ.*, doi:10.1111/j.1365-3040.2010.02230.x, 2010.
- Scheidegger, Y., Saurer, M., Bahn, M., and Siegwolf, R.: Linking stable oxygen and carbon isotopes with stomatal conductance and photosynthetic capacity: A conceptual model, *Oecologia*, 125, 350–357, 2000.
- Schmidt, H. L.: Fundamentals and systematics of the non-statistical distributions of isotopes in natural compounds, *Naturwissenschaften*, 90, 537–552, 2003.
- Schmidt, H. L. and Gleixner, G.: Carbon isotope effect on key reactions in plant metabolism and  $^{13}\text{C}$ -patterns in natural compounds, in: *Stable isotopes, integration of biological, ecological and geochemical processes*, edited by: Griffiths, H., 13–25, Bios scientific publishers, 1998.
- Schmidt, H. L., Werner, R. A., and Rossmann, A.: O-18 pattern and biosynthesis of natural plant products, *Phytochemistry*, 58, 9–32, 2001.
- Schmidt, M., Maseyk, K., Lett, C., Biron, P., Richard, P., Bariac, T., and Seibt, U.: Concentration effects on laser-based  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  measurements and implications for the calibration of

2706

- vapour measurements with liquid standards, *Rapid Commun. Mass Spectrom.*, 24, 3553–3561, 2010.
- Snyder, K. and Williams, D. G.: Water sources used by riparian trees varies among stream types on the San Pedro river, Arizona. *Agricul. For. Meteo.*, 105, 227–240, 2000.
- 5 Seibt, U., Wingate, L., Berry, J., and Lloyd, J.: Non-steady-state effects in diurnal  $^{18}\text{O}$  discrimination by *Picea sitchensis* branches in the field, *Plant Cell Environ.*, 29, 928–939, 2006.
- Seibt, U., Wingate, L., Lloyd, J., and Berry, J. A.: Nocturnal stomatal conductance effects on the  $\delta^{18}\text{O}$  of foliage gas exchange observed in two forest ecosystems, *Tree Physiol.*, 27, 585–595, 2007.
- 10 Shimoyama, K., Hiyama, T., Fukushima, Y. and Inoue, G.: Controls on evapotranspiration in a west siberian bog, *J. Geophys. Res.*, 109, D08111, 1–12, doi:10.1029/2003JD004114, 2004.
- Sidorova, O. V., Siegwolf, R. T. W., Saurer, M., Shashkin, A. V., Knorre, A. A., Prokushkin, A. S., Vaganov, E. A., and Kiryanov, A. V.: Do centennial tree-ring and stable isotope trends of *Larix gmelinii* (Rupr.) Rupr. indicate increasing water shortage in the Siberian north? *Oecologia*, 161, 825–835, 2009.
- 15 Sjoström, D. and Welker, J. M.: Storm-track trajectories control the isotope geochemistry of precipitation in the Eastern US, *Exploration Geology-Isobalance, Special Issue*, 102, 103–112, 2009.
- Stratton, L. C., Goldstein, G., and Meinzer, F. C.: Temporal and spatial partitioning of water resources among eight woody species in a Hawaiian dry forest, *Oecologia*, 124, 309–317, 2000.
- Sternberg, L., Pinzon, M. C., Anderson, W. T., and Jahren, A. H.: Variation in oxygen isotope fractionation during cellulose synthesis: intramolecular and biosynthetic effects, *Plant Cell Environ.*, 29, 1881–1889, 2006.
- 25 Sturm, P. and Knohl, A.: Water vapor  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  measurements using off-axis integrated cavity output spectroscopy, *Atmos. Meas. Tech.*, 3, 67–77, doi:10.5194/amt-3-67-2010, 2010.
- Sullivan, P. F. and Welker, J. M.: Variation in leaf physiology of *Salix arctica* within and across ecosystems in the High Arctic: test of a dual  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  conceptual model, *Oecologia*, 151, 372–386, 2007.
- 30 Tans, P., Berry, J. A., and Keeling, R.: Oceanic  $^{13}\text{C}/^{12}\text{C}$  observations: A new window on ocean  $\text{CO}_2$  uptake, *Global Biogeochem. Cy.*, 7, 353–368, 1993.

2707

- Tcherkez, G.: Do metabolic fluxes matter for interpreting isotopic respiratory signals? *New Phytologist* 186, 566–568, 2010.
- Tcherkez, G., Nogues, S., Bleton, J., Cornic, G., Badeck, F., and Ghashghaie, J.: Metabolic origin of carbon isotope composition of leaf dark-respired  $\text{CO}_2$  in French bean, *Plant Physiol.*, 131, 237–244, 2003.
- 5 Tcherkez, G., Farquhar, G., Badeck, F., and Ghashghaie, J.: Theoretical considerations about carbon isotope distribution in glucose of C-3 plants, *Funct. Plant Biol.*, 31, 857–877, 2004.
- Tcherkez, G., Schäufele, R., Nogués, S., Piel, C., Boom, A., Lanigan, G., Barbaroux, C., Mata, C., Elhani, S., Hemming, D., Maguas, C., Yakir, D., Badeck, F., Griffiths, H., Schnyder, H., and Ghashghaie, J.: On the  $^{13}\text{C}/^{12}\text{C}$  isotopic signal of day and night respiration at the mesocosm level, *Plant Cell Environ.*, 33, 900–913, 2010.
- 10 Terwilliger, V. J., Kitajima, K., Le Roux-Swarthout, D. J., Mulkey, S., and Wright, S. J.: Intrinsic water-use efficiency and heterotrophic investment in tropical leaf growth of two neotropical pioneer tree species as estimated from  $\delta^{13}\text{C}$  values, *New Phytol.*, 152, 267–281, 2001.
- 15 Tian L., Yao, T., MacClune, K., White, J. W. C., Schilla, A., Vaughn, B., Vachon, R., and Ichiyangi, K.: Isotopic variations in West China: a consideration of moisture sources, *J. Geophys. Res.*, 112, D10112, doi:10.1029/2006JD007718, 2007.
- Tu, K. P. and Dawson, T. E.: Partitioning ecosystem respiration using stable carbon isotope analyses of  $\text{CO}_2$ , in: *Stable Isotopes and Biosphere-Atmosphere Interactions: Processes and Biological Controls*, edited by: Flanagan, L. B., Ehleringer, J. R., and Pataki, D. E., Elsevier Academic Press, San Diego, CA, 125–153, 2005.
- 20 Tu, K. P. and Dawson, T. E.: Partitioning ecosystem respiration between plant and microbial sources using natural abundance stable carbon isotopes: A study of four California ecosystems, *Oecologia*, in review, 2011.
- 25 Tu, K. P., Brooks, P. D., and Dawson, T. E.: Using septum capped vials with continuous flow isotope ratio mass spectrometric analysis of atmospheric  $\text{CO}_2$  for Keeling plot applications, *Rapid Commun. Mass Spect.*, 15, 952–956, 2001.
- Tuzson, B., Henne, S., Brunner, D., Steinbacher, M., Mohn, J., Buchmann, B., and Emmenegger, L.: Continuous isotopic composition measurements of tropospheric  $\text{CO}_2$  at Jungfraujoch (3580 m a.s.l.), Switzerland: real-time observation of regional pollution events, *Atmos. Chem. Phys. Discuss.*, 10, 24563–24593, doi:10.5194/acpd-10-24563-2010, 2010.
- 30 Unger, S., Máguas, C., Pereira, J. S., Aires, L. M., David, T. S., and Werner, C.: Partitioning carbon fluxes in a Mediterranean oak forest to disentangle changes in ecosystem sink strength

2708

- during drought, *Agr. Forest Meteorol.*, 149, 949–961, doi:10.1016/j.agrformet.2008.11.013, 2009.
- Unger, S., Máguas, C., Pereira, J. S., Aires, L. M., David, T. S. and Werner, C.: Disentangling drought-induced variation in ecosystem and soil respiration by stable isotope partitioning, *Oecologia*, 163, 1043–1057, doi:10.1007/s00442-010-1576-6, 2010a.
- 5 Unger, S., Máguas, C., Pereira, J. S., David, T. S., and Werner, C.: The influence of precipitation pulses on soil respiration – Assessing the “Birch effect” by stable carbon isotopes, *Soil Biol. Biochem.*, 42, 1800–1810, doi:10.1016/j.soilbio.2010.06.01, 2010b.
- Vachon, R., White, J. W., Gutmann, E., and Welker, J. M.: Amount-weighted annual isotopic ( $\delta^{18}\text{O}$ ) values are affected by the seasonality of precipitation: A sensitivity study, *Geophys. Res. Lett.*, 34, L21707, doi:10.1029/2007GL030547, 2007.
- 10 Vachon, R. W., Welker, J. M., White, J. W. C., and Vaughn, B. H.: Monthly precipitation isoscapes ( $\delta^{18}\text{O}$ ) of the United States: Connections with surface temperatures, moisture source conditions, and air mass trajectories, *J. Geophys. Res.*, 115, D21126, doi:10.1029/2010JD014105, 2010a.
- Vachon, R. W., Welker, J. M., White, J. W., and Vaughn, B. H.: Moisture source temperatures and precipitation  $\delta^{18}\text{O}$ -temperature relationships across the US, *Water Resour. Res.*, 46, W07523, 1–14, doi:10.1029/2009WR008558, 2010b.
- Van Bel, A. J. E. and Hess, P. H.: Hexoses as phloem transport sugars: the end of a dogma? *J. Exp. Bot.*, 59, 261–272, 2008.
- 20 Warren, C. R. and Adams, M. A.: Internal conductance does not scale with photosynthetic capacity: implications for carbon isotope discrimination and the economics of water and nitrogen use in photosynthesis, *Plant Cell Environ.*, 29, 192–201, 2006.
- Wegener, F., Beyschlag, W., and Werner, C.: The magnitude of diurnal variation in carbon isotopic composition of leaf dark respired  $\text{CO}_2$  correlates with the difference between  $\delta^{13}\text{C}$  of leaf and root material, *Funct. Plant Biol.*, 37, 849–858, doi:10.1071/FP09224, 2010.
- 25 Welker, J. M.: Isotopic ( $\delta^{18}\text{O}$ ) characteristics of weekly precipitation collected across the United States: An initial analysis with application to water source studies, *Hydrol. Process.*, 14, 1449–1464, 2000.
- 30 Welker, J. M., Jonsdottir, I., and Fahenstock, J. T.: Isotopic ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) characteristics of *Carex* plants and populations along the Eurasian Coastal Arctic, *Polar Biol.*, 27, 29–37, 2003.
- Welker, J. M., Rogers, M., Cable, J., Cohn, B., Bogan, D., and Arp, C.: The Alaska Water


2709

- Isotope Network, International Isotope Symposium, Fairbanks, AK, Proceedings, 2010.
- Welp, L. R., Lee, X., Kim, K., Griffis, T. J., Billmark, K. A., and Baker, J. M.:  $\delta^{18}\text{O}$  of water vapour, evapotranspiration and the sites of leaf water evaporation in a soybean canopy, *Plant Cell Environ.*, 31(9), 1214–1228, 2008.
- 5 Wen, X. F., Sun, X. M., Zhang, S. C., Yu, G. R., Sargent, S. D., and Lee, X.: Continuous measurement of water vapor D/H and  $^{18}\text{O}/^{16}\text{O}$  isotope ratios in the atmosphere, *J. Hydrol.*, 349, 489–500, 2008.
- Werner, C.: Do isotopic respiratory signals trace changes in metabolic fluxes? *New Phytol.*, 186, 569–571, doi:10.1111/j.1469-8137.2010.03248.x, 2010.
- 10 Werner, C. and Máguas, C.: Carbon isotope discrimination as a tracer of functional traits in a Mediterranean macchia plant community, *Funct. Plant Biol.*, 37, 467–477, doi:10.1071/FP09081, 2010.
- Werner, C. and Gessler, A.: Diel variations in the carbon isotope composition of respired  $\text{CO}_2$  and associated carbon sources: a review of dynamics and mechanisms, *Biogeosciences Discuss.*, 8, 2183–2233, doi:10.5194/bgd-8-2183-2011, 2011.
- 15 Werner, C., Maia, R., and Máguas, C.: Are carbon and oxygen isotope compositions of bulk leaf material reliable predictors of water use efficiency in slow-growing drought-adapted species? PS2001 Proceedings, CSIRO Publishing, Melbourne, Australia, 2001.
- Werner, C., Unger, S., Pereira, J. S., Maia, R., Kurz-Besson, C., David, T. S., David, J. S., and Máguas, C.: Importance of short-term dynamics in carbon isotope ratios of ecosystem respiration ( $\delta^{13}\text{C}_R$ ) in a Mediterranean oak woodland and linkage to environmental factors, *New Phytol.*, 172, 330–346, doi:10.1111/j.1469-8137.2006.01836.x, 2006.
- 20 Werner, C., Hasenbein, N., Maia, R., Beyschlag, W., and Máguas, C.: Evaluating high time-resolved changes in carbon isotope ratio of respired  $\text{CO}_2$  by a rapid in-tube incubation technique, *Rapid Commun. Mass Sp.*, 21, 1352–1360, doi:10.1002/rcm.2970, 2007.
- 25 Werner, C., Wegener, F., Unger, S., Nogués, S., and Priault, P.: Short-term dynamics of isotopic composition of leaf respired  $\text{CO}_2$  upon darkening: measurements and implications, *Rapid Commun. Mass Sp.*, 23, 2428–2438, doi:10.1002/rcm.4036, 2009.
- West, A. G., Goldsmith, G. R., Brooks, P. D., and Dawson, T. E.: Discrepancies between isotope ratio infrared spectroscopy and isotope ratio mass spectrometry for the stable isotope analysis of plant and soil waters, *Rapid Commun. Mass Spectrom.*, 24, 1948–1954, 2010.
- 30 Wingate, L., Seibt, U., Moncrieff, J., Lloyd, J., and Berry, J.: Variations in  $^{13}\text{C}$  discrimination during  $\text{CO}_2$  exchange in *Picea sitchensis* branches in the field, *Plant Cell Environ.*, 30, 600–

2710


- 616, doi:10.1111/j.1365-3040.2007.01647.x, 2007.
- Wingate, L., Seibt, U., Maseyk, K., Ogée, J., Almeida, P., Yakir, D., Pereira, J. S., and Mencuccini, M.: Evaporation and carbonic anhydrase activity recorded in oxygen isotope signatures of net CO<sub>2</sub> fluxes from a mediterranean soil, *Glob. Change Biol.*, 14, 2178–2193, doi:10.1111/j.1365-2486.2008.01635.x, 2008.
- Wingate, L., Ogée, J., Cuntz, M., Genty, B., Reiter, I., Seibt, U., Yakir, D., Maseyk, K., Pendall, E. G., Barbour, M. M., Mortazavi, B., Burrell, R., Peylin, P., Miller, J., Mencuccini, M., Shim, J. H., Hunt, J., and Grace, J.: The impact of soil microorganisms on the global budget of  $\delta^{18}\text{O}$  in atmospheric CO<sub>2</sub>, *P. Natl. Acad. Sci. USA*, 106, 22411–22415, 2009.
- Wingate, L., Ogée, J., Burrell, R., and Bosc, A.: Strong seasonal disequilibrium measured between the oxygen isotope signals of leaf and soil CO<sub>2</sub> exchange, *Glob. Change Biol.*, 16(11), 3048–3064, doi:10.1111/j.1365-2486.2010.02186.x, 2010a.
- Wingate, L., Ogée, J., Burrell, R., Bosc, A., Devaux, M., Grace, J., Loustau, D., and Gessler, A.: Photosynthetic carbon isotope discrimination and its relationship to the carbon isotope signals of stem, soil and ecosystem respiration, *New Phytol.*, 188, 576–589, 2010b.
- Wittmer, M. H. O. M., Auerwald, K., Schönbach, P., Schäufele, R., Müller, K., Yang, H., Bai, Y. F., Susenbeth, A., Taube, F., and Schnyder, H.: Do grazer hair and faeces reflect the carbon isotope composition of semi-arid C3/C4 grassland? *Basic Appl. Ecol.*, 11, 83–92, 2010.
- Wu, H. J. and Lee, X.: Short-term effects of rain on soil respiration in two New England forests, *Plant Soil*, 338, 329–342, 2011.
- Xu, L., Baldocchi, D. D., and Tang, J.: How soil moisture, rain pulses, and growth alter the response of ecosystem respiration to temperature, *Glob. Biogeochem. Cycles*, 18, GB4002, doi:10.1029/2004GB002281, 1–10, 2004.
- Yakir, D.: Oxygen-18 of leaf water: a crossroad for plant-associated isotopic signals, in: *Stable Isotopes: Integration of Biological and Geochemical Processes*, edited by: Griffiths, H., BIOS Scientific Publishers, Oxford, 147–168, 1998.
- Yakir, D.: The stable isotopic composition of atmospheric CO<sub>2</sub>, *Treatise on Geochemistry*, 4, 175–212, 2003.
- Yakir, D. and Sternberg, L.: The use of stable isotopes to study ecosystem gas exchange, *Oecologia*, 123, 297–311, 2000.
- Yoneyama, T., Handley, L. L., Scrimgeour, C. M., Fisher, D. B., and Raven, J. A.: Variations of the natural abundances of nitrogen and carbon isotopes in *Triticum aestivum*, with special reference to phloem and xylem exudates, *New Phytol.*, 137, 205–213, 1997.

2711




**Fig. 1.** Use of carbon and water stable isotopes in different compounds/biomarkers to reveal information on carbon and water cycles at different spatial and temporal scales. GPP – Gross primary productivity, Re – respiration, ET – evapotranspiration, DOC – dissolved organic carbon.

2712



**Fig. 2.** Relationship between carbon isotope discrimination ( $\Delta^{13}\text{C}$ ) of leaf organic matter and the ratio of leaf internal to external  $\text{CO}_2$  concentration ( $c_i/c_a$ ) determined from gas exchange measurements for three Mediterranean species: the evergreen sclerophyllous *Quercus coccoifera* L. (circles), the drought semi-deciduous *Cistus albidus* L. (triangles) and the herbaceous *Vinca difformis* Pourret (diamonds). Species were subjected to increasing drought treatment (light symbols) for 12 weeks and compared to well-watered controls (dark symbols) under semi-natural conditions in late spring at the University of Lisbon. Pooled data were collected after week one, six and 11 of the experiments. The line indicates the theoretical relationship from Farquhar et al. (1989). From Unger unpublished.

2713



**Fig. 3.** Principal component analysis (PCA) of eleven co-occurring Mediterranean macchia species based on mean leaf bulk  $\Delta^{13}\text{C}$  of current and of previous year grown leaves. The first axis separated species by annual mean  $\Delta^{13}\text{C}$ , while the second axis was loaded with seasonal changes in  $\Delta^{13}\text{C}$ . Analyzed macchia species from Serra da Arrabida, Southwest Portugal, were drought semi-deciduous shrubs *Cistus albidus* (CA), *Cistus monspeliensis* (CM) and *Rosmarinus officinalis* (RO), the sclerophyllous evergreens *Quercus coccoifera* (QC), *Arbutus unedo* (AU), *Olea europaea* (OE), *Phillyrea latifolia* (PL), *P. angustifolia* (PA), *Pistacia lentiscus* (PiL), *Erica arborea* (EA) and the gymnosperm *Juniperus phoenicea* (JP). Adapted from Werner and Máguas (2010).

2714







Fig. 5.

2717

**Fig. 5** Temporal analysis  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  variations: examples of different samples and archives. **(A)** Tree ring  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  analysis of bulk wood (triangles) and extracted cellulose (circle), data from Sidorova et al. (2009), with two different corrections for the  $^{13}\text{C}$  values in tree rings one according to Francey et al. (1999), corrected for the Süess-Effect (lines), the other according to Loader et al. (2010) (lined symbols), which also takes the  $\text{CO}_2$  response of photosynthesis into account. **(B)** Carbon isotope discrimination ( $\Delta^{13}\text{C}$ ) derived from horns of Alpine ibex (*Capra ibex*) from Augstmatthorn – Brienzler Rothorn between 1938 and 2006, data from Barbosa et al. (2010). **(C)** The monthly  $^{18}\text{O}$  precipitation values of a site in eastern Washington in comparison to the Pacific Decadal Oscillation (PDO) index: monthly means values two-days running averages. Note periods of close and fractured correspondence; data from Cohn et al. (2009). Please note different scales on x- and y-axes.

2718



**Fig. 6.** Changes in the monthly spatial distribution of  $\delta^{18}\text{O}$  of precipitation (isoscapes) across the US. Note the strong depletion of precipitation in the northern Rocky Mountain region. Data from Vachon et al. (2010b).