



**HAL**  
open science

# Methanogens and methanogenesis in the rumens and ceca of lambs fed two different high-grain-content diets

Milka Popova, Diego Morgavi, Cécile Martin

► **To cite this version:**

Milka Popova, Diego Morgavi, Cécile Martin. Methanogens and methanogenesis in the rumens and ceca of lambs fed two different high-grain-content diets. *Applied and Environmental Microbiology*, 2013, 79 (6), pp.1777-1786. 10.1128/AEM.03115-12 . hal-02646467

**HAL Id: hal-02646467**

**<https://hal.inrae.fr/hal-02646467>**

Submitted on 29 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Methanogens and Methanogenesis in the Rumens and Ceca of Lambs Fed Two Different High-Grain-Content Diets

M. Popova, D. P. Morgavi and C. Martin  
*Appl. Environ. Microbiol.* 2013, 79(6):1777. DOI:  
10.1128/AEM.03115-12.  
Published Ahead of Print 14 December 2012.

---

Updated information and services can be found at:  
<http://aem.asm.org/content/79/6/1777>

---

**SUPPLEMENTAL MATERIAL**

*These include:*

[Supplemental material](#)

**REFERENCES**

This article cites 41 articles, 16 of which can be accessed free  
at: <http://aem.asm.org/content/79/6/1777#ref-list-1>

**CONTENT ALERTS**

Receive: RSS Feeds, eTOCs, free email alerts (when new  
articles cite this article), [more»](#)

---

---

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>  
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

---

# Methanogens and Methanogenesis in the Rumens and Ceca of Lambs Fed Two Different High-Grain-Content Diets

M. Popova, D. P. Morgavi, C. Martin

INRA, UMR1213 Herbivores, Saint-Genès-Champanelle Clermont Université, VetAgro Sup, UMR Herbivores, Clermont-Ferrand, France

The amount and nature of dietary starch are known to influence the extent and site of feed digestion in ruminants. However, how starch degradability may affect methanogenesis and methanogens along the ruminant's digestive tract is poorly understood. This study examined the diversity and metabolic activity of methanogens in the rumen and cecum of lambs receiving wheat or corn high-grain-content diets. Methane production *in vivo* and *ex situ* was also monitored. *In vivo* daily methane emissions ( $\text{CH}_4$  g/day) were 36% ( $P < 0.05$ ) lower in corn-fed lambs than in wheat-fed lambs. *Ex situ* methane production ( $\mu\text{mol/h}$ ) was 4-fold higher for ruminal contents than for cecal contents ( $P < 0.01$ ), while methanogens were 10-fold higher in the rumen than in the cecum (*mcrA* copy numbers;  $P < 0.01$ ). Clone library analysis indicated that *Methanobrevibacter* was the dominant genus in both sites. Diet induced changes at the species level, as the *Methanobrevibacter millerae*-*M. gottschalkii*-*M. smithii* clade represented 78% of the sequences from the rumen of wheat-fed lambs and just about 52% of the sequences from the rumen of the corn-fed lambs. Diet did not affect *mcrA* expression in the rumen. In the cecum, however, expression was 4-fold and 2-fold lower than in the rumen for wheat- and corn-fed lambs, respectively. Though we had no direct evidence for compensation of reduced rumen methane production with higher cecum methanogenesis, the ecology of methanogens in the cecum should be better considered.

Global warming, caused by increasing atmospheric concentrations of greenhouse gases, is a major worldwide environmental, economic, and social threat, and it is well documented that livestock production contributes to this problem (1). Enteric methane ( $\text{CH}_4$ ) is a natural byproduct arising from microbial fermentation of feeds within the rumen and, to some extent, in the cecum (2). Microbial populations in the digestive tract of ruminants ferment carbohydrates, proteins, and, to a lesser extent, lipids to produce mainly volatile fatty acids ([VFA]; acetate, propionate, and butyrate as the main ones), dihydrogen ( $\text{H}_2$ ), and carbon dioxide ( $\text{CO}_2$ ). Methanogenic *Archaea* mainly use  $\text{H}_2$  to reduce  $\text{CO}_2$  to  $\text{CH}_4$ .

Diet manipulation is an effective way of decreasing enteric methane emissions in ruminant production systems (3). In intensive production, it is well established that high-cereal diets decrease enteric methane production. In parallel, these diets have been shown to decrease the diversity (4, 5) of methanogenic *Archaea* in the rumen of sheep. Concerning the effect of the nature of the cereal on methane production, few direct comparisons have been carried out. Beauchemin and McGinn (6) reported lower methane emissions in feedlot cattle fed corn (slowly degradable starch) than in animals receiving barley (readily degradable starch) finishing diets. Those authors suggested that this was mediated by the lower ruminal pH observed with the corn diet. In the present study, we hypothesized that the inclusion of readily degradable starch in the diet would shift plant cell wall degradation from the rumen to the cecum (7), providing more organic matter for cecal fermentation.

The cecum of ruminants harbors a complex microbial community (8, 9), but information about its members and particularly the methanogenic *Archaea* is scarce. Moreover, to our knowledge, there are no studies reporting the effect of dietary starch on the community of methanogens in the cecum.

The aim of this work was to compare the methanogenic communities in terms of diversity and metabolic activity in the rumen and cecum of growing lambs fed either corn (slowly degradable

starch) or wheat (readily degradable starch) diets. By using these contrasting starch cereals, we sought to bring out differences between the rumen and cecum microbiota. We also measured *in vivo* methane emissions by lambs. In order to compare the two digestive compartments, we measured the *ex situ* methanogenic potential of the contents of the rumen and cecum, as well as their capacity to degrade wheat or corn *in vitro*.

## MATERIALS AND METHODS

**Animals, diet, and experimental design.** The study was conducted at the animal experimental facilities of INRA's Herbivore Research Unit in Saint-Genès Champanelle (France). Animal procedures were in accordance with the guidelines for animal research of the French Ministry of Agriculture and applicable European guidelines and regulations for animal experimentation ([http://www2.vet-lyon.fr/ens/expa/acc\\_regl.html](http://www2.vet-lyon.fr/ens/expa/acc_regl.html)). The experimentation protocol was approved by the Regional Ethics Committee on Animal Experimentation, approval number CE2-10.

Twelve INRA 401 newborn male lambs were separated in two homogeneous (age, weight at birth) groups. Lambs were kept with their mothers until weaning at the age of 9 weeks. In each group, ewes were adapted progressively to high-grain-content diets based on wheat or corn, so from birth lambs were in contact with only one type of cereal. After weaning, lambs on each group were housed and fed together until the end of fattening (~24 weeks). Three weeks before slaughter, lambs were housed in individual pens for *in vivo* methane production and digestion measurements. Through the whole experimental period, animals were fed twice daily at 0800 h and 1300 h. Diets contained 550 g of cereal grain completed with 700 g of barley hay and 10 g of bicarbonate. Ingredients and chemical

Received 11 October 2012 Accepted 11 December 2012

Published ahead of print 14 December 2012

Address correspondence to M. Popova, milka.popova@clermont.inra.fr.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AEM.03115-12>.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.03115-12

composition of the experimental diets as fed are presented in Table S1 in the supplemental material.

**Intake, total tract digestibility, and *in vivo* methane production.** For practical reasons, animals were separated into two experimental groups, each one composed of 3 corn-fed and 3 wheat-fed lambs having similar live weights. There was a 3-week period between measurements of intake, digestibility, and methane production in each experimental group; also, lambs in the first group were slaughtered 3 weeks before those in the second one.

Intake, total tract digestibility, and daily methane production were measured during a 5-day period at the end of fattening while the lambs were in individual pens. Feed intake andorts were measured and recorded daily during the sample collection period to calculate dry matter intake (DMI). Total tract digestibility was determined from the total collection of feces during the measurement period. Fecal samples were weighed and mixed before a 10% aliquot was sampled. Feed and fecal samples from the 5 days were pooled by animal; one aliquot was used for dry matter determination (103°C for 24 h) and another for chemical analysis as previously reported (10). The neutral detergent fiber (NDF) and acid detergent fiber (ADF) contents were determined by sequential procedures after pretreatment with amylase and were expressed inclusive of residual ash (11). A polarimetric method was used for starch quantification (12).

Methane production was determined using the sulfur hexafluoride (SF<sub>6</sub>) tracer technique (13), as described by Martin et al. (10). A calibrated permeation tube was introduced into the rumen of each lamb through the mouth. Mean permeation rates of SF<sub>6</sub> from the brass permeation tubes were 703 ± 128 ng/min. Background concentrations of methane and SF<sub>6</sub> were also measured on ambient air samples collected every day in the barn during the sampling period. Gas samples were analyzed by gas chromatography as described elsewhere (10).

**Rumen content sampling for fermentation and microbial parameters.** Lambs were slaughtered at INRA-Theix's experimental abattoir at an average age of 171 ± 5.7 days and average body weight of 28.2 ± 0.94 kg. Lambs were last fed the morning of the day before slaughter. The entire gastrointestinal tract was removed as soon as possible after slaughter, and representative samples of homogenized ruminal and cecal contents were collected and weighed along with whole ruminal and cecal contents. Rumen pH was measured with a portable pH meter (CG840, electrode Ag/AgCl; Schott Geräte, Hofheim, Germany) at 5 locations immediately after opening the rumen. Cecum pH was measured only on the liquid phase. Aliquots of ruminal (~250-g) and cecal (~50-g) contents were strained through polyester monofilament fabric (250-µm-mesh aperture) to separate the liquid phase. For VFA analysis, ruminal and cecal filtrates were sampled as already described (14). VFA were analyzed by gas chromatography (15) using crotonic acid as the internal standard on a CP 9002 gas chromatograph (Chrompack, Middelburg, Germany). For enumeration of protozoa, rumen fluid was mixed with MFS solution (35 ml/liter formaldehyde, 0.14 M NaCl, 0.92 mM methyl green) and stored in the dark at room temperature until microscope counting was performed as already described (16).

Another aliquot of ruminal and cecal contents (~30 g each) was diluted with 15 ml ice-cold phosphate-buffered saline (PBS) (pH 6.8) and homogenized, for three 1-min cycles with 1-min intervals on ice, using a Polytron grinding mill (Kinematica GmbH, Steinhofhalde, Switzerland). Approximately 0.5 g was transferred to a 2-ml Eppendorf tube and mixed with 1 ml of RNALater tissue collection RNA stabilization solution (Applied Biosystems, Austin, TX). Tubes were kept at 4°C overnight to allow the solution to thoroughly penetrate the cells, as suggested by the manufacturer, and stored at -80°C until molecular analyses were performed.

**Rumen and cecum content sampling for *ex situ* and *in vitro* fermentations.** Total ruminal and cecal contents, as well as corresponding strained fluids, were used as the inoculum for *ex situ* and *in vitro* incubations, respectively. *Ex situ* fermentations were performed to measure the methanogenic potential of gastrointestinal contents. For that, 100-ml serum vials containing 30 ml of anaerobic buffer solution (17), kept at 39°C under an O<sub>2</sub>-free CO<sub>2</sub> gas headspace, were inoculated with 30 g of total

ruminal or cecal contents from each animal in duplicate. For *in vitro* fermentations, vials contained buffer as described above and 0.5 g of ground corn or wheat as the substrate. These vials were inoculated with 15 ml of strained rumen or cecum fluid (one sample per animal) in order to assess their capacity to ferment the starchy feed substrates. Vials were capped with butyl rubber stoppers and incubated at 39°C for 24 h. Vials without inoculum were used as a negative control. Gas production was measured at 6 h and 24 h of incubation with the aid of a pressure transducer, and samples were collected for analysis of constituents by gas chromatograph (Micro GC 3000A; Agilent Technologies, France). At the end of incubation, vials were opened, pH was measured, and samples for VFA analysis were taken as described above.

**tNA extraction and cDNA synthesis.** Total nucleic acids (tNA) were extracted as described by Popova et al. (18). Each tNA sample was divided into two equal fractions; DNA in one of the fractions was digested with DNase provided in a total RNA isolation kit (Macherey-Nagel, France) using a modified protocol (18).

The yield and the purity of the extracted tNA and RNA samples were assessed by optical density measurements using a NanoQuant Plate on an Infinity spectrophotometer (Tecan, Switzerland). RNA integrity was estimated with an Agilent RNA 6000 Nano kit on an Agilent 2100 bioanalyzer (Agilent Technologies, Santa Clara, CA) according to the manufacturer's recommendations. Results were analyzed using 2100 Expert software, version B.02.07.SI482 (Beta) (Agilent Technologies, Waldbronn, Germany). Indicators of RNA integrity were the RNA integrity number (RIN; maximum value, 10), the 23S rRNA/16S rRNA ratio (optimum value, 2), and the absence of degradation (low baseline) (19). Total RNA (1 µg) was reverse transcribed in cDNA using random primers (Promega, Madison, WI) (0.5 µg/µg RNA), as described by Popova et al. (16).

**Microbial community quantification and gene expression.** Real-time quantitative PCR (qPCR) was carried out using a StepOne system (Applied Biosystems, Courtabeuf, France). Assays were performed in triplicate using SYBR Premix *Ex Taq* (TaKaRa Bio Inc., Otsu, Japan) and used 20 ng of tNA or 2 µl of total cDNA. Negative controls without DNA template were run with every assay to assess overall specificity.

Primers used to quantify *mcrA* and *rrs* DNA or cDNA and reaction setup and temperature cycles were as described by Popova et al. (16). The amplification efficiency, slope, and R<sup>2</sup> of qPCR assays targeting the *mcrA* gene were 91.8%, -3.53, and 0.999, respectively. The amplification efficiency, slope, and R<sup>2</sup> of qPCR assays targeting the *rrs* gene were 95%, -3.35, and 0.999, respectively.

Absolute quantification involved the use of standard curves that had been prepared with PCR products corresponding to a partial sequence of the *mcrA* gene of *Methanobrevibacter ruminantium* DSM 1093 and almost the entire sequence of the *rrs* gene of *Prevotella bryantii* B14 (DSM 11371). Standard curves were created using triplicate, 10-fold dilutions series ranging from 2.5 × 10<sup>2</sup> to 2.5 × 10<sup>8</sup> copies for the *mcrA* gene and from 1 × 10<sup>2</sup> to 1 × 10<sup>8</sup> copies for the bacterial *rrs* gene. Gene copies in ruminal and cecal content samples were quantified, and results were expressed as numbers of gene copies per g dry matter (DM) of content or numbers of gene copies in total ruminal or cecal DM contents.

Expression of the functional *mcrA* gene was assessed using relative quantification by the threshold cycle (C<sub>T</sub>) of the qPCR. Levels of expression of the *mcrA* gene were compared in animals fed different diets using the ΔC<sub>T</sub> (see below) and 2<sup>-ΔC<sub>T</sub></sup> values (20) for the corn and wheat diets. Fold change in *mcrA* gene expression in the contents of the cecum compared to the contents of the rumen of the same animal was computed using the 2<sup>-ΔΔC<sub>T</sub></sup> method as follows:

$$\Delta C_T = (C_T mcrA - C_T rrs) \quad (1)$$

$$\Delta\Delta C_T = (C_T mcrA - C_T rrs)_{\text{cecum}} - (C_T mcrA - C_T rrs)_{\text{rumen}} \quad (2)$$

Statistical analysis of gene expression was done on the ΔC<sub>T</sub> and 2<sup>-ΔC<sub>T</sub></sup> values as indicated by Schmittgen and Livak (20).

**Microbial diversity analysis by PCR-DGGE.** The *mcrA* and *rrs* genes were used as molecular markers to assess, respectively, methanogen and bacterial diversity by PCR-denaturing gradient gel electrophoresis (PCR-DGGE). For PCR amplification, 50 ng of total nucleic acids and 5  $\mu$ l of cDNA samples were used. PCR amplifications and DGGE analysis were performed as described before (16).

Images of DGGE gels were acquired using an calibrated optical-density scanner (Epson, France) at a spatial resolution of 600 dots per inch (dpi). Images were analyzed with GelCompar II version 4.0 (Applied Maths, Kortrijk, Belgium). Banding profiles were quantified within each profile by determining the total number of bands, the peak surface of each band, and the sum of all the peak surfaces of all bands (21). This was used to calculate the community biodiversity using three indices, the Shannon index, the dominance index, and the evenness index, according to Fromin et al. (21). To normalize for differences among gels, *mcrA* or *rrs* PCR products of a sample taken from a wether outside the study were run on every gel and used as standards for comparisons of gels. The similarities of standards between gels were 88% for the *mcrA* gene and 91% for the bacterial *rrs* gene.

DGGE profiles were compared using hierarchical clustering to join similar profiles into groups (21). To this end, all images of DGGE gels were matched using the standards and bands were quantified after local-background subtraction. A tolerance in the band position of 1% was applied. The similarity among profiles was calculated with the Pearson product-moment correlation coefficient, and the clustering was done with the unweighted-pair group method using average linkages (UPGMA).

**Clone library construction, sequencing, and analysis of nucleotide sequences.** Methanogen *mcrA* gene libraries were constructed by cloning PCR-amplified products from tNA samples pooled by digestive compartment (rumen or cecum) and diet (corn or wheat). Four clone libraries (rumen-corn [RC], rumen-wheat [RW], cecum-corn [CC], and cecum-wheat [CW]) were created using pGEM-T Easy Vector System II (Promega, Madison, WI), according to the manufacturer's recommendations. Clone libraries were sent to LGC Genomics GmbH (Germany) sequencing services for plasmid purification and DNA sequencing using the M13 forward primer.

Twenty-one completed genomes of methanogenic *Archaea* were retrieved from the NCBI genome database (<http://www.ncbi.nlm.nih.gov/sites/genome>; accessed 22 September 2011). Strains belonged to all six known orders of methanogens, representing a total of 8 families. There were 56 sequences corresponding to *rrs* (average length, 1,468 bp) and 21 sequences corresponding to *mcrA* and MCR (methyl coenzyme M reductase alpha subunit) (average lengths, 1,673 bp and 557 amino acids). *mcrA* sequences were truncated to a fragment limited by a pair of widely used primers (22) (named *mcrA*-Luton); this DNA fragment was *in silico* translated to MCR-Luton amino acid sequences. Muscle (23) implemented in the SeaView version 4 program (24) was used to align *rrs*, *mcrA*-Luton, and MCR-Luton fragments. MEGA 5 (25) was used to compute distance matrixes for both *mcrA*-Luton sequences (using the Kimura 2-parameter algorithm) and MCR-Luton sequences (using the Dayhoff model). We used different species cutoff limits (unique and from 0.01 to 0.06) for *rrs* in MOTHUR and confirmed for this data set that 0.02 (98% similarity) was the best-adapted value to define archaeal species as previously reported (26). Pairwise comparisons of *mcrA*-Luton and MCR-Luton fragments and *rrs* sequences were performed to calculate the nucleotide or amino acid difference and then the percent sequence identity. The similarities of the 21 *mcrA*-Luton, MCR-Luton, and *rrs* sequences were linearly correlated and used to define distance cutoff values for methanogenic species-level operational taxonomic units (OTUs) (see Fig. S1 in the supplemental material). The 0.02 species cutoff value based on the *rrs* gene corresponded with cutoff values of 0.13 (i.e., 87% identity) ( $R^2 = 0.79$ ) and 0.06 (i.e., 94% identity) ( $R^2 = 0.79$ ) for *mcrA*-Luton and MCR-Luton, respectively.

Sequences obtained in this study were compared with entries in the GenBank nr (nonredundant) database using BLASTn. Chimeras were

TABLE 1 Methane output from lambs fed wheat- or corn-based diets<sup>a</sup>

Parameter	Methane output for indicated diet		SEM	P value <sup>b</sup>
	Wheat	Corn		
g/day	12.84	8.19	1.254	<0.05
g/kg DM intake	15.17	11.97	0.763	<0.05
g/kg NDF intake	41.16	32.93	4.212	NS
g/kg starch intake	31.56	25.32	2.852	NS
g/kg digested DM	25.28	20.92	1.683	0.10
g/kg digested NDF	101.16	108.29	23.730	NS
g/kg digested starch	35.84	27.78	3.370	NS

<sup>a</sup> Values represent means of the results determined with 6 lambs.

<sup>b</sup> NS, not significant.

identified using Bellerophon (27). After elimination of nonmethanogen and chimeric sequences, there were 68, 83, 77, and 71 sequences in the RW, RC, CW, and CC clone libraries, respectively. Sequences were aligned with Muscle (23) implemented in the SeaView version 4 program (24), and evolutionary distance matrices were calculated according to the Kimura 2-parameter algorithm implemented in MEGA 5 (25). Sequences were assigned to OTUs on the basis of 87% (*mcrA*) sequence identity using the furthest-neighbor algorithm in mothur v.1.21.1 (28). Rarefaction curves were constructed to check the sampling effort. Community structures of the libraries were compared using the libshuff command of mothur. For phylogenetic reconstruction, representative sequences, one from each OTU, were combined with *mcrA* sequences retrieved from GenBank representing major archaeal phylogenetic groups. Sequences were aligned with Muscle (23), and phylogenetic trees were constructed in MEGA 5 (25) using the neighbor-joining method; trees were bootstrap reassembled 1,000 times.

**Calculations and statistical analysis.** Hydrogen recoveries for the *ex situ* and *in vitro* assays of both rumen and cecum inocula (total contents without substrate and strained fluids with substrate) were calculated as described by Demeyer et al. (29).

All data were analyzed using PROC MIXED of SAS version 9.1.2 (30). To determine effects on intake, total tract digestion, counts of protozoa, and *in vivo* methane output, the model included the diet (D) and experimental group (EG) as fixed effects with animal within diet as a random effect. The EG effect was not significant, so it is not presented here. The model for statistical analysis of fermentation parameters, molecular quantification data, and data from *ex situ* incubations of total contents included the diet (D) and fermentative compartment (FC) and their interaction (D  $\times$  FC) as fixed effects with animal within diet as a random effect. For data from *in vitro* incubations of strained fluids, the model included the D, FC, and substrate (S) and their interactions (D  $\times$  FC, D  $\times$  S, FC  $\times$  S, and D  $\times$  FC  $\times$  S) as fixed effects with animal within diet as a random effect. In the latter case, interactions were not significant and they are not presented in this paper. Effects were declared significant at  $P \leq 0.05$  and trends were declared for  $P \leq 0.10$ .

**Nucleotide sequence accession numbers.** The sequences from this study have been deposited in the GenBank database under the accession numbers JX488056 to JX488138 and JX894252 to JX894470.

## RESULTS

### Dry matter intake, total tract digestibility, and methane output.

Diet had no effect on intake or total tract digestion of DM, NDF, and starch ( $P > 0.05$ ; data not shown). Mean daily weight gains were similar between the two diets (0.19 kg/day for corn versus 0.17 kg/day for wheat diet,  $P > 0.05$ ). In contrast, lambs fed the corn-based diet had lower daily methane emissions (CH<sub>4</sub> g/day) and methane yield (CH<sub>4</sub> g/kg DMI) (36% [ $P < 0.05$ ] and 21% [ $P < 0.05$ ], respectively) than wheat-fed animals (Table 1).

TABLE 2 Characteristics of digestive contents and *in vivo* fermentation parameters in lambs fed wheat- or corn-based diets<sup>a</sup>

Parameter	Value for indicated diet <sup>b</sup>				SEM	<i>P</i> value <sup>c</sup>		
	Wheat		Corn			D	FC	D × FC
	Rumen	Cecum	Rumen	Cecum				
<b>Digestive content</b>								
Total wet wt (g)	3,023.30a	430.45b	3,921.70a	273.33c	20.336	<0.01	<0.01	<0.01
Total dry wt (g)	423.24a	76.13b	622.23c	55.14b	45.105	0.087	<0.01	<0.05
Dry matter (%)	14.00	17.80	15.85	20.67	1.147	NS	<0.01	NS
<b>Fermentation</b>								
pH	6.66	6.79	6.68	6.52	0.088	NS	NS	NS
Total VFA (mM)	66.91	46.91	68.64	57.68	4.997	NS	<0.01	NS
Acetate (%) <sup>d</sup>	69a	78b	68a	76b	15.2	NS	<0.01	NS
Propionate (%) <sup>d</sup>	17	16	19	13	1.6	NS	0.06	0.10
Butyrate (%) <sup>d</sup>	12a	1b	8a	10a	14.1	NS	<0.05	<0.01
Minor VFA (%) <sup>d,e</sup>	3	2	4	2	0.3	NS	<0.01	NS
Acetate/propionate ratio	4.43	5.03	3.75	7.12	0.851	NS	NS	NS

<sup>a</sup> Values represent means of the results determined with 6 lambs.

<sup>b</sup> Different letters (a, b, and c) within the same row indicate significant differences ( $P < 0.05$ ).

<sup>c</sup> D, diet; FC, fermentative compartment (rumen or cecum); D × FC, interaction between diet and fermentative compartment; NS, not significant.

<sup>d</sup> Data represent % mM of individual VFA per mM of total VFA produced.

<sup>e</sup> Minor VFA are valerate, isovalerate, isobutyrate, and caproate.

***In vivo* fermentation patterns.** The ruminal and cecal contents of corn-fed lambs were, respectively, 1.4-fold heavier ( $P < 0.01$ ) and 1.6-fold lighter ( $P < 0.01$ ) than the corresponding contents of wheat-fed lambs (Table 2). Irrespective of the diet, the VFA concentration in the cecum was about 30% of that found in the rumen but with a higher proportion of acetate ( $P < 0.01$ ).

***Ex situ* and *in vitro* fermentation patterns.** Diet did not influence *ex situ* fermentation potential, with the exception of total

gas, which tended to be higher ( $P = 0.101$ ) in digestive contents from corn-fed lambs (Table 3). The main differences were observed between compartments, with cecal contents producing ~4-fold less methane and ~20% less VFA ( $P < 0.01$ ) than equal volumes of ruminal contents. The hydrogen recovery was also significantly lower with cecum inoculates ( $P < 0.01$ ).

The capacity of ruminal and cecal strained fluids to degrade readily fermentable substrate *in vitro* is summarized in Table 4.

TABLE 3 *Ex situ* methanogenic and fermentative potential of total rumen and cecal contents from lambs receiving wheat- or corn-based diets<sup>a</sup>

Parameter	Diet				SEM	<i>P</i> value <sup>b</sup>	
	Wheat		Corn			D	FC
	Rumen	Cecum	Rumen	Cecum			
<b>Gas production at 6 h of incubation</b>							
Total (μmol/h)	288.53	272.69	343.61	309.36	26.703	0.101	NS
Methane (μmol/h)	36.64	6.89	33.30	6.93	3.379	NS	<0.01
Methane (%) <sup>c</sup>	11.18	4.09	9.68	3.16	0.0001	NS	<0.01
<b>Gas production at 24 h of incubation</b>							
Total (μmol/h)	146.44	154.30	188.25	164.54	12.155	<0.05	NS
Methane (μmol/h)	28.03	7.43	26.46	6.39	2.71	NS	<0.01
Methane (%) <sup>c</sup>	14.86	6.77	13.63	5.54	0.673	NS	<0.01
Hydrogen recovery (%)	73.45	56.72	77.71	47.85	4.510	NS	<0.01
<b>VFA production at 24 h of incubation</b>							
Total (mM)	110.84	91.15	115.83	88.49	6.697	NS	<0.01
Acetate (%) <sup>d</sup>	61.10	64.00	61.00	61.00	1.928	NS	NS
Propionate (%) <sup>d</sup>	20.75	21.75	23.00	25.33	2.119	NS	NS
Butyrate (%) <sup>d</sup>	11.50	9.56	10.50	10.00	1.486	NS	NS
Acetate/propionate ratio	3.03	3.43	2.66	2.49	0.543	NS	NS
Minor VFA (%) <sup>d,e</sup>	6.34	4.89	5.50	4.00	0.518	NS	<0.05
pH	5.89	6.03	5.56	5.79	0.138	NS	NS

<sup>a</sup> Values represent the means of the results of 12 observations.

<sup>b</sup> D, diet; FC, fermentative compartment. The D × FC interaction was never significant ( $P > 0.05$ ). NS, not significant.

<sup>c</sup> μmol of CH<sub>4</sub> per μmol of total gas produced.

<sup>d</sup> % mM of individual VFA per mM of total VFA produced.

<sup>e</sup> Minor VFA are valerate, isovalerate, isobutyrate, and caproate.

TABLE 4 Capacity of strained rumen and cecum liquids from lambs fed wheat- or corn-based diets to degrade readily fermentable substrates *in vitro*<sup>a</sup>

Parameter	Value for indicated diet and substrate								SEM	P value <sup>b</sup>		
	Wheat				Corn							
	Rumen		Cecum		Rumen		Cecum			D	FC	S
	Wheat	Corn	Wheat	Corn	Wheat	Corn	Wheat	Corn				
Gas production at 6 h of incubation												
Total ( $\mu\text{mol/h}$ )	440.40	357.05	382.55	302.66	367.64	264.83	379.96	279.24	45.872	NS	NS	<0.01
Methane ( $\mu\text{mol/h}$ )	49.75	37.82	25.90	19.52	36.00	23.57	9.03	6.61	5.344	<0.01	<0.01	<0.05
Methane (%) <sup>c</sup>	11.18	10.46	6.28	6.10	9.47	8.43	2.60	2.40	0.873	<0.01	<0.01	<0.05
Gas production at 24 h of incubation												
Total ( $\mu\text{mol/h}$ )	548.47	493.79	465.47	398.28	485.38	397.97	458.20	378.63	40.417	NS	= 0.06	<0.05
Methane ( $\mu\text{mol/h}$ )	660.60	645.18	315.22	322.35	567.47	498.07	156.25	164.74	55.042	<0.05	<0.01	NS
Methane (%) <sup>c</sup>	13.33	13.12	7.62	8.11	12.28	11.73	4.92	5.44	0.001	0.06	<0.01	NS
Hydrogen recovery (%)	68.11	74.10	50.43	53.24	72.57	70.66	45.36	47.67	3.500	NS	<0.01	0.101
VFA production at 24 h of incubation												
Total (mM)	122.18	119.47	118.48	111.03	96.96	104.12	96.47	86.59	5.205	<0.01	NS	NS
Acetate (%) <sup>d</sup>	58.04	54.17	63.10	60.48	58.90	53.03	59.75	59.30	7.455	NS	<0.05	<0.01
Propionate (%) <sup>d</sup>	23.10	25.18	17.17	16.84	21.37	22.07	25.20	24.95	7.59	NS	NS	NS
Butyrate (%) <sup>d</sup>	13.70	15.04	15.26	17.34	13.63	19.30	11.25	11.65	9.85	NS	NS	<0.01
Acetate/butyrate ratio	2.76	2.30	4.72	4.09	2.97	2.50	2.38	2.38	0.642	NS	NS	NS
Minor VFA (%) <sup>d,e</sup>	6.21	6.70	5.39	5.88	5.94	5.91	3.63	3.55	1.124	NS	NS	NS

<sup>a</sup> Values represent the means of the results of 6 observations.

<sup>b</sup> D, diet; FC, fermentative compartment; S, substrate. The double and triple interactions between D, FC, and S were never significant ( $P > 0.05$ ). NS, not significant.

<sup>c</sup>  $\mu\text{mol}$  of  $\text{CH}_4$  per  $\mu\text{mol}$  of total gas produced.

<sup>d</sup> % mM of individual VFA per mM of total VFA produced.

<sup>e</sup> Minor VFA are valerate, isovalerate, isobutyrate, and caproate.

Methane production ( $\mu\text{mol/h}$ ) was higher ( $P < 0.01$ ) with strained ruminal or cecal fluids from wheat-fed lambs than with that from corn-fed lambs. Incubation of cecal contents always yielded less methane and a poorer hydrogen recovery than incubation of ruminal contents ( $P < 0.01$ ). Similar to the methane results, total VFA concentrations were lower ( $P < 0.01$ ) in vials inoculated with inocula from corn-fed lambs (Table 4). Acetate molar proportion was higher ( $P < 0.05$ ) in cecal than in ruminal contents.

**Microbial community structure.** Total protozoal numbers tended to increase ( $P = 0.06$ ) in wheat-fed lambs compared to lambs fed the corn diet (Table 5). Vestibuliferida (*Isostricha* and *Dasytricha*) increased with wheat. Bacterial numbers (*rrs* copy numbers) were similar ( $P > 0.05$ ) in the rumens of wheat- and corn-fed lambs and were higher ( $P < 0.05$ ) in the cecum of wheat-fed lambs than in the cecum of corn-fed lambs (Table 5). Diet did not induce changes in the bacterial community diversity as measured using PCR-DGGE. In contrast, PCR-DGGE analysis partially separated rumen and cecum samples (Fig. 1). Cecum samples grouped more closely than did rumen samples. The Shannon index, the dominance index, and the evenness index values were similar between diets and digestive compartments (data not shown).

Methanogens (*mcrA* copies), as for protozoa, were higher ( $P < 0.01$ ) in digestive contents of animals fed wheat relative to lambs fed corn. Whatever the diet, the concentration with respect to *mcrA* gene copy numbers (number of copies/g DM contents) and the total numbers of *mcrA* and *rrs* copies were higher ( $P < 0.01$ ) in ruminal relative to cecal contents (Table 5).

The expression of the bacterial *rfs* gene remained constant in ruminal and cecal contents, as assessed by a Student's test on the *rfs*  $2^{-\Delta CT}$  values (data not shown), allowing us to use it as a calibrator for an *mcrA* expression study (31). The *mcrA*  $\Delta C_T$  values were higher ( $P < 0.05$ ) in the cecum than in the rumen for both groups of animals. The *mcrA*  $2^{-\Delta CT}$  values were also higher, but the difference was not significant (Table 5). Nonetheless, the  $2^{-\Delta CT}$  values in cecal contents were 2 times higher in corn-fed lambs than in wheat-fed lambs. In addition, expression of the *mcrA* gene decreased by more than 2 times ( $2^{-\Delta \Delta CT} = -2.167$ ) in the cecum of corn-fed lambs compared to their rumen. In wheat-fed lambs, *mcrA* expression in the cecum was 4 times lower than in the rumen ( $2^{-\Delta \Delta CT} = -4.091$ ).

Cluster analysis of the *mcrA* DNA sequences from PCR-DGGE profiles partially separated the rumen from the cecum samples (Fig. 2). Three main clusters were formed, one grouping 7 rumen samples, another grouping 6 cecum samples, and the third formed with 4 rumen and 4 cecum samples. Although, within each cluster, samples tended to group by diet, the separation was not clear. Neither the number of bands nor any of the diversity indices calculated from the PCR-DGGE profiles presented significant differences related to diet or fermentative compartment (data not shown). However, several bands were uniquely found either in the ruminal or the cecal contents.

We constructed clone libraries of the *mcrA* gene to investigate the phylogeny of methanogenic *Archaea* in ruminal and cecal contents of wheat- and corn-fed lambs. Rarefaction curves for all four clone libraries tended to reach the horizontal asymptote for the 87% species cutoff value, which we calculated was equivalent to

TABLE 5 Counts of protozoa and absolute and relative qPCR quantification results<sup>a</sup>

Parameter	Value for indicated diet <sup>b</sup>				SEM	P value <sup>c</sup>		
	Wheat		Corn			D	FC	D × FC
	Rumen	Cecum	Rumen	Cecum				
Total protozoa (log cells/ml)	5.89	/ <sup>d</sup>	5.55	/	0.108	0.06	/	/
Small (<100- $\mu$ m) entodiniomorphs (log cells/ml)	5.89	/	5.54	/	0.102	<0.05	/	/
Large (>100- $\mu$ m) entodiniomorphs (log cells/ml)	3.84	/	3.44	/	0.263	NS	/	/
<i>Dasytricha</i> (log cells/ml)	3.93	/	2.72	/	0.146	<0.01	/	/
<i>Isotricha</i> (log cells/ml)	4.25	/	2.81	/	0.187	<0.01	/	/
Gene quantification								
<i>mcrA</i> (log copies/g DM)	8.81	8.34	8.46	7.66	0.088	<0.01	<0.01	0.08
<i>rrs</i> (log copies/g DM)	12.10a	12.18a	12.04a	11.77b	0.078	<0.05	NS	<0.05
<i>mcrA</i> (log total no.)	11.41a	10.20b	11.26a	9.40c	0.095	<0.01	<0.01	<0.05
<i>rrs</i> (log total no.)	14.70a	14.06b	14.82a	13.51c	0.074	<0.05	<0.01	<0.01
<i>mcrA/rrs</i> ( $\times 10^{-2}$ )	5.16a	1.42b	2.89c	0.80b	2.3	<0.01	<0.01	<0.05
<i>mcrA</i> gene expression								
$\Delta CT$	10.45	12.79	11.21	12.01	0.874	NS	<0.05	NS
$2^{-\Delta CT}$ ( $\times 10^{-5}$ )	11.1	4.40	17.1	9.30	5.55	NS	NS	NS

<sup>a</sup> Values represent the means of the results determined with 6 lambs.

<sup>b</sup> Values with different letters (a, b, and c) in the same row are significantly different.

<sup>c</sup> D, diet; FC, fermentative compartment; D × FC, interaction between diet and fermentative compartment; NS, not significant.

<sup>d</sup> /, the population of protozoa was not counted in the cecal contents.

the 98% species cutoff value of the *rrs* gene. For higher cutoff values, the asymptote was not reached (see Fig. S2 in the supplemental material). Pairwise comparisons by the libshuff test yielded significance values < 0.0043, attesting that our four sequence collections were all different. We used an 87% species cutoff value for *mcrA* sequences, and the 299 sequences used in this study grouped into 16 distinct OTUs. Clones were unevenly distributed between OTUs; the majority of the sequences from each library grouped into a single OTU closely related to the *Methanobrevibacter millerae*-*M. gottschalkii*-*M. smithii* clade (Fig. 3). Close relatives to the *M. ruminantium* clade represented, respectively, 29%, 12%, 18%, and 10% of the RC, RW, CC, and CW

libraries. In contrast, members of the *Methanomicrobiales* order were detected only in CW and RC libraries (14% and 12% of the sequences, respectively). All four libraries contained sequences that clustered with uncultured archaeal clones belonging to rumen cluster C.

## DISCUSSION

We hypothesized that readily degradable starch from wheat would reduce overall feed degradation in the rumen, increase organic matter arrival and fermentation in the cecum, and thus would induce changes in the two digestive microbiota. The aim of this study was to describe the ruminal and cecal microbiota, particu-




FIG 1 PCR-DGGE profiles of *rrs* DNA sequences of rumen bacteria from two fermentative compartments (FC) of lambs (animal numbers shown) fed a wheat- or corn-based diet. The similarity among profiles was calculated with the Pearson product-moment correlation coefficient, and the clustering was done with the unweighted-pair group method using average linkages (UPGMA).






FIG 2 PCR-DGGE profiles of *mcrA* DNA sequences of rumen methanogens from two fermentative compartments (FC) of lambs (animal numbers shown) fed a wheat- or corn-based diet. The similarity among profiles was calculated with the Pearson product-moment correlation coefficient, and the clustering was done with the unweighted-pair group method using average linkages (UPGMA).

larly, methanogenic *Archaea*, of growing lambs fed two contrasting diets in term of starch degradability. In order to link microbial community structure and function, we measured *in vivo* methane production and followed VFA patterns in the rumen and cecum. *Ex situ* and *in vitro* methane production was used to compare, respectively, methanogenic potential of ruminal and cecal contents and their capacity to degrade readily fermentable substrates.

Methanogenic *Archaea* are usually studied by targeting two main molecular markers—the *rrs* gene coding for the 16S rRNA and *mcrA* gene coding for a subunit of MCR (22), a crucial enzyme for the terminal step of methanogenesis (32). Targeting a gene which codes for an enzyme also offers the potential for development of activity-based detection methods by analysis of mRNA. Thus, molecular biology tools that target the *mcrA* gene in DNA and RNA extracts are useful to simultaneously highlight the diversity and activity of methanogens in different environments (16, 33).

Diet did not influence animal performances (intake, growth), but methane production and methane yield were higher in wheat-fed than in corn-fed lambs. Numbers of rumen protozoa tended to increase in the rumen of lambs fed the wheat diet, likely due to the greater amount of ruminal readily degradable starch (16, 34). Protozoal density is positively related to enteric methane emissions (reviewed in reference 35) and may help explain the higher methane produced with the wheat diet. In contrast, the concentration of methanogens in the rumen was not affected by diet. The lack of correlation between numbers of methanogens in the rumen and methane production was also observed with other mitigation strategies, such as removal of protozoa or lipid supply (reviewed in reference 35). A reduction in numbers of methanogens associated with methane reduction was instead reported with compounds toxic to methanogens (36, 37). When no changes in methanogen abundance are observed, higher methanogenesis could be attributed to increases in the metabolic activity of methanogens. A positive correlation between methanogenesis and *mcrA* gene expression has already been reported *in vivo* for beef

cattle fed different high-grain-content diets (16) and *in vitro* with a tea saponin supplementation (11). In the current trial,  $2^{-\Delta CT}$  values of ruminal contents of wheat-fed lambs were 1.5 times higher than  $2^{-\Delta CT}$  values in corn-fed lambs. This difference was not statistically significant, but it should be noted that rumen contents were sampled at slaughter and that animals were last fed the morning of the day before slaughter. Methanogenic activity under these conditions would be at its lowest.

In our study, *Methanobacteriales*-related sequences were dominant in both the ruminal and the cecal contents. *Methanobacteriales* have been reported to be the predominant methanogens in the bovine (38, 39) and ovine (40) rumen. No dominant archaeal group has been identified in other sampling locations of the gastrointestinal tract in these ruminants (38). Clone library analysis suggested the presence of diet-induced changes of the methanogenic community composition in the rumen. The majority of the sequences from the RW library (78%) belonged to the *M. millerae*-*M. gottschalkii*-*M. smithii* clade. In addition, it should be noted that this library did not present sequences related to the *Methanomicrobiales* order. In contrast, the RC library had only 52% of the sequences belonging to the *M. millerae*-*M. gottschalkii*-*M. smithii* clade. The other half were distributed between the *M. ruminantium* clade, the order of *Methanobacteriales*, and rumen cluster C. On the other hand, it seems that diet had no effect on the composition of the community of methanogens in the cecum of growing lambs, as sequences from the CC and the CW library had similar phylogenetic distributions. However, *Methanomicrobiales*-like sequences were retrieved only in the CW library.

Finally, we observed no diet-induced changes in the rumen bacterial community structure: *rrs* copy concentrations and total numbers and *rrs* PCR-DGGE profiles from the rumens were similar in the two groups of animals. This is in accordance with the resemblances of VFA profiles between the rumens of corn- and wheat-fed lambs. It has to be noted that PCR-DGGE detects large




FIG 3 Phylogenetic analysis of partial *mcrA* sequences derived from RW (rumen wheat), RC (rumen corn), CW (cecum wheat), and CC (cecum corn) clone libraries. Evolutionary analyses were conducted in MEGA 5. There were a total of 390 positions in the final data set. The evolutionary history was inferred using the neighbor-joining method. The tree was bootstrap resampled 1,000 times; only bootstrap values > 50 are shown. The scale bar equals an average of 0.5 nucleotide substitutions per 100 positions. The GenBank accession numbers for reference nucleotide sequences are given between vertical bars. Numbers in parentheses indicate the number of sequences grouped in the same OTU at 87% species cutoff. *Methanopyrus kandleri* was used as an outgroup to root the tree.

changes in the community whereas subtle variations are unnoticed. The differences between diets observed in the proportions of methanogens might reflect variations in the availability of substrates for methanogenesis and also variations in the microbes producing them. For instance, increases in the rumen cluster C members might be due to a higher availability of methanol, a required substrate for this group (41).

We reported that cecal contents were 7 and 14 times lighter than ruminal contents in wheat- and corn-fed lambs, respectively. In consequence, concentrations of molecular markers (numbers of gene copies/g DM) as well as total numbers of the bacterial and methanogenic populations were also lower. This is consistent with a recent work reporting that microbial numbers are reduced by 8 orders of magnitude from the rumen, through the abomasum, and into the duodenum of lactating dairy cows (42). Interestingly, despite these differences in microbial concentrations between rumen and cecal contents, we did not observe any differences in their *ex situ* total gas production. However, the lower VFA concentrations and the lower methanogenic potential in the cecal contents suggested lower fermentation rates than in the ruminal contents. These observations are in accordance with the lower mean  $2^{-\Delta\Delta CT}$

values, showing that *mcrA* expression was lower in the cecum of both groups of lambs than in their rumen.

In both *ex situ* and *in vitro* trials, we observed higher molar proportions of acetate with cecal than with ruminal inocula. The enhanced production of acetate may have been partly due to reductive acetogenesis. This hydrogen sink pathway of acetate production is an alternative to methanogenesis that was reported in the cecum (43). Lower values for hydrogen recoveries in the cecum (both for total contents and strained-fluid incubations) suggest that reductive acetogenesis was a substantial source of acetate in our cecal samples (43).

Because of the existence of reductive acetogenesis and the absence of hydrogen-producing protozoa in the cecum, we expected differences in the diversity of ruminal and cecal methanogenic *Archaea* community. Cluster analysis of DGGE profiles showed differences in methanogenic communities between fermentative compartments that were confirmed by comparisons of clone libraries using Libshuff. In another study, targeting the archaeal *rrs* gene, the richness index of the archaeal community in the rumen was significantly higher than in the feces (44). In the study of Romero-Pérez et al. (9), the sample type (rumen liquor or rectal

fecal samples) had a larger effect on bacterial communities than diet. Also, the bacterial communities, assessed by capillary electrophoresis single-stranded conformation polymorphism profiles of *rrs* genes, showed different structures between the forestomach and fecal contents of cows (8). These observations could be explained by environmental conditions such as pH and the presence of oxygen but also by the source and amount of the substrate, which differ greatly between the rumen and cecum.

**Conclusions.** Data presented here showed that the nature of dietary starch greatly influenced methane production by growing lambs. Differences in methane emissions between lambs fed corn- or wheat-based diet are probably due to differences in the ruminal digestion of substrates. However, effects of diet on the microbial ecosystem of the rumen and of the cecum were less obvious. Our study is among the first aiming to characterize methanogenic community structure (in terms of numbers and diversity) and activity in the two microbial fermentation compartments of the ruminant digestive tract. Our results suggest that several archaeal species would be better adapted to either the rumen or cecum environment. Further research using high-throughput sequencing would be required for detailed studies of this community. Also, the contribution of other microbial populations of the cecum to the nutrition and health of ruminants is seldom addressed but warrants further investigation.

## ACKNOWLEDGMENTS

We thank the staff of the animal experimental facilities and the slaughterhouse of the INRA's Herbivores Research Unit for animal care, help in methane measurements, and sampling, as well as D. Graviou, G. Gentes, L. Genestoux, P. Savajols, and Y. Rochette for help in sample analysis.

M. Popova was the recipient of an INRA-Région Auvergne scholarship.

## REFERENCES

- O'Mara FP. 2011. The significance of livestock as a contributor to global greenhouse gas emissions today and in the near future. *Anim. Feed Sci. Technol.* 166:7–15.
- Murray RM, Bryant AM, Leng RA. 1976. Rates of production of methane in the rumen and large intestine of sheep. *Br. J. Nutr.* 36:1–14.
- Beauchemin KA, McAllister TA, McGinn SM. 2009. Dietary mitigation of enteric methane from cattle. *CAB Rev.* 4:1–18.
- Wright A-DG, Williams AJ, Winder B, Christophersen CT, Rodgers SL, Smith KD. 2004. Molecular diversity of rumen methanogens from sheep in western Australia. *Appl. Environ. Microbiol.* 70:1263–1270.
- Wright ADG, Kennedy P, O'Neill CJ, Toovey AF, Popovski S, Rea SM, Pimm CL, Klein L. 2004. Reducing methane emissions in sheep by immunization against rumen methanogens. *Vaccine* 22:3976–3985.
- Beauchemin KA, McGinn SM. 2005. Methane emissions from feedlot cattle fed barley or corn diets. *J. Anim. Sci.* 83:653–661.
- Martin C, Philippeau C, Michalet-Doreau B. 1999. Effect of wheat and corn variety on fiber digestion in beef steers fed high-grain diets. *J. Anim. Sci.* 77:2269–2278.
- Michelland RJ, Monteils V, Zened A, Combes S, Cauquil L, Gidenne T, Hamelin J, Fortun-Lamothe L. 2009. Spatial and temporal variations of the bacterial community in the bovine digestive tract. *J. Appl. Microbiol.* 107:1642–1650.
- Romero-Pérez GA, Ominski KH, McAllister TA, Krause DO. 2011. Effect of environmental factors and influence of rumen and hindgut biogeography on bacterial communities in steers. *Appl. Environ. Microbiol.* 77:258–268.
- Martin C, Rouel J, Jouany JP, Doreau M, Chilliard Y. 2008. Methane output and diet digestibility in response to feeding dairy cows crude linseed, extruded linseed, or linseed oil. *J. Anim. Sci.* 86:2642–2650.
- Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74:3583–3597.
- AFNOR. 1985. Dosage de l'amidon. Méthode polarimétrique, p 123–125. *In* *Aliments des animaux. Méthodes d'analyses françaises et communautaires*, 2nd ed. Association Française de Normalisation, Paris, France.
- Johnson K, Huyler M, Westberg H, Lamb B, Zimmerman P. 1994. Measurement of methane emissions from ruminant livestock using a sulfur hexafluoride tracer technique. *Environ. Sci. Technol.* 28:359–362.
- Lettat A, Nozière P, Silberberg M, Morgavi DP, Berger C, Martin C. 2010. Experimental feed induction of ruminal lactic, propionic, or butyric acidosis in sheep. *J. Anim. Sci.* 88:3041–3046.
- Morgavi DP, Boudra H, Jouany JP, Graviou D. 2003. Prevention of patulin toxicity on rumen microbial fermentation by SH-containing reducing agents. *J. Agric. Food Chem.* 51:6906–6910.
- Popova M, Martin C, Eugène M, Mialon MM, Doreau M, Morgavi DP. 2011. Effect of fibre- and starch-rich finishing diets on methanogenic *Archaea* diversity and activity in the rumen of feedlot bulls. *Anim. Feed Sci. Technol.* 166:113–121.
- Goering HK, Van Soest PJ. 1970. Forage fiber analysis (apparatus, reagents, procedures, and some applications). U.S. Agricultural Research Service, Washington, DC.
- Popova M, Martin C, Morgavi DP. 2010. Improved protocol for high-quality coextraction of DNA and RNA from rumen digesta. *Folia Microbiol. (Praha)* 55:368–372.
- Schroeder A, Mueller O, Stocker S, Salowsky R, Leiber M, Gassmann M, Lightfoot S, Menzel W, Granzow M, Ragg T. 2006. The RIN: an RNA integrity number for assigning integrity values to RNA measurements. *BMC Mol. Biol.* 7:3. doi:10.1186/1471-2199-7-3.
- Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative Ct method. *Nat. Protoc.* 3:1101–1108.
- Fromin N, Hamelin J, Tarnawski S, Roesti D, Jourdain-Miserez K, Forestier N, Teysier-Cuvelle S, Gillet F, Aragno M, Rossi P. 2002. Statistical analysis of denaturing gel electrophoresis (DGE) fingerprinting patterns. *Environ. Microbiol.* 4:634–643.
- Luton PE, Wayne JM, Sharp RJ, Riley PW. 2002. The *mcrA* gene as an alternative to 16S rRNA in the phylogenetic analysis of methanogen populations in landfill. *Microbiology* 148:3521–3530.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32:1792–1797.
- Gouy M, Guindon S, Gascuel O. 2010. SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* 27:221–224.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28:2731–2739.
- Wright A-DG, Northwood KS, Obispo NE. 2009. Rumen-like methanogens identified from the crop of the folivorous South American bird, the hoatzin (*Opisthocomus hoazin*). *ISME J.* 3:1120–1126.
- Huber T, Faulkner G, Hugenholtz P. 2004. Bellerophon: a program to detect chimeric sequences in multiple sequence alignments. *Bioinformatics* 20:2317–2319.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75:7537–7541.
- Demeyer DI, Fiedler D, De Graeve KG. 1996. Attempted induction of reductive acetogenesis into the rumen fermentation in vitro. *Reprod. Nutr. Dev.* 36:233–240.
- SAS Institute, Inc. 2003. Statistical analysis system, Windows version 9.1.2. SAS Institute, Inc., Cary, NC.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta Ct}$  method. *Methods* 25:402–408.
- Thauer RK. 1998. Biochemistry of methanogenesis: a tribute to Marjory Stephenson. 1998 Marjory Stephenson Prize Lecture. *Microbiology* 144(Pt 9):2377–2406.
- Freitag TE, Prosser JI. 2009. Correlation of methane production and functional gene transcriptional activity in a peat soil. *Appl. Environ. Microbiol.* 75:6679–6687.
- Mackie RI, Gilchrist FMC, Robberts AM, Hannah PE, Schwartz HM. 1978. Microbiological and chemical changes in the rumen during the stepwise adaptation of sheep to high concentrate diets. *J. Agric. Sci.* 90:241–254.

35. Morgavi DP, Forano E, Martin C, Newbold CJ. 2010. Microbial ecosystem and methanogenesis in ruminants. *Animal* 4:1024–1036.
36. Denman SE, Tomkins NW, McSweeney CS. 2007. Quantitation and diversity analysis of ruminal methanogenic populations in response to the antimethanogenic compound bromochloromethane. *FEMS Microbiol. Ecol.* 62:313–322.
37. Knight T, Ronimus RS, Dey D, Tootill C, Naylor G, Evans P, Molano G, Smith A, Tavendale M, Pinares-Patiño CS, Clark H. 2011. Chloroform decreases rumen methanogenesis and methanogen populations without altering rumen function in cattle. *Anim. Feed Sci. Technol.* 166:101–112.
38. Lin C, Raskin L, Stahl DA. 1997. Microbial community structure in gastrointestinal tracts of domestic animals: comparative analyses using rRNA-targeted oligonucleotide probes. *FEMS Microbiol. Ecol.* 22:281–294.
39. Zhou M, Hernandez-Sanabria E, Guan LL. 2009. Assessment of the microbial ecology of ruminal methanogens in cattle with different feed efficiencies. *Appl. Environ. Microbiol.* 75:6524–6533.
40. Wright A-D, Ma X, Obispo N. 2008. *Methanobrevibacter* phylotypes are the dominant methanogens in sheep from Venezuela. *Microb. Ecol.* 56:390–394.
41. Paul K, Nonoh J, Mikulski L, Brune A. 2012. “*Methanoplasmatales*,” *Thermoplasmatales*-related archaea in termite guts and other environments, are the seventh order of methanogens. *Appl. Environ. Microbiol.* 78:8245–8253.
42. Frey JC, Pell AN, Berthiaume R, Lapierre H, Lee S, Ha JK, Mendell JE, Angert ER. 2010. Comparative studies of microbial populations in the rumen, duodenum, ileum and faeces of lactating dairy cows. *J. Appl. Microbiol.* 108:1982–1993.
43. Demeyer D. 1991. Differences in stoichiometry between rumen and hind-gut fermentation. *Adv. Anim. Physiol. Anim. Nutr.* 22:50–61.
44. Michelland RJ, Monteils V, Combes S, Cauquil L, Gidenne T, Fortun-Lamothe L. 2010. Comparison of the archaeal community in the fermentative compartment and faeces of the cow and the rabbit. *Anaerobe* 16:396–401.