



HAL
open science

GH10 xylanase D from *Penicillium funiculosum*: biochemical studies and xylooligosaccharide production

Mickael Lafond, Alexandra Tauzin, Veronique Desseaux, Estelle Bonnin,
El-Hassan Ajandouz, Thierry Giardina

► To cite this version:

Mickael Lafond, Alexandra Tauzin, Veronique Desseaux, Estelle Bonnin, El-Hassan Ajandouz, et al.. GH10 xylanase D from *Penicillium funiculosum*: biochemical studies and xylooligosaccharide production. *Microbial Cell Factories*, 2011, 10, 10.1186/1475-2859-10-20 . hal-02647803

HAL Id: hal-02647803

<https://hal.inrae.fr/hal-02647803>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH

Open Access

GH10 xylanase D from *Penicillium funiculosum*: biochemical studies and xylooligosaccharide production

Mickael Lafond¹, Alexandra Tauzin¹, Véronique Desseaux¹, Estelle Bonnin², El-Hassan Ajandouz¹ and Thierry Giardina^{1*}

Abstract

Background: The filamentous fungus *Penicillium funiculosum* produces a range of glycoside hydrolases (GH). The *XynD* gene, encoding the sole *P. funiculosum* GH10 xylanase described so far, was cloned into the pPICZ α A vector and expressed in methylotrophic yeast *Pichia pastoris*, in order to compare the results obtained with the *P. funiculosum* GH11 xylanases data.

Results: High level expression of recombinant XynD was obtained with a secretion of around 60 mg.L⁻¹. The protein was purified to homogeneity using one purification step. The apparent size on SDS-PAGE was around 64 kDa and was 46 kDa by mass spectrometry thus higher than the expected molecular mass of 41 kDa. The recombinant protein was N- and O-glycosylated, as demonstrated using glycoprotein staining and deglycosylation reactions, which explained the discrepancy in molecular mass. Enzyme-catalysed hydrolysis of low viscosity arabinoxylan (LVAX) was maximal at pH 5.0 with $K_{m(\text{app})}$ and $k_{\text{cat}}/K_{m(\text{app})}$ of 3.7 ± 0.2 (mg.mL⁻¹) and 132 (s⁻¹mg⁻¹.mL), respectively. The activity of XynD was optimal at 80°C and the recombinant enzyme has shown an interesting high thermal stability at 70°C for at least 180 min without loss of activity. The enzyme had an endo-mode of action on xylan forming mainly xylobiose and short-chain xylooligosaccharides (XOS). The initial rate data from the hydrolysis of short XOS indicated that the catalytic efficiency increased slightly with increasing their chain length with a small difference of the XynD catalytic efficiency against the different XOS.

Conclusion: Because of its attractive properties XynD might be considered for biotechnological applications. Moreover, XOS hydrolysis suggested that XynD possess four catalytic subsites with a high energy of interaction with the substrate and a fifth subsite with a small energy of interaction, according to the GH10 xylanase literature data.

Keywords: xylanase GH10, *Penicillium funiculosum*, xylooligosaccharides

1. Introduction

Xylanases (endo-1,4- β -xylanases; EC 3.2.1.8) catalyze the hydrolysis of β -1,4 bonds of xylan, the major hemicellulose component of the plant cell wall. According to the sequence-based glycoside hydrolase (GH) classification (CAZy classification) [1], xylanases are mainly grouped into families 10 (GH10) and 11 (GH11) but they were also found in glycoside hydrolase families 5, 8 and 43 <http://www.cazy.org>. A GH7 enzyme from *Penicillium*

funiculosum with xylanase/cellobiohydrolase activities was also reported and called XynA. GH10 family contains plant, fungal and bacterial enzymes whereas the structurally unrelated GH11 family only includes fungal and bacterial enzymes [1-3]. Xylanases are used in many food and feed sectors such as additives for broiler feed, bread making, in separation processes of gluten from starch in wheat or other cereals gluten from starch, production of juice from fruits or vegetables, extracting more fermentable sugar from barley for brewing, etc [4-6]. Recently, the potential applications of xylanases for bioconversion of hemicellulosic materials to xylo-oligosaccharides (XOS) have gained popularity instead of acid hydrolysis which is

* Correspondence: thierry.giardina@univ-cezanne.fr

¹Université Paul Cézanne, ISM2/Biosciences UMR CNRS 6263 (Université Aix Marseille III/CNRS), service 342, Faculté des Sciences et Techniques Saint-Jérôme, 13397 Marseille Cedex 20, France

Full list of author information is available at the end of the article

not environmentally feasible [7-9]. XOS have been shown to impact on human health by reducing cholesterol level and precancerous lesions, improving the biological availability of calcium and exhibiting a positive effect on gut microbiota [10-12]. The latter positive effect should be due to the enhancement of the production of different prebiotic molecules like short chain fatty acids known to promote the growth of beneficial bacteria such as *Lactobacillus* or *Bifidobacteria* and to lowering the cecal pH value [13-15].

In addition to the health promoting effects, XOS are moderately sweet and stable over a wide range of pH and temperatures and have organoleptic characteristics that make them suitable for incorporation into foods [16,17].

Lignocellulosic biomass degradation used for biotechnological XOS production, needs thermal treatments coupled to thermostable xylanases. Moreover, a large amount of hemicellulosic fraction is insoluble and need action of appropriate enzymes. So, researches concerning new xylanases acting on high temperatures and able to hydrolyze insoluble substrate remain topical.

The filamentous fungi *Penicillium funiculosum* secretes a lot of glycosidases and the recent sequencing of the *P. funiculosum* genome, allowed discriminating about 200 genes encoding proteins with potential glycosidase activity [18]. Among the glycosidases which could be used during biotechnological processes, different types of xylanases were purified and their genes have been cloned [19,20]. XynB (GH11) and XynC (GH11) were recently extensively studied [21,22] whereas little is known concerning XynD the only GH10 member family found to date in this microorganism [20]. GH10 xylanases typically exhibit a molecular weight ≥ 30 kDa and a low *pI* [23]. The crystal structures of several GH10 enzymes showed that the catalytic domain is an 8-fold α/β -barrel forming a 'salad bowl'. Often, one or more extra domains are present, corresponding to carbohydrate binding module (CBM1) which binds to cellulose [24].

The activity of GH10 xylanases on AXs produces shorter XOS than those produced by GH11 xylanases. The former can act near the substituted xylose residue whereas the latter are bothered by additional groups like 4-O-methyl-D glucuronate, acetate and α -L-arabinofuranose, thus restricting the access to the β -1,4-linkages in the xylan backbone [25]. It is worth noting that in the GH10 family, the members are able to produce different end-products due to the difference in their catalytic site [26].

Here we report the molecular characterization of XynD from *Penicillium funiculosum* encoding a 41 kDa GH10 xylanase (XynD). The cDNA cloning and

heterologous expression of XynD in *Pichia pastoris* was performed, and biochemical properties of the recombinant protein were determined, notably showing a natural thermal resistance. The enzyme was also characterized under kinetic viewpoint using various short and long substrates, especially regarding the XOS liberation profile.

2. Materials and methods

Materials

The pPICZ α A expression vector and the *P. pastoris* expression kit used, including the *P. pastoris* strain X-33, zeocin, oligonucleotides and all restriction DNA modifying enzymes (except DNA polymerase) were from Invitrogen (Groningen, Netherlands). PrimeSTARTM HS DNA polymerase for polymerase chain reactions (PCR) was from Takara (Madison, USA). *Escherichia coli* DH5 α (*supE44*, *hsdR17*, *recA1*, *endA1*, *gyrA96*, *thi-1*, and *relA1*) was used for the DNA procedures (Invitrogen). Sodium phosphate and citric acid was from Sigma-Aldrich and Sephacryl S200 HR 26/60 chromatographic columns were from Amersham-Pharmacia Biotech (Uppsala, Sweden). UltracelTM system and UltracelTM PES membrane were from Millipore (Billerica, USA). The XynD concentration was determined by the Bradford method (1976) using the "Protein Assay" Reagent from Bio-Rad (Marnes-La-Coquette, France) [27].

Cloning, expression and purification of XynD

cDNA encoding XynD was synthesized by Geneart (Germany) using the GenBank AJ634957.1 accession number. The pPICZ α A-derived *Pichia pastoris* expression plasmid with the cDNA insert encoding XynD was constructed using standard procedures. The insert was purified using the QIAquick[®] (QIAGEN) purification kit and ligated into the XhoI/XbaI pPICZ α A plasmid, which was digested with PmeI to linearize the DNA for integration into *P. pastoris* genome. The linearized DNA was used for electroporation into *P. pastoris* strain X-33 using a Multiporator[®] (Eppendorf) at 1500 V for 5 ms. The transformants were selected using Minimal Methanol and Minimal Dextrose plates. Finally, in order to select multicopy vector strain transformants YPDS plates containing 100, 200, 500 and 1000 μ g/mL zeocin[®] were used.

Large-scale expression was carried out as previously described [21]. The culture supernatant was concentrated using UltracelTM PES ultrafiltration membrane (3 kDa) and subjected to gel filtration at a flow rate of 1.5 mL.min⁻¹ on a Sephacryl S-200 column connected to FPLC[®] equipment (Akta Purifier 10, GE Healthcare, Piscataway, USA) and eluted with 50 mM Na-phosphate buffer, pH 7.2, containing 150 mM NaCl. The fractions containing xylanases activity were pooled.

Polyacrylamide gel electrophoresis, glycosylation staining, N-terminal sequencing, molecular mass determination, and mass spectrometry analysis

SDS-PAGE was performed in 10 or 12% (w/v) polyacrylamide gel in the presence of 2% SDS under reducing or non-reducing conditions [28]. Proteins were visualized by Coomassie blue R-250 staining. For glycosylation determination, SDS-polyacrylamide gel were stained for carbohydrate with periodic acid-Schiff (PAS) using a SIGMA glycoprotein staining kit[®] according to the manufacturer's protocol.

N-terminal amino acid sequencing of the Ponceau red stained protein after electro-transfer on a polyvinylidene difluoride membrane was performed by automated repetitive Edman degradation on a model 494 Procise[™] Protein Sequencer (Applied Biosystems Division) [29]. Molecular mass determination was performed by Matrix-assisted laser desorption ionization/mass spectrometry (Ettan Maldi-Tof Pro, GE Healthcare, Uppsala, Sweden) as previously described [21].

Deglycosylation assay

Purified XynD was N-deglycosylated using endo- β -acetlyglucosaminidase H (Endo-H, Sigma Chemical Co.) and/or α -mannosidase from *Canavalia ensiformis* (Sigma Chemical Co.). Treatment with Endo-H was carried out in a 10 mM sodium citrate buffer, pH 5.0, for 6 h at 37°C, with a ratio of 1000 U: 77.5 μ g native and denaturated enzyme following the supplier recommendations. For O-deglycosylation, N-deglycosylated XynD was treated with α -mannosidase in a 50 mM sodium phosphate buffer, pH 6.0 for 16 h at 37°C, with a ratio of 14.2 U: 60 μ g protein.

Enzyme assays and effect of temperature and pH on enzyme activity

Xylanase activity was measured using the dinitrosalicylic acid (DNS) assay. Aliquot of the xylanase (10 μ L) was mixed with 10 mg.mL⁻¹ low viscosity AX from wheat (LVAX) (Megazyme) in McIlvaine's buffer, pH 5.0 (total volume reaction volume 250 μ l) at 70°C for 2 min. The reaction was stopped by the addition of 250 μ l DNS reagent before boiling the mixture during 5 min. The reactions were cooled and centrifuged for 3 min at 17 000 g and 200 μ L was transferred to a microplate and the absorption was measured at 545 nm. One unit (U) of xylanase activity was defined as the amount of protein that released 1 μ mol of xylose min⁻¹, based on xylose calibration curve. Optimal pH for xylanase activity was estimated using the DNS assay with LVAX (10 mg.mL⁻¹) in McIlvaine's buffer in a pH range of 2.0 to 8.0. Optimal temperature was also estimated using the same xylanase assay with LVAX (10 mg.mL⁻¹) in McIlvaine's buffer at pH 5.0 and temperatures ranging from 20°C to 100°C.

Thermal stability was performed at pH 5.0 and 70°C or 80°C for different time periods before performing activity assay.

Determination of substrate specificity and kinetic parameters

The substrate specificity of purified recombinant XynD was evaluated using the following substrates at 10 mg.mL⁻¹ in McIlvaine's buffer, pH 5.0: LVAX, medium viscosity AX (MVAX, Megazyme), high viscosity AX (HVAX, Megazyme), water-insoluble AX (IAX, Megazyme), Birchwood Xylan (Sigma), Beechwood Xylan (Sigma), xyloglucan from tamarind (XGT, Megazyme). Water-extractable wheat AX with different ratio arabinose:xylose were prepared by sequential precipitation with 30%, 50% and 60% ethanol (AXMF30, AXMF50, AXMF60), [30]. Their A/X values were 0.33, 0.53 and 0.73, respectively. Activity was measured by the DNS assay as described above.

Determinations of kinetic parameters were realized using the same condition described previously with LVAX as the substrate. Six concentrations of LVAX (2.5-20 mg.mL⁻¹) were used. The final concentration of XynD was 1.6 nM. The reaction was stopped at appropriate time intervals (2, 4, 6, 8 and 10 min) and the initial rates of hydrolysis of LVAX have been determined accordingly and expressed as μ M xylose/min. All experiments were repeated three times. Kinetic analyses were performed using the Michaelis-Menten, Lineweaver-Burk, Eadie-Hofstee and Dixon plots.

XOS analysis and quantification

Xylooligosaccharides and their products generated after XynD hydrolysis were analysed by high performance anion exchange chromatography coupled with a pulsed amperometric detection (HPAEC-PAD; Dionex, Sunnyvale, CA, USA) equipped with a Carbo-Pac PA-100 analytical column (250 \times 4 mm), a GP40 gradient pump, and a AS3500 auto-sampler (Thermo-Electron). The protocol used is the same as described by Cervera-Tison *et al*, in 2009 excepted for the composition of buffer A (NaOAc, 5 mM; NaOH, 80 mM) [31]. To evaluate the activity of the XynD on XOS (xylohexaose, xylopentaose, xylotetraose and xylotriose), initial slopes of progress curves were used to determine the catalytic efficiency (k_{cat}/K_m) of the reaction following the equation of Matsui as previously described by Cervera-Tison *et al*, in 2009 [31]. All assays were carried out in duplicate.

3. Results and discussion

Characterization of XynD gene

The *XynD* cDNA is predicted to encode a modular mature protein of 390 amino acid residues, which includes a 317 amino acid catalytic module, a 38 amino acid Ser/Thr rich linker and a 35 amino acid carbohydrate binding module

with significant similarity to the carbohydrate binding module characteristic of fungal CBM1 [24]. Deduced molecular mass and pI of XynD were calculated to be 41540 Da and 5.02, respectively. One N-glycosylation site was detected, at position N101, using NetNGlyc 1.0 Server and the DictyOGlyc server proposed several potential O-glycosylation sites (Figure 1).

A BLAST search showed that XynD shares highest amino acid sequence identity of 87 and 86% with family 10 xylanases from *Talaromyces stipitatus* and *Penicillium marneffeii*, respectively (data not shown). This result together with biochemical characteristics, suggested that XynD, belongs to GH10 family, confirming the results from Furniss et al. in 2005 [20].

Expression, purification and biochemical characterization of XynD

In order to study the biochemical properties of the xylanase from *P. funiculosum*, the *XynD* cDNA was cloned into the pPICZαA plasmid and expressed in *Pichia pastoris* as a recombinant protein using the native *Saccharomyces cerevisiae* α-factor secretion signal sequence under the control of the AOX1 promoter. After an induction step with methanol, a major protein band with an apparent molecular mass of 64 kDa, which was not present in the supernatant of untransformed cells, was observed in the culture filtrate after SDS-PAGE analysis (data not shown). The amount of the 64 kDa protein increased with time of induction (data not shown). Only trace amounts of other proteins were present in the culture supernatant. XynD was purified to homogeneity as described in the Materials and Methods section, giving a single peak after chromatography on a Sephacryl S200 gel containing 21% of the supernatant activity. Thus, the enzyme was purified 1.9 fold with a specific activity of 33.9 U.mg⁻¹.

The N-terminal sequence EAEAQAQ was in complete agreement with that expected from the recombinant protein, EAEA corresponding to the C-terminal end of the α-factor signal peptide of pPICZαA vector after efficient peptidase signal digestion (Kex2), followed by QAQ from XynD N-terminal mature sequence. The purified

```

QAQLNTAAKAAGLLYFGTAVDNPDLSDSKYLVNLETADFG
QITPANAMKWQPTSPQGSYFTTQGDQIASLAKSNNDYLRC
HNLVWYNQLPSYITSGSWTN*ATLIAALKEHINGVVTHYKGG
QCYAWDVVNEALNEDGTYRQNVFYQHIGEA YIPIAFAAAA
AADPNAKLYNDYNI EYAGAKATGAQGIVKLIQAAGGRIDG
VGLQSHFIVGQTPSLATQKANMAAFTALGVDVAITELDIR
MTLPDTSALQTQSTDYQTTTTACVQTKGCVGITLWDYTD
KYSWVPGTFSGQGDACPWDSNYNKKPAYYGILAGLQSGSGS
SSSTSSITLITTTPTASSSTTSATTSATSGAAHWGQCGGIG
WSGPTICVSPYTCQVLNPPYSQCL
    
```

Figure 1 XynD peptidic sequence. N*: N-glycosylation putative site, ST: O-glycosylation putative sites are underline. Identified peptides of N- and O-deglycosylated protein after tryptic digestion and mass spectrometry appear in bold.

recombinant enzyme migrated in SDS-PAGE give a single band around 64 kDa (Figure 2, lane 2). The estimated size on electrophoresis gel and the 46 kDa molecular mass determinate by mass spectrometry, thus much higher than the theoretical size of 41540 Da, suggesting that the recombinant XynD was glycosylated. The presence of sugars linked to the protein was confirmed by periodic acid Schiff staining (Figure 2, lane 6). One N-glycosylation site is present in XynD amino acid sequence at position N101, suggesting that N-glycosylation is responsible for the higher molecular mass. To confirm this change was due to glycosylation, deglycosylation analysis was performed. After treatment with Endo H to remove N-linked carbohydrate moieties, only a single band located at about 60 kDa was observed (Figure 2, lane 3) which confirms that XynD was N-glycosylated. However, the molecular mass was still higher than that expected and periodic acid Schiff staining still positive, suggesting the presence of O-glycosylated moieties (Figure 2, lane 7). Actually, linker peptide connecting catalytic and carbohydrate-binding modules in fungal glycoside hydrolases are rich in Ser and Thr residues, and they are often O-glycosylated [32].

Thus, α-mannosidase treatment was performed to remove O-glycosylated sugars and a band with a molecular mass around 43 kDa appeared, corresponding to theoretical molecular mass of XynD and the deglycosylated enzyme was not stained by periodic acid Schiff (Figure 2, lane 4 and 8). Tryptic digestion and mass spectrometry analysis of the N- and O-deglycosylated 43 kDa protein confirmed that enzyme is XynD (Figure 1). A large amount of polymeric α-mannosidase which gave 2





Figure 2 SDS-PAGE (10%) of XynD and deglycosylated XynD (1-5: Coomassie R250 coloration, 6-8: Schiff reagent coloration). (1) Molecular weight; (2) Native XynD; (3) XynD N-deglycosylated; (4) *: XynD N- and O-deglycosylated; (5) α-mannosidase; (6) Native XynD; (7) XynD N-deglycosylated; (8) XynD N- and O-deglycosylated.

different bands around 44 and 66 kDa (Figure 2, Lane 5) was necessary to totally O-deglycosylated XynD.

The experimental pI determined by isoelectrofocusing was 5.9 (data not shown). This result and the experimentally determined molecular mass were in good agreement with data collected in the literature. It is worth noting that almost all basic xylanases having a molecular mass of less than 30 kDa belong to GH11 family, whereas acid xylanases have a molecular mass higher than 30 kDa and belong to GH10 family [33].

Effect of temperature and pH on XynD activity

The effects of pH and temperature on the enzymatic activity were investigated on the recombinant enzyme. XynD displayed optimum activity in the 4.0 to 5.5 pH range whereas the activity was lost at pH 2.5 (Figure 3A).



At pH 5.0, the optimal enzymatic activity was observed at 80°C whereas the activity drastically decreased above 90°C (Figure 3B). The thermal stability of the enzyme was determined at 70 and 80°C at the same pH value (Figure 3C). XynD enzymatic activity decreases about 80% after 10 min at 80°C in contrast XynD is stable at least one hour at 70°C. The high stability of XynD at 70°C should be highly useful for industrial purposes. This could be due to post-translational modifications of xylanase during excretion process, such as glycosylation. However, deglycosylated enzyme treated as above was found to be as stable as the glycosylated enzyme, suggesting that glycosylation had no effect on thermal stability.

Substrate specificity of XynD from *P. funiculosum*

The hydrolytic activity of XynD from *P. funiculosum* on various substrates was determined (Table 1). The enzyme showed a high specificity and a insensitivity towards different soluble xylans tested with specific activities ranging between $50.5 \pm 0.5 \text{ U.mg}^{-1}$ for Birchwood Xylan and $114.3 \pm 1.0 \text{ U.mg}^{-1}$ for HVAX. The rate of insoluble xylan degradation was similar to that of soluble xylan degradation and XynD seemed to be a few sensitive to the xylan substitution degree with specific activities ranging between $80.6 \pm 1.8 \text{ U.mg}^{-1}$ for AXMF60 and $120.8 \pm 1.6 \text{ U.mg}^{-1}$ for AXMF30. The equivalent results were previously observed for the GH10 xylanase from *Aspergillus aculeatus* [34]. Because it is well-known that GH10 xylanases are preferentially active against soluble substrates, this result is very interesting, suggesting an important role of the CBM to hydrolyze the insoluble and viscous xylans during XynD action [35]. This would be highly useful for degradation of cell wall polysaccharides allowing easier access to starchy materials in cereals or other starch-rich grains. Finally, low activity was detected using xyloglucan as substrate.

Table 1 Specific activities (U.mg^{-1}) of XynD on different substrates

Substrates	Specific Activity (U.mg^{-1})	SD
LVAX	97.9	8.5
MVAX	106.4	0.2
HVAX	114.3	1.0
IAX	85.7	5.7
Birchwood X	50.5	0.5
Beechwood X	60.1	0.5
AXMF30	120.8	1.6
AXMF50	89.4	2.8
AXMF60	80.6	1.8
XGT	0.9	0.1

SD: standard deviation.

Table 2 Kinetic values for XynD with LVAX as substrate and comparison with the *P. funiculosus* family 11 xylanases

Enzyme	Plot	$K_{m(app)}$ (mg.mL ⁻¹)	k_{cat} (s ⁻¹)	$k_{cat}/K_{m(app)}$ (s ⁻¹ mg ⁻¹ .mL)	References
XynD	M.-M.	3.7 ± 0.2	399	108	This study
XynD	L.-B.	5.0 ± 0.2	833	167	"
XynD	E.-H.	3.9 ± 0.2	523	134	"
XynB	M.-M.	40.0 ± 3.0	5405	135	[21]
XynC	M.-M.	14.4 ± 0.7	2939	204	[22]

For XynD, the parameters were determined from primary plots of Michaelis-Menten (M.-M), Lineweaver-Burk (L.B) and Eadie-Hofstee (E.-H) plots.

Determination of kinetic parameters

The XynD-catalysed hydrolysis of LVAX was measured with the optimal temperature (70°C) and pH 5 conditions (Table 2). The experimental initial rates were used for determination of k_{cat} and $K_{m(app)}$ and calculation of the catalytic efficiency, $k_{cat}/K_{m(app)}$ according the kinetic models of Michaelis-Menten, Lineweaver-Burk and Eadie-Hofstee (plot not shown). The $K_{m(app)}$ of XynD with LVAX as the substrate is 3 to 8 fold higher than those of *P. funiculosus* GH11 xylanases but the catalytic efficiency ($k_{cat}/K_{m(app)}$) were similar [21,22].

In order to evaluate the mode of action of XynD, the products generated using highly polymeric substrates and on short substrates ranging from xylo-tetraose to xylo-hexaose were analysed by HPAEC-PAD. XynD displayed mainly an endo-activity against wheat AX (LVAX) because the XOS generated during the initial stages of hydrolysis were progressively degraded yielding principally xylobiose and to a lower extent of xylo-triose and xylose (Figure 4). This characteristic is obviously similar to GH11 xylanases that yield xylobiose and xylo-triose as main end reaction products [22,31].




Figure 4 HPAEC analysis of wheat LVAX hydrolysis by XynD. XynD (25.0 nM) was incubated with 5 mg.mL⁻¹ wheat AX in McIlvaine's buffer of pH 5.0 at 50°C. The reaction was stopped after 0, 30, 60 and 1024 min and subjected to HPAEC analysis. Xylose (X1), xylobiose (X2), xylo-triose (X3), xylo-tetraose (X4), xylo-pentaose (X5), and xylo-hexaose (X6).

XynD exhibited an exclusive endo-mode of action against XOS. Indeed, xylohexaose was hydrolysed to xylo-triose, xylo-tetraose and xylobiose; xylopentaose to xylobiose and xylo-triose; whereas xylo-tetraose was cleaved to produce exclusively xylobiose (Figure 5). No significant amount of xylose was generated during hydrolysis of XOS. The catalytic efficiency (k_{cat}/K_m) increased slightly with increasing chain length of XOS up to n = 6 (Table 3) and the relative value on xylo-tetraose, xylo-pentaose and xylo-hexaose, was 1:1.3:8.8. Xylanase C from *P. funiculosus*, a GH11 xylanase family, showed different catalytic efficiency (k_{cat}/K_m) results on the same substrates with the following relative values: 1:2.1:39.3 [22]. Finally, the most important




Figure 5 Progress curves of the XOS generated by XynD after hydrolysis of xylo-tetraose (A), xylo-pentaose (B), and xylo-hexaose (C). Recombinant enzymes were incubated with 100 μM of XOS in McIlvaine's buffer pH 5.0 at 70°C. The quantity of xylose (X1), xylobiose (X2), xylo-triose (X3), xylo-tetraose (X4), xylo-pentaose (X5), and xylo-hexaose (X6) produced during the course of the reaction is indicated. The concentrations of enzymes used were 640 nM (A), 16.1 nM (B), and 6.4 nM (C).

Table 3 Catalytic efficiencies of XynD compared to 3 *Penicillium sp.* GH11 xylanases

Enzymes	Substrates (100 μM)	k_{cat}/K_m ($\text{min}^{-1} \cdot \text{M}^{-1}$)	References
XynD	X6	5.9.10 ⁶	This study
	X5	4.4.10 ⁶	"
	X4	1.5.10 ⁶	"
XynC	X6	6.5.10 ⁶	[22]
	X5	3.1.10 ⁶	"
	X4	1.8.10 ⁵	"
XynA	X6	2.6.10 ⁶	[22]
	X5	4.6.10 ⁵	"
	X4	3.0.10 ⁴	"
XynB	X6	8.0.10 ⁵	[31]
	X5	7.4.10 ⁴	"
	X4	6.2.10 ³	"

Xylohexaose (X6), xylopentaose (X5) and xyloetraose (X4) were used as substrates.

difference was found with XynB from *P. griseofulvum* 1:12:129 [31]. Both the studies concluded that these xylanases potentially contain six subsites in their active site. Xylohexaose degradation produced only xylotriase, indicating that the minimum length of the substrate is four residues. Moreover, the small difference of the XynD catalytic efficiency against xyloetraose and xylopentaose and against xylopentaose and xylohexaose suggest that XynD possess four catalytic subsites with a high energy of interaction with the substrate and a fifth subsite with a small energy of interaction. However this hypothesis will be confirmed by a modeling study.

4. Conclusion

Filamentous fungus *P. funiculosum* is known for its capacity to produce xylanases with different capacities that may provide the fungus with the maximum potential to degrade xylans from different sources. In this study, the *P. funiculosum* GH10 unique gene encoding a xylanase was cloned and express in *P. pastoris* yeast. Biochemical characterization and enzymatic analysis showed that XynD presented a high temperature optimal for hydrolysis activity, was thermostable and it was a versatile xylanase in term of activities and catalytic efficiencies than the others *P. funiculosum* xylanases. Because of its attractive properties XynD might be considered for biotechnological applications.

Abbreviations

AX: arabinoxylan; AXMF: AXs with different ratio A/X; CBM: carbohydrate binding module; DNS: dinitrosalicylic acid; GH: glycoside hydrolase; HAVX: high viscosity AXs; HPAEC-PAD: high performance anion exchange chromatography; IAX: insoluble AXs; LVAX: low viscosity AXs; MAVX: medium viscosity AXs; PAS: periodic acid Schiff; PBS: phosphate-buffered saline; PES: polyether sulfone; XGT: xyloglucan; XOS: xylooligosaccharides.

Author details

¹Université Paul Cézanne, ISM2/BiosSciences UMR CNRS 6263 (Université Aix Marseille III/CNRS), service 342, Faculté des Sciences et Techniques Saint-Jérôme, 13397 Marseille Cedex 20, France. ²INRA, UR 1268 - Biopolymères - Interactions - Assemblages, 44316 Nantes Cedex 03, France.

Authors' contributions

ML was involved in study design, data interpretation and manuscript writing; AT was involved in biochemical and enzymatic analysis; VD was involved in biochemical analysis design and data interpretation; EB was involved in the determination of the activity assays and in the data interpretation; EHA was involved in data interpretation and manuscript writing; TG was the coordinator of the study design and was involved in data interpretation and manuscript writing and editing. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 15 December 2010 Accepted: 5 April 2011

Published: 5 April 2011

References

- Henrissat B: A Classification of Glycosyl Hydrolases Based on Amino-Acid-Sequence Similarities. *Biochem J* 1991, **280**:309-316.
- Henrissat B, Bairoch A: New Families in the Classification of Glycosyl Hydrolases Based on Amino-Acid-Sequence Similarities. *Biochem J* 1993, **293**:781-788.
- Henrissat B, Bairoch A: Updating the sequence-based classification of glycosyl hydrolases. *Biochem J* 1996, **316**:695-696.
- Subramaniyan S, Prema P: Biotechnology of microbial xylanases: Enzymology, molecular biology, and application. *Crit Rev Biotechnol* 2002, **22**:33-64.
- Beg QK, Kapoor M, Mahajan L, Hoondal GS: Microbial xylanases and their industrial applications: a review. *Appl Microbiol Biot* 2001, **56**:326-338.
- Techapun C, Poosaran N, Watanabe M, Sasaki K: Thermostable and alkalinetolerant microbial cellulase-free xylanases produced from agricultural wastes and the properties required for use in pulp bleaching bioprocesses: a review. *Process Biochem* 2003, **38**:1327-1340.
- Wong KKY, Saddler JN: Trichoderma Xylanases, Their Properties and Application. *Crit Rev Biotechnol* 1992, **12**:413-435.
- Bajpai P: Application of enzymes in the pulp and paper industry. *Biotechnol Progr* 1999, **15**:147-157.
- Niehaus F, Bertoldo C, Kahler M, Antranikian G: Extremophiles as a source of novel enzymes for industrial application. *Appl Microbiol Biot* 1999, **51**:711-729.
- Okazaki M, Fujikawa S, Matsumoto N: Effect of xylo-oligosaccharide on the growth of *bifidobacteria*. *Bifidobact Microflora* 1990, **9**:77-86.
- Campbell JM, Fahey GC, Wolf BW: Selected indigestible oligosaccharides affect large bowel mass, cecal and fecal short-chain fatty acids, pH and microflora in rats. *J Nutr* 1997, **127**:130-136.
- Madhukumar MS, Muralikrishna G: Structural characterisation and determination of prebiotic activity of purified xylo-oligosaccharides obtained from Bengal gram husk (*Cicer arietinum L.*) and wheat bran (*Triticum aestivum*). *Food Chem* 2010, **118**:215-223.
- Roberfroid MB: Health benefits of non-digestible oligosaccharides. In *Dietary Fiber in Health and Disease*. Edited by: Kritchevsky D, Bonfield Ch. New York: Plenum Press; 1997:211-219.
- Hsu CK, Liao JW, Chung YC, Hsieh CP, Chan YC: Xylo-oligosaccharides and fructooligosaccharides affect the intestinal microbiota and precancerous colonic lesion development in rats. *J Nutr* 2004, **134**:1523-1528.
- Moure A, Gullon P, Dominguez H Parajo JC: Advances in the manufacture, purification and applications of xylo-oligosaccharides as food additives and nutraceuticals. *Process Biochem* 2006, **41**:1913-1923.
- Gibson GR, Roberfroid MB: Dietary Modulation of the Human Colonic Microbiota - Introducing the Concept of Prebiotics. *J Nutr* 1995, **125**:1401-1412.
- Barreteau H, Delattre C, Michaud P: Production of oligosaccharides as promising new food additive generation. *Food Technol Biotech* 2006, **44**:323-333.

18. Guais O, Borderies G, Pichereaux C, Maestracci M, Neugnot V, Rossignol M, Francois JM : Proteomics analysis of "Rovabio™ Excel", a secreted protein cocktail from the filamentous fungus *Penicillium funiculosum* grown under industrial process fermentation. *J Ind Microbiol Biot* 2008, **35**:1659-1668.
19. Furniss CSM, Belshaw NJ, Alcocer MJC, Williamson G, Elliott GO, Gebruers K, Haigh NP, Fish NM, Kroon PA: A family 11 xylanase from *Penicillium funiculosum* is strongly inhibited by three wheat xylanase inhibitors. *Biochim Biophys Acta* 2002, **1598**:24-29.
20. Furniss CSM, Williamson G, Kroon PA: The substrate specificity and susceptibility to wheat inhibitor proteins of *Penicillium funiculosum* xylanases from a commercial enzyme preparation. *J Sci Food Agr* 2005, **85**:574-582.
21. Brutus A, Villard C, Durand A, Tahir T, Furniss C, Puigserver P, Juge N, Giardina T: The inhibition specificity of recombinant *Penicillium funiculosum* xylanase B towards wheat proteinaceous inhibitors. *Biochim Biophys Acta* 2004, **1701**:121-128.
22. Berrin JG, Ajandouz EH, Georis J, Arnaut F, Juge N: Substrate and product hydrolysis specificity in family 11 glycoside hydrolases: an analysis of *Penicillium funiculosum* and *Penicillium griseofulvum* xylanases. *Appl Microbiol Biot* 2007, **74**:1001-1010.
23. Torronen A, Rouvinen J: Structural and functional properties of low molecular weight endo-1,4-beta-xylanases. *J Biotechnol* 1997, **57**:137-149.
24. Levasseur A, Navarro D, Punt PJ, Belaich JP, Asther M, Record E: Construction of engineered bifunctional enzymes and their overproduction in *Aspergillus niger* for improved enzymatic tools to degrade agricultural by-products. *Appl Environ Microb* 2005, **71**:8132-8140.
25. Beaugrand J, Chambat G, Wong VWK, Goubet F, Rémond C, Paës G: Impact and efficiency of GH10 and GH11 thermostable endoxylanases on wheat bran and alkali-extractable arabinoxylans. *Carbohydr Res* 2004, **339**:2529-2540.
26. Pell G, Szabo L, Charnock SJ, Xie HF, Gloster TM, Davies GJ, Gilbert HJ: Structural and biochemical analysis of *Cellvibrio japonicus* xylanase 10C - How variation in substrate-binding cleft influences the catalytic profile of family GH-10 xylanases. *J Biol Chem* 2004, **279**:11777-11788.
27. Bradford MM: Rapid and Sensitive Method for Quantitation of Microgram Quantities of Protein Utilizing Principle of Protein-Dye Binding. *Anal Biochem* 1976, **72**:248-254.
28. Laemmli UK: Cleavage of Structural Proteins During Assembly of Head of Bacteriophage-T4. *Nature* 1970, **227**:680-685.
29. Matsudaira P: Sequence from picomole quantities of proteins electroblotted on to polyvinylidene difluoride membranes. *J Biol Chem* 1987, **262**:10035-10038.
30. Dervilly G, Saulnier L, Roger P, Thibault JF: Isolation of homogeneous fractions from wheat water-soluble arabinoxylans. Influence of the structure on their macromolecular characteristics. *J Agr Food Chem* 2000, **48**:270-278.
31. Cervera-Tison MC, Andre-Leroux G, Lafond M, Georis J, Juge N, Berrin J-G : Molecular determinants of substrate and inhibitor specificities of the *Penicillium griseofulvum* family 11 xylanases. *Biochim Biophys Acta* 2009, **1794**:438-445.
32. Goto M: Protein O-glycosylation in *fungi*: Diverse structures and multiple functions. *Biosci Biotech Bioch* 2007, **71**:1415-1427.
33. Wong KKY, Tan LUL, Saddler JN: Multiplicity of Beta-1,4-Xylanase in Microorganisms - Functions and Applications. *Microbiol Rev* 1988, **52**:305-317.
34. Bonnin E, Daviet S, Sorensen JF, Sibbesen O, Goldson AJ, Juge N, Saulnier L: Behaviour of family 10 and 11 xylanases towards arabinoxylans with varying structure. *J Sci Food Agr* 2006, **86**:1618-1622.
35. Maes C, Vangeneugden B, Delcour JA: Relative activity of two endoxylanases towards water-unextractable arabinoxylans in wheat bran. *J Cereal Sci* 2004, **39**:181-186.

doi:10.1186/1475-2859-10-20

Cite this article as: Lafond et al.: GH10 xylanase D from *Penicillium funiculosum*: biochemical studies and xylooligosaccharide production. *Microbial Cell Factories* 2011 **10**:20.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

