



HAL
open science

Stress-induced GSK3 regulates the redox stress response by phosphorylating glucose-6-phosphate dehydrogenase in Arabidopsis

Sylvia Dal Santo, Hansjörg Stampfl, Julia Krasensky, Stefan Kempa, Yves Y. Gibon, Elena Petutschnig, Wilfried Rozhon, Alexander Heuck, Tim Clausen, Claudia Jonak

► To cite this version:

Sylvia Dal Santo, Hansjörg Stampfl, Julia Krasensky, Stefan Kempa, Yves Y. Gibon, et al.. Stress-induced GSK3 regulates the redox stress response by phosphorylating glucose-6-phosphate dehydrogenase in Arabidopsis. *The Plant cell*, 2012, 24 (8), pp.3380-3392. 10.1105/tpc.112.101279 . hal-02647870

HAL Id: hal-02647870

<https://hal.inrae.fr/hal-02647870>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Stress-Induced GSK3 Regulates the Redox Stress Response by Phosphorylating
Glucose-6-Phosphate Dehydrogenase in *Arabidopsis***

Silvia Dal Santo, Hansjörg Stampfl, Julia Krasensky, Stefan Kempa, Yves Gibon, Elena Petutschnig,
Wilfried Rozhon, Alexander Heuck, Tim Clausen and Claudia Jonak
Plant Cell 2012;24;3380-3392; originally published online August 10, 2012;
DOI 10.1105/tpc.112.101279

This information is current as of October 2, 2013

Supplemental Data	http://www.plantcell.org/content/suppl/2012/08/02/tpc.112.101279.DC1.html http://www.plantcell.org/content/suppl/2012/08/21/tpc.112.101279.DC2.html
References	This article cites 64 articles, 19 of which can be accessed free at: http://www.plantcell.org/content/24/8/3380.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm

Stress-Induced GSK3 Regulates the Redox Stress Response by Phosphorylating Glucose-6-Phosphate Dehydrogenase in *Arabidopsis*

Silvia Dal Santo,^{a,1,2} Hansjörg Stampfl,^{a,1} Julia Krasensky,^a Stefan Kempa,^{a,3} Yves Gibon,^b Elena Petutschnig,^{a,4} Wilfried Rozhon,^{a,5} Alexander Heuck,^c Tim Clausen,^c and Claudia Jonak^{a,6}

^aGregor Mendel Institute of Molecular Plant Biology, Austrian Academy of Sciences, 1030 Vienna, Austria

^bInstitut National de la Recherche Agronomique-Bordeaux, La Grande Ferrade, 33883 Villenave d'Ornon, France

^cResearch Institute of Molecular Pathology, 1030 Vienna, Austria

Diverse stresses such as high salt conditions cause an increase in reactive oxygen species (ROS), necessitating a redox stress response. However, little is known about the signaling pathways that regulate the antioxidant system to counteract oxidative stress. Here, we show that a Glycogen Synthase Kinase3 from *Arabidopsis thaliana* (ASK α) regulates stress tolerance by activating Glc-6-phosphate dehydrogenase (G6PD), which is essential for maintaining the cellular redox balance. Loss of stress-activated ASK α leads to reduced G6PD activity, elevated levels of ROS, and enhanced sensitivity to salt stress. Conversely, plants overexpressing ASK α have increased G6PD activity and low levels of ROS in response to stress and are more tolerant to salt stress. ASK α stimulates the activity of a specific cytosolic G6PD isoform by phosphorylating the evolutionarily conserved Thr-467, which is implicated in cosubstrate binding. Our results reveal a novel mechanism of G6PD adaptive regulation that is critical for the cellular stress response.

INTRODUCTION

High soil salinity is a major environmental constraint for plant growth and development and negatively affects agricultural productivity (Boyer, 1982). Salinity imposes a water deficit and ion stress, which cause effects such as inhibition of essential enzymes, destabilization of cell membranes, a decrease in nutrient supply, and overproduction of reactive oxygen species (ROS) (Hasegawa et al., 2000; Zhu, 2001; Munns, 2002). ROS production is a universal feature of aerobic metabolism. Although recent evidence has uncovered a role for ROS as a signaling molecule and key physiological regulator of diverse cellular processes (Mittler et al., 2011), excess levels generated as a consequence of an insult cause oxidative damage and can

ultimately lead to cell death (Møller et al., 2007). Therefore, tight regulation of excess ROS is an essential protective mechanism in all organisms.

Glc-6-phosphate dehydrogenase (G6PD; EC 1.1.1.49) catalyzes a key step of the oxidative pentose phosphate pathway (OPPP) that provides NADPH for reductive biosynthesis and maintenance of the cellular redox state. In animals, G6PD is the principal enzyme providing NADPH and is of central importance for cellular redox regulation in response to stress. For example, mouse embryonic stem cells in which G6PD activity is disrupted are extremely sensitive to oxidative stress (Pandolfi et al., 1995; Filosa et al., 2003). In addition, exogenous stresses such as pathogen infection, certain drugs, or ingestion of fava beans (*Vicia faba*) can trigger hemolytic anemia in humans with G6PD deficiency, which is the most common enzymopathy. Similarly, yeast mutants defective in G6PD show an increased susceptibility to oxidative stress and fail to induce adaptation (Juhnke et al., 1996; Izawa et al., 1998). In plants, G6PD activity is present in the plastids and cytosol (Debnam and Emes, 1999). G6PD activity has been positively correlated with environmental stresses such as salt stress (Valderrama et al., 2006; Wang et al., 2008), aluminum toxicity (Slaski et al., 2006), drought (Scharte et al., 2009), and infection by viral and fungal pathogens (Sindelár and Sindelárová, 2002; Scharte et al., 2009); by contrast, a decrease in chloroplastic G6PD activity has been shown to be beneficial for withstanding oxidative stress (Debnam et al., 2004). Thus, the role of G6PD in the stress response is currently unclear.

The activity of G6PD is tightly regulated. In addition to regulation at the transcriptional level, redox regulation and the cellular NADPH/NADP⁺ ratio modulate the activity of the different G6PD isoforms (Wendt et al., 2000; Debnam et al., 2004; Schürmann

¹ These authors contributed equally to this work.

² Current address: Department of Biotechnology, Strada Le Grazie 15, 37134 Verona, Italy.

³ Current address: Berlin Institute for Medical Systems Biology, Robert Rössle Strasse 10, 13092 Berlin, Germany.


⁴ Current address: Albrecht-von-Haller-Institute of Plant Sciences, Georg-August-University, Untere Karspuele 2, 37073 Goettingen, Germany.


⁵ Current address: Center for Life and Food Sciences Weihenstephan, Technische Universität München, Liesel-Beckmann-Straße 1, 85354 Freising, Germany.

⁶ Address correspondence to claudia.jonak@gmi.oeaw.ac.at.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Claudia Jonak (claudia.jonak@gmi.oeaw.ac.at).

 Some figures in this article are displayed in color online but in black and white in the print edition.

 Online version contains Web-only data.

 Open Access articles can be viewed online without a subscription.

www.plantcell.org/cgi/doi/10.1105/tpc.112.101279

and Buchanan, 2008). G6PD is phosphorylated in animals and plants, which has been correlated with both enhanced (Ramnanan and Storey, 2006; Dieni and Storey, 2010; Gupte et al., 2011) and reduced activity (Hauschild and von Schaewen, 2003; Xu et al., 2005; Zhang et al., 2000). While different signaling pathways, including protein kinase C, protein kinase A, and Src kinase, have been implicated in regulating G6PD phosphorylation in animals under a variety of different conditions (Xu et al., 2005; Gupte et al., 2009, 2011), the upstream regulator(s) of G6PD phosphorylation in plants have yet to be identified.

Glycogen Synthase Kinase3 (GSK3) constitutes a class of evolutionarily conserved Ser/Thr protein kinases. Originally identified in mammals as a cytoplasmic modulator of glycogen metabolism, GSK3 is now recognized as a central regulator of an array of cellular events, including cell fate determination, microtubule function, cell cycle regulation, apoptosis, and inflammatory responses (Cohen and Frame, 2001; Doble and Woodgett, 2003; Jope and Johnson, 2004; Martin et al., 2005). In plants, GSK3/shaggy-like kinases (GSKs) are encoded by a gene family that directs different physiological responses (Jonak and Hirt, 2002; Saidi et al., 2012). In alfalfa (*Medicago sativa*), several GSKs have emerged as regulatory components in stress signaling. For example, wound-induced GSK is posttranslationally activated by wounding (Jonak et al., 2000), Msk1 is involved in innate immunity by limiting the severity of virulent bacterial infections (Wrzaczek et al., 2007), and Msk4 regulates high salt tolerance by adjusting carbohydrate metabolism in response to environmental stress (Kempa et al., 2007). In *Arabidopsis thaliana*, overexpression of AtGSK1/ASK α enhanced salt tolerance (Piao et al., 2001), and in rice (*Oryza sativa*), knockout mutants of GSK1 showed increased tolerance to salt stress (Koh et al., 2007). However, the mechanisms by which these stress-related GSKs act, including their direct targets for phosphorylation, remain elusive.

In a changing environment, cellular metabolism needs to be altered to enable an adapted physiological response. Protein phosphorylation represents an important means of fine-tuning the activity of metabolic enzymes. However, our knowledge of how signal transduction is linked to redox regulation is limited. Here, we provide direct evidence that a stress-activated GSK3 from *Arabidopsis*, ASK α , is a key regulatory component protecting cells against oxidative stress. Moreover, we identify G6PD as an *in vivo* target of ASK α and present a novel molecular mechanism for the regulation of G6PD activity.

RESULTS

ASK α Is Involved in *Arabidopsis* Salt Stress Resistance

To investigate the role of *Arabidopsis* GSKs (ASKs) in stress tolerance, we screened ASK activity mutants (i.e., *ask* knockout and ASK overexpressor lines) for their performance under abiotic stress conditions. This systematic study revealed that ASK α activity mutants resembled the wild type under normal growth conditions but displayed altered tolerance to high salt conditions. High soil salinity stress, imposed by watering soil-grown plants with 150 mM NaCl solution over 3 weeks, severely affected the growth of *ask* α T-DNA insertion knockout plants (see

Supplemental Figures 1A and 1B online) compared with wild-type plants (Figure 1A). No difference was observed between the different genotypes in untreated plants (see Supplemental Figure 1C online).

We next analyzed whether ASK α was also important for tolerance to salt stress during early seedling development. On medium supplemented with 100 mM NaCl, both germination efficiency and primary root length were significantly reduced in *ask* α mutants compared with wild-type plants grown under the same high salt stress conditions (Figures 1B and 1C). By contrast, plants overexpressing ASK α from the strong constitutive 35S promoter (see Supplemental Figure 1B online) germinated equally well under normal and salt stress conditions, and root length was only slightly affected by high salt stress. In high salt conditions, both germination efficiency and root length of plants overexpressing ASK α were significantly increased compared with wild-type plants. Thus, ASK α plays an important role in sensitivity to salt stress, both during early seedling development and in adult plants.

ASK α Is Activated by High Salinity Stress

To assess whether ASK α protein kinase activity is modulated by salt stress, *in vivo* ASK α kinase activity was first determined from long-term salt stressed and nonstressed plants expressing ASK α -myc. Immunokinase assays using anti-myc antibodies and myelin basic protein (MBP) as a general substrate showed basal ASK α -myc kinase activity in nonstressed plants and enhanced kinase activity in plants exposed to salt stress (Figure 1D). To study the effect of salt stress on the activity of endogenous ASK α , an ASK α -specific peptide antibody (see Supplemental Figures 1D and 1E online) was used for the immunokinase assays. Consistent with the previous result, wild-type Columbia-0 (Col-0) plants grown under high salt conditions contained enhanced endogenous ASK α activity compared with plants grown under normal conditions (Figure 1E).

Plants respond to high soil salinity by a sequence of temporally successive cellular and physiological responses. To study the kinetics of ASK α activation at the onset of stress, ASK α immunokinase assays were performed on plants exposed to salt stress for 6 h, 12 h, 1 d, or 3 d. In control plants, ASK α activity was low. However, high soil salinity induced ASK α activity rapidly and persistently over an experimental period of 3 d (Figure 1F, top panel), suggesting that ASK α activation is part of a primary stress response. By contrast, the ASK α steady state protein levels remained constant (Figure 1F, bottom panel). This indicates that there is a posttranslational mechanism of ASK α activation upon salt stress.

Altered G6PD Activity and ROS Levels of ASK α Activity Mutants under Salt Stress Conditions

Our previous work indicated a role for the GSK MSK4 in metabolic regulation in alfalfa (Kempa et al., 2007), so we hypothesized that this salt stress induction of ASK α might impact cellular metabolism by regulating the activity of specific metabolic enzymes. We therefore performed a high-throughput robot-based screen for multiple enzyme activities (Gibon et al., 2004) in




Figure 1. *ASKα* Is Activated by High Salinity and Modulates *Arabidopsis* Salt Stress Resistance.

(A) Phenotype of *ASKα* activity mutants upon prolonged salt stress. Four-week-old soil-grown *Arabidopsis* wild-type Col-0, *askα*, and *ASKα* over-expressor lines OE#4 and OE#5 were watered with 150 mM NaCl for 3 weeks. An image of single representative individuals of the phenotypes' mean of 30 plants is shown.

(B) and (C) Salt tolerance of *ASKα* activity mutants during early seedling development.

(B) Germination efficiency on half-strength MS plates and on plates supplemented with 100 mM NaCl. Data are means \pm SD of three independent biological replicates with $n = 100$ each. Asterisks indicate a significant difference (** $P < 0.01$) using Student's *t* test for pairwise comparison to Col-0 under stress conditions.

(C) Primary root growth on control medium and on medium supplemented with 100 mM NaCl. Root length was determined 7 d after sowing. Data are means \pm SD of three independent biological replicates. Asterisks indicate a significant difference (** $P < 0.01$) using Student's *t* test for pairwise comparison to Col-0 under stress conditions.

(D) and (E) *ASKα* in vivo kinase activity upon prolonged salt stress. Four-week-old soil-grown *Arabidopsis* wild type and *ASKα* activity mutants were either irrigated with water or with a 150 mM NaCl solution for 3 weeks. Twenty plants per line were pooled and used for immunokinase assays. *ASKα* was either immunoprecipitated from 100 μg leaf protein extracts with anti-myc antibodies (D) or with *ASKα*-specific antibodies (E). Subsequent kinase assays were performed with [γ - 32 P]ATP and MBP as a substrate.

(F) Kinetics of *ASKα* kinase activation. Soil-grown *Arabidopsis* Col-0 plants were exposed to high soil salinity for 6 h, 12 h, 24 h, or 3 d. Plant material was pooled from rosette leaves of 10 plants per time point. Top panel: Immunokinase assays were performed on 100 μg of leaf protein extracts. *ASKα*-specific antibodies were used for immunoprecipitation. Subsequently, *ASKα* kinase activity was determined in kinase assays with [γ - 32 P]ATP and MBP as a substrate. Bottom panel: *ASKα* protein levels were determined by immunoprecipitation of *ASKα* from 200 μg total protein extract and subsequent immunoblot analysis with *ASKα*-specific antibodies. ctrl., control.

All experiments in this figure were performed at least twice and showed similar results.

[See online article for color version of this figure.]

askα plants to identify potential downstream effectors of the stress response. This analysis revealed reduced G6PD activity in stressed *askα* plants. Analysis of total G6PD activity showed that prolonged salt stress enhanced G6PD activity in wild-type plants (Figure 2A). However, in *askα*, G6PD activity was strongly reduced by salinity. By contrast, plants overexpressing ASKα showed a stress-induced increase in G6PD activity that was higher than that in the wild type (Figure 2A).

G6PD activity provides reducing power important for ROS detoxification. We thus explored levels of the ROS hydrogen peroxide (H_2O_2) in leaves upon salt stress. Consistent with the reduced total G6PD activity, H_2O_2 levels were significantly elevated in *askα* plants exposed to high soil salinity. In stressed ASKα overexpressor plants, H_2O_2 levels were lower than in the wild type (Figure 2B), indicating that ASKα activity is involved in modulating stress-induced ROS accumulation.

ASKα Phosphorylates and Activates G6PD6 in Vitro

To study the mechanism by which ASKα regulates G6PD, we first analyzed total G6PD levels in leaves which were similar in all genotypes under normal and salt stress conditions (Figure 2C). Next, we assessed the possibility that ASKα might regulate G6PD activity directly by phosphorylation. The major G6PD activity in leaves originates from cytosolic G6PD isoforms (Debnam and Emes, 1999). The lack of a potential plastid targeting sequence in ASKα prompted us to focus our analyses on the two cytosolic G6PD isoforms, G6PD5 and G6PD6 (Wakao et al., 2008).

In an in vitro kinase assay with recombinant proteins, ASKα was able to phosphorylate G6PD5 and G6PD6 (Figure 3A). Subsequent analysis of G6PD activity revealed that phosphorylation by active ASKα stimulated the activity of G6PD6 (Figure 3B), whereas a kinase-dead version of ASKα (ASKα K98R) could not phosphorylate G6PD6 and also failed to modify its activity. Interestingly, ASKα did not enhance G6PD5 activity, indicating that the two G6PD isoforms might be regulated by distinct mechanisms.

Thr-467 Is Necessary for G6PD6 Activation by ASKα

To investigate the molecular mechanism of G6PD6 activation by ASKα, liquid trap quadrupole mass spectrometry analysis was performed on recombinant G6PD phosphorylated by ASKα in vitro. Thr-467 at the C terminus of G6PD6 was identified as the ASKα target site (see Supplemental Figure 2 online). Thr-467, and the adjacent amino acids, are highly conserved in eukaryotes (see Supplemental Figure 3 online), suggesting an important role for this residue in regulating G6PD6 activity.

To understand the molecular effects of Thr-467 phosphorylation, we analyzed the homologous crystal structure of human G6PD (Au et al., 2000; Kotaka et al., 2005). The corresponding human Thr residue (Thr-466) is exposed on the G6PD surface and thus is readily accessible by the kinase (Figure 4A). Active G6PD exists in a dimer-tetramer equilibrium (Au et al., 2000). Human Thr-466 is located remotely from the dimer and tetramer interfaces of the G6PD molecule, suggesting that this residue would not affect the oligomeric state (Figure 4A). Interestingly,




Figure 2. G6PD Activity and H_2O_2 Levels Are Altered in ASKα Activity Mutants upon Salt Stress.

Four-week-old soil-grown *Arabidopsis* plants were watered with 150 mM NaCl for 3 weeks. The same plant material was used for (A) to (C). Analyses were performed with pools of 20 plants on three independent salt watering experiments. Data are means \pm relative sd. Controls for each genotype were normalized to one [(A) and (B)].

(A) Total G6PD activity in leaf extracts of wild-type Col-0, *askα*, and ASKα overexpressor lines OE#4 and OE#5 grown under normal and high salinity conditions.

(B) H_2O_2 levels. H_2O_2 content was assessed spectrophotometrically in Col-0 and ASKα activity mutants.

(C) Total G6PD levels. Top lane: Total G6PD was immunoprecipitated from 50 μ g protein extract and subsequently detected by immunoblot analysis using a polyclonal G6PD antibody. Bottom lane: CBB-stained gel.

human Thr-466 is in close proximity (3 Å) to the β E- α E loop, which is part of the NADP binding site (Figure 4B). Phosphorylation of Thr-467 in G6PD6 might thus introduce structural changes in the coenzyme binding site connected with an enhanced G6PD activity.




Figure 3. ASK α Phosphorylates and Thereby Activates G6PD6 in Vitro.

(A) Phosphorylation of G6PD5 and G6PD6 by ASK α . Kinase assays were performed with purified GST-ASK α and GST-G6PD5 or GST-G6PD6. ASK α did not phosphorylate GST alone.

(B) Enzymatic activities of G6PD5 and G6PD6. G6PD activity was quantified after a phosphorylation reaction of GST-G6PD5 and GST-G6PD6 without kinase or with GST-ASK α or with GST-ASK α K98R. Data are means \pm relative sd. The assay was performed three times using fresh proteins from independent purifications. Asterisks indicate a significant difference (*** P < 0.005) tested by Student's t test for pairwise comparison to the unphosphorylated control.

To explore the significance of Thr-467 modification for G6PD6 activity further, the residue was mutated either to Ala (T467A), which cannot be phosphorylated, or to Glu (T467E), mimicking the phosphorylation event. In ASK α kinase assays, phosphorylation of the T467A (nonphosphorylatable) mutant was reduced compared with wild-type protein, suggesting that Thr-467 is a central target site for ASK α phosphorylation (Figure 4C). The observed residual phosphorylation of G6PD6 T467A and G6PD6 T467E indicates that ASK α phosphorylates additional sites, at

least in vitro. Importantly, consistent with the results of the kinase assays, analyses of the impact of ASK α on the activity of G6PD6 mutant versions revealed that the activity of G6PD6 T467A could no longer be enhanced by ASK α , whereas the G6PD6 T467E phosphomimic mutant displayed constitutively




Figure 4. Thr-467 Is Necessary for G6PD6 Activation by ASK α in Vitro.

(A) and (B) The crystal structure of human G6PD suggests that Thr-467 phosphorylation influences coenzyme binding.

(A) Crystal structure of human G6PD with G6PD dimers are colored in red and gray and Thr-466 highlighted in ball representation (green).

(B) Human Thr-466 (stick representation with carbon atoms colored in yellow) is positioned in close proximity to the β E- α E loop (in green), which is part of the NADP binding region (NADP highlighted in stick representation with carbon atoms colored in white). Representations were generated with PyMOL (DeLano, 2002) using the PDB codes 2BH9 and 1QKI (Au et al., 2000; Kotaka et al., 2005).

(C) In vitro kinase assay with GST-ASK α and GST-G6PD6, GST-G6PD6 T467A, or GST-G6PD6 T467E. The experiment was repeated three times, showing comparable results. WT, wild type.

(D) Enzymatic activity of G6PD6, G6PD6 T467A, and G6PD6 T467E. G6PD6 activity was quantified after a phosphorylation reaction with ASK α or ASK α K98R, or without kinase. Data are means \pm relative sd. The assay was performed four times using fresh proteins from independent purifications. Asterisks indicate a significant difference (*** P < 0.005) tested by Student's t test for pairwise comparison to the unphosphorylated control.




Figure 5. ASK α Activates G6PD6 in Vivo.

(A) Phosphorylation of G6PD6 by immunoprecipitated ASK α . Immunokinase assay using anti-myc antibodies (Santa Cruz) to purify ASK α -myc from nonstressed and high salinity-stressed ASK α -myc-expressing plants (same plant material as in Figure 1D) using GST-G6PD6 as a substrate.

(B) G6PD6 is an in vivo phosphoprotein. *Arabidopsis* protoplasts were transformed with G6PD6-HA or cotransformed with G6PD6-HA and ASK α -myc or ASK α K98R-myc and labeled with 32 P-orthophosphate. The autoradiogram shows G6PD6 immunoprecipitated from 200 μ g of protein extracts with HA antibodies. ASK α K98R-myc and untransformed protoplasts were used as specificity controls for phosphorylation and immunoprecipitation.

(C) In vivo activity of G6PD6, G6PD6 T467A, and G6PD6 T467E. G6PD activity (top) and protein levels (bottom) of *Arabidopsis* protoplasts transformed solely with G6PD6-HA, G6PD6-HA T467A, and G6PD6-HA T467E (lanes 1, 4, and 7, respectively), or cotransformed either with

elevated enzyme activity (Figure 4D). Thus, phosphorylation of G6PD6 on Thr-467 is necessary for stimulating G6PD6 activity.

G6PD6 Is Phosphorylated and Activated by ASK α in Vivo

We have shown that G6PD activity is modified in ASK α activity mutants upon salt stress. To assess whether G6PD6 might be a direct target of ASK α in vivo, we first tested whether ASK α activity immunoprecipitated from control and stressed plants could phosphorylate recombinant G6PD6. Consistent with the data shown in Figure 1, ASK α immunoprecipitated from salt-stressed ASK α -overexpressing plants showed a higher kinase activity toward G6PD6 than ASK α immunoprecipitated from plants grown under normal conditions (Figure 5A).

We next determined whether G6PD6 exists as an in vivo phosphoprotein. Hemagglutinin (HA)-tagged G6PD6 was transiently expressed in radiolabeled *Arabidopsis* protoplast cells and immunoprecipitated with anti-HA antibodies. As shown in Figure 5B, G6PD6 is phosphorylated in vivo. Cotransformation of G6PD6-HA with ASK α further enhanced G6PD6 phosphorylation, also compared with coexpression of the kinase-dead mutant ASK α K98R.

To verify whether ASK α might modulate G6PD6 activity in vivo, G6PD activity was quantified in protoplast cells transformed with G6PD6 in the presence or absence of ASK α (Figure 5C). Indeed, G6PD activity was enhanced by ASK α but not by ASK α K98R in cells expressing G6PD6. However, when protoplasts were transformed with the nonphosphorylatable mutant G6PD6 T467A, ASK α was unable to stimulate G6PD activity. Cells expressing the phosphomimicking mutant G6PD6 T467E showed constitutively high G6PD activity, which could not be further stimulated by ASK α . Collectively, these results suggest that Thr-467 phosphorylation by ASK α is necessary and sufficient for activation of G6PD6 in vivo.

Loss of G6PD6 Alters the Cellular Redox State and Renders Plants More Sensitive to High Salt Stress

Given that G6PD activity is modulated by high salt conditions, we postulated that it plays a direct role in stress tolerance. To test this, *g6pd6.2* knockout plants (see Supplemental Figures 4A and 4B online) were assayed for their stress sensitivity. Wild-type seeds germinated nearly as well on medium supplemented with 100 mM NaCl as on normal growth medium. However, the germination efficiency of *g6pd6.2* was significantly reduced under high salinity conditions (Figure 6A). Similarly, root growth of *g6pd6.2* was more strongly affected by NaCl compared with the wild type (Figure 6B). Thus, G6PD6 is indeed important for tolerance to high salt.

Glutathione is a key marker of the intracellular redox state. Consistent with a role of G6PD6 in redox regulation, levels of

ASK α -myc (lanes 2, 5, and 8) or with ASK α K98R-myc (lanes 3, 6 and 9). Data are means \pm sd. The assay was repeated three times. Asterisks indicate a significant difference (* P < 0.05) using Student's *t* test for pairwise comparison to cells solely transformed with wild-type G6PD6-HA. WT, the wild type.




Figure 6. *g6pd6.2* Shows an Altered Redox Status and Enhanced Sensitivity to Salt Stress.

(A) Germination efficiency of wild-type Col-0 and *askα* and *g6pd6.2* mutants under normal and salt stress conditions. Germination rate on control plates and on plates supplemented with 100 mM NaCl. Data are means \pm SD of three independent biological replicates with $n = 100$ each. Asterisks indicate a significant difference (** $P < 0.01$) using Student's t test for pairwise comparison to Col-0 under stress conditions.

(B) Wild-type Col-0, *askα*, and *g6pd6.2* were exposed to salt stress during early seedling development. Primary root growth under control conditions and on medium supplemented with 100 mM NaCl. Root length was determined 7 d after sowing. Data are means \pm SD of three independent experiments. Asterisks indicate a significant difference (** $P < 0.01$) using Student's t test for pairwise comparison to wild-type Col-0 under stress conditions.

(C) and **(D)** Glutathione redox status. GSSG levels **(C)** and the ratio of GSH to GSSG **(D)** in leaf extracts from Col-0, *askα*, and *g6pd6.2* grown in soil under normal or high salinity conditions. Analyses were performed with pools of 20 plants on three independent salt watering experiments. Data are means \pm relative SD. Asterisks indicate a significant difference (** $P < 0.01$) using Student's t test for pairwise comparison to wild-type Col-0 under stress conditions.

GSSG were elevated (Figure 6C) and the ratio of reduced GSH to oxidized glutathione was altered in *g6pd6.2* under salt stress conditions (Figure 6D). Early seedling development and the glutathione redox status of *askα* plants were more strongly affected by high salinity conditions than in *g6pd6.2* plants, suggesting that ASK α also regulates other targets in addition to G6PD6 for successful acclimation to stress.

We also analyzed a previously described G6PD6 T-DNA insertion line, which was unaffected by salt stress (Wakao et al., 2008). In this T-DNA insertion line, G6PD6 is disrupted downstream of the ASK α phosphorylation site Thr-467, underlining the significance of Thr-467 for adaptive regulation of G6PD.

Finally, to assess whether the effect of ASK α on stress tolerance is indeed mediated by phosphorylation of G6PD6, we used the *askα* knockout plants, which are hypersensitive to salt stress. This hypersensitivity was restored by expression of the constitutively phosphorylated mutant G6PD6 T467E (Figure 7A; see Supplemental Figure 4C online). In addition, *askα g6pd6.2*

double knockout plants showed a similar sensitivity to salt as *askα* and *g6pd6.2* (Figure 7B), suggesting that they are indeed part of the same pathway. Thus, our results support a role for T467 phosphorylation of G6PD6 by ASK α in protecting plants against salt stress.

DISCUSSION

A change in ROS levels is a key phenomenon in response to abiotic and biotic stresses (Mittler, 2002; Apel and Hirt, 2004). ROS can act as a physiological signal, but when overproduced causes oxidative damage; therefore, ROS formation and removal must be tightly regulated. Plants, like other organisms, have the potential to adapt to environmental stress, and successful protection against oxidative stress is a major determinant in coping with stressful conditions. In this study, we identified the *Arabidopsis* GSK3, ASK α , as a regulator of G6PD6 activity, glutathione redox status, and ROS levels and showed that ASK α is important for acclimation to salt stress.




Figure 7. ASK α Regulates Salt Stress Tolerance by Phosphorylating G6PD6.

(A) Phosphorylation of G6PD6 is important for salt stress tolerance. Germination efficiency of Col-0, *askα*, and *askα* mutants transformed with either G6PD6 or G6PD6 T467E on half-strength MS plates supplemented with 100 mM NaCl. Data are means \pm relative SD of three independent biological replicates with at least 75 plants each. Asterisks indicate a significant difference compared with *askα* under stress conditions (* $P < 0.05$, ** $P < 0.01$, and *** $P < 0.005$) using Student's *t* test for pairwise comparison. WT, wild type.

(B) ASK α and G6PD6 function in the same signaling pathway. Survival rate of Col-0, *askα*, *g6pd6*, and *askα g6pd6* double knockout plants on medium supplemented with 200 mM NaCl. Data are means \pm relative SD of three independent biological experiments with at least 75 plants each.

(C) Model: ASK α -mediated phosphorylation of G6PD6 contributes to maintaining cellular redox balance under high salinity conditions.

ASK α is a novel component regulating the high salinity response. Plant GSK3s are encoded by a gene family, and different GSK3 proteins have diverse functions (Jonak and Hirt, 2002; Saidi et al., 2012). In addition to ASK α , other GSK3 family members have been implicated in the response of plants to salinity. Salt stress-induced transcriptional upregulation has been reported for some GSK3 family members in different plant species (Piao et al., 1999; Charrier et al., 2002; Chen et al., 2003; Koh et al., 2007). Overexpression of *Arabidopsis* ASK α and alfalfa Msk4 enhanced salt stress tolerance (Piao et al., 2001; Kempa et al., 2007), whereas knockout mutants of rice OsGSK1 showed increased tolerance to salt stress (Koh et al., 2007), suggesting that different GSK3s have distinct, either positive or negative, regulatory functions in salt stress tolerance. Furthermore, different subcellular localization of stress-related GSK3s indicates that distinct GSK3s might be involved in different aspects of the high-salinity response.

ASK α has previously been implicated in flower development (Dornelas et al., 2000); however, using *askα* knockout and ASK α -overexpressing lines, we did not observe the flower phenotype described in ASK α antisense plants, which might be due to the

different approaches used to generate plants deficient in ASK α . Our results suggest that ASK α is activated by salt stress at the posttranslational level. The activity of mammalian GSK3 is modulated by various mechanisms, including phosphorylation and differential protein complex formation (Cohen and Frame, 2001; Woodgett, 2001; Jope and Johnson, 2004). We have shown that ASKs are phosphorylated in vivo and that Tyr-230 phosphorylation is necessary for ASK α activity (de la Fuente van Bentem et al., 2008). An important role of Tyr phosphorylation for ASK activity was confirmed in studies on BIN2/ASK η . BIN2 is phosphorylated at the corresponding Tyr (Tyr-200). Brassinosteroid-triggered signaling led to inhibition of active BIN2 via dephosphorylation by BSU1 phosphatase (Kim et al., 2009). Although Tyr phosphorylation is essential for ASK α activity (de la Fuente van Bentem et al., 2008), preliminary data indicate that salinity-induced activation of ASK α involves other mechanisms in addition to Tyr-230 phosphorylation. Future studies will aim to identify the molecular mechanism of ASK α activation during stress.

In mammals, GSK3 is a central regulatory component modulating a multitude of cellular targets (Cohen and Frame, 2001; Woodgett, 2001; Jope and Johnson, 2004). Similarly, the localization of ASK α

to the cytoplasm and the nucleus implies that ASK α might act on different organelle-specific targets. G6PD6 appears to be one of the cytosolic *in vivo* targets of ASK α . Notably, ASK γ , the closest homolog of ASK α , was unable to enhance G6PD6 activity in cells (see Supplemental Figure 5 online), thus emphasizing the specificity of ASK α on G6PD6 *in vivo* regulation.

G6PD is subject to complex control, and regulation of G6PD transcript and protein levels has been reported (Salvemini et al., 1999; Nemoto and Sasakuma, 2000; Scharte et al., 2009). Additionally, redox regulation and product inhibition participate in modulating G6PD activity (Wendt et al., 2000; Debnam et al., 2004; Schürmann and Buchanan, 2008). It is interesting to note that G6PD6 was shown to be inactivated by oxidation (Wakao and Benning, 2005) and that total G6PD activity was reduced in salt-stressed *ask* α mutants. Thus, in salt-stressed plants, ASK α might play a crucial role in counterbalancing oxidative inhibition of G6PD6 by phosphorylating and thereby enhancing G6PD6 activity.

Previous studies in plants suggested a negative regulation of G6PD by phosphorylation (Zhang et al., 2000; Hauschild and von Schaewen, 2003; Xu et al., 2005). However, we provide direct evidence that G6PD6 activity can be enhanced by phosphorylation. Consistent with our data, aestivation-induced phosphorylation of G6PD in the snail *Olata lactea* also positively correlated with G6PD activity (Ramnanan and Storey, 2006).

The ASK α phosphorylation site Thr-467 of G6PD6 is conserved in eukaryotes. Interestingly, structural analyses show that Thr-467 is in close proximity to the β E- α E loop that is part of the NADP binding region. Phosphorylation of Thr-467 might thus introduce defined structural rearrangements in the active site cleft required for proper cosubstrate binding. Consistently, mutations in the β E- α E loop were shown to affect the catalytic activity of G6PD *in vitro* and *in vivo* (Roos et al., 1999). Notably, changing Thr-467 of G6PD5 to Glu enhanced G6PD activity to a similar extent to that of G6PD6 (see Supplemental Figure 6B online), suggesting that Thr-467 phosphorylation might be a common mechanism regulating G6PD activity. Since ASK α does not phosphorylate G6PD5 on this residue (see Supplemental Figure 6A online), future studies are aimed at identifying the protein kinase targeting G6PD5.

In animals and yeast, the pentose phosphate pathway is the main source of NADPH for ROS removal (Pandolfi et al., 1995; Juhnke et al., 1996); however, in plants, the role of the OPPP during salt stress is unclear. Under normal growth conditions, photosynthesis supplies reducing equivalents; however, under stress conditions, when photosynthesis may be impaired, the OPPP might deliver reducing power for maintenance of the cellular redox balance. Several lines of evidence indicate that the OPPP is important during salt stress responses in plants. The activity of G6PD, a key enzyme of the OPPP, increased in *Arabidopsis* plants upon high salinity stress. Consistently, G6PD activity was reported to be induced by NaCl treatment in other plant species (Valderrama et al., 2006; Wang et al., 2008). Importantly, loss of G6PD6 impinged on the redox status of glutathione and rendered plants more sensitive to salt stress. These data, together with the observation that ectopic overexpression of a plastidic G6PD isoform in the cytosol enhanced drought tolerance of tobacco (*Nicotiana tabacum*; Scharte et al., 2009),

not only establish an important (positive) role of the OPPP in stress defense in plants but also provide strong evidence for a function for the cytoplasmic OPPP in stress acclimation.

In summary, our data indicate a model in which ASK α is an important regulator of ROS detoxification and, thus, acclimation to salt stress (Figure 7C). High salinity activates ASK α , which in turn phosphorylates G6PD6 on Thr-467, thereby stimulating its activity. Enhanced G6PD activity provides NADPH for the antioxidant system to remove excess ROS. Reduction of H₂O₂ to water can then be mediated by the glutathione peroxidase cycle or by the ascorbate-glutathione cycle.

G6PD is a major determinant of cellular redox homeostasis, which plays a pivotal role in determining cellular responsiveness to stress. ROS generation and redox imbalance are closely linked to aging and a wide range of diseases, including inflammation, cancer, and neurodegenerative disorders. Our data not only provide novel mechanistic insights into the regulation of G6PD6 activity by phosphorylation, but also offer a starting point for future studies beyond metabolic adaptation of plants to adverse environments.

METHODS

Plant Growth and Stress Treatments

Arabidopsis thaliana ecotype Col-0 was germinated on half-strength Murashige and Skoog (MS) medium (Duchefa). After 10 d, seedlings were transferred to soil and cultivated in a 16-h-light/8-h-dark regime at 150 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ light intensity and 60% relative humidity. For high soil salinity stress, plants were watered with a 150 mM NaCl solution. For germination and root length assay under high salt conditions, seeds were allowed to germinate on half-strength MS or half-strength MS supplemented with 100 mM NaCl in vertical plates in a 16-h-light/8-h-dark regime. Seeds used for one experiment were propagated in the same growth chamber at the same time.

Plant Material and Plasmid Constructs

The *ask* α mutant (SAIL_1055_F02) was genotyped using primers ASK α 1 5' and ASK α 9ex 3' (see Supplemental Table 1 online). The *g6pd6.2* mutant (Gabi_KAT_142G07) was genotyped using primers G6PD6 9ex and G6PD6 13ex (see Supplemental Table 1 online). The cDNA of ASK α -Myc and ASK α K98R-Myc were amplified by PCR using the primers ASK α cDNA forward and reverse (see Supplemental Table 1 online) and were cloned as *Bam*I fragments into the expression vector pGreenII0029 under the control of the 35S promoter. G6PD6-HA and its mutated variants were cloned into pGWR8 (Rozhon et al., 2010). *Arabidopsis* Col-0 plants were transformed using the floral dipping method (Clough and Bent, 1998).

Gene Expression Analysis

Total RNA was extracted from plant material using the RNeasy plant mini kit (Qiagen) and treated with RNase-free DNase following the manufacturer's protocol. RT-PCR experiments were performed with cDNAs generated from 2 μg of total RNA using oligo(dT) primers and M-MuLV reverse transcriptase (Q-BIOgene). The RT-PCR exponential phase was determined on 22 to 30 cycles. Oligonucleotide primers used for RT-PCR were designed based on the 3' untranslated region of control and selected genes (see Supplemental Table 1 online).

For real-time PCR, first-strand cDNA synthesis from 1 μg of total RNA was done using qScript cDNA SuperMix (Quanta). Samples were then diluted 1:5 with water. Five microliters of the diluted samples was used for

real-time PCR reactions using the SensiMix SYBR and Fluorescein kit (Peqlab) on an IQ5 multicolor real-time PCR detection system (Bio-Rad). All experiments were performed three times with independent RNA samples under the following cycling conditions: a 95°C hold for 10 min followed by 45 cycles at 95°C for 15 s, 58°C for 15 s, and 72°C for 15 s. Nonspecific PCR products were identified by dissociation curves. Relative expression values were calculated using the $2^{-\Delta\Delta CT}$ (for cycle threshold) method (Livak and Schmittgen, 2001). Primer efficiencies were calculated by relative standard curves. PP2A was used as a normalization control as described by Czechowski et al. (2005). Normalized gene expression was represented relative to a wild-type control. The oligonucleotide primers used were designed based on the 3' untranslated region for G6PD6 and G6PD5, or in case of ASK α , spanning the junction between exon 11 and 12 (see Supplemental Table 1 online).

Total G6PD Activity Measurement

Fifty milligrams of plant material was extracted in 0.5 mL of enzyme extraction buffer (10% glycerol, 0.25% BSA, 0.1% Triton X-100, 50 mM HEPES-KOH, pH 7.5, 10 mM MgCl₂, 1 mM EDTA, 1 mM EGTA, 1 mM benzamidine, 1 mM 6-aminocaproic acid, 1 mM phenylmethylsulfonyl fluoride, and 10 μ M leupeptin). G6PD total activity was determined as described (Stanton et al., 1991) with some modifications. Five microliters of enzyme extract was used in a 100- μ L total reaction volume. Absorbance was measured with a microplate reader spectrophotometer (GeniosPro; Tecan). In vivo G6PD activities from plants were calculated in nmol NADPH/(min*mg fresh weight) and from protoplast cells in nmol NADPH/(min* μ g protein). A minimum of four technical replicas was performed per experiment. For determination of G6PD activities from recombinant proteins, several enzyme concentrations were tested per measurement and G6PD activities were calculated in nmol NADPH/(min* μ g protein).

H₂O₂ Level Measurement

H₂O₂ levels were measured according to Wolff (1994). Briefly, 50 mg of freshly pulverized plant material was extracted with 0.5 mL of 25 mM H₂SO₄. Samples were centrifuged at 4°C, and 100 μ L of clear supernatant was added to 900 μ L of reagent solution [0.1 mM Xylenol Orange tetrasodium salt, 0.25 mM ammonium-iron(II)sulfate, 100 mM sorbitol, and 25 mM H₂SO₄]. Reactions were incubated for 1 h at room temperature in the dark. Absorbance was measured at 580 nm with a single beam spectrophotometer (Ultrospec 3100 pro; Amersham Biosciences). A calibration curve with dilutions from 0 to 10 nmol H₂O₂ in 100 μ L of 25 mM H₂SO₄ was recorded and used for quantification. To avoid oxidation of plant material, no more than 12 samples were measured at a time.

Glutathione Measurement

Oxidized and reduced forms of glutathione were determined with a spectrophotometric plate reader assay as described by Queval and Noctor (2007). Assays were performed on a BioTek's Synergy 4 Multi-Mode Microplate Reader using Corning 96-well UV-transparent plates. Standards and sample extracts were assayed as triplicates.

Arabidopsis Protoplast Transformation

Protoplasts from an *Arabidopsis* cell suspension culture were transformed as described (Cardinale et al., 2002).

In Vivo Labeling Assay

HA-tagged G6PD6 was transiently expressed in *Arabidopsis* protoplasts. One hour after transformation, [³²P] orthophosphoric acid (0.1 mCi/mL) (carrier free; Hartmann Analytic) was added and protoplasts were incubated

overnight in the dark. Immunoprecipitation was performed using HA antibodies (Santa-Cruz).

Expression and Purification of Recombinant GST-Fusion Proteins

ASK α cDNA was cloned as a *Bam*HI fragment into pGEX5x3, and cDNAs from G6PD5 and G6PD6 were cloned as *Eco*RI fragments into pGEX4T1. Recombinant proteins were expressed as GST fusion proteins in the *Escherichia coli* BL21codon plus strain. Proteins were purified using the Sepharose beads affinity method (Glutathione Sepharose 4B; GE Healthcare).

In Vitro Mutagenesis

PCR was performed using 2.5 U of Pfu Ultra (Stratagene) with the primers listed in Supplemental Table 1 online. PCR mixtures were digested with the enzyme *Dpn*I for 2 h at 37°C and transformed into *E. coli*. In vitro mutagenesis was verified by sequencing.

Antibody Production and Specificity Tests

A rabbit polyclonal ASK α -specific antibody was raised against the synthetic peptide (PNPGARDSTGVDKL) from the N terminus of ASK α . Crude serum antibody was used for immunoprecipitation of the kinase. An antibody specificity test for immunoprecipitation was performed using ASK α , and its two closest homologs, ASK γ and ASK ϵ in vitro translated with the T3/T7 coupled translation kit using the T7 primer (Amersham Biosciences). Kinases were immunoprecipitated with the ASK α antibody as described for immunokinase assays.

For the peptide-competition assay, 10 μ g of ASK α antibody was incubated with 5 μ g of synthetic ASK α peptide at 4°C overnight. After immunoprecipitation of ASK α with ASK α -specific antibodies from 20 μ g total protein extract, immunoblot analysis was performed with ASK α -specific antibodies with and without prior peptide incubation.

Immunokinase and in Vitro Kinase Assay

Immunokinase assays were performed as described (Jonak et al., 2000). G6PD or MBP was used as the specific or general substrate, respectively (Jonak et al., 2000; De Rybel et al., 2009). In vitro kinase assays using recombinant proteins were performed in a total volume of 20 μ L of kinase buffer (20 mM HEPES, pH 7.5, 15 mM MgCl₂, and 5 mM EGTA). The reaction was started with 2 μ Ci [γ -³²P]ATP and incubated at room temperature for 30 min. The reaction was stopped by the addition of 5 μ L of 4 \times SDS loading buffer. Proteins were resolved by 8% SDS-PAGE. The gel was dried and exposed overnight to a phosphor imager screen.

Mass Spectrometry and Phosphopeptide Analysis

Following the in vitro kinase assay between ASK α and G6PD6, proteins were separated by SDS-PAGE. Polyacrylamide gels were stained with a colloidal Coomassie Brilliant Blue (CBB) staining solution (Roti-Blue; Carl Roth). Excised bands were processed and putative phosphopeptides were identified by liquid chromatography-tandem mass spectrometry analysis as previously described (Penkner et al., 2009). Raw spectra were interpreted by Discoverer 1.3 (Thermo Scientific). Peptide tolerance was set to 2 D, and fragment ion tolerance was set to 0.8 D. Carbamidomethylcysteine was set as static, and oxidation of Met residues, phosphorylation of Ser, Thr, and Tyr residues as variable modifications. Phospho RS 2.0 was used to evaluate phosphorylation sites, and the spectra of possibly phosphorylated peptides were evaluated manually. An in-house generated FASTA database was used for the search containing the sequences of the target proteins, common contaminants, and proteolytic enzymes. Peptides were filtered according to the XCorr/Charge state values; false positive rate was set to 5%.

Protein Extraction and Immunoblot Analysis

Total proteins from *Arabidopsis* leaves and protoplasts were extracted and blotted on polyvinylidene fluoride membranes as described (Wrzaczek et al., 2007). Protein concentration was assessed using the Bradford method. Membranes were probed with a 1:5000 dilution of anti-ASK α , anti-Myc (Santa-Cruz), or anti-HA (Santa-Cruz) antibodies. To detect total G6PD protein levels, a 1:6000 dilution of polyclonal antibody against *Saccharomyces cerevisiae* G6PD (Sigma-Aldrich) was used as described for plants (Valderrama et al., 2006). Alkaline phosphatase-conjugated goat anti-rabbit IgG (Santa-Cruz) was used as the secondary antibody. The reaction was detected using the CDP-Star detection reagent (GE Healthcare).

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative under the following accession numbers: ASK α , At5g26751; G6PD6, AT5G40760; G6PD5, AT3G27300; β -tubulin, At5g12250; GAPDH, AT1G13440; and PP2A, AT1G13320.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure 1. *ask* α T-DNA Insertion Mutant and Specificity of the Anti-ASK α Antibody.

Supplemental Figure 2. Mass Spectrometry Analysis of G6PD6 Phosphorylation by ASK α .

Supplemental Figure 3. G6PD6 Thr-467 Is Evolutionary Conserved.

Supplemental Figure 4. ASK α and G6PD6 Function in the Same Signaling Pathway.

Supplemental Figure 5. ASK γ Does Not Enhance G6PD6 Activity in Vivo.

Supplemental Figure 6. Characterization of Thr-467 of G6PD5.

Supplemental Table 1. List of Primers.

ACKNOWLEDGMENTS

We thank Anneliese Auer, Andrey Belokourow, and Bettina Dekrout for technical assistance, Edina Csaszar for mass spectrometry, and Manuela Baccarini for critical comments on the article. This work was supported by grants from the Austrian Science Foundation (P20375-B03) and the Vienna Science and Technology Fund (LS 123).

AUTHOR CONTRIBUTIONS

S.D.S. and H.S. analyzed the stress tolerance and performed the protein kinase and enzyme activity assays. S.D.S. identified the phosphorylation site. H.S. and J.K. analyzed the cellular redox status. S.K. and Y.G. performed the robot-based screen for enzyme activities. E.P. and W.R. generated transgenic plants. A.H. and T.C. performed the structural analyses. C.J. conceived, designed, and evaluated the experiments and wrote the article.

Received June 6, 2012; revised July 6, 2012; accepted July 12, 2012; published August 10, 2012.

REFERENCES

- Apel, K., and Hirt, H.** (2004). Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **55**: 373–399.

Au, S.W., Gover, S., Lam, V.M., and Adams, M.J. (2000). Human glucose-6-phosphate dehydrogenase: The crystal structure reveals a structural NADP(+) molecule and provides insights into enzyme deficiency. *Structure* **8**: 293–303.

Boyer, J.S. (1982). Plant productivity and environment. *Science* **218**: 443–448.

Cardinale, F., Meskiene, I., Ouaked, F., and Hirt, H. (2002). Convergence and divergence of stress-induced mitogen-activated protein kinase signaling pathways at the level of two distinct mitogen-activated protein kinase kinases. *Plant Cell* **14**: 703–711.

Charrier, B., Champion, A., Henry, Y., and Kreis, M. (2002). Expression profiling of the whole Arabidopsis shaggy-like kinase multigene family by real-time reverse transcriptase-polymerase chain reaction. *Plant Physiol.* **130**: 577–590.

Chen, G.P., Ma, W.S., Huang, Z.J., Xu, T., Xue, Y.B., and Shen, Y.Z. (2003). Isolation and characterization of TaGSK1 involved in wheat salt tolerance. *Plant Sci.* **165**: 1369–1375.

Clough, S.J., and Bent, A.F. (1998). Floral dip: A simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**: 735–743.

Cohen, P., and Frame, S. (2001). The renaissance of GSK3. *Nat. Rev. Mol. Cell Biol.* **2**: 769–776.

Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K., and Scheible, W.R. (2005). Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiol.* **139**: 5–17.

Debnam, P.M., and Emes, M.J. (1999). Subcellular distribution of enzymes of the oxidative pentose phosphate pathway in root and leaf tissues. *J. Exp. Bot.* **50**: 1653–1661.

Debnam, P.M., Fernie, A.R., Lisse, A., Golding, A., Bowsher, C.G., Grimshaw, C., Knight, J.S., and Emes, M.J. (2004). Altered activity of the P2 isoform of plastidic glucose 6-phosphate dehydrogenase in tobacco (*Nicotiana tabacum* cv. Samsun) causes changes in carbohydrate metabolism and response to oxidative stress in leaves. *Plant J.* **38**: 49–59.

de la Fuente van Bentem, S., et al. (2008). Site-specific phosphorylation profiling of Arabidopsis proteins by mass spectrometry and peptide chip analysis. *J. Proteome Res.* **7**: 2458–2470.

DeLano, W.L. (2002). The PyMOL Molecular Graphics System. (Palo Alto, CA: DeLano Scientific).

De Rybel, B., et al. (2009). Chemical inhibition of a subset of *Arabidopsis thaliana* GSK3-like kinases activates brassinosteroid signaling. *Chem. Biol.* **16**: 594–604.

Dieni, C.A., and Storey, K.B. (2010). Regulation of glucose-6-phosphate dehydrogenase by reversible phosphorylation in liver of a freeze tolerant frog. *J. Comp. Physiol. B* **180**: 1133–1142.

Doble, B.W., and Woodgett, J.R. (2003). GSK-3: Tricks of the trade for a multi-tasking kinase. *J. Cell Sci.* **116**: 1175–1186.


Dornelas, M.C., Van Lammeren, A.A., and Kreis, M. (2000). *Arabidopsis thaliana* SHAGGY-related protein kinases (AtSK11 and 12) function in perianth and gynoecium development. *Plant J.* **21**: 419–429.

Filosa, S., Fico, A., Paglialunga, F., Balestrieri, M., Croke, A., Verde, P., Abrescia, P., Bautista, J.M., and Martini, G. (2003). Failure to increase glucose consumption through the pentose-phosphate pathway results in the death of glucose-6-phosphate dehydrogenase gene-deleted mouse embryonic stem cells subjected to oxidative stress. *Biochem. J.* **370**: 935–943.

Gibon, Y., Blaessing, O.E., Hannemann, J., Carillo, P., Höhne, M., Hendriks, J.H., Palacios, N., Cross, J., Selbig, J., and Stitt, M. (2004). A Robot-based platform to measure multiple enzyme activities in *Arabidopsis* using a set of cycling assays: Comparison of changes of enzyme activities and transcript levels during diurnal cycles and in prolonged darkness. *Plant Cell* **16**: 3304–3325.

- Gupte, R.S., Ata, H., Rawat, D., Abe, M., Taylor, M.S., Ochi, R., and Gupte, S.A.** (2011). Glucose-6-phosphate dehydrogenase is a regulator of vascular smooth muscle contraction. *Antioxid. Redox Signal.* **14**: 543–558.
- Gupte, R.S., Floyd, B.C., Kozicky, M., George, S., Ungvari, Z.I., Neito, V., Wolin, M.S., and Gupte, S.A.** (2009). Synergistic activation of glucose-6-phosphate dehydrogenase and NAD(P)H oxidase by Src kinase elevates superoxide in type 2 diabetic, Zucker fa/fa, rat liver. *Free Radic. Biol. Med.* **47**: 219–228.
- Hasegawa, P.M., Bressan, R.A., Zhu, J.K., and Bohnert, H.J.** (2000). Plant cellular and molecular responses to high salinity. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **51**: 463–499.
- Hauschild, R., and von Schaewen, A.** (2003). Differential regulation of glucose-6-phosphate dehydrogenase isoenzyme activities in potato. *Plant Physiol.* **133**: 47–62.
- Izawa, S., Maeda, K., Miki, T., Mano, J., Inoue, Y., and Kimura, A.** (1998). Importance of glucose-6-phosphate dehydrogenase in the adaptive response to hydrogen peroxide in *Saccharomyces cerevisiae*. *Biochem. J.* **330**: 811–817.
- Jonak, C., Beisteiner, D., Beyerly, J., and Hirt, H.** (2000). Wound-induced expression and activation of WIG, a novel glycogen synthase kinase 3. *Plant Cell* **12**: 1467–1475.
- Jonak, C., and Hirt, H.** (2002). Glycogen synthase kinase 3/SHAGGY-like kinases in plants: An emerging family with novel functions. *Trends Plant Sci.* **7**: 457–461.
- Joep, R.S., and Johnson, G.V.** (2004). The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem. Sci.* **29**: 95–102.
- Juhnke, H., Krems, B., Kötter, P., and Entian, K.D.** (1996). Mutants that show increased sensitivity to hydrogen peroxide reveal an important role for the pentose phosphate pathway in protection of yeast against oxidative stress. *Mol. Gen. Genet.* **252**: 456–464.
- Kempa, S., Rozhon, W., Samaj, J., Erban, A., Baluska, F., Becker, T., Haselmayer, J., Schleiff, E., Kopka, J., Hirt, H., and Jonak, C.** (2007). A plastid-localized glycogen synthase kinase 3 modulates stress tolerance and carbohydrate metabolism. *Plant J.* **49**: 1076–1090.
- Kim, T.W., Guan, S., Sun, Y., Deng, Z., Tang, W., Shang, J.X., Sun, Y., Burlingame, A.L., and Wang, Z.Y.** (2009). Brassinosteroid signal transduction from cell-surface receptor kinases to nuclear transcription factors. *Nat. Cell Biol.* **11**: 1254–1260.
- Koh, S., Lee, S.-C., Kim, M.-K., Koh, J.H., Lee, S., An, G., Choe, S., and Kim, S.-R.** (2007). T-DNA tagged knockout mutation of rice OsGSK1, an orthologue of Arabidopsis BIN2, with enhanced tolerance to various abiotic stresses. *Plant Mol. Biol.* **65**: 453–466.
- Kotaka, M., Gover, S., Vandeputte-Rutten, L., Au, S.W., Lam, V.M., and Adams, M.J.** (2005). Structural studies of glucose-6-phosphate and NADP+ binding to human glucose-6-phosphate dehydrogenase. *Acta Crystallogr. D Biol. Crystallogr.* **61**: 495–504.
- Livak, K.J., and Schmittgen, T.D.** (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C(T)}$ method. *Methods* **25**: 402–408.
- Martin, M., Rehani, K., Joep, R.S., and Michalek, S.M.** (2005). Toll-like receptor-mediated cytokine production is differentially regulated by glycogen synthase kinase 3. *Nat. Immunol.* **6**: 777–784.
- Mittler, R.** (2002). Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* **7**: 405–410.
- Mittler, R., Vanderauwera, S., Suzuki, N., Miller, G., Tognetti, V.B., Vandepoele, K., Gollery, M., Shulaev, V., and Van Breusegem, F.** (2011). ROS signaling: The new wave? *Trends Plant Sci.* **16**: 300–309.
- Møller, I.M., Jensen, P.E., and Hansson, A.** (2007). Oxidative modifications to cellular components in plants. *Annu. Rev. Plant Biol.* **58**: 459–481.
- Munns, R.** (2002). Comparative physiology of salt and water stress. *Plant Cell Environ.* **25**: 239–250.
- Nemoto, Y., and Sasakuma, T.** (2000). Specific expression of glucose-6-phosphate dehydrogenase (G6PDH) gene by salt stress in wheat (*Triticum aestivum* L.). *Plant Sci.* **158**: 53–60.
- Pandolfi, P.P., Sonati, F., Rivi, R., Mason, P., Grosveld, F., and Luzzatto, L.** (1995). Targeted disruption of the housekeeping gene encoding glucose 6-phosphate dehydrogenase (G6PD): G6PD is dispensable for pentose synthesis but essential for defense against oxidative stress. *EMBO J.* **14**: 5209–5215.
- Penkner, A.M., Fridkin, A., Gloggnitzer, J., Baudrimont, A., Machacek, T., Woglar, A., Csaszar, E., Pasierbek, P., Ammerer, G., Gruenbaum, Y., and Jantsch, V.** (2009). Meiotic chromosome homology search involves modifications of the nuclear envelope protein Matefin/SUN-1. *Cell* **139**: 920–933.
- Piao, H.L., Lim, J.H., Kim, S.J., Cheong, G.W., and Hwang, I.** (2001). Constitutive over-expression of AtGSK1 induces NaCl stress responses in the absence of NaCl stress and results in enhanced NaCl tolerance in Arabidopsis. *Plant J.* **27**: 305–314.
- Piao, H.L., Pih, K.T., Lim, J.H., Kang, S.G., Jin, J.B., Kim, S.H., and Hwang, I.** (1999). An Arabidopsis GSK3/shaggy-like gene that complements yeast salt stress-sensitive mutants is induced by NaCl and abscisic acid. *Plant Physiol.* **119**: 1527–1534.
- Queval, G., and Noctor, G.** (2007). A plate reader method for the measurement of NAD, NADP, glutathione, and ascorbate in tissue extracts: Application to redox profiling during Arabidopsis rosette development. *Anal. Biochem.* **363**: 58–69.
- Ramnanan, C.J., and Storey, K.B.** (2006). Glucose-6-phosphate dehydrogenase regulation during hypometabolism. *Biochem. Biophys. Res. Commun.* **339**: 7–16.
- Roos, D., et al.** (1999). Molecular basis and enzymatic properties of glucose 6-phosphate dehydrogenase volendam, leading to chronic nonspherocytic anemia, granulocyte dysfunction, and increased susceptibility to infections. *Blood* **94**: 2955–2962.
- Rozhon, W., Mayerhofer, J., Petutschnig, E., Fujioka, S., and Jonak, C.** (2010). ASKtheta, a group-III Arabidopsis GSK3, functions in the brassinosteroid signalling pathway. *Plant J.* **62**: 215–223.
- Saidi, Y., Hearn, T.J., and Coates, J.C.** (2012). Function and evolution of 'green' GSK3/Shaggy-like kinases. *Trends Plant Sci.* **17**: 39–46.
- Salvemini, F., Franzé, A., Iervolino, A., Filosa, S., Salzano, S., and Ursini, M.V.** (1999). Enhanced glutathione levels and oxidoresistance mediated by increased glucose-6-phosphate dehydrogenase expression. *J. Biol. Chem.* **274**: 2750–2757.
- Scharte, J., Schön, H., Tjaden, Z., Weis, E., and von Schaewen, A.** (2009). Isoenzyme replacement of glucose-6-phosphate dehydrogenase in the cytosol improves stress tolerance in plants. *Proc. Natl. Acad. Sci. USA* **106**: 8061–8066.
- Schürmann, P., and Buchanan, B.B.** (2008). The ferredoxin/thioredoxin system of oxygenic photosynthesis. *Antioxid. Redox Signal.* **10**: 1235–1274.
- Sindelár, L., and Sindelárová, M.** (2002). Correlation of viral RNA biosynthesis with glucose-6-phosphate dehydrogenase activity and host resistance. *Planta* **215**: 862–869.
- Slaski, J.J., Zhang, G., Basu, U., Stephens, J.L., and Taylor, G.J.** (2006). Aluminum resistance in wheat (*Triticum aestivum*) is associated with rapid, Al-induced changes in activities of glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase in root apices. *Physiol. Plant.* **98**: 477–487.
- Stanton, R.C., Seifert, J.L., Boxer, D.C., Zimmerman, E., and Cantley, L.C.** (1991). Rapid release of bound glucose-6-phosphate dehydrogenase by growth factors. Correlation with increased enzymatic activity. *J. Biol. Chem.* **266**: 12442–12448.

- Valderrama, R., Corpas, F.J., Carreras, A., Gómez-Rodríguez, M.V., Chaki, M., Pedrajas, J.R., Fernández-Ocaña, A., Del Río, L.A., and Barroso, J.B.** (2006). The dehydrogenase-mediated recycling of NADPH is a key antioxidant system against salt-induced oxidative stress in olive plants. *Plant Cell Environ.* **29**: 1449–1459.
- Wakao, S., Andre, C., and Benning, C.** (2008). Functional analyses of cytosolic glucose-6-phosphate dehydrogenases and their contribution to seed oil accumulation in Arabidopsis. *Plant Physiol.* **146**: 277–288.
- Wakao, S., and Benning, C.** (2005). Genome-wide analysis of glucose-6-phosphate dehydrogenases in Arabidopsis. *Plant J.* **41**: 243–256.
- Wang, X., Ma, Y., Huang, C., Wan, Q., Li, N., and Bi, Y.** (2008). Glucose-6-phosphate dehydrogenase plays a central role in modulating reduced glutathione levels in reed callus under salt stress. *Planta* **227**: 611–623.
- Wendt, U.K., Wenderoth, I., Tegeler, A., and Von Schaewen, A.** (2000). Molecular characterization of a novel glucose-6-phosphate dehydrogenase from potato (*Solanum tuberosum* L.). *Plant J.* **23**: 723–733.
- Wolff, S.P.** (1994). Ferrous ion oxidation in presence of ferric ion indicator xenol orange for measurement of hydroperoxides. *Methods Enzymol.* **233**: 182–189.
- Woodgett, J.R.** (2001). Judging a protein by more than its name: GSK-3. *Sci. STKE* **2001**: re12.
- Wrzaczek, M., Rozhon, W., and Jonak, C.** (2007). A proteasome-regulated glycogen synthase kinase-3 modulates disease response in plants. *J. Biol. Chem.* **282**: 5249–5255.
- Xu, Y., Osborne, B.W., and Stanton, R.C.** (2005). Diabetes causes inhibition of glucose-6-phosphate dehydrogenase via activation of PKA, which contributes to oxidative stress in rat kidney cortex. *Am. J. Physiol. Renal Physiol.* **289**: F1040–F1047.
- Zhang, Z., Apse, K., Pang, J., and Stanton, R.C.** (2000). High glucose inhibits glucose-6-phosphate dehydrogenase via cAMP in aortic endothelial cells. *J. Biol. Chem.* **275**: 40042–40047.
- Zhu, J.K.** (2001). Plant salt tolerance. *Trends Plant Sci.* **6**: 66–71.



Supplemental Figure 1. The *aska* T-DNA insertion mutant and specificity of the anti-ASKα antibody.

(A) Schematic diagram of *ASKα* showing the site of T-DNA insertion in *aska*. Black boxes indicate coding and white boxes intron regions.

(B) qRT-PCR analysis of wild-type Col 0, *aska* and ASKα overexpressor lines OE#4 and OE#5 showing the absence of a full-length *ASKα* transcript in *aska* plants and enhanced *ASKα* transcript levels in ASKα overexpressor lines. Data are means \pm relative SD of three independent experiments.


(C) Four-week-old *A. thaliana* wild-type Col 0, *aska* and ASKα-overexpressor lines OE#4 and OE#5 grown under control conditions in soil.

(D) ³⁵S-methionine-labelled *in vitro*-translated proteins ASKα, ASKγ and ASKε (lanes 1 to 3, respectively) and immunoprecipitations (IP) of the *in vitro*-translated proteins ASKα, ASKγ and ASKε with anti-ASKα antibody (lanes 4 to 6, respectively).

(E) Immunoprecipitation blot analysis of *Arabidopsis* leaf protein extract. ASKα was immunoprecipitated with ASKα-specific antibodies from 100 μg of total protein extract. Subsequent analysis was performed with an anti-ASKα antibody with (+) or without (-) prior blocking of the antibody with the N-terminal ASKα peptide.


A

CID MS/MS spectrum of VAWEIF#TPLLHR (m/z 781.53 Da); Xcorr:3.41; phospho RS score:114



B

CID MS/MS/MS spectrum of VAWEIF#TPLLHR (m/z 781.53 Da); Xcorr:2.18



C

MS/MS of m/z 781.53 Da

#1	b ⁺	b ²⁺	Seq.	y ⁺	y ²⁺	#2
1	100.07570	50.54149	V			12
2	171.11282	86.06005	A	1462.72432	731.86580	11
3	357.19214	179.09971	W	1391.68720	696.34724	10
4	486.23474	243.62101	E	1205.60788	603.30758	9
5	599.31881	300.16304	I	1076.56528	538.78628	8
6	746.38723	373.69725	F	963.48121	482.24424	7
7	927.40124	464.20426	T-Phospho	816.41279	408.71003	6
8	1024.45401	512.73064	P	635.39878	318.20303	5
9	1137.53808	569.27268	L	538.34601	269.67664	4
10	1250.62215	625.81471	L	425.26194	213.13461	3
11	1387.68106	694.34417	H	312.17787	156.59257	2
12			R	175.11896	88.06312	1

D

MS/MS/MS of m/z 781.53 Da

#1	b ⁺	b ²⁺	Seq.	y ⁺	y ²⁺	#2
1	100.07570	50.54149	V			12
2	171.11282	86.06005	A	1462.72432	731.86580	11
3	357.19214	179.09971	W	1391.68720	696.34724	10
4	486.23474	243.62101	E	1205.60788	603.30758	9
5	599.31881	300.16304	I	1076.56528	538.78628	8
6	746.38723	373.69725	F	963.48121	482.24424	7
7	927.40124	464.20426	T-Phospho	816.41279	408.71003	6
8	1024.45401	512.73064	P	635.39878	318.20303	5
9	1137.53808	569.27268	L	538.34601	269.67664	4
10	1250.62215	625.81471	L	425.26194	213.13461	3
11	1387.68106	694.34417	H	312.17787	156.59257	2
12			R	175.11896	88.06312	1

Supplemental Figure 2. Mass spectrometry (MS) analysis of G6PD6 phosphorylation by ASK α . (A) CID (collision induced decay) MS/MS and (B) CID MS/MS/MS spectrum of VAWEIF#TPLLHR. (C, D) List of fragment ions of MS/MS (C) and MS/MS/MS (D) of m/z 781.53 Da. Detected fragment ions are labeled red or blue for b or y ions, respectively.


```

AtG6PD6      QHFVRRDELKVAWEIFTPLLLHRIDKGE-VKSIPIYKPGSRGPKEADQLLEKAGYLQTHGYI 509
AtG6PD5      QHFVRRDELKAWEIFTPLLLHRIDKGE-VKSVPYKQSGRGPAEADQLLLKAGYMQTHGYI 510
Potato       QHFVRRDELKAWEIFTPLLLHRIDNGE-VKPIPYKPSGRGPAEADELLQNAGYVQTHGYI 505
Poplar       QHFVRRDELKAWEIFTPLLLHRIDNGE-MKPKEYQPSGRGPVEADELLAKAGYVQTHGYI 505
Grapevine    QHFVRRDELKAWEIFTPLLLHRIDNGE-MKPIPYKPSGRGPSEADELLSKSGYVQTHGYI 510
Rice         QHFVRRDELKAAWQIFTPLLLHDIDEGK-VKSIPYQPSGRGPKEADELSERVGYMQTHGYI 498
Wheat        QHFVRRDELKAAWQIFTPLLLHDIDAGK-LKAVSYKPSGRGPKEADELSEKVGYMQTHGYI 502
Medicago    QHFVRRDELKASWQIFTPLLLHKIDRGE-LKPVPYKPSGRGPAEADELLEKAGYVQTHGYI 509
Pea          QHFVRRDELKASWEIFTPLLLHKIDRGE-LKPIPYKPSGRGPAEADELLEKAGYVQTHGYI 511
Chlamydomonas QHFVRRDELRAAWAIFTPLLLHAIDAGA-VPLHPYPYGAR----- 286
Physcomitrella QHFVRRDELKVAWEIFTPLLLHRIDVGK-LELIPYKESGRGPAEADELNARVGYRRTEGYC 520
Yeast        SNFVRDELDISWGIEFTPLLLKHIERPDGTPEIYPYGSRGPKGLKEYMQKHKYVMPEKHP 491
Drosophila   MHFVRSDELREAWRIFTPLLLHKIEHER-IPPIPYYGSRGPTEADRKCVENNFIYSASYK 514
Mouse        MHFVRIDELREGWRIFTPLLLHKIEREK-PQFPPYVYGSRGPTEADELMRRVGFQYKGTYK 508
Human        MHFVRSDELREAWRIFTPLLLQIELEK-PKPIPIYYGSRGPTEADELMKRVGFQYEGTYK 508
:*** ** . * *****: * : * * *

```

Supplemental Figure 3. G6PD6 threonine 467 is conserved.


Partial protein alignment of eukaryotic G6PDs was performed with ClustalW using Blosum as protein weight matrix, 10 as GAP Open and 0,1 as Gap Extension. Thr467 of G6PDs is boxed. Asterisks denote identical residues, colons indicate conserved amino acid substitutions and periods designate semi-conservative substitutions. AtG6PD6 and AtG6PD5, *Arabidopsis thaliana*, Q9FJ15 and Q9LK23, respectively; Potato, *Solanum tuberosum*, P37830; Poplar, *Populus trichocarpa*, Q2L9V8; Grapevine, *Vitis vinifera*, D7UBH2; Rice, *Oryza sativa* subsp. japonica, Q8RY51; Wheat, *Triticum aestivum*, Q9LR19; *Medicago truncatula*, G7KS28; Pea, *Pisum sativum*, A6XIG0; *Chlamydomonas reinhardtii*, A8J6N3; *Physcomitrella patens* subsp. patens, A9TA54; Yeast, *Saccharomyces cerevisiae*, P11412; *Drosophila*, *Drosophila mojavensis*, B4L7Z3; Mouse, *Mus musculus*, P97324; Human, *Homo sapiens*, P11413.




Supplemental Figure 4. ASK α and G6PD6 function in the same signaling pathway.

(A, B) *g6pd6.2* T-DNA insertion mutant. Schematic diagram of G6PD6 showing the site of T-DNA insertion in the *g6pd6.2* mutant (A). Black boxes indicate coding and white boxes indicate non-coding regions. The position of Thr467 is depicted in red. RT-PCR analysis of wild-type (WT) and *g6pd6.2* (B). No full length G6PD6 transcript was present in *g6pd6.2* plants in three independent biological replicas.

(C) Wild-type G6PD6-HA and G6PD6T467E-HA protein levels in *askα* plants. Upper: Wild-type G6PD6-HA and G6PD6T467E-HA protein levels were detected by western blot analysis using an anti-HA antibody. Lower: Coomassie stained gel.



Supplemental Figure 5. ASK γ does not enhance G6PD6 activity *in vivo*. *In vivo* G6PD activity of *Arabidopsis* protoplasts transformed with G6PD6-HA or co-transformed with G6PD6-HA and ASK γ -myc. Data are means \pm SD with n = 3. The assays were performed three times independently and showed similar results.



Supplemental Figure 6. Characterization of Thr467 of G6PD5.

(A) *In vitro* kinase assay with GST-ASK α and GST-G6PD5, GST-G6PD5 T467A or GST-G6PD5 T467E. The experiment was repeated three times showing comparable results.

(B) G6PD5 T467E displays enhanced enzymatic activity. Enzymatic activity of G6PD5, G6PD5 T467A and G6PD5 T467E. G6PD6 activity was quantified after a phosphorylation reaction with ASK α , ASK α K98R or without kinase. Data are means \pm relative SD. The assay was performed four times using fresh proteins from independent purifications. Asterisk indicates a significant difference (* P <0.05) tested by Student's t-test for pairwise comparison to the un-phosphorylated control.

Gene	Primer Name	Primer Sequence	Purpose
At-ASK α	qRT_ASK α _11/12ex	(F) 5'-AAGAGACCCAAACGCACGTCTACC-3'	qRT-PCR
		(R) 5'-ACACCTTTCAGCTCGTGAGGCTT-3'	qRT-PCR
	ASK α cDNA	(F) 5'-GGATCCATTATGGCGTCAGTGGGTATAGCTCC-3'	Cloning
		(R) 5'-GGATCCGCGCGCCGCAAACCGAGCCAAGGACACTGC-3'	Cloning
	ASK α K98R	(F) 5'-GTTGCGATAAGGAAAGTTTTACAAGATAGG-3'	Mutagenesis
		(R) 5'-CCTATCTTGTAATAACTTTCTTATCGCAAC-3'	Mutagenesis
	ASK α 1 5'	(F) 5'-TGGCTGAGCGTGTGTTGGTCA-3'	Genotyping
	ASK α 9ex 3'	(R) 5'-GAATTTGAATTCCTGTAGTTTGG-3'	Genotyping
At-G6PD6	G6PD6 cDNA	(F) 5'-GCATGGGATCTGGTCAATGGCACGTTGAG-3'	Cloning pGEX4T1
		(R) 5'-TAGTGTAGGAGGGATCCAGATATAGCC-3'	Cloning pGEX4T1
	G6PD6 cDNA1	(F) 5'-GGAGGTGGATCCATGGGATCTGGTCAATGG-3'	Cloning pGWR8
		(R) 5'-ACCTCCGGATCC TAGTGTAGGAGGAATCCAG-3'	Cloning pGWR8
	G6PD6 T467A	(F) 5'-GCGTGGGAGATCTTCGCGCCGCTACTCCACAGG-3'	Mutagenesis
		(R) 5'-CCTGTGGAGTAGCGGCGCGAAGATCTCCACGC-3'	Mutagenesis
	G6PD6 T467E	(F) 5'-GCGTGGGAGATCTTCGAGCCGCTACTCCACAGG-3'	Mutagenesis
		(R) 5'-CCTGTGGAGTAGCGGCTCGAAGATCTCCACGC-3'	Mutagenesis
G6PD6 9ex	(F) 5'-TTGCCATGGAGAAACCAATATCTC-3'	Genotyping	
G6PD6 13ex	(R) 5'-ACCTTGATACCGTTGCCCATACG-3'	Genotyping	
At-G6PD5	G6PD5 cDNA	(F) 5'-GCATGGGTTCTGGTCAATGGCATATGGAG-3'	Cloning pGEX4T1
		(R) 5'-CAATGTAGGAGGGATCCAAATGTAGCC-3'	Cloning pGEX4T1
	G6PD5 cDNA1	(F) 5'-GGAGGTGGATCCATGGGTTCTGGTCAAT-3'	Cloning pGWR8
		(R) 5'-ACCTCCGGATCCCAATGTAGGAGGAATCC-3'	Cloning pGWR8
At- β -Tubulin	RT-TUB	(F) 5'-ACCACTCCTAGCTTTGGTGATCTG-3'	RT-PCR
		(R) 5'-AGGTTCACTGCGAGCTTCCTCA-3'	RT-PCR
At-PP2A	qRT-PP2A	(F) 5'-TAACGTGGCCAAATGATGC-3'	qRT-PCR
		(R) 5'-GTTCTCCACAACCGCTTGGT-3'	qRT-PCR
At-GAPDH	qRT-GAPDH	(F) 5'-TTGGTGACAACAGGTCAAGCA-3'	qRT-PCR
		(R) 5'-AAACTTGTCGCTCAATGCAATC-3'	qRT-PCR
Gabi T-DNA LB	GK_LB	(R) 5'-CCCATTGGACGTGAATGTAGACAC-3'	Genotyping
Sail T-DNA LB	SAIL_LB1	(R) 5'-TAGCATCTGAATTTCCATAACCAATCTCGA-3'	Genotyping

Supplemental Table 1. List of primers.

Mutated nucleotides are depicted in red. The BamHI site is shown in pink, the NotI site in green, and in blue nucleotides added to maintain the right frame.