



Annotated genetic linkage maps of *Pinus pinaster* Ait. from a Central Spain population using microsatellite and gene based markers

Marina de Miguel, Nuria de Maria, Luis Diaz, Enrique Sáez-Laguna, David Sánchez-Gómez, Emilie E. Chancerel, Ismael Aranda, Carmen Collada, Christophe Plomion, José-Antonio Cabezas, et al.

► To cite this version:

Marina de Miguel, Nuria de Maria, Luis Diaz, Enrique Sáez-Laguna, David Sánchez-Gómez, et al.. Annotated genetic linkage maps of *Pinus pinaster* Ait. from a Central Spain population using microsatellite and gene based markers. BMC Genomics, 2012, 13, 14 p. 10.1186/1471-2164-13-527 . hal-02648778

HAL Id: hal-02648778

<https://hal.inrae.fr/hal-02648778>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access

Annotated genetic linkage maps of *Pinus pinaster* Ait. from a Central Spain population using microsatellite and gene based markers

Marina de Miguel^{1,2}, Nuria de Maria^{1,2}, M Ángeles Guevara^{1,2}, Luis Díaz^{1,2}, Enrique Sáez-Laguna^{1,2}, David Sánchez-Gómez¹, Emilie Chancerel^{4,5}, Ismael Aranda¹, Carmen Collada^{2,3}, Christophe Plomion^{4,5}, José-Antonio Cabezas^{1,2} and María-Teresa Cervera^{1,2*}

Abstract

Background: *Pinus pinaster* Ait. is a major resin producing species in Spain. Genetic linkage mapping can facilitate marker-assisted selection (MAS) through the identification of Quantitative Trait Loci and selection of allelic variants of interest in breeding populations. In this study, we report annotated genetic linkage maps for two individuals (C14 and C15) belonging to a breeding program aiming to increase resin production. We use different types of DNA markers, including last-generation molecular markers.

Results: We obtained 13 and 14 linkage groups for C14 and C15 maps, respectively. A total of 211 and 215 markers were positioned on each map and estimated genome length was between 1,870 and 2,166 cM respectively, which represents near 65% of genome coverage. Comparative mapping with previously developed genetic linkage maps for *P. pinaster* based on about 60 common markers enabled aligning linkage groups to this reference map. The comparison of our annotated linkage maps and linkage maps reporting QTL information revealed 11 annotated SNPs in candidate genes that co-localized with previously reported QTLs for wood properties and water use efficiency.

Conclusions: This study provides genetic linkage maps from a Spanish population that shows high levels of genetic divergence with French populations from which segregating progenies have been previously mapped. These genetic maps will be of interest to construct a reliable consensus linkage map for the species. The importance of developing functional genetic linkage maps is highlighted, especially when working with breeding populations for its future application in MAS for traits of interest.

Keywords: *Pinus pinaster*, Genetic linkage map, Functional annotation, Microsatellites, SNPs

Background

Maritime pine (*Pinus Pinaster* Ait.) is one of the most important species in the Mediterranean region for its ecology and wood productiveness. As other conifers, this long lived species dominates different landscapes and can withstand severe environmental conditions [1]. Several studies have revealed high levels of phenotypic variation [2-4] and genetic diversity [5-7] in maritime pine. This species has a fragmented geographic distribution

that could be subdivided into different meta-populations based on its high level of genetic differentiation [8-10]. In the Iberian Peninsula different patterns of local adaptation have been identified [11]. Besides its ecological value, maritime pine is also a significant species for its economic importance. Particularly, *P. pinaster* is a major resin producing species in the Iberian Peninsula [12]. The resin is at the basis of many manufactured products such as turpentine, oils, varnishes, sealing wax, plastics and others. In 1990s resin tapping was reintroduced in Spain after a drastic reduction in 1970s due to the international crisis in this sector [13]. Many of the abandoned stands have been tapped again. In particular,

* Correspondence: cervera@inia.es

¹INIA-CIFOR, Departamento de Ecología y Genética Forestal, Carretera de la Coruña Km7.5, Madrid 28040, Spain

²Unidad Mixta de Genómica y Ecofisiología Forestal, INIA/UPM, Madrid, Spain
Full list of author information is available at the end of the article

natural stands of Central Spain are one of the most important resin tapping regions [14]. As resin production shows high heritability [15] a breeding program is a useful strategy to improve productiveness [16]. Consequently several breeding programs have been implemented for resin production in maritime pine [17-19].

Genetic linkage mapping can facilitate marker-assisted selection (MAS) as it allows the identification of quantitative trait Loci (QTL) [20-23]. Furthermore, as genome organization is well conserved in conifers, comparative mapping is a useful strategy to find homologous chromosomal segments involved in the genetic control of economical and adaptive traits [24,25].

Traditional molecular makers, such as proteins, RFLPs (Restriction Fragment Length Polymorphisms), RAPDs (Random Amplified Polymorphic DNAs), AFLPs (Amplified Fragment Length Polymorphisms) and nSSRs (nuclear Simple Sequence Repeats) have helped to build a first generation genetic linkage maps in forest trees [26,27]. The use of RAPDs and AFLPs, randomly distributed in the genome [28,29], has allowed the construction of genetic linkage maps from species with large genome sizes like conifers [30-32]. An alternative for species with extremely large genomes or for populations with low levels of polymorphism are SAMPL markers (Selective Amplification of Microsatellite Polymorphic Loci). SAMPL combines the advantages of AFLPs and microsatellites resulting in higher percentage of polymorphic markers per assay and higher repeatability between assays [33].

In recent years, efforts have focused in sequencing genes of interest to build genetic linkage maps with direct functional information [34]. Functional genetic linkage maps have experienced a revolution with the availability of new sets of markers from coding regions such as: EST-Ps (Expressed Sequence Tags Polymorphisms), EST-SSRs (EST derived microsatellites) and SNPs (Single Nucleotide Polymorphisms) [35-37]. Functional genetic linkage maps based on annotated genes allow to assess redundant and paralogous EST markers and further improve the quality and utility of genetic maps [38]. Specifically, SNPs have several advantages for their use as molecular makers because they are very abundant in the genome, they show higher stability than SSRs, are usually bi-allelic and codominant [39,40]. Moreover, new technologies have been developed for high throughput detection and genotyping of SNPs reducing the cost of assays [41,42]. Thus, highly saturated genetic linkage maps can be constructed even for species with large and un-sequenced genomes like conifers [21,43-46].

As other pines, *P. pinaster* is a diploid organism characterized by a large and complex genome with high low-copy fraction [47,48]. Particularly, maritime pine has $2n=24$ chromosomes and its genome size is estimated between 51–62 pg/2C [49,50]. Several genetic linkage

maps have been developed for maritime pine based on proteins [51-54], RAPDs [54-57], AFLPs [44,49,54,58], SSRs [44,58-60], EST-Ps [44,58,61] and SNPs [44]. Also, comparative mapping have been performed with *Pinus taeda* L. [44,61]. None of the genetic linkage maps available for *P. pinaster* has been derived from individuals belonging to Spanish populations. These populations show high levels of genetic divergence with the French populations used to design mapping progenies in previous genetic linkage maps [8]. As maritime pine shows a fragmented geographic distribution with high levels of population genetic structure and variation [6,8] it is important to explore the genetic organization in a representative population from the Castilian Plateau (Central Spain) and thus better cover the natural distribution of the species.

Thus, the main objective of this work was to construct saturated genetic linkage maps for *P. pinaster* using controlled crosses between two trees that take part in a breeding program for resin production in a natural population from Central Spain, as a first step to the genetic dissection of this trait. Combining different kind of molecular markers we aim to construct a map with annotated gene functions and homologous markers with previous maps for contributing to the development of a consensus map for the species. A second objective was to identify candidate genes overlapping with QTL already detected in this species [62-64].

Methods

Mapping populations

Two outbred full-sibs families of *P. pinaster* were used for genetic linkage mapping. Progenies were originated from two reciprocal controlled crosses between two progenitors (C14 and C15) belonging to a natural population in Coca (Segovia) located in Central Spain (41° 12' N 4° 31' W). Previous studies on this population have showed a differential genetic structure when compared with other populations of the natural distribution of the species [8]. Progenitors took part in a breeding program for resin production started in 1994 and they were selected for their contrasting resin production, low for C14 and higher for C15. Controlled crosses were carried out in 1999 for C14xC15 and in 2000 for C15xC14. F₁ seeds were collected and germinated in controlled conditions at Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, INIA (Madrid, Spain). Then they were planted in semi-controlled conditions at Dirección Nacional de Biodiversidad– Madrid (40° 27' N 3° 44' W). A paternity test analysis was performed with 13 SSRs. Finally, once the contaminants were removed, the mapping population comprised 161 individuals: 106 from family C14xC15 and 55 individuals from C15xC14.

Molecular markers

Genomic DNA was extracted from needles using a modified protocol from Dellaporta et al. [65] for all marker analyses, but for the 1536 and 384 GoldenGate assays (Illumina Inc., San Diego, CA, USA), for which a commercial *Invisorb DNA plants HTS 96kit* (Invitex GmbH, Berlin, Germany) was used. Four types of molecular marker were used for genotyping the mapping populations: nSSRs, EST-Ps, SAMPLs and SNPs.

nSSRs: Forty seven primer pairs designed for amplification of nSSR loci in *P. pinaster* and *P. taeda* [60,66,67] were tested for segregation in the mapping populations. Thirteen loci were polymorphic, 27 were monomorphic, and seven resulted in multi-banding or non-clear patterns. Amplification of A6F03, A5B01, A5A11, A6F10, A6D04, A5B07 loci was performed as in Guevara et al. [66]. Amplification of NZPR823, NZPR413, NZPR114, NZPR544, SsrPt_ctg64, SsrPt_ctg275 loci was performed as described by Chagné et al. [60] and the amplification of PtTX3116 followed the protocol described by Auckland et al. [68] with modified touchdown profile, using 55°C and 45°C as starting and final temperatures [69]. A Perkin-Elmer GenAmp 9700 thermal cycler (Perkin Elmer Inc., Waltham, Massachusetts, USA) was used to carry out PCR reactions. Amplified products were separated in denaturing gels containing 6% acrylamide / bisacrylamide (19:3), 7 M urea and 1x TBE. Amplified products were visualized in a DNA Analyzer System (4300, LI-COR Biosciences, Lincoln, NE, USA). Fragments were scored visually as codominant markers.

EST-Ps: EST-P genotyping was carried out by Tilling (Targeting Induced Local Lesions in Genomes) as described by Till et al. [70]. This technique allows detection of multiple SNP sites heterozygous in the same progenitor [71]. A set of 14 EST-P primer pairs (PtIFG_893, PtIFG_9136, PtIFG_9034, PtIFG_1955, PtIFG_8429, PtIFG_8702, PtIFG_3C8E, PtIFG_22B8, PtIFG_1CA6C, PtIFG_9044, PtIFG_2253, PtIFG_8436, PtIFG_8887, PtIFG_C6H11) derived from cDNA sequences of *P. taeda* and *P. pinaster* [61,72,73] were tested, in order to identify the most informative markers. A total of 11 EST-P primer pairs generated 25 polymorphic markers. PCRs were performed in 10 µl containing 10 ng of DNA; 1x PCR reaction buffer (Fermentas, Ontario, Canada), 0.2 mM of each dNTP, 2 mM MgSO₄, 0.25U Pfu DNA polymerase (Fermentas, Ontario, Canada), 0.2 µM of each primer (forward primers were labeled on its 5' end with IRDye 700 and reverse primers with IRDye 800). A Perkin-Elmer GenAmp 9700 thermal cycler (Perkin Elmer Inc., Waltham, Massachusetts, USA) was used to carry out PCR reactions. Thermocycler parameters were: 94°C 2 min, 10 touchdown cycles of 94°C 20s, (Tm + 3)°C, 45 s (-0.8°C/cycle), 72°C 1 min;

45 cycles of 94°C 20s, (Tm-5)°C 45 s, 72°C 1 min and final extension step of 72°C for 7 min. Amplification products were visualized on 1% agarose gels to verify amplification. PCR products were digested with CEL I nuclease purified as described by Till et al. [8]. Previously, the concentration of nuclease added, was screened to optimize the detection of heteroduplex between heterozygous sites. Partial DNA digestions were stopped by the addition of 5 µl of 0.5 M EDTA. The mixture were transferred to 96-well Sephadex G50 spin plates (GE Healthcare, Waukesha, WI, USA) for cleaning up by centrifugation into formamide solution and heated at 70°C to reduce the volume to 8 µl. DNA fragments were separated in denaturing gels containing 8% Long Ranger polyacrylamide (Cambrex, East Rutherford, NJ, USA), 7 M urea and 1x TBE. Fragments detection was carried out on a DNA Analyzer System (4300, LI-COR Biosciences, Lincoln, NE, USA). Fragments were scored as dominant markers. Polymorphism was inferred from the resulting fragment pattern and confirmed by sequencing independently undigested amplified products from four haploid megagametophyte DNAs for each progenitor.

SAMPLs: SAMPL genotyping was performed as indicated by Vos et al. [28] with several modifications [74]. Preamplifications were carried out using three primer combinations (*EcoRI* + A/ *MseI* + G; *EcoRI* + A/ *MseI* + C; *EcoRI* + A/ *MseI* + T). For the selective amplification a SAMPL primer [CATA: (CA)₈(TA)₂; GATA: (GA)₈(TA)₂ [75]], was used in combination with an *EcoRI* + 3 primer. In order to select the most informative combinations (those with a higher level of polymorphism) different combinations were tested using template DNA from the parental lines and 9 offspring. Progenitor C14 revealed lower levels of polymorphisms than C15 (see *Results* section), thus primer combinations were chosen in order to equilibrate the number of markers segregating from each progenitor. A total of 31 CATA/*EcoRI* and 26 GATA/*EcoRI* primer combinations were used for the selective amplification. Selective PCR reaction were performed in 10 µl of 1x PCR Buffer (10 mM Tris-HCl, 50 mM KCl, pH 8.3), 0.1 mM of each dNTP, 2.5 mM MgCl₂ (Roche, Basel, Switzerland), 3 ng IRDye 800 5'end labeled CATA or GATA primers, 15 ng *EcoRI* + 3 primer, 0.2U Taq DNA polymerase (Invitrogen, Grand Island, NY, USA) and 5 µl of 10-fold diluted pre-amplification DNA fragments using classical AFLP cycling parameters [12]. Samples were loaded into denaturing gels containing 8% Long Ranger polyacrylamide (Cambrex, East Rutherford, NJ, USA), 7 M urea and 1x TBE. Fragments detection was carried out on a DNA Analyzer System (4300, LI-COR Biosciences, Lincoln, NE, USA). Fragments were scored visually as dominant markers.

SNPs: two SNP genotyping assays were used in this study; a 1,536 BeadArray™ and a 384 BeadXpress®

Golden Gate assays (Illumina Inc., San Diego, CA, USA). SNPs selected for 1,536 Golden Gate assay corresponded to three different sets (see Chancerel et al. [44] for further details): *in vitro* polymorphisms from 35 candidate genes for cell wall formation and drought stress resistance; *in silico* SNPs from a maritime pine EST assembly; and *in silico* polymorphism from re-sequenced amplicons of the species. In this genotyping assay, 95 DNA samples of the mapping progenies were genotyped (73 for C14xC15 and 22 for C15xC14). In order to increase the number of genotyped individuals for a set of genes of interest, another genotyping assay was developed. This genotyping assay (384 SNPlex) consisted in a subsample of SNPs selected from the 1,536 genotyping assay and 14 additional SNPs from candidate genes for drought resistance [76]. It was carried out at Center for Genomic Regulation (CRG, Barcelona, Spain) for a total of 119 DNA samples (79 for C14xC15 and 40 for C15xC14). Both genotyping assays were realized according to the manufacturer's instructions (Illumina Inc., San Diego, CA, USA) and SNPs clusters revised manually with Illumina Bead Studio v2.0 Software. When the same SNP was successfully genotyped in both assays priority was given for the 384 Vera Code data because of the higher number of DNA samples genotyped in this assay. Contig and gene sequences containing the polymorphic SNPs are presented in Additional file 1.

Linkage map construction

For each progenitor we assembled three different linkage maps belonging to datasets of C14xC15 (106 individuals), C15xC14 (55 individuals) and a dataset with the information of the individuals of both reciprocal crosses (161 individuals). Since no relevant differences were found as a consequence of merging both progenies (see *Results* section), further linkage analyses were developed using only the data set with the merged information of both progenies. Parental maps were constructed using the "two-way-pseudo-testcross" mapping strategy [77]. Markers with more than 70% of missing data were excluded from further analysis. Linkage analyses and map estimations were performed using the regression mapping algorithm implemented in the software JoinMap v4.0 [78] with the CP population type and using a recombination fraction < 0.35 and a LOD > 3 as mapping parameters. Map distances were calculated using Kosambi mapping function [79]. When difficulties in estimating marker order are found, two additional maps are constructed (map2 and map3). In map2, new markers are added because more pairwise data are available. In map3, the remaining loci are added by decreasing statistical support. In these cases we kept map2 for further analyses. When a pair of markers was considered identical, only one of the markers was selected for

mapping. In order to assign unlinked loci to selected linkage groups (LG), the strongest cross link was employed with a LOD value of 3 (JoinMap command "assign ungrouped loci to SCL-groups"). Segregation ratios were tested using χ^2 test ($P \leq 0.01$).

Evaluation of homogeneity of recombination rate between female and male meiosis

In order to evaluate whether the male and female gametes presented different levels of recombination, we tested departure from homogeneity of recombination fraction following Plomion et al. [57]. Since the statistical power of homogeneity depends largely on the sample size, the test was performed for all markers pairs in common in the three genetic maps for each progenitor and having a recombination fraction lower than 0.1 (Additional file 2).

Comparative mapping

The linkage maps of both progenitors were compared based on common markers. Besides genetic maps were compared with previously developed *P. pinaster* maps [44] based on common SSRs, ESTPs and SNPs. LGs were named according to Chancerel et al. [44] using loblolly pine nomenclature, as it is the reference pine species.

Genome length and map coverage

Total genome length was calculated as the sum of all mapped marker intervals. Estimated genome length (G_e), was determined from the partial linkage data according to Hulbert et al. [80] modified by Chakravarti et al. [81] (Method 3). A minimum LOD score of three was chosen to estimate genome length using framework maps constructed following the methodology previously described in order to avoid overestimation of genome size because of clustered markers. Observed map coverage was calculated as the ratio of total genome length to estimated genome length [82].

Marker distribution

To evaluate whether markers were randomly distributed, we tested the procedure explained in Echt et al. [38]. A Kolmogorov-Smirnov test for two populations was implemented to compare the observed marker distribution frequencies with expected distribution frequencies under the assumption of randomness. SAMPLs and SNPs distribution were also analyzed by calculating Pearson correlation coefficient between the number of SAMPLs and SNPs in the LGs and the size of the LGs as in Cervera et al. [82].

Heterozygosity levels

The average heterozygosity was estimated for each progenitor and for each molecular marker type independently. Heterozygosity levels based on SSRs were calculated as the ratio between polymorphic and total number of tested SSRs, discarding those with multi-banding and non-clear patterns. Three SSR primer pairs resulted in the amplification of two different loci with clearly different segregation patterns and were scored as different markers, but they were considered as only one for heterozygosity estimations. Heterozygosity levels based on SNPs were calculated as the ratio of polymorphic SNPs and total number of SNPs successfully genotyped. Heterozygosity estimates for SAMPLs were calculated for the first primer combination tested; since the following ones were selected in order to maximize the number of polymorphic markers in C14 (see *Molecular markers* subsection). Heterozygosity based on EST-Ps was not calculated because we only analyzed markers that had been found polymorphic in previous studies in other pine species [61,72,73], therefore a bias could be introduced.

Functional annotation

Functional annotation of the mapped SNP-based genes was carried out using sequence information from the Oligo Pool Assay-OPA (60 nucleotides in length at both sides of the SNP position). In order to obtain homology with longer sequences a BLAST-N search was performed using the pine Gene Index [83] and GeneBank [84]. We retained sequences showing the highest homology (e-value lower than 10^{-20} were considered significant). Then, these longer sequences were annotated using Blast2GO software [85]. Gene Ontology (GO) annotation terms for molecular function at ontology level equal to 3 were placed in the map in order to search for clusters of genes with similar function. For sequences where GO annotation for level 3 was not available we selected the GO annotation terms for level 2. To evaluate whether similar GO terms were clustered or randomly distributed along the genome we performed for each GO term at level 2 a Kolmogorov-Smirnov test for two populations as explained in *Marker Distribution* subsection.

In order to detect interesting co-localizations between candidate genes and QTLs the linkage maps developed in this study were aligned with maps previously constructed for *P. pinaster* containing enough number of orthologous markers to detect homologous LGs and the respective position of QTLs for different traits [44,61,86].

Marker nomenclature

Marker nomenclature for SSRs and SNPs were maintained according to their original publications (see

Molecular Markers subsection). EST-Ps also conserved original nomenclature, but the size of the amplified band was added to the marker name. SAMPLs were named with the differential selective nucleotide used in the preamplification (C, G or T), followed by the targeted microsatellite (CATA or GATA), and the selective *EcoRI*+3 primer employed, ending by the size of the amplified band fragment.

Results and discussion

The paternity test analysis revealed seven contaminants for C14xC15 and three for C15xC14 that were removed for further analyses. The final number of individuals per progeny, 106 for C14xC15 and 55 for C15xC14, was in the limit for reliable estimations and as we did not observe significant differences in recombination fraction between female and male meiosis (see next subsection) we constructed the genetic linkage maps by pooling all individuals of both reciprocal crosses.

Evaluation of homogeneity of recombination between female and male meiosis

Ninety-six marker pairs for C14 and 42 for C15, with a recombination fraction below 0.1, were available in all three maps (C14xC15, C15xC14 and pool map) (see *Methods* section). Eight marker pairs out of the 96 analyzed, showed significant differences between female and male meiosis for C14 (data not shown). Five of them showed a higher recombination rate for male meiosis and three for female meiosis. No marker pair resulted in significant differences in recombination rate for C15. The low level of differences detected in recombination fraction between female and male meiosis supports the merging of both progenies in order to obtain a higher number of offspring in the mapping population and thereby establishes more precise parental maps. No evidence of heterogeneity of recombination was previously reported for *P. pinaster* [56] and other conifer species [87]. However, Plomion and O'Malley [57] suggested that recombination fraction could be higher in male meiosis for *P. pinaster*. The important differences in number of individuals between our mapping progenies (106 versus 55) compelled us to perform the analyses with a narrow window of markers (only those with a recombination fraction lower than 0.1), while in Plomion and O'Malley [57] analyses were performed with a wider window (markers pairs with a recombination fraction lower than 0.3). This could explain the difference in results obtained. Nevertheless, further research in testing homogeneity of recombination between female and male meiosis is needed to clarify whether or not female and male gametes exhibit similar recombination rate, which can have some implications for MAS.

Individual linkage maps and comparative mapping

Previous analysis of the mapping population with four AFLP primer combinations revealed very low levels of polymorphism (data not shown). Therefore, we decided to use SAMPL technique to increase the number of polymorphic fragments. SAMPL analysis was performed using the most informative primer combinations (see *Methods* section). This result validates the use of SAMPLs as an alternative for genotyping low polymorphic populations.

Out of the total set of molecular markers available (Table 1), four and five markers were excluded from C14 and C15 datasets respectively, because of their identical segregation profiles with other markers. All of them were SNPs belonging to the same gene or contig. Markers with more than 70% of missing data were also excluded. Most of them were SAMPLs genotyped only in the C15xC14 pedigree. In addition, in a "two-way-pseudo-test-cross" mapping strategy, intercross markers i.e. markers with the same heterozygous allelic configuration in both progenitors, are less informative. Because of that, several SAMPLs, SNPs and one microsatellite marker were excluded. However, when it was possible we kept a number of intercross markers because they allow to align homologous LGs between both parental maps.

Near 5% of markers used for linkage analysis were unlinked, which is in the same range of what has been observed in other conifer maps [24,45]. Most of them presented more than 35% of missing data and corresponded to SAMPLs genotyped only in the C14xC15 pedigree. Several SNPs were also unlinked. Near 94% of the markers could be assigned to LGs and 60% could be positioned in the final maps (Table 1, Figure 1, Figure 2 and Additional file 3). The lower percentage of markers positioned when compared with other highly saturated maps [46] is due to the use of SAMPLs only scoring in one of the mapping progenies, as revealed by the low percentage of positioned SAMPLs (Table 1). When we discard SAMPLs scored only in one or the two mapping progenies, the percentage of SAMPL markers positioned in the parental maps increases to 65.3% for C14 and 72.1% for C15. These results are very similar to those obtained with positioned SNPs (Table 1) indicating that both type of markers are suitable for the construction of linkage maps. Even more, for a complete coverage of the genome it is interesting to use markers with different target sequences, since coding and non-coding regions seems not to be randomly distributed along the genome [87,88].

In a first phase, before aligning on the reference *P. pinaster* linkage map, 22 LGs were obtained for C14 and 20 for C15 (Table 1, Figure 1 and Figure 2). The smallest LGs were similar in size between C14

and C15 maps. However, the largest LG were higher in C15 than in C14. Also, average size of LGs was slightly higher for C15 than for C14. This was explained because average distance and maximum distance between two adjacent markers was larger in C15 than in C14. Thirty-one intercross markers between both parental maps allowed the identification of homologous LGs (Table 2). Eleven markers with segregation 1:2:1 (same heterozygous combination in both parents) could only be positioned in one parental map (Table 2). Five of them could not be mapped in the other parent because they were ungrouped and the remaining six markers because of the increase in the goodness of fit calculated for the order of markers when were included in the map.

The alignment with maps described by Chancerel et al. [44], based on common SSRs, EST-PS and SNPs (Table 2), made it possible to bring together some LGs resulting into 13 LGs for C14 and 14 LGs for C15 (Table 1), close to the 12 chromosomes of the haploid *P. pinaster* genome [89]. In general, similar size of LGs for parental maps was obtained except for LGs 1, 3 and 10 that were larger in C15 map and LG12 that was larger in C14 map. The fact that we could not assemble the markers in 12 LGs, the differences in size of homologous LGs and the presence of common markers only positioned in one parental map, are probably related with the presence of homozygous regions in the genome of these individuals that prevent mapping markers in these areas. This effect was partially expected because previous studies of the population of origin of both parental trees, Coca, revealed a high coefficient of endogamy [90]. As a result of endogamy we would expect a loss of polymorphisms in the individuals coming from this population, strictly confirmed by the low levels of polymorphism detected by AFLPs genotyping (data not shown) and the low levels of heterozygosity found in the parental trees (Table 3) compared with observed heterozygosity in other provenances of *P. pinaster* [5].

In this respect, it is important to point out to the difference in heterozygosity between C14 and C15 parental trees as revealed by the estimation obtained from SAMPLs (Table 3). This difference was overcome by further genotyping using selected SAMPL primer combinations with a higher number of polymorphic markers in C14 (see *Methods* section). Percentage of heterozygosity calculated from SSRs and SNPs yielded lower values than those obtained from SAMPLs and differences in heterozygosity between C14 and C15 could not be appreciated. One possible explanation is that analyzed SNPs were selected from coding regions where the level of polymorphism is lower than in non coding regions [91].

Table 1 Mapping parameters of parental linkage maps constructed by merging two reciprocal crosses: C14xC15 and C15xC14

Mapping parameter	C14	C15
Total number of available markers	402	410
Number of SSRs loci	11	13
Number of ESTP loci ^a	13	12
Number of SAMPL loci	228	237
Number of SNP loci ^b	150	148
Total number of distorted (p ≤ 0.01) markers	39	52
Number of excluded markers ^c	62	72
Number of SSRs loci	1	1
Number of ESTP loci	0	0
Number of SAMPL loci	46	60
Number of SNP loci	15	11
Number of markers not excluded	340	338
Number of assigned markers ^d	321	319
Number of SSRs loci	9	10
Number of ESTP loci	11	11
Number of SAMPL loci	174	166
Number of SNP loci	127	132
Number of positioned markers ^e	215 (63.2%)	211 (62.4%)
Number of SSR loci	6 (60%)	7 (58.3%)
Number of ESTP loci	10 (76.9%)	7 (58.3%)
Number of SAMPL loci	98 (53.8%)	98 (55.4%)
Number of SNP loci	101 (74.8%)	99 (72.3%)
Number of distorted (p ≤ 0.01) positioned markers	14	14
Unlinked markers (%) ^f	19(6.3%)	19 (5.6%)
Number of LG >3 before making alignments	22	20
Number of LG >3 after making alignments	13	14
Smallest LG (cM) before making alignments	17.5	13.4
Largest LG (cM) before making alignments	81.1	155.3
Average length (cM) LG ± SD before alignments	53.7 ± 20.6	69 ± 35.6
Smallest LG (cM) after making alignments	52	42.3
Largest LG (cM) after making alignments	142.2	155.3
Average length (cM) of a LG ± SD after alignments	90.8 ± 29.14	98.5 ± 38
Maximum distance (cM) between 2 adjacent markers	24.9	35.3
Average distance (cM) between 2 adjacent markers ± SD	6.12 ± 5.8	7.22 ± 6.4

Table 1 Mapping parameters of parental linkage maps constructed by merging two reciprocal crosses: C14xC15 and C15xC14 (Continued)

Observed map length (cM)	1180.4	1379.5
Estimated map length (cM)	1870.2	2166.6
Observed map coverage	63%	64%

^a The 25 ESTP-s correspond to 11 gene loci.

^b The SNPs markers correspond to 47 gene loci and 143 contigs.

^c Markers with more than 70% of missing data (see *Methods* section) and identical markers.

^d Assigned markers correspond to markers linked with more than 2 other markers.

^e Unpositioned markers correspond to markers with a recombination frequency higher than 0.35 with the nearest linked marker (unlinked markers) or markers which position could not be reliably estimated. Percentage of positioned markers was calculated over the number of not excluded markers.

^f Percentage of unlinked markers was calculated over the number of not excluded markers.


SD Standard deviation.

Twenty four contigs with several SNPs (from two to seven) were included in the linkage maps. SNPs belonging to the same gene or contig mapped always in the same position or less than 3 cM away (Figure 1 and Figure 2), except *m682* and *m127* (LG 5), separated by 26.8 cM. Marker *m682* was distorted at the 0.1% significance level, which could affect the accurateness of its position. Alternatively, both SNPs may be associated to different loci at the same LG. The fact that nearly all SNPs belonging to the same contig were mapped in the same position supports the accuracy of the genotyping method used, as previously reported [41,44].

Alignments with the linkage maps developed by Chancerel et al. [44] pointed out that marker order was highly conserved, excepting small inversions of less than 5 cM (data not shown). The only major inconsistency in data was found for marker PtIFG_8436_200, which was amplified using the same primer combination as in Chancerel et al. [44], but subjected to different detection techniques, tilling versus SSCP (Figure 2). This EST-P marker was mapped in LG 10 in our mapping progeny in agreement with previous developed maps in *P. taeda* [72]. However, in other published linkage maps of *P. pinaster* this gene was mapped in LG 7 [44,61]. Chagné et al. [61] discussed the possibility that PtIFG_8436 in *P. pinaster* targeted a paralogous gene as they found low similarity at the DNA sequence level. Our result suggests the existence of an orthologous sequence between *P. pinaster* and *P. taeda* genomes for the region amplified by PtIFG_8436 marker in LG 10 and a paralogous sequence in LG 7 of *P. pinaster* genome.

Segregation distortion

A χ^2 test (d.f. = 1) was performed to test Mendelian segregation of each marker. We detected 9.7% of markers showing distorted segregation ratios at 1% significance level for C14 and 12% for C15 linkage maps (Table 1).



These results are very similar to those obtained in other pine [92,93] and conifer species [36]. The number of distorted markers excluded and unlinked was similar to the number of distorted markers finally assigned to LGs (Additional file 4). Besides, among the unlinked and excluded loci the number of distorted markers was not higher than those showing no segregation distortion (Table 1). Thus, in this case, unlinked and excluded loci seem not to be the result of segregation distortion, as previously reported in other linkage studies [24]. Distorted markers assigned to a LG were randomly distributed (Additional file 4). Only 13 distorted markers could be positioned in each map indicating the difficulty to estimate an accurate position for these distorted markers. Distorted markers positioned in the maps did not appear clustered in specific regions of the genome (Table 1, Figure 1 and Figure 2) suggesting that segregation distortion was probably related with genotyping errors rather than the effect

of pre or post-zygotic selection. As they were not clustered they did not compromise map structure [93].

Marker distribution

Markers were randomly distributed along the genome as no significant differences were found between distribution of markers along the LGs and expected distribution under the hypothesis of randomness (Kolmogorov-Smirnov for two populations, $D = 0.55$, $p\text{-value} = 0.124$ for C14 map and $D = 0.5$, $p\text{-value} = 0.474$ for C15 map), in accordance to other conifer maps [24,32]. Besides, largest LG had more SNPs (Pearson correlation, $r = 0.53$, $p\text{-value} = 0.01$ for C14 map and $r = 0.62$, $p\text{-value} = 0.003$ for C15 map) than smaller LG. Same results were obtained for SAMPLs (Pearson correlation, $r = 0.69$, $p\text{-value} < 0.001$ for C14 map and $r = 0.75$, $p\text{-value} < 0.001$ for C15 map) indicating that they are also randomly distributed along the genome, as expected for this kind of multiband markers [28].




Figure 2 Genetic linkage maps: LGs 7 to 15. Bars on the left represent the LGs obtained for C14 and the bars on the right the LGs obtained for C15. Common markers between both maps are in bold and connected with a solid line. Markers in italics are in common with maps of Chancerel et al. [44] and the homologous LG in this study is indicated with brackets. Markers showing any special feature (see Results section) are underlined. Markers in color are candidate genes that co-localize with QTLs reported in previously published maps for wood properties (green), isotopic composition of C¹³ (violet) and ring growth (blue). SNPs belonging to the same contig are surrounded by a solid line and when they were too far from each other they are connected by a solid line in the left of the chromosome bar. Markers showing significant distorted segregation ratios are indicated with asterisks (** means significant at 0.01 p-value, *** at 0.005, **** at 0.001, ***** at 0.0005 and *****) at 0.0001). Annotations of SNPs are indicated by the term GO and a numeric code. Numeric codes for molecular function annotation level 2: 1 - binding; 2 - catalytic activity; 3 - structural molecule activity; 4 - transporter activity; 5 - enzyme regulator activity. Numeric codes for molecular function level 3: 1.1 - nucleic acid binding; 1.2 - nucleotide binding; 1.3 - protein binding; 1.4 - carbohydrate binding; 1.5 - lipid binding; 2.1 - hydrolase activity; 2.2 - transferase activity; 5.1 - sequence specific DNA binding; 6.1 - signal transducer activity.

Table 2 Markers used for comparative mapping within the species

Marker	C14	C15
Common markers between both parental maps		33
Markers segregating in both parents positioned only in one parental map	5	6
Common markers with Chancerel et al. [44]	65	57
Common SSR loci	2	3
Common ESTP loci	7	3
Common SNP loci	56	51
Number of LGs without common markers with Chancerel et al. [44]	1	2

Genome length and map coverage

Observed genome length ranged from 1,180.4 (C14) to 1,379.5 cM (C15), 200 cM larger for C15 map than for C14 map (Table 1). In other *P. pinaster* maps observed genome length ranged from 869 to 1,860 cM depending on the density of markers [44,55,58]. The higher genome length observed in C15 map agrees with its higher heterozygosity estimation compared to C14 map (Table 3). Estimated genome length ranged from 1,870.2 to 2,166.6 cM, in line with what has been obtained in previous *P. pinaster* maps (1,223 to 3,252 cM depending on the method of estimation [44,56,88]). The last generation maps estimated *P. pinaster* genome size to be 2,500 cM [44], a value which is near our estimates and close to other pine species [94]. Estimated genome length was higher for C15

Table 3 Heterozygosity

Marker	C14			C15		
	Poly.	Mono.	Heterozygosity (%)	Poly.	Mono.	Heterozygosity (%)
SSR	9	31	0.23	10	30	0.25
SAMPL	133	191	0.41	251	191	0.57
SNP	150	672	0.18	96	726	0.12

Percentage of heterozygosity calculated as the ratio of polymorphic markers segregating in each parental map over total (polymorphic -Poly.- and monomorphic-Mono.-) markers.

Heterozygosity estimates for SAMPLs were calculated for the first primer combination tested in the family.

linkage map although the observed map coverage (near 65%, Table 1) was similar for both parental maps. High density genetic linkage maps usually report map coverage over 90% [46,95]. However, previously published *P. pinaster* genetic linkage maps also reported map coverage near 65% [44,57], indicating the difficulty to achieve a complete coverage for such a large and complex genome.

Functional annotation

We validated and improved the functional annotation information for mapped SNPs. Significant sequence homology was found in pine Gene Index database for 160 out of the 171 mapped SNPs in both parental linkage maps [83] (Additional file 5). Sequence homology was found for several species with the top hit homologies for *Picea sitchensis*, *Pinus taeda*, *Pinus radiata*, *Vitis vinifera* and *Picea glauca* (Additional file 5). Nine sequences over the 171 sequences showed no match with the InterPro database [96] and we did not find any GO term for seven sequences. Thus, a total of 144 sequences were annotated, 132 of them for molecular function and 101 sequences with annotation for molecular function levels 2 or 3. Most of the mapped

SNPs were associated to cDNA belonging to the GO terms: *binding*, *catalytic activity* and *hydrolase activity* (Additional file 6). As expected, SNPs belonging to the same contig reported identical GO annotation terms. However, our results could not confirm statistically (Kolmogorov-Smirnov test for two populations not significant, data not shown) if neighboring SNPs belonging to different genes or contigs exhibited GO terms for the same molecular function. Denser genetic maps with deeper functional annotation are required to evaluate if genes with similar functions are clustered or not.

The comparison of our annotated linkage maps and linkage maps reporting QTL information revealed candidate genes for several QTLs for wood properties or isotopic composition of C¹³ (δC^{13}) [61,62,64]. δC^{13} is a character closely related with water use efficiency [97]. In our study, SNPs annotated for water-stress inducible proteins, AQUAPORINs and DEHYDRINs were positioned in the same region as QTLs for δC^{13} [62] (Table 4). This outcome reinforces the hypothesis that the genomic regions identified by QTL analysis [62] might play a key role in the genetic control of water use efficiency. Also SNPs associated with a *CELLULOSE SYNTHASE CESA3*, a *PEROXIDASE* (enzyme involved

Table 4 Co-localizations of SNPs and QTLs

SNP ID	Sequence description	e-value	LG	Trait QTL	Reference
m157	CELLULOSE SYNTHASE	1.5e ⁻⁶⁸	3	wood	Pot et al. [64]
m264	PEROXIDASE	2.9e ⁻¹⁴⁷	8	wood	
m941	ENDO- BETA-XYLANASE A-LIKE	0	4	wood	
m1542	WATER-STRESS INDUCIBLE PROTEIN 1	9.5e ⁻²⁶	2	δC^{13}	Brendel et al. [62]
m1543	WATER-STRESS INDUCIBLE PROTEIN 3	7.3e ⁻³⁴	2	δC^{13}	
m426	WATER DEFICIT INDUCIBLE LP3-LIKE PROTEIN	3.2e ⁻⁶²	3	δC^{13}	
m295	AQUAPORIN	9.4e ⁻³	5	δC^{13}	
m965	THYLAKOID LUMENAL 19 KDA PROTEIN	7.1e ⁻¹¹⁹	5	ring growth	
m712	DEHYDRIN 9 PROTEIN	1.2e ⁻⁷¹	8	δC^{13}	
m716	DEHYDRIN 2	1.9e ⁻⁴³	8	δC^{13}	
m859	THYLAKOID LUMENAL PROTEIN	9.8e ⁻¹⁴⁵	12	δC^{13}	

Co-localization of mapped SNPs with QTLs detected in previously published maps of *P. pinaster*. δC^{13} stands for isotopic composition of C¹³. *Wood* stands for wood chemical composition and fibre properties.

in lignin polymerization [98]) and a *ENDO-1,4BETA-XYLANASE A-LIKE* gene (Additional file 5) co-localized with QTLs for wood chemical composition and fiber properties [64] (Table 4). This result increases the evidence of function assigned to these genes and has special relevance when we consider that orthologous QTLs for wood properties were also found in other *Pinus* species [61]. This finding highlights the importance of developing functional genetic linkage maps to be used as useful tools to look for favorable allelic variants to be implemented in MAS.

Conclusions

Our study demonstrates the importance of developing genetic linkage maps from different populations representing different genetic backgrounds in order to generate an accurate consensus linkage map of the same species. Comparative mapping is a key process to facilitate the understanding of genome organization and evolution in conifers. For that purpose it is essential to correctly identify orthologous versus paralogous genes. New efforts in detecting orthologous markers as well as progress in sequencing conifer genomes will improve comparative mapping studies in the future. Here we also confirm the importance of developing functional genetic linkage maps, especially when working with breeding populations for its future application in MAS for traits of interest.

Additional files

Additional file 1: Sequence of contigs and genes containing the analyzed SNPs. Nucleotide sequences in fasta format of contigs and genes containing the polymorphic SNPs identified in the mapping progenies. All sequences are the consensus sequence obtained from different individuals except for genes *LP3-1* and *LP3-3* which sequences obtained for one individual.

Additional file 2: Standard error of recombination frequency. Representation of the standard error of recombination frequency for markers segregating 1:1 in the F_1 mapping populations C14xC15 (N=106) and C15xC14 (N=55). Standard error calculated following Ritter et al. [99].

Additional file 3: Polymorphic SNPs. List of the polymorphic SNPs in mapping progenies and associated features.

Additional file 4: Distorted markers. Number of total (assigned, ungrouped and excluded) markers showing distortion in their segregation rates at $p \leq 0.01$ per LG obtained for C14 and C15 linkage maps constructed by merging two reciprocal crosses: C14xC15 and C15xC14.

Additional file 5: BLAST analysis of SNP-sequences. Results of BLAST-N search in pine databases of the Gene Index Project and GeneBank for mapped SNPs.

Additional file 6: GO annotation for mapped SNPs. Sequence distribution of GO terms for molecular function.

Authors' contributions

MdM: Analyzed the quality of SNP genotyping results, constructed the linkage maps and wrote the first draft of the manuscript. NdM, MAG and LD: Carried out SSRs, EST-PS and SAMPL analyses. DSG: Handled and managed plant material. EC and CP: helped MdM in analyzing SNPs from de 1,536 SNP array. ESL: Carried out DNA extraction. CC: Performed functional annotation. JAC: Participated in the construction of linkage maps. MTC, CP and IA: Designed experiments and coordinated projects. All authors contributed to writing the article and approved the final manuscript.

Acknowledgements

MdM was supported by an FPI-INIA PhD fellowship granted by the "Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria". ES was supported by an FPI PhD fellowship granted by the "Ministerio de Ciencia e Innovación" (MICINN). DS-G was supported by the "Juan de la Cierva" program granted by MICINN. This work was supported by funding from the Spanish MICINN (AGL2009-10496, MAPINSEQ), the transnational Plant-KBBE program (PLE2009-0016, SUSTAINPINE), and the European Commission (FP7-211868, NOVELTREE). The authors are very grateful to S. González-Martínez for designing the 384 SNPlex and discussing quality of SNP genotyping results; Ricardo Alía and Salustiano Iglesias for providing seeds and land resources to install the mapping progenies, respectively. We also would like to thank two anonymous reviewers for their helpful comments that have improved the manuscript.

Author details

¹INIA-CIFOR, Departamento de Ecología y Genética Forestal, Carretera de la Coruña Km7.5, Madrid 28040, Spain. ²Unidad Mixta de Genómica y Ecofisiología Forestal, INIA/UPM, Madrid, Spain. ³ETSIM, Departamento de Biotecnología, Ciudad Universitaria s/n, Madrid 28040, Spain. ⁴INRA, UMR1202 BIOGECO, Cestas F-33610, France. ⁵Université de Bordeaux, UMR1202 BIOGECO, Talence 33170, France.

Received: 9 June 2012 Accepted: 26 September 2012

Published: 4 October 2012

References

- Blanco E, Casado MA, Costa M, Escribano R, García-Anton M, Génova M, Gómez-Manzanque A, Gómez-Manzanque F, Moreno JC, Morla C, Regato P, Sainz H: *Los bosques ibéricos. Una interpretación geobotánica*. 4th edition. Barcelona: Planeta; 2005.
- Fernández M, Gil L, Pardos JA: Effects of water supply on gas exchange in *Pinus pinaster* Ait. provenances during their first growing season. *Ann For Sci* 2000, **57**:9-16.
- Sánchez-Gómez D, Majada J, Alía R, Feito I, Aranda I: Intraspecific variation in growth and allocation patterns in seedlings of *Pinus pinaster* Ait. submitted to contrasting watering regimes: can water availability explain regional variation? *Ann For Sci* 2010, **67**:505.
- Aranda I, Alía R, Ortega U, Dantas AK, Majada J: Intra-specific variability in biomass partitioning and carbon isotopic discrimination under moderate drought stress in seedlings from four *Pinus pinaster* populations. *Tree Genet Genomes* 2010, **6**:169-178.
- Mariette S, Chagné D, Lézier C, Pastuszka P, Raffin A, Plomion C, Kremer A: Genetic diversity within and among *Pinus pinaster* populations: comparison between AFLP and microsatellite markers. *Heredity* 2001, **86**:469-479.
- Ribeiro M, Mariette S, Vendramin G, Szmidt A, Plomion C, Kremer A: Comparison of genetic diversity estimates within and among populations of maritime pine using chloroplast simple-sequence repeat and amplified fragment length polymorphism data. *Mol Ecol* 2002, **11**:869-877.
- González-Martínez SC, Mariette S, Ribeiro MM, Burban C, Raffin A, Chambel MR, Ribeiro CAM, Aguiar A, Plomion C, Alía R, Gil L, Vendramin GG, Kremer A: Genetic resources in maritime pine (*Pinus pinaster* Aiton): molecular and quantitative measures of genetic variation and differentiation among maternal lineages. *For Ecol Manag* 2004, **197**:103-115.
- Eveno E, Collada C, Guevara MA, Léger V, Soto A, Díaz L, Léger P, González-Martínez SC, Cervera MT, Plomion C, Garnier-Géré P: Contrasting patterns of selection at *Pinus pinaster* Ait. drought stress candidate genes as revealed by genetic differentiation analyses. *Mol Biol Evol* 2008, **25**:417-437.

Competing interests

The authors declare that they have no competing interests.

9. Vendramin G, Anzidei M, Madaghiale A, Bucci G: **Distribution of genetic diversity in *Pinus pinaster* Ait. as revealed by chloroplast microsatellites.** *Theor Appl Genet* 1998, **97**:456–463.
10. Bucci G, González-Martínez SC, Le Provost G, Plomion C, Ribeiro MM, Sebastiani F, Alía R, Vendramin GG: **Range-wide phylogeography and gene zones in *Pinus pinaster* Ait. revealed by chloroplast microsatellite markers.** *Molec Ecol* 2007, **16**:2137–2153.
11. González-Martínez SC, Alía R, Gil L: **Population genetic structure in a Mediterranean pine (*Pinus pinaster* Ait.): a comparison of allozyme markers and quantitative traits.** *Heredity* 2002, **89**:199–206.
12. Farjon A: *A natural history of Conifers*. Portland: Timber Press, Inc.; 2008.
13. Tadesse W, Nanos N, Aunon F, Arrabal C, García C, Gil L, Alía R, Pardos J: **Genetic improvement of resin yield from maritime pine in Spain.** *Forest Chemicals Review* 2001, **11**:11.
14. Nanos N, Tadesse W, Montero G, Gil L, Alía R: **Modelling resin production distributions for *Pinus Pinaster* Ait using two probability functions.** *Ann For Sci* 2000, **57**:369–377.
15. Mergen F, Hoekstra P, Echols RM: **Genetic control of oleoresin yield and viscosity in Slash Pine.** *For Sci* 1955, **1**:19–30.
16. White TL, Adams WT, Neale D: **Tree Improvement Programs-Structure, Concepts and Importance.** In *Forest Genetics*. Edited by White TL, Adams WT, Neale D. Oxfordshire: CABI Publishing; 2007:285–302.
17. Arrabal C, García-Vallejo MC, Cadahia E, Cortijo M, de Simón BF: **Characterization of two chemotypes of *Pinus pinaster* by their terpene and acid patterns in needles.** *Plant Syst Evol* 2012, **298**:511–522.
18. Tadesse W, Nanos N, Alía R, Aunon F, Gil L: **Evaluation of high resin yielders of *Pinus pinaster* Ait.** *International Journal of Forest Genetics* 2001, **8**:271–278.
19. Arrabal C, Cortijo M, De Simon BF, García Vallejo MC, Cadahia E: **Differentiation among five Spanish *Pinus pinaster* provenances based on its oleoresin terpenic composition.** *Biochem Syst Ecol* 2005, **33**:1007–1016.
20. Gerber S, Lascoux M, Kremer A: **Relation between protein markers and quantitative traits in maritime pine (*Pinus pinaster* AIT.).** *Silvae Genet* 1997, **46**:286–291.
21. Pelgas B, Bousquet J, Meirmans P, Ritland K, Isabel N: **QTL mapping in white spruce: gene maps and genomic regions underlying adaptive traits across pedigrees, years and environments.** *BMC Genomics* 2011, **12**:145.
22. Devey ME, Carson SD, Nolan ME, Matheson AC, Te Riini C, Hohepa J: **QTL associations for density and diameter in *Pinus radiata* and the potential for marker-aided selection.** *Theor Appl Genet* 2004, **108**:516–524.
23. Plomion C, Yani A, Marpeau A: **Genetic determinism of delta 3-carene in maritime pine using RAPD markers.** *Genome* 1996, **39**:1123–1127.
24. Pelgas B, Bousquet J, Beauseigle S, Isabel N: **A composite linkage map from two crosses for the species complex *Picea marianax Picea rubens* and analysis of synteny with other Pinaceae.** *Theor Appl Genet* 2005, **111**:1466–1488.
25. Krutovsky KV, Troggio M, Brown GR, Jermstad KD, Neale DB: **Comparative mapping in the Pinaceae.** *Genetics* 2004, **168**:447–461.
26. Cervera MT, Plomion C, Malpica C: **Molecular markers and genome mapping in woody plants.** In *Molecular Biology of Woody Plants*. 1st edition. Edited by Jain SM, Minocha SC. Dordrecht: Kluwer Academic Publishers; 2000:375–394.
27. Devey ME, Sewell MM, Uren TL, Neale DB: **Comparative mapping in loblolly and radiata pine using RFLP and microsatellite markers.** *Theor Appl Genet* 1999, **99**:656–662.
28. Vos P, Hogers R, Bleeker M, Reijans M, Lee T, Hornes M, Friters A, Pot J, Paleman J, Kuiper M: **AFLP: a new technique for DNA fingerprinting.** *Nucleic Acids Res* 1995, **23**:4407.
29. Williams JGK, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV: **DNA polymorphisms amplified by arbitrary primers are useful as genetic markers.** *Nucleic Acids Res* 1990, **18**:6531–6535.
30. Gosselin I, Zhou Y, Bousquet J, Isabel N: **Megagametophyte-derived linkage maps of white spruce (*Picea glauca*) based on RAPD, SCAR and ESTP markers.** *Theor Appl Genet* 2002, **104**:987–997.
31. Ritland K, Krutovsky KV, Tsumura Y, Pelgas B, Isabel N, Bousquet J: **Genetic mapping in conifers.** In *Genetics, Genomics and Breeding of Conifers*. Edited by Plomion C, Bousquet J. Edenbridge: Science Publishers; 2011:196–238.
32. Remington D, Whetten R, Liu BH, O'malley D: **Construction of an AFLP genetic map with nearly complete genome coverage in *Pinus taeda*.** *Theor Appl Genet* 1999, **98**:1279–1292.
33. Paglia G, Morgante M: **PCR-based multiplex DNA fingerprinting techniques for the analysis of conifer genomes.** *Mol Breed* 1998, **4**:173–177.
34. González-Martínez SC, Krutovsky KV, Neale DB: **Forest-tree population genomics and adaptive evolution.** *New Phytol* 2006, **170**:227–238.
35. Komulainen P, Brown GR, Mikkonen M, Karhu A, García-Gil MR, O'Malley D, Lee B, Neale DB, Savolainen O: **Comparing EST-based genetic maps between *Pinus sylvestris* and *Pinus taeda*.** *Theor Appl Genet* 2003, **107**:667–678.
36. Tani N, Takahashi T, Iwata H, Mukai Y, Ujino-Ihara T, Matsumoto A, Yoshimura K, Yoshimaru H, Murai M, Nagasaka K, Tsumura Y: **A consensus linkage map for Sugi (*Cryptomeria japonica*) from two pedigrees, based on microsatellites and Expressed Sequence Tags.** *Genetics* 2003, **165**:1551–1568.
37. Plomion C, Hurme P, Frigerio J, Ridolfi M, Pot D, Pionneau C, Avila C, Gallardo F, David H, Neutelings G: **Developing SSCP markers in two *Pinus* species.** *Mol Breed* 1999, **5**:21–31.
38. Echt C, Saha S, Krutovsky K, Wimalanathan K, Erpelding J, Liang C: **An annotated genetic map of loblolly pine based on microsatellite and cDNA markers.** *BMC Genet* 2011, **12**:17.
39. Neale DB: **Genomics to tree breeding and forest health.** *Curr Opin Genet Dev* 2007, **17**:539–544.
40. Cho RJ, Mindrinos M, Richards DR, Sapolsky RJ, Anderson M, Drenkard E, Dewdney J, Reuber TL, Stammers M, Federspiel N: **Genome-wide mapping with biallelic markers in *Arabidopsis thaliana*.** *Nat Genet* 1999, **23**:203–207.
41. Lepointevin C, Frigerio JM, Garnier-Géré P, Salin F, Cervera MT, Vornam B, Harvengt L, Plomion C: **In Vitro vs In Silico Detected SNPs for the Development of a Genotyping Array: What Can We Learn from a Non-Model Species?** *PLoS ONE* 2010, **5**:e11034.
42. Pavy N, Parsons L, Paule C, MacKay J, Bousquet J: **Automated SNP detection from a large collection of white spruce expressed sequences: contributing factors and approaches for the categorization of SNPs.** *BMC Genomics* 2006, **7**:174.
43. Eckert AJ, Pande B, Ersoz ES, Wright MH, Rashbrook VK, Nicolet CM, Neale DB: **High-throughput genotyping and mapping of single nucleotide polymorphisms in loblolly pine (*Pinus taeda* L.).** *Tree Genet Genomes* 2009, **5**:225–234.
44. Chancerel E, Lepointevin C, Le Provost G, Lin Y-C, Jaramillo-Correa J, Eckert A, Wegryn J, Zelenika D, Boland A, Frigerio J-M, Chaumeil P, Garnier-Gere P, Boury C, Grivet D, Gonzalez-Martinez S, Rouze P, Van de Peer Y, Neale D, Cervera M, Kremer A, Plomion C: **Development and implementation of a highly-multiplexed SNP array for genetic mapping in maritime pine and comparative mapping with loblolly pine.** *BMC Genomics* 2011, **12**:368.
45. Pavy N, Pelgas B, Beauseigle S, Blais S, Gagnon F, Gosselin I, Lamothe M, Isabel N, Bousquet J: **Enhancing genetic mapping of complex genomes through the design of highly-multiplexed SNP arrays: application to the large and unsequenced genomes of white spruce and black spruce.** *BMC Genomics* 2008, **9**:21.
46. Moriguchi Y, Ujino-Ihara T, Futamura N, Saito M, Ueno S, Matsumoto A, Tani N, Taira H, Shinohara K, Tsumura Y: **The construction of a high-density linkage map for identifying SNP markers that are tightly linked to a nuclear-recessive major gene for male sterility in *Cryptomeria japonica* D. Don.** *BMC Genomics* 2012, **13**:95.
47. Guevara MA, Cervera MT, Soto A, Collada C, Plomion C, Savolainen O, Neale DB, González-Martínez SC: **Genomics applied to the study of adaptation in pine species.** *Investigacion Agraria: Sistemas y Recursos Forestales* 2005, **14**:292–306.
48. Elisk C, Williams C: **Retroelements contribute to the excess low-copy-number DNA in pine.** *Mol Gen Genet* 2000, **264**:47–55.
49. Chagné D, Lalanne C, Madur D, Kumar S, Frigério JM, Krier C, Decroocq S, Savouré A, Bou-Dagher-Kharrat M, Bertocchi E, Brach J, Plomion C: **A high density genetic map of maritime pine based on AFLPs.** *Ann For Sci* 2002, **59**:627–636.
50. Grotkopp E, Rejmánek M, Sanderson MJ, Rost TL: **Evolution of genome size in pines (*Pinus*) and its life-history correlates: supertree analyses.** *Evolution* 2004, **58**:1705–1729.
51. Bahrman N, Damerval C: **Linkage relationships of loci controlling protein amounts in maritime pine (*Pinus pinaster* Ait).** *Heredity* 1989, **63**:267–274.

52. Gerber S, Rodolphe F, Bahrman N, Baradat P: **Seed-protein variation in maritime pine (*Pinus pinaster* Ait) revealed by two-dimensional electrophoresis: genetic determinism and construction of a linkage map.** *Theor Appl Genet* 1993, **85**:521–528.
53. Plomion C, Costa P, Bahrman N: **Genetic analysis of needle protein in Maritime pine. Mapping dominant and codominant protein markers assayed on diploid tissue, in a haploid-base genetic map.** *Silvae Genet* 1997, **46**:161–165.
54. Costa P, Pot D, Dubos C, Frigerio JM, Pionneau C, Bodenès C, Bertocchi E, Cervera M-T, Remington DL, Plomion C: **A genetic map of Maritime pine based on AFLP, RAPD and protein markers.** *Theor Appl Genet* 2000, **100**:39–48.
55. Plomion C, Bahrman N, Durel CE, O'Malley DM: **Genomic mapping in *Pinus pinaster* (maritime pine) using RAPD and protein markers.** *Heredity* 1995, **74**:661–668.
56. Plomion C, O'Malley D, Durel C-E: **Genomic analysis in maritime pine (*Pinus pinaster*) comparison of two RAPD maps using selfed and open-pollinated seeds of the same individual.** *Theor Appl Genet* 1995, **90**:1028–1034.
57. Plomion C, O'Malley D: **Recombination rate differences for pollen parents and seed parents in pine.** *Heredity* 1996, **77**:341–350.
58. Ritter E, Aragones A, Markussen T, Achere V, Espinel S, Fladung M, Wrobel S, Faivre-Rampant P, Jeandroz S, Favre JM: **Towards construction of an ultra high density linkage map for *Pinus pinaster*.** *Annals of Forest Sciences* 2002, **59**:637–643.
59. Mariette S, Chagné D, Decroocq S, Vendramin GG, Lalanne C, Madur D, Plomion C: **Microsatellite markers for *Pinus pinaster* Ait.** *Ann For Sci* 2001, **58**:203–206.
60. Chagné D, Chaumeil P, Ramboer A, Collada C, Guevara MA, Cervera MT, Vendramin GG, García V, Frigerio J-M, Echt C, Richardson T, Plomion C: **Cross-species transferability and mapping of genomic and cDNA SSRs in pines.** *Theor Appl Genet* 2004, **109**:1204–1214.
61. Chagné D, Brown GR, Lalanne C, Madur D, Pot D, Neale D, Plomion C: **Comparative genome and QTL mapping between maritime and loblolly pines.** *Mol Breed* 2003, **12**:185–195.
62. Brendel O, Pot D, Plomion C, Rozenberg P, Guehl JM: **Genetic parameters and QTL analysis of delta C-13 and ring width in maritime pine.** *Plant Cell Environ* 2002, **25**:945–953.
63. Markussen T, Fladung M, Achere V, Favre JM, Faivre-Rampant P, Aragones A, DA Silva Pérez D, Havengt L, Ritter E: **Identification of QTLs controlling growth, chemical and physical wood property traits in *Pinus pinaster*, Ait.** *Silvae Genet* 2003, **52**:8–15.
64. Pot D, Rodrigues J, Rozenberg P, Chantre G, Tibbits J, Cahalan C, Pichavant F, Plomion C: **QTLs and candidate genes for wood properties in maritime pine (*Pinus pinaster* Ait.).** *Tree Genet Genomes* 2006, **2**:10–24.
65. Dellaporta SL, Wood J, Hicks JB: **A plant DNA miniprep: version II.** *Plant Mol Biol Report* 1983, **1**:19–21.
66. Guevara MA, Chagné D, Almeida H, Byrnes M, Collada C, Favre JM, Havengt L, Jeandroz S, Orazio C, Plomion C, Ramboer A, Rocheta M, Sebastiani F, Soto A, Vendramin GG, Cervera MT: **Isolation and characterization of nuclear microsatellite loci in *Pinus pinaster* Ait.** *Mol Ecol Notes* 2005, **5**:57–59.
67. Elisk CG, Williams CG: **Low copy microsatellite recovery from a conifer genome.** *Theor Appl Genet* 2001, **103**:1189–1195.
68. Auckland L: **Conifer microsatellite handbook.** Raleigh: Texas A & M University; 2002.
69. González-Martínez SC, Robledo-Arnuncio JJ, Collada C, Díaz A, Williams CG, Alía R, Cervera MT: **Cross-amplification and sequence variation of microsatellite loci in Eurasian hard pines.** *Theor Appl Genet* 2004, **109**:103–111.
70. Till BJ, Reynolds SH, Greene EA, Codomo CA, Enns LC, Johnson JE, Burtner C, Odden AR, Young K, Taylor NE, Henikoff JG, Comai L, Henikoff S: **Large-Scale Discovery of Induced Point Mutations With High-Throughput TILLING.** *Genome Res* 2003, **13**:524–530.
71. Rungis D, Hamberger B, Bérubé Y, Wilkin J, Bohlmann J, Ritland K: **Efficient genetic mapping of single nucleotide polymorphisms based upon DNA mismatch digestion.** *Mol Breed* 2005, **16**:261–270.
72. Brown G, Kadel E III, Bassoni D, Kiehne K, Temesgen B, Van Buijtenen J, Sewell M, Marshall K, Neale D: **Anchored reference loci in loblolly pine (*Pinus taeda* L.) for integrating pine genomics.** *Genetics* 2001, **159**:799–809.
73. Temesgen B, Brown GR, Harry DE, Kinlaw CS, Sewell MM, Neale DB: **Genetic mapping of expressed sequence tag polymorphism (ESTP) markers in loblolly pine (*Pinus taeda* L.).** *Theor Appl Genet* 2001, **102**:664–675.
74. Cervera M, Gusmao J, Steenackers M, Peleman J, Storme V, Vanden Broeck A, Van Montagu M, Boerjan W: **Identification of AFLP molecular markers for resistance against *Melampsora larici-populina* in *Populus*.** *Theor Appl Genet* 1996, **93**:733–737.
75. Schmidt A, Doudrick R, Heslop-Harrison J, Schmidt T: **The contribution of short repeats of low sequence complexity to large conifer genomes.** *Theor Appl Genet* 2000, **101**:7–14.
76. Grivet D, Sebastiani F, Alía R, Bataillon T, Torre S, Zabal-Aguirre M, Vendramin GG, González-Martínez SC: **Molecular footprints of local adaptation in two Mediterranean conifers.** *Mol Biol Evol* 2011, **28**:101.
77. Grattapaglia D, Sederoff R: **Genetic linkage maps of *Eucalyptus grandis* and *Eucalyptus urophylla* using a pseudo-testcross mapping strategy and RAPD markers.** *Genetics* 1994, **137**:1121–1137.
78. Van Ooijen JW: **In Joinmap 4, software for the calculation of genetic maps in experimental populations.** Edited by Kiazma BV. Wageningen; 2006.
79. Kosambi D: **The estimation of map distances from recombination values.** *Annals Eugen* 1944, **12**:172–175.
80. Hulbert S, Iltott T, Legg E, Lincoln S, Lander E, Michelmore R: **Genetic analysis of the fungus, *Bremia lactucae*, using restriction fragment length polymorphisms.** *Genetics* 1988, **120**:947.
81. Chakravarti A, Lasher LK, Reefer JE: **A maximum likelihood method for estimating genome length using genetic linkage data.** *Genetics* 1991, **128**:175–182.
82. Cervera M-T, Storme V, Ivens B, Gusmao J, Liu BH, Hostyn V, Van Slycken J, Van Montagu M, Boerjan W: **Dense Genetic Linkage Maps of Three *Populus* Species (*Populus deltoides*, *P. nigra* and *P. trichocarpa*) Based on AFLP and Microsatellite Markers.** *Genetics* 2001, **158**:787–809.
83. **The Gene Index Project.** <http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/Blast/index.cgi>.
84. **Gene Bank.** <http://www.ncbi.nlm.nih.gov/genbank/>.
85. Conesa A, Götts S, García-Gómez JM, Terol J, Talon M, Robles M: **Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research.** *Bioinformatics* 2005, **21**:3674–3676.
86. Pot D, Chantre G, Rozenberg P, Rodrigues JC, Jones GL, Pereira H, Hannrup B, Cahalan C, Plomion C: **Genetic control of pulp and timber properties in maritime pine (*Pinus pinaster* Ait.).** *Ann For Sci* 2002, **59**:563–575.
87. Scotti I, Burelli A, Cattonaro F, Chagné D, Fuller J, Hedley PE, Jansson G, Lalanne C, Madur D, Neale D, Plomion C, Powell W, Troglio M, Morgante M: **Analysis of the distribution of marker classes in a genetic linkage map: a case study in Norway spruce (*Picea abies* Karst).** *Tree Genet Genomes* 2005, **1**:93–102.
88. Gerber S, Rodolphe F: **An estimation of the genome length of maritime pine (*Pinus pinaster* Ait.).** *Theor Appl Genet* 1994, **88**:289–292.
89. Sax K, Sax HJ: **Chromosome number and morphology in the conifers.** *J Arnold Arbor* 1933, **14**:356–375.
90. González-Martínez S: **Estructura poblacional y flujo genético de *Pinus pinaster* Aiton en el noroeste de la Península Ibérica.** PhD Thesis. Universidad Politécnica de Madrid, ETSIM; 2001.
91. Nordborg M, Hu TT, Ishino Y, Jhaveri J, Toomajian C, Zheng H, Bakker E, Calabrese P, Gladstone J, Goyal R, Jakobsson M, Kim S, Morozov Y, Padhukasahasram B, Plagnol V, Rosenberg NA, Shah C, Wall JD, Wang J, Zhao K, Kalbfleisch T, Schulz V, Kreitman M, Bergelson J: **The pattern of polymorphism in *Arabidopsis thaliana*.** *PLoS Biol* 2005, **3**: e196.
92. Yin TM, Wang XR, Andersson B, Lerceteau-Köhler E: **Nearly complete genetic maps of *Pinus sylvestris* L. (Scots pine) constructed by AFLP marker analysis in a full-sib family.** *Theor Appl Genet* 2003, **106**:1075–1083.
93. Kubisiak TL, Nelson CD, Nance W, Stine M: **RAPD linkage mapping in a longleaf pine x slash pine F1 family.** *Theor Appl Genet* 1995, **90**:1119–1127.
94. Echt C, Nelson C: **Linkage mapping and genome length in eastern white pine (*Pinus strobus* L.).** *Theor Appl Genet* 1997, **94**:1031–1037.
95. Yin T, DiFazio SP, Gunter LE, Riemenschneider D, Tuskan GA: **Large-scale heterospecific segregation distortion in *Populus* revealed by a dense genetic map.** *Theor Appl Genet* 2004, **109**:451–463.
96. **InterPro protein sequence analysis & classification.** <http://www.ebi.ac.uk/interpro/>.

97. Farquhar GD, Richards RA: **Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes.** *Aust J Plant Physiol* 1984, **11**:539–552.
98. Boerjan W, Ralph J, Baucher M: **Lignin biosynthesis.** *Annu Rev Plant Biol* 2003, **54**:519–546.
99. Ritter E, Gebhardt C, Salamini F: **Estimation of recombination frequencies and construction of RFLP linkage maps in Plants from crosses between heterozygous parents.** *Genetics* 1990, **125**:645–654.

doi:10.1186/1471-2164-13-527

Cite this article as: de Miguel *et al.*: Annotated genetic linkage maps of *Pinus pinaster* Ait. from a Central Spain population using microsatellite and gene based markers. *BMC Genomics* 2012 **13**:527.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

