



HAL
open science

Competitive interactions in plant-parasitic nematode communities affecting organic vegetable cropping systems

Thierry T. Mateille, Johannes Tavoillot, Claire Goillon, Laure Pares, Amélie Lefevre, Hélène Védie, Mireille Navarrete, Alain Palloix, Anne-Marie Sage-Palloix, Philippe Castagnone-Sereno, et al.

► To cite this version:

Thierry T. Mateille, Johannes Tavoillot, Claire Goillon, Laure Pares, Amélie Lefevre, et al.. Competitive interactions in plant-parasitic nematode communities affecting organic vegetable cropping systems. *Crop Protection*, 2020, 135, pp.105206. 10.1016/j.cropro.2020.105206 . hal-02649096

HAL Id: hal-02649096

<https://hal.inrae.fr/hal-02649096>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Title**

2 Competitive interactions in plant-parasitic nematode communities affecting organic vegetable
3 cropping systems.

4
5 **Authors**

6 Thierry Mateille ^{a*}, Johannes Tavoillot ^a, Claire Goillon ^b, Laure Pares ^c, Amélie Lefèvre ^c,
7 Hélène Védie ^d, Mireille Navarrete ^e, Alain Palloix ^{f,†}, Anne-Marie Sage-Palloix ^f, Philippe
8 Castagnone-Sereno ^g, Ariane Fazari ^g, Nathalie Marteu ^g, Caroline Djian-Caporalino ^g

9 **Affiliations**

10 ^a CBGP, IRD, CIRAD, INRA, Montpellier SupAgro, Univ. Montpellier, Montpellier, France.

11 johannes.tavoillot@ird.fr, thierry.mateille@ird.fr

12 ^b APREL (Provence Association for Vegetable Research and Experimentation), Saint-Rémy

13 de Provence. goillon@aprel.fr

14 ^c INRA, Alénia Roussillon Experimental Station, Alénia, France. _amelie.lefevre@inra.fr,

15 laure.pares@inra.fr

16 ^d GRAB (Organic Agriculture Research Group), Avignon, France. _helene.vedie@grab.fr

17 ^e ECODEVELOPPEMENT, INRA, 84000 Avignon, France. mireille.navarrete@inra.fr

18 ^f INRA UR 1052, Genetics and Breeding of Fruit and Vegetables, Montfavet, France. anne-

19 marie.sage-palloix@inra.fr

20 ^g INRA, Université Côte d'Azur, CNRS, ISA, France. caroline.caporalino@inra.fr,

21 philippe.castagnone@inra.fr, _ariane.fazari@inra.fr, _nathalie.marteu@inra.fr

22

23 * **Corresponding author:** thierry.mateille@ird.fr

24 **ABSTRACT**

25 Plant-parasitic nematodes (PPN) are detected everywhere as mixed-species communities.
26 Most non-chemical control strategies of PPN only target some species, thus raising questions
27 about the consequences that this specificity may have on the residual community. In this
28 respect, the long-term ecological sustainability of such strategies is challenged. In order to
29 evaluate the impacts of agronomical practices on PPN communities, two four-year
30 experiments that differed by the presence or absence of root-knot nematodes (RKN -
31 *Meloidogyne* spp.) were carried out under cold shelters in the south of France, under native
32 field conditions of vegetable cropping systems that included a nematicidal sorghum green
33 manure and a pepper variety carrying a RKN resistance gene. At the site with RKN, RKN
34 populations developed on susceptible vegetables. But they were controlled by the green
35 manure but not by the R-pepper, and were also vulnerable to low soil temperatures. At the site
36 without RKN, Paratylenchidae populations developed on susceptible vegetables, but were
37 controlled by both the green manure and the R-pepper, and not by low temperatures. At each
38 site, populations of Telotylenchidae exhibited dynamics suggesting competition with RKN or
39 Paratylenchidae. Hypotheses about competition models are discussed according to the
40 specific life traits of the PPN involved, including ecto- vs. endoparasitism and sedentary vs.
41 free-living behaviour, and to the antagonist mechanisms of the cover and resistant crops that
42 must be introduced in vegetable cropping systems.

43

44 **Keywords:** Community; Green manure; Plant-parasitic nematode; Plant resistance; Species
45 competition; Vegetables.

46

47 **1. Introduction**

48 Plant parasitic nematodes (PPNs) are responsible for great yield losses estimated
49 at US\$100 billion annually worldwide (Abd-Elgawad and Askary, 2015). Chemical
50 nematicides used for the management of plant-parasitic nematodes (PPNs) can have broad
51 and unintended effects. They act not only on PPNs, but also on all free-living nematode
52 species (Chitwood, 2003). Moreover, they are able to kill a wide range of other soil-borne
53 organisms and may negatively impact soil biodiversity (Rich et al., 2004). On the other hand,
54 the natural plant-protection alternatives against PPNs, such as service plants, plant resistance
55 and biocontrol agents, are more species-specific, and better preserve soil functions involved in
56 soil health and plant production (Doran and Zeiss, 2000; Timper, 2014).

57 In Mediterranean regions, RKN are most destructive to vegetable farms (Djian-
58 Caporalino, 2012). Despite the promising results obtained with these alternatives for
59 controlling RKN populations in vegetable cropping systems, questions arise concerning the
60 effect of such strategies on the entire PPN community (including RKNs) and, more precisely,
61 the effects that these strategies may have on other members of the PPN community, especially
62 competing one. Indeed, all natural control alternatives developed in agriculture focus on a few
63 target species compared to the whole PPN diversity encountered in specific sites (Jones et al.,
64 2013). As an example, most developed methods on vegetable crops concern only a few RKN
65 species (Nyczepir and Thomas, 2009). Thus, when bearing in mind that PPNs occur
66 everywhere as mixed-species communities, most natural plant protection alternatives would
67 induce long-term richness erosion, community rearrangements, increased development of
68 minor species, etc. (Mateille et al., 2008). Furthermore, the sustainability of soil
69 suppressiveness (i.e. capacity of soils to suppress plant diseases even in the presence of a
70 virulent soil-borne pest and a susceptible host) should not only be considered in terms of
71 success over time in relation to emblematic crop-specific nematode species, but in terms of
72 biodiversity and long-term soil health. In that sense, a promising option is the combination of

73 control methods targeted against the main pathogenic species plus non-specific cropping
74 practices aimed at promoting suppressiveness such as by organic amendments (Evans et al.,
75 1993; Luc et al., 2005).

76 PPN community assemblages are subjected to various types of constraints: nematode
77 evolution, intra- and interspecific interactions, climate, plant diversity and attractiveness, soil
78 characteristics and functions, land-use changes and cropping practices, etc. (Hodda et al.,
79 2009). In that way, PPN community assemblages would be shifted when applying control
80 alternatives in cropping systems (Wang et al., 2016). Previous works have been shown that
81 some sorghum species and varieties are able to reduce PPN populations such as RKNs,
82 *Helicotylenchus dihystera*, *Pratylenchus penetrans* and *Rotylenchus reniformis* (La Mondia et
83 al., 2002; Wang et al., 2004; Asmus et al., 2008; Stapleton et al., 2010; Navarrete et al.,
84 2016). However, the same sorghum species can lead to an increase of other taxa such as
85 *Belonolaimus longicaudatus*, *Mesocriconema* sp., *Paratrichodorus minor*, *Pratylenchus* spp.
86 and *Tylenchorhynchus* spp. (Rhoades, 1983; McSorley and Dickson, 1995; Crow et al., 2001;
87 Bhan et al., 2010; Villenave et al., 2010; Fraedrich et al., 2012). In southeast France, RKNs
88 occur now over 40% of the vegetable production area since the ban of methyl bromide in
89 Europe (Djian-Caporalino, 2012). As soon as alternative strategies were implemented to
90 control RKNs in this area, especially using Sudan grass as green manure, the expansion of
91 *Paratylenchus* spp. infestations was observed (Mateille and Tavoillot, 2019), even leading to
92 worrisome crop damage in some cases as seen elsewhere (Faulkner, 1964; Wang et al., 2016).

93 Usually, because PPN species utilize the same trophic source, they are under
94 competition either by exploitation (use of the same resource in limited quantities) or by
95 interference (reciprocal disturbances generated by the search for this resource when it is not in
96 limited quantities) (Begon et al., 2006). Thus, exploitation produces the elimination of
97 species, either by direct exclusion or by moving or reducing their niche until coexistence
98 becomes possible (Connell, 1980). The coexistence between PPN species depends partly on

99 their different life traits: (i) they exhibit all reproduction modalities (amphimixis, mitotic and
100 meiotic parthenogenesis, hermaphroditism), and different modalities can be found among the
101 same genus, such as RKN or *Pratylenchus* (Chitwood and Perry, 2006); (ii) they exhibit
102 different parasitism strategies (Bird and Bird, 2001): PPN may feed on plant tissues from
103 outside the plant (i.e., ectoparasites) or inside the tissues (i.e., endoparasites) and they can
104 move through plant tissues (migratory species) or can become swollen and permanently
105 immobile (sedentary species).

106 Therefore, considering the PPN diversity associated with vegetable cropping systems,
107 this study focused on the unbalanced development of PPN populations in communities
108 subjected to management techniques targeting RKNs especially. Moreover, two sites differing
109 by the presence/absence of RKNs were chosen in order to better understand how PPN species
110 compete.

111

112 **2. Materials and methods**

113 *2.1. Field survey designs*

114 Two field trials were performed each on two separate commercial organic farms
115 located in southern France, with Mediterranean climate, from 2012 to 2016. The trials were
116 carried out under 40 m x 8 m x 3.5 m plastic cover plots that were previously cropped with
117 RKN susceptible vegetables (salad and melon) since several seasons. One trial was performed
118 near Lambesc (43.65N, 5.21E). The sandy-silty soil (37.5% sand, 22.3% silt, 10.7% clay,
119 3.5% organic matter, pH 8.4) was heavily infested with RKN. During the study period, the
120 soil temperature at a depth of 15 cm varied from 5°C in winter to 30°C in summer. In order to
121 explore deeper interactions between species, the other trial was performed on a site free of
122 RKNs, located at the INRA Experimental Centre near Alénia (42.64N, 2.97E). The soil was
123 sandy-silty (33.7% sand, 48.1% silt, 18.1% clay, 1.8% organic matter, pH 7.5) and its
124 temperature reported at a depth of 10 cm varied from 4.7°C in winter to 26°C in summer.

125 In both sites, a nematicidal green manure (hybrid sorghum ‘270911’ = three-way
126 hybrid from *S. bicolor* spp. *bicolor* x Sudan grass cvs. ‘Almuden’, ‘Bihar’, ‘Magno’ and
127 ‘Artis’ cross developed by UPL France SASTM) was used for biofumigation. Four weeks after
128 sowing, the plants were cut, and then grounded and rotavated. The soil was rolled and left
129 uncovered for one month to allow biofumigation (Goillon et al., 2019). Moreover, a resistant
130 pepper crop (*Me-3* DLL R-pepper) was cultivated in spring and summer. This resistant pepper
131 is *Capsicum annuum* sweet pepper grafted on a resistant pepper rootstock carrying the major
132 *Meloidogyne* R-gene *Me-3* in the susceptible genetic background Doux-Long-des-Landes
133 (BC1-S1 [(DH149 × DLL) × DLL]). The rootstock was provided by the Genetic Resources
134 Centre for Vegetable Species (CRB-Leg) at INRA Montfavet, and the sweet pepper variety
135 was grafted by Scea Meffre PlantsTM.

136 The green manure and the resistant crops were included into crop sequences that
137 alternated RKN-susceptible vegetables such as melon (*Cucumis melo*) in spring, and lettuce
138 (*Lactuca sativa*) or Swiss chard (*Beta vulgaris* subsp. *vulgaris*) in winter.

139

140 2.2. Nematode analyses

141 Soil samples (eight random replicates in each plot at each sampling date) were
142 collected from the top 20 cm soil layer (500 ml for each replicate) at before the experiment
143 and after each susceptible crop was terminated. The samples were systematically taken from
144 the same core site, to minimize the effects of heterogeneity in the distribution of nematodes
145 over the plot. PPNs were extracted from a 250-ml aliquot of each soil replicate using the
146 elutriation procedure (Seinhorst, 1962). They were identified first to genus (Mai and Mullin,
147 1996) and counted in 5 mL aliquots sampled from 25 mL suspensions under a
148 stereomicroscope at 60x magnification (Merny and Luc, 1969). Then, nematode suspensions
149 were fixed (De Grisse, 1969) and one hundred PPN specimens at least were mounted onto
150 slides and morphologically identified to species level according to specific keys (Van

151 Bezooijen, 2006). PPN levels were expressed as the number of individuals per dm³ of fresh
152 soil. RKN populations were identified using the SCAR-PCR procedure (Zijlstra, 2000;
153 Zijlstra et al., 2000) and their avirulence was assessed on *Mi-1*-tomato and *Me3*-peppers in
154 controlled conditions (Djian-Caporalino et al., 2011).

155

156 2.3. Data analyses

157 In each site, population dynamics were monitored for each taxa. Mean population
158 levels were analysed and compared between taxa (ANOVA and Wilcoxon-Mann-Whitney
159 test with P<0.05) at each sampling date. In order to analyse competitions between PPN taxa,
160 the whole PPN data (individuals from each family /100mL of soil) gathered during the
161 experiment were analysed with normalized Principal Component Analyses (PCA). The PCA
162 were performed by using the *ade-4* package provided in R software (Chessel et al., 2004; R
163 Core Team, 2016). Then we monitored the regression kinetics of pairs of taxa adapted from
164 the Lotka-Volterra model (Begon et al., 2006) by displaying the population levels of each pair
165 of taxa at each sampling date.

166

167 3. Results

168 3.1. Dynamics of plant-parasitic nematodes in communities

169 3.1.1. At the 'Lambesc' site

170 The PPN taxa detected were *Ditylenchus acutus*, *Nothotylenchus acutus* and *N. thornei*
171 (Anguinidae), *Helicotylenchus canadensis* (Hoplolaimidae), *Meloidogyne arenaria*, *M.*
172 *incognita* (Meloidogynidae), *Mesocriconema* spp. (Criconematidae), *Paratylenchus nanus*
173 (Paratylenchidae), *Pratylenchus thornei* (Pratylenchidae), *Histotylenchus* sp., *Merlinius*
174 *microdorus* and *Tylenchorhynchus clarus* (Telotylenchidae), *Xiphinema pachtaicum*
175 (Longidoridae), and Tylenchidae species (*Basiria tumida*, *Boleodorus thylactus*, *Filenchus*
176 *hamatus*, *F. misellus*, *Ottolenchus facultativus*, *Psilenchus aestuarius*, *P. hilarulus*). *M.*

177 *arenaria* and *M. incognita* populations were identified as *Mi-1* and *Me-3* avirulent.

178 Only nematode families with a total abundance > 1% were considered for dynamics:
179 Hoplolaimidae, Meloidogynidae, Telotylenchidae and Tylenchidae. The Hoplolaimidae and
180 Tylenchidae nematodes remained in low abundance throughout the experiment and were not
181 impacted by either the green manure or by the R-pepper (Fig. 1). Meloidogynidae (98% *M.*
182 *arenaria* and 2% *M. incognita*) were the most dominant with $1,883 \pm 398$ individuals/100 mL
183 of soil at the beginning of the experiment. The Meloidogynidae abundance was strongly
184 reduced after the sorghum green manure (-94.4% in 2012 and -81.8% in 2014). At the same
185 time, the Telotylenchidae population (73.5% *T. clarus*) was enhanced (+81.9% in 2012 and
186 +57.5% in 2014), respectively. Unexpectedly, the Meloidogynidae population increased on
187 the resistant pepper crop (+99.8% in 2013 and +91.6% in 2015), while the Telotylenchidae
188 population decreased in 2013 (-44.3%) and increased in 2015 (+7.5%). Furthermore,
189 Meloidogynidae decreased on susceptible vegetables when cultivated in winter (-99% on
190 Swiss chard in 2012; -77.9% on lettuce in 2014 and -92.4% in 2015; -70.1% on lettuce in
191 2016). In contrast, the dynamics of Telotylenchidae depended on the crop succession:
192 populations declined on Swiss chard (-51.9% in 2013) and on lettuce (-48.8% in 2015)
193 following the green manure. On the other hand, they increased on lettuce following resistant
194 pepper (+17.9% in 2014 and +64.9% in 2016). The melon crop multiplied both
195 Meloidogynidae and Telotylenchidae populations increased on the melon following lettuce in
196 2014 (+85.7% and +32.3%, respectively). From the beginning to the end of the experiment,
197 the alternation of sorghum green manure and resistant pepper resulted in an overall 11.7%
198 reduction of the Meloidogynidae population, and an overall 8.7% increase of the
199 Telotylenchidae populations.

200

201 3.1.2. At the 'Alénya' site

202 The PPN taxa detected were *N. acutus* and *N. geraerti* (Anguinidae), *Lelenchus*

203 *leptosoma* (Ecphyadophoridae), *P. nanus* (Paratylenchidae), *P. thornei* (Pratylenchidae),
204 *Histotylenchus* sp. and *T. clarus* (Telotylenchidae), and Tylenchidae species (*B. tumida*, *F.*
205 *hamatus*, *O. facultativus*, *P. aestuarius*).

206 As indicated above, only the most representative nematode families were considered
207 (total abundance > 1%): Paratylenchidae, Telotylenchidae and Tylenchidae. The Tylenchidae
208 populations were the most abundant at this site at the beginning of the experiment (Fig. 2),
209 but they decreased just after the first lettuce crop and remained at a very low level throughout
210 the experiment. The two applications of green manure in 2013 and 2015 did not prevent the
211 Telotylenchidae populations (73% *T. clarus*) from increasing on lettuce and melon (+88.6%
212 and +48.6%, respectively). In the same time, the Paratylenchidae populations decreased (-
213 46.4% and -89.5%, respectively). Both Telotylenchidae and Paratylenchidae populations were
214 able to multiply on all vegetables, except when lettuce was cultivated after a four-month bare
215 period (2013), but they declined on the resistant pepper crop in 2014 (-49.7% and -94.4%,
216 respectively). Nevertheless, from the beginning to the end of the experiment, the alternation
217 of sorghum green manure and resistant pepper resulted in an overall increase of the
218 Telotylenchidae and Paratylenchidae populations (+5.4% and +28.9%, respectively).

219

220 3.2. Interactions between nematode taxa

221 When modelling the correspondence between all the nematode families by using the
222 whole PPN data gathered during the experiment at the 'Lambesc' site, the PCA analysis
223 revealed the major contribution of the Meloidogynidae (Me) and the Telotylenchidae (Te)
224 variables, and their opposite position on the first PCA axis (Fig. 3A). Moreover, when
225 modelling the kinetic regression between RKN and Telotylenchidae populations throughout
226 the experiment, we observed, with few exceptions, that Meloidogynidae populations
227 decreased when Telotylenchidae populations increased, and vice versa (Fig. 4A). In addition,
228 there was a constant increase of the Telotylenchidae at the expense of the Meloidogynidae

229 because the regression kinetic moved as a spiral according to a long-term reversal of the
230 Meloidogynidae /Telotylenchidae ratios in favour of Telotylenchidae nematodes.

231 The PCA analysis modelled on all of the nematode family data at the 'Alénya' site
232 revealed the major contribution of Paratylenchidae (Pa) and Telotylenchidae (Te) variables
233 and their opposite position on the first PCA axis (Fig. 3B). The regression modelled between
234 these two families showed a cyclic kinetic, meaning that populations of Telotylenchidae
235 decreased when populations of Paratylenchidae increased, and vice versa (Fig. 4B), but with a
236 long-term increase of both PPN families until the end of the experiment.

237

238 **4. Discussion**

239 Since RKN are usually dominant in vegetable soils, the resort to the two sites, one
240 highly infested with RKN and one free of RKN, should make it possible to analyse contrasted
241 communities and their dynamics when submitted to similar cropping systems. We have
242 chosen two different sites because RKN infestation and non-infestation conditions cannot be
243 found on the same site. Therefore, the strict comparison of the two sites cannot be performed
244 "all other factors being equal" (with/without RKN, crop sequence, climatic conditions, etc.)
245 and the interpretation should take into consideration all the different agro-environmental
246 conditions. In addition, we deliberately conducted this study under native farm conditions, i.e.
247 on large enough plots to manage all practices as in real conditions, even if it makes the
248 analysis more difficult. Therefore, replicate plots within each site could not be designed, and
249 the individual sampling points were considered as replicates.

250

251 *4.1. Response of PPN populations to seasons*

252 Looking at vegetable cropping periods, low soil temperatures that occurred during
253 each winter (5 to 10°C) decreased Meloidogynidae populations at the 'Lambesc' site,
254 although they were able to infest lettuces that are highly susceptible. *Meloidogyne arenaria*

255 and *M. incognita* can survive poorly in soils below 10°C (Evans and Perry, 2009). Low
256 temperatures are known to affect several functions. They reduce the mobility of free RKN
257 juveniles in the soil, which impacts their penetration into the roots, hence their survival.
258 Juveniles that have infested roots fail to reproduce and the embryogenesis is slowed down by
259 curbing eggs in tardicultus states (Evans and Perry, 2009). At the ‘Alénya’ site, winter periods
260 did not affect the development of *Paratylenchus nanus* that increased on winter lettuce,
261 except when the lettuce followed the 4 month-bare period in 2013. In fact, *Paratylenchus*
262 species are widespread under all types of climates and *P. nanus* is able to develop at high
263 altitudes and in cold countries (Talavera and Navas, 2002; Ryss et al., 2005). That could
264 explain why *P. nanus* was not disturbed by cold periods. Telotylenchidae nematodes that
265 were dominated by *Tylenchorhynchus clarus* exhibited contrasted behaviours. Their
266 populations declined during the winter periods at the ‘Lambesc’ site, except in 2015, whereas
267 they were enhanced or maintained at the ‘Alénya’ site, except after the bare period in 2013.
268 Like the other Telotylenchidae, *T. clarus* is cosmopolitan and is not that sensitive to
269 temperature (Noel and Lownsbery, 1978). Consequently, Meloidogynidae species were
270 obviously susceptible to low temperatures while Paratylenchidae nematodes were tolerant,
271 and Telotylenchidae were indifferently affected or not. On the other hand, the
272 Meloidogynidae, Paratylenchidae and Telotylenchidae nematodes reproduced during hot
273 periods (spring and summer), especially on melon, while it is well known that RKN are very
274 aggressive on Cucurbitaceae, and Paratylenchidae and Telotylenchidae have been shown to
275 cause damage to several vegetables (Potter and Olthof, 1993; Faske, 2013).

276

277 4.2. Response of PPN populations to R-pepper

278 Looking at the R-pepper crop that was introduced in spring and summer (soil
279 temperature up to 28°C), the low efficiency of the resistance of the *Me3*-DLL variety towards
280 RKN was confirmed at the ‘Lambesc’ site. The *Me-3* gene induces early root-cell necrosis

281 around the second stage juveniles in the upper root layers (epidermis and cortex), preventing
282 many of the juveniles from reaching their feeding site on the vascular cylinder and continuing
283 their life cycle and reproducing. This gene is weakened because it is introgressed by
284 backcross in a highly susceptible genetic background, which favours the development of
285 RKN when submitted to a high-inoculation pressure (Barbary et al., 2014). At the ‘Alénya’
286 site, the *P. nanus* populations strongly decreased during the R-pepper crop, whereas this crop
287 was installed during a hot period. Therefore, it is hypothesized that either *Capsicum annuum*
288 sweet pepper is not a good host plant for *P. nanus*, or that the *Me-3* gene may have some
289 effect on the reproduction of a nematode species other than RKN. The Telotylenchidae
290 populations were either reduced (‘Lambesc’ 2013 and ‘Alénya’ 2014) or enhanced
291 (‘Lambesc’ 2015), meaning that population dynamics would be under outer drivers (soil,
292 climate, etc.) and then that this pepper is probably also a host plant for this nematode family
293 (Santos et al., 2005).

294

295 4.3. Response of PPN populations to sorghum

296 RKN populations were significantly reduced by the sorghum hybrid ‘270911’. *P.*
297 *nanus* was also reduced but to a lesser extent, despite the fact that sorghum is a good host for
298 Paratylenchidae (Siddiqi et al., 1993). On the other hand, Telotylenchidae populations
299 reproduced at both experimental sites. Sorghum is a good host for *Tylenchorhynchus* species
300 (Fraedrich et al., 2012). However, this nematicidal hybrid did not affect them. It is therefore
301 surprising that the three nematode families did not react in the same way to sorghum
302 ‘270911’. The same observations were previously made with other sorghum varieties: in PPN
303 communities, ring (*Mesocriconema* spp.) and lesion (*Pratylenchus* spp.) nematodes
304 multiplied, whereas root-knot nematodes (*M. incognita*) were controlled (Bhan et al., 2010);
305 Sudan grass growth was reduced by *B. longicaudatus*, whereas the abundance of *M. incognita*
306 was kept constant (Crow et al., 2001). Considering that no PPN species should be immune to

307 the HCN released after burial of Sudan grass, it was postulated (i) that HCN activity would be
308 very short-lived, or (ii) that HCN would not be uniformly distributed in roots and soil
309 (McGuidwin and Layne, 1995). We suspect that the effectiveness of the sorghum ‘270911’
310 depends on the parasitic behaviour of each nematode group. Second-stage *Meloidogyne*
311 juveniles are free in the soil, but they immediately infest roots after hatching, and the next
312 stages concerning the females are endoparasite and sedentary. *Paratylenchus* species are
313 ectoparasites, but they become sedentary when feeding on cortical root cells with their long
314 stylet. Telotylenchidae are ectoparasites, feeding on epidermal cells and root hairs. Thus,
315 since only the species exhibiting a sedentary behaviour (RKN and *Paratylenchus*) were
316 reduced with sorghum ‘270911’ and not the free species in the soil (*T. clarus*), we
317 hypothesize that the deep feeding of sedentary species in toxic roots of living sorghum plants
318 would be more efficient than the toxicity on the nematode free-living stages of the HCN
319 released in the soil after sorghum incorporation. This hypothesis is supported by another
320 experiment done at ‘Lambesc’ and ‘Alenya’ with the Sudan grass ‘Piper’, commonly used as
321 green manure in France and with low leaf dhurrin content (compared to root content), thus
322 less HCN release. The same results as those obtained with sorghum ‘270911’ were described
323 in the field, and an additional experiment in controlled conditions showed that both sorghums
324 were very poor RKN hosts, not supporting reproduction of RKNs (Djian-Caporalino et al.,
325 2019).

326

327 4.4. Competitive interactions

328 Consequently, the response of the different PPN towards the practices introduced into
329 complex vegetable cropping systems depends on their species diversity, but the plant-
330 nematode interaction is not the only interaction involved. Indeed, as an example, the
331 Telotylenchidae populations once increased and again decreased on the same crop (e.g., on
332 lettuce and R-pepper) while the soil sample replicates were removed from the same core

333 places avoiding possible bias due to the aggregated distribution of PPN. This probably means
334 that other interactions occurred, such as PPN-PPN competition. It is known that the intensity
335 of interspecific competition is directly related to the overlap level of the ecological niches of
336 species (Pianka, 1978). We observed that Paratylenchidae populations were not able to
337 develop when Meloidogynidae were present, as observed at the ‘Lambesc’ site. In this case,
338 the almost total exclusion of *P. nanus* would be due to RKN juveniles, perhaps because they
339 occupy the same cortical parenchyma niche, at least temporarily, either by moving between
340 cortical cells before reaching their feeding site on the vascular cylinder (for RKN) or by
341 feeding on the cortical cells (for *P. nanus*). Moreover, the strong opposition between
342 Telotylenchidae and either Meloidogynidae (‘Lambesc’ site) or Paratylenchidae (‘Alénya’
343 site) revealed by PCA and time regression analyses confirmed competition in PPN
344 communities. However, the competition differed according to the species involved. At the
345 ‘Lambesc’ site, the cropping system that is targeted for controlling Meloidogynidae led to the
346 long-term replacement of RKN by Telotylenchidae. This would mean that when a crop
347 succession is susceptible to RKN, the Telotylenchidae are excluded by competition. On the
348 contrary, in a crop sequence that reduces Meloidogynidae, the competition is lessened in
349 favour of the Telotylenchidae. Thus, the replacement of Meloidogynidae by Telotylenchidae
350 would be due to the long-term control of the Meloidogynidae. At the ‘Alénya’ site, the
351 competition between Paratylenchidae and Telotylenchidae seems to be more cyclic with a
352 long-term increase of both PPN families, meaning less dependence on the cropping system.
353 The cyclic competition between Paratylenchidae and Telotylenchidae would be due to their
354 ectoparasitic behaviour (i.e., competition for root surface). It seems that competition between
355 Telotylenchidae and RKN, on one hand, and Telotylenchidae and Paratylenchidae, on the
356 other, would correspond to hierarchic and cyclic models, respectively (Daly et al., 2015).
357 Nevertheless, more research should be conducted in order to (i) understand how competition
358 occurs between PPN species and contributes in up and down kinetics (microcosm

359 experiments), and (ii) predict how changing cropping practices would regulate PPN diversity
360 with competition (Groselj et al., 2015).

361

362 **5. Conclusion**

363 From an agronomic point of view, these competitive interactions raise the question of
364 the risk due to a residual PPN community, once a target species (e.g., RKN) is controlled
365 (Ferris et al., 2004; De Araujo Filho et al., 2016). Even if some cropping strategies have been
366 shown to control RKN populations, Paratylenchidae and Telotylenchidae are known to be
367 pathogenic on vegetables (Khan et al., 1986; Potter and Olthof, 1993). In these experiments,
368 the replacement of RKN by Telotylenchidae nematodes and the extensive development of
369 Paratylenchidae nematodes raise questions about the sustainable reduction of the global
370 pathogenicity of the PPN community. It is therefore necessary to reconsider long-term
371 unyielding soil suppressiveness strategies such as managing the diversity of the PPN
372 communities rather than focusing on controlling targeted species. This requires a more
373 holistic approach associating several scientific disciplines such as soil ecology, nematology,
374 breeding and agronomy.

375

376 **Acknowledgements**

377 The authors are especially grateful to O. Arnaud (organic vegetable producer, Ecocert
378 certification, Lambesc, France) and to all the technicians of the INRA experimental station
379 (French National Institute for Agricultural Research, Alénya, France) who performed all the
380 cropping operations. This work was part of the GEDUNEM project (Varietal and technical
381 innovations for the sustainable and integrated management of root-knot nematodes in
382 protected vegetable cropping systems) supported by the INRA metaprogramme, SMaCH
383 (Sustainable Management of Crop Health - www.smach.inra.fr), and the GIS PicLeg (French
384 scientific interest group on integrated vegetable production - www.picleg.fr), labelled

385 TERRALIA (French competitiveness cluster for the South-East agricultural and food sector -
386 www.pole-terralia.com) and managed by Dr. C. Djian-Caporalino (INRA, France).

387

388 **References**

389 Abd-Elgawad, M.M.M., Askary, T.H., 2015. Impact of phytonematodes on agriculture
390 economy. In: Biocontrol agents of phytonematodes. CABI, pp. 3-49.

391 Asmus, G.L., Inomoto, M.M., Cargnin, R.A., 2008. Cover crops for reniform nematode
392 suppression in cotton: greenhouse and field evaluations. Trop. Pl. Pathol. 23, 85-89.

393 Barbary, A., Palloix, A., Fazari, A., Marteu, N. Castagnone-Sereno, P., Djian-Caporalino, C.,
394 2014. The plant genetic background affects the efficiency of the pepper major nematode
395 resistance genes *Me1* and *Me3*. Theor. Appl. Genet. 127, 499-507.

396 Begon, M., Townsend, C.R., Harper, J.L., 2006. Ecology: from individuals to ecosystems.
397 Blackwell.

398 Bhan, M., McSorley, R., Chase, C.A., 2010. Effect of cropping system complexity on plant-
399 parasitic nematodes associated with organically grown vegetables in Florida. Nematropica
400 40, 53-70.

401 Bird, D.K., Bird, A.F., 2001. Plant-parasitic nematodes. In: Parasitic nematodes. Molecular
402 biology, biochemistry and immunology. CABI, pp. 139-166.

403 Chessel, D., Dufour, A.B., Thioulouse, J., 2004. The ade4 package-I- One-table methods. R
404 News 4, 5-10.

405 Chitwood, D.J., 2003. Nematicides. In: Encyclopedia of Agrochemicals (Vol. 3). John Wiley,
406 pp. 1104-1115.

407 Chitwood, D.J., Perry, R.N., 2006. Reproduction, physiology and biochemistry. In: Plant
408 nematology. CABI, pp. 182-200.

409 Connell, J.H., 1980. Diversity and the coevolution of competitors, or the ghost of competition
410 past. Oikos 35, 131-138.

411 Crestin, J.M., Vannier, S., 2007. Les engrais verts en maraîchage. APREL / CA84 / Agence
412 de l'Eau Méditerranée & Corse.

413 Crow, W.T., Weingartner, D.P., Dickson, D.W., McSorley, R., 2001. Effect of sorghum
414 Sudan grass and velvetbean cover crops on plant-parasitic nematodes associated with
415 potato production in Florida. J. Nematol. 33, 285-288.

416 Daly, A.J., Baetens, J.M., De Baets, B., 2015. The impact of initial evenness on biodiversity
417 maintenance for a four-species in silico bacterial community. J. Theor. Biol. 387, 189-205.

418 De Araujo Filho, J.V., Machado, A.C.Z., Dallagnol, L.J., Camargo, L.E.A., 2016. Root-knot
419 nematodes (*Meloidogyne* spp.) parasitizing resistant tobacco cultivars in southern Brazil.
420 Pl. Dis. 100, 1222-1231.

421 De Grisse, A.T., 1969. Redescription ou modifications de quelques techniques utilisées dans
422 l'étude des nématodes phytoparasites. Int. S. Crop 34, 351-369.

423 Djian-Caporalino, C., 2012. Root-knot nematodes (*Meloidogyne* spp.), a growing problem in
424 French vegetable crops. EPPO Bull. 42, 127-137.

425 Djian-Caporalino, C., Mateille T., Bailly-Bechet, M., Marteu, N., Fazari, A., Bautheac, P.,
426 Raptopoulo, A., Van Duong, L., Tavoillot, J., Martiny, B., Goillon, C., Castagnone-Sereno,
427 P., 2019. Evaluating sorghums as green manure against root-knot nematodes. Crop Prot.
428 122, 142-150.

429 Djian-Caporalino, C., Molinari, S., Palloix, A., Ciancio, A., Fazari, A., Marteu, N., Ris, N.,
430 Castagnone-Sereno, P., 2011. The reproductive potential of the root-knot nematode
431 *Meloidogyne incognita* is affected by selection for virulence against major resistance genes
432 from tomato and pepper. Eu. Pl. Pathol. 131, 431-440.

433 Doran, J.W., Zeiss, M.R., 2000. Soil health and sustainability: managing the biotic component
434 of soil quality. Appl. Soil Ecol. 15, 3-11.

435 Evans, A.A.F, Perry, R.N., 2009. Survival mechanisms. In: Root-knot nematodes. CABI, pp.
436 201-222.

437 Evans, K., Trudgill, D.L., Webster, J.M., 1993. Plant parasitic nematodes in temperate
438 agriculture. Oxford University.

439 Fasje, T.R., 2013. Penetration, post-penetration development, and reproduction of
440 *Meloidogyne incognita* on *Cucumis melo* var. *Texanus*. J. Nematol. 45, 58-65.

441 Faulkner, L.R., 1964. Pathogenicity and population dynamics of *Paratylenchus hamatus*
442 Thorne and Allen, on *Mentha* spp. Phytopathol. 54, 344-348.

443 Ferris, H., McKenry, M.V., Jaffee, B.A., Anderson, C.E., Juurma, A., 2004. Population
444 characteristics and dosage trajectory analysis for *Mesocriconema xenoplax* in California
445 Prunus orchards. J. Nematol. 36, 505-516.

446 Fraedrich, S.W., Cram, M.M., Handoo, Z.A., Zarnoch, S.J., 2012. Influence of
447 *Tylenchorhynchus ewingi* on growth of loblolly pine seedlings, and host suitability of
448 legumes and small grains. Nematol. 14, 417-425.

449 Groselj, D., Jenko, F., Frey, E., 2015. How turbulence regulates biodiversity in systems with
450 cyclic competition. Phys. Rev. E 91, no033009.

451 Hodda, M., Peters, L., Traunspurger, W., 2009. Nematode diversity in terrestrial, freshwater
452 aquatic and marine systems. In: Nematodes as environmental indicators. CABI, pp. 45-93.

453 Jones, J.T., Haegeman, A., Danchin, E.G., Gaur, H.S., Helder, J., Jones, M.K.G., Kikuchi, T.,
454 Manzanilla-López, R., Palomares-Rius, J.E., Wesemael, W., Perry, R.N., 2013. Top 10
455 plant-parasitic nematodes in molecular plant pathology. Mol. Pl. Pathol. 14, 946-961.

456 Khan, R. M., Khan, A.M., Khan, M.W., 1986. Interaction between *Meloidogyne incognita*,
457 *Rotylenchulus reniformis* and *Tylenchorhynchus brassicae* on tomato. Rev. Nématol. 9,
458 245–250.

459 LaMondia, J.A., Elmer, W.H., Mervosh, T.L., Cowles, R.S., 2002. Integrated management of
460 strawberry pests by rotation and intercropping. Crop Prot. 21, 837-846.

461 Luc, M., Sikora, R.A., Bridge, J., 2005. Plant parasitic nematodes in subtropical and tropical
462 agriculture. CABI.

463 McGuidwin, A.E., Layne, T.L., 1995. Response of nematode communities to Sudangrass and
464 Sorghum-Sudangrass hybrids grown as green manure crops. *J. Nematol.* 27, 609-616.

465 McSorley, R., Dickson, D.W., 1995. Effect of tropical rotation crops on *Meloidogyne*
466 *incognita* and other plant-parasitic nematodes. *J. Nematol.* 27, 535-544.

467 Mai, W.F., Mullin P.G., 1996. Plant-parasitic nematodes: a pictorial key to genera, 5th ed.
468 Comstock Publishing Associates, Cornell University Press.

469 Mateille, T., Cadet, P., Fargette, M., 2008. Control and management of plant-parasitic
470 nematode communities in a soil conservation approach. In: Integrated management and
471 biocontrol of vegetable and grain crops nematodes. Springer, pp. 79-97.

472 Mateille, T., Tavoillot, J., 2019. [http://ephytia.inra.fr/fr/C/7624/Info-Bioagresseurs-](http://ephytia.inra.fr/fr/C/7624/Info-Bioagresseurs-Paratylenchus)
473 *Paratylenchus*.

474 Merny, G., Luc, M., 1969. Les techniques d'évaluation des populations dans le sol. In:
475 Problèmes d'écologie, l'échantillonnage des peuplements animaux dans les milieux
476 terrestres. Masson, pp. 257-292.

477 Navarrete, M., Djian-Caporalino, C., Mateille, T., Palloix, A., Sage-Palloix, A.M., Lefèvre,
478 A., Fazari, A., Marteu, N., Tavoillot, J., Dufils, A., Furnio, C., Pares, L., Forest, I., 2016. A
479 resistant pepper as a trap cover crop in vegetable producing strongly decreases root-knot
480 nematode infestation in soil. *Agron. Sustain. Dev.* 36, 68.

481 Noel, G.R., Lownsbery, B.F., 1978. Effects of temperature on the pathogenicity of
482 *Tylenchorhynchus clarus* to Alfalfa and observations on feeding. *J. Nematol.* 10, 195-198.

483 Nyczepir, A.P., Thomas, S.H., 2009. Current and future management strategies in intensive
484 crop production systems. In: Root-knot nematodes. CABI, pp: 412-443.

485 Pianka, E.R., 1978. Evolutionary ecology. Harper and Row.

486 Potter, J.W., Olthof, T.H.A., 1993. Nematode pests of vegetable crops. In: Plant parasitic
487 nematodes in temperate agriculture. CABI, pp: 171-207.

488 R Core Team, 2016. R: a language and environment for statistical computing: R Foundation

489 for Statistical Computing. www.Rproject.org.

490 Rich, J.R., Dunn, R.A., Noling, J.W., 2004. Nematicides: past and present uses. In:

491 Nematology: advances and perspectives, Vol. 2 Nematode management and utilization.

492 CABI, pp: 1179-1200.

493 Rhoades, H.L., 1983. Effects of cover crops and fallowing on populations of *Belonolaimus*

494 *longicaudatus* and *Meloidogyne incognita* and subsequent crop yields. *Nematropica* 13, 9-

495 16.

496 Ryss, A., Boström, S., Sohlenius, B., 2005. Tylenchid nematodes found on the Nunatak

497 Basen, East Antarctica. *Ann. Zool.* 55, 315-324.

498 Santos, B.M., Gilreath, J.P., Motis, T.N., 2005. Managing nutsedge and stunt nematode in

499 pepper with reduced methyl bromide plus chloropicrin rates under virtually impermeable

500 films. *Hort Technol.* 15, 596-599.

501 Seinhorst, J.W., 1962. Modifications of the elutriation method for extracting nematodes from

502 soil. *Nematologica* 8, 117-128.

503 Siddiqi, M.R., Baujard, P., Mounport, D., 1993. Descriptions of *Paratylenchus*

504 *pernoxius* sp.n. and *Paralongidorus duncani* sp.n. from Senegal, and the synonymization

505 of *Longidoroides* with *Paralongidorus*. *Afro-Asian J. Nematol.* 3, 81-89.

506 Stapleton, J.J., Summers, C.G., Mitchell, J.P., Prather, T.S., 2010. Deleterious activity of

507 cultivated grasses (Poaceae) and residues on soilborne fungal, nematode and weed pests.

508 *Phytoparasitica* 38, 61-69.

509 Talavera, M., Navas, A., 2002. Incidence of plant-parasitic nematodes in natural and semi-

510 natural mountain grassland and the host status of some common grass species. *Nematology*

511 4, 541-552.

512 Timper, P., 2014. Conserving and enhancing biological control of nematodes. *J. Nematol.* 46,

513 75-89.

514 Van Bezooijen, J., 2006. Methods and techniques for nematology. Wageningen University.

515 Villenave, C., Saj, S., Pablo, A.L., Sall, S., Djigal, D., Chotte, J.L., Bonzi, M., 2010.
516 Influence of long-term organic and mineral fertilization on soil nematofauna when growing
517 *Sorghum bicolor* in Burkina Faso. Biol. Fert. Soils 46, 659-670.

518 Wang, K., Li, Y., Xie, H., Wu, W.J., Xu, C.L., 2016. Pin nematode slow decline of
519 *Anthurium andraeanum*, a new disease caused by the pin nematode *Paratylenchus*
520 *shenzhenensis*. Pl. Dis. 100, 940-945.

521 Wang, K.H., McSorley, R., Gallaher, R.N., 2004. Winter cover crops and nematode densities.
522 J. Nematol. 36, 517-523.

523 Zijlstra, C., 2000. Identification of *Meloidogyne chitwoodi*, *M. fallax* and *M. hapla* based on
524 SCAR-PCR, a powerful way of enabling reliable identification of populations or
525 individuals that share common traits. Eur. J. Pl. Pathol. 106, 283-290.

526 Zijlstra, C., Donkers-Venne, D.T., Fargette, M., 2000. Identification of *Meloidogyne*
527 *incognita*, *M. javanica* and *M. arenaria* using sequence characterised amplified region
528 (SCAR) based PCR assays. Nematology 2, 847-853.

529

530 **Legends for figures and tables**

531

532 **Figure 1.** Kinetics of plant-parasitic nematode populations all along the cropping schedule
533 assayed at the ‘Lambesc’ site. S = nematicidal *Sorghum* hybrid ‘270911’; R pepper =
534 *Capsicum annuum* rootstock *Me3* DLL. Unnamed periods = bare periods. Stars indicate
535 significant differences between population levels at each sampling date ($P < 0.05$).

536 **Figure 2.** Kinetics of plant-parasitic nematode populations all along the cropping schedule
537 assayed at the ‘Alénia’ site. S = nematicidal *Sorghum* hybrid ‘270911’; R pepper =
538 *Capsicum annuum* rootstock *Me3* DLL. Unnamed periods = bare periods. Stars indicate
539 significant differences between population levels at each sampling date ($P < 0.05$).

540 **Figure 3.** Covariation among taxa in plant-parasitic nematode communities at the ‘Lambesc’
541 (A) and ‘Alénia’ (B) sites. Normalized PCA loading plot of the plant-parasitic nematode
542 families (Ho = Hoplolaimidae; Me = Meloidogynidae; Pa = Paratylenchidae; Te =
543 Telotylenchidae; Ty = Tylenchidae).

544 **Figure 4.** Covariation among taxa in plant-parasitic nematode communities at the ‘Lambesc’
545 (A) and ‘Alénia’ (B) sites. Time regression between nematode families (1 to 11 =
546 sampling dates; dotted lines = tendency line).

547

Figure 1




Figure 3




Figure 4

