



Duplication and partitioning in evolution and function of homoeologous Q loci governing domestication characters in polyploid wheat

Zengcui Zhanga, Harry H. Belcram, Piotr Gornickic, Mathieu Charles, Jérémy Just, Cécile C. Huneau, Ghislaine Magdelenat, Arnaud Couloux, Sylvie Samaind, Bikram S. Gille, et al.

► To cite this version:

Zengcui Zhanga, Harry H. Belcram, Piotr Gornickic, Mathieu Charles, Jérémy Just, et al.. Duplication and partitioning in evolution and function of homoeologous Q loci governing domestication characters in polyploid wheat. Proceedings of the National Academy of Sciences of the United States of America, 2011, 108 (46), pp.18737 - 18742. 10.1073/pnas.1110552108 . hal-02650358

HAL Id: hal-02650358

<https://hal.inrae.fr/hal-02650358>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Correction

EVOLUTION

Correction for “Duplication and partitioning in evolution and function of homoeologous *Q* loci governing domestication characters in polyploid wheat,” by Zengcui Zhang, Harry Belcram, Piotr Gornicki, Mathieu Charles, Jérémy Just, Cécile Huneau, Ghislaine Magdelenat, Arnaud Couloux, Sylvie Samain, Bikram S. Gill, Jack B. Rasmussen, Valérie Barbe, Justin D. Faris, and Boulos Chalhouh, which appeared in issue 46, November 15, 2011, of *Proc Natl Acad Sci USA* (108:18737–18742; first published October 31, 2011; 10.1073/pnas.1110552108).

The authors note that, due to a printer’s error, on page 18737, right column, first paragraph, line 13, “*T. turgidum*” should instead appear as “*T. aestivum*.” Additionally, on page 18738, left column, second full paragraph, line 7, “*T. turgidum*” should instead appear as “*T. aestivum*.” Lastly, on page 18739, left column, third full paragraph, line 4, “*T. turgidum*” should instead appear as “*T. aestivum*.”

www.pnas.org/cgi/doi/10.1073/pnas.1121066109




Fig. 1. Comparison of effects of 5AQ on morphological and domestication-related traits of hexaploid wheat cv. CS spikes. (A) Presence of 5AQ confers a square-shaped spike (Left), and the absence of 5AQ (in the deletion mutant CS del143) (Right) results in a speltoid spike characterized by a much longer and spear-shaped spike compared with the spike of CS primarily due to the elongation of rachis segments. (B) Seed of CS with 5AQ (Left) is free threshing because the seed is loosely held and abscission occurs at the base of the glumes, leaving the rachis intact. CS lacking the 5AQ gene (Right) is not free threshing because the glumes hold the seed more tightly and abscission occurs within the rachis, causing disarticulation of the entire spikelet with a portion of the rachis left attached to the base of the spikelet. (C) Detailed illustration of a single spikelet of CS when 5AQ is present (Left) and absent (Right). Both spikelets have three fertile florets (first, second, and third) and each spikelet has a pair of glumes (1). Each floret has a lemma (2), seed (3), and palea (4). Note the looseness of the seeds due to the difference in the position of the glumes in 5AQ compared with the mutant lacking 5AQ. The abscission zones are indicated by the blue arrows (the lower spikelet in the 5AQ mutant was removed so that the abscission zone could be observed).

tween homoeoalleles, or other changes caused by polyploidization, are essential as well.

Here, we evaluate the structure, evolution, and function of the homoeologous *Q/q* loci to establish a molecular basis for their contribution to the domestication phenotype in the genetic background of diploid and allopolyploid wheat.

Results

Organization of the *Q/q* Gene Locus in Different Wheat Genomes and Species. The sequences of 11 homologous and homoeologous genomic regions spanning the *Q/q* loci from five wheat species (A, B, S, and D genomes and three ploidy levels) were compared: the A-genome locus from the diploid *T. urartu* (URA) accession TMU138, the tetraploid *T. turgidum* cv. Langdon (LDN) and the hexaploid *T. turgidum* cvs. Renan (RE) and Chinese Spring (CS); the B-genome locus from LDN, RE, and CS; the S-genome locus from the diploid *Ae. speltooides* (SPE) accession SPELT1; and the D-genome locus from the diploid *Ae. tauschii* (TAU) accession AL8/78, as well as from RE and CS (Fig. 2 and Fig. S2).

Q/q was the only gene identified in the sequenced fragments of the B- and S-genome loci. A 23S ribosomal protein-like gene was identified 40 kb downstream of *5Dq* in the D genomes and 100 kb




Fig. 2. Schematic presentation of the 11 sequenced homologous and homoeologous genomic regions, spanning the *Q/q* loci from five wheat species. Line diagrams illustrate sequenced DNA segments. Color-coded block diagrams illustrate results of pairwise sequence comparisons. Additional information is provided in Fig. S2.

downstream of *5Aq* in the A genome of LDN. This gene is located 55 kb from the *5Aq* gene in *Triticum monococcum* and 9 kb from the *q* ortholog in rice chromosome 3 (27).

Eighty percent of the sequenced genomic regions consist of nested transposable elements (TEs) interspersed with stretches of unassigned sequences (Fig. 2 and Fig. S2). TEs are different when comparing homoeologous loci, i.e., between A, B/S, and D loci, highlighting their dynamic nature. Most TEs are conserved among A, B, and D homologous loci where 19 large genome-specific indels and breaks in genome colinearity due to recent events such as transposon insertions were revealed (Fig. 2 and Fig. S2). TE annotation and insertion dates estimated for 52 retrotransposons with complete long-terminal repeats (LTRs) are presented in Fig. S2 and Table S1.

All *Q/q* homoeoalleles Are Transcriptionally Active, but *5Bq* Is a Pseudogene That Does Not Encode a Full-Length *q* Protein. The in-

tron/exon structure of the *5AQ* gene (Fig. 3) was previously established (23). Sequences of the *5AQ* coding regions of LDN and CS were identical and differed from the RE sequence by only two substitutions in introns. The coding sequence of URA *5aq* differed from *5AQ* coding sequence of CS, RE, and LDN by five substitutions including two nonsynonymous substitutions, one being the V329I mutation (numbering as in *5AQ*) proposed to be responsible for the phenotypic differences between *5AQ* and *5aq* alleles (23).

For *5Bq*, two transcript variants, *5Bq.1* and *5Bq.2*, were identified for CS (Fig. 3). *5Bq.1* contains only six exons, corresponding to exons 1 and 5–10 of *5AQ*. The first exon ends 45 bp upstream and then continues from 79 bp upstream of the corresponding 3'- and 5'-splice sites of exons 1 and 5 of *5AQ*, respectively, due to a 531-bp deletion including sequences corresponding to exons 2–4 as well as parts of exon 1 and intron 4 of *5AQ*. *5Bq.2* has the same structure as *5Bq.1*, except that the sequence corresponding to intron 7 of *5AQ* is not spliced out. *5Bq.1* and *5Bq.2* encode truncated 354-bp ORFs with a stop codon located at a position equivalent to nucleotides 63–61 upstream of the corresponding 5'-splice site of *5AQ* exon 5.

The 531-bp deletion was not found in *5Bq* of LDN and RE, but a frameshift caused by a 2-bp deletion in predicted exon 2 leads to a stop codon 69 bp from the beginning of predicted exon 4 and a 519-bp truncated ORF (Fig. 3). The 2-bp deletion is not found in CS *5Bq* because of the larger 531-bp deletion. The *5Bq* 2-bp de-




Fig. 3. An overview of the intron/exon structure of the *Q/q* homoeoalleles determined by RT-PCR cloning and sequencing. Exons conforming to the splicing pattern found for the *5AQ* gene are shown in green (deduced from cDNA sequences) and gray (predicted), alternative exons are shown in orange, and introns are shown in yellow. 5'- and 3'-UTR lengths (L) by RACE are shown for CS mRNAs. Gene sizes shown are from the translation start to the stop codon. Exons encoding the AP2 domains and the position of the codon for amino acid residue 329 in *5AQ* (I), and the corresponding codons in *5AQ* (V) and *5Dq* (L) are also shown.

letion is present in hexaploid wheats and in most domesticated tetraploids, but it is much less prevalent among wild tetraploids (*T. turgidum* ssp. *dicoccoides*), suggesting that it occurred after tetraploidization and was perhaps selected upon domestication. The 531-bp deletion was only found in some hexaploid wheat lines, suggesting that it occurred after allohexaploidization.

For LDN, one *5Bq* transcript type with the *5AQ* splicing pattern was confirmed (Fig. 3). For RE, five different transcript types were detected by RT-PCR cloning and sequencing. These transcripts include two to four additional exons spliced out of a 9,535-bp helitron inserted two bp upstream of the 5'-splice site of the predicted exon 5 of the *5Bq* gene (Fig. 3). The entire helitron sequence was presumably transcribed. Some *5Bq* transcripts from RE lacked the canonical exon 5 or used an alternative 3' splicing site for intron 4. One transcript lacked the canonical exon 9. The RE sequence differed from CS by one synonymous substitution and they both differed from LDN by two more synonymous and one nonsynonymous substitutions, if the *5AQ* splicing pattern was assumed and the frameshift-causing deletions were ignored.

5Bq of LDN, CS, and RE are transcriptionally active, but do not encode full-length functional *q* proteins. Therefore, *5Bq* is a pseudogene in all three species.

The 9,536-bp helitron transposon in RE is the first complete helitron identified in wheat (more information is provided in Fig. S3). The helitron insertion was traced in the RE pedigree to Mironovskaya 808, a *T. turgidum* winter wheat cultivar of Russian origin. This insertion has not been found in any other hexaploid nor in any wild or domesticated tetraploid wheat line (Table S2), suggesting that it occurred only very recently.

Two transcript variants were identified for the CS *5Dq* gene. *5Dq.1* results from splicing of 10 exons matching the splicing pattern of *5AQ*, but for transcript *5Dq.2*, retention of the first intron leads to a premature stop codon and a 495-bp truncated

ORF (Fig. 3). The coding parts of *5Dq* of CS and REN are identical and differ from the TAU sequence by only one synonymous substitution and three intron substitutions. *5Sg* in SPE encodes a full-length *q* protein, on the basis of the splicing pattern of *5AQ* (Fig. 3).

A *q* Gene Duplication in the Wheat Ancestral Genome Was Probably Followed by Differential Loss of One Copy. Full-length *Triticeae Q/q* coding sequences were used in the phylogenetic analysis (Fig. 4). For the *5Bq* pseudogenes, the frameshifts caused by the deletions were adjusted to generate coding sequences corresponding to full-length ORFs found in other *Q/q* genes, even though such mRNAs and corresponding proteins are not made. The null hypothesis of equal evolutionary rate (third codon positions) throughout the tree was not rejected under any of several models tested. The molecular clock was calibrated using 11.6 million years ago (Mya) for the divergence time between barley and wheat (12).

It has been shown previously that *5AQ* and *5Aq* alleles and corresponding phenotypes are correlated with a V329I mutation (23). This mutation occurred very recently and is present in the A genome of only some wheats (Fig. 4). V is the ancestral state at residue 329 and it mutated to L in the B/D/S *q* lineage.

The 2.6 ± 0.8 Mya divergence time between the D and B/S genomes (S genome appears to be more closely related to B than to D genome) (Fig. 4) is consistent with our previous estimates (12, 13, 28). The divergence time of *5Aq* of *T. monococcum* from *5AQ/q* of *T. urartu* and polyploid wheats was estimated at 2.6 ± 0.7 Mya, an earlier date than similar estimates for the *Acc-1* gene (13). In the absence of any evidence of an accelerated substitution rate, an ancestral *q* gene duplication some 5.8 Mya (Fig. 4) followed by a selective loss of one of the copies in the A-genome progenitor and the other one in the progenitor of the B, D, and S genomes is the




Fig. 4. Phylogenetic relationships and divergence times of the *Q/q* genes. The phylogenetic tree for synonymous substitutions was constructed by the neighbor-joining method. Branch lengths are shown by the number of substitutions. Bootstrap values (%) are shown in square brackets. The molecular clock was calibrated using 11.6 Mya for the time of divergence between barley and wheat (12). Species with the Q allele carrying I₃₂₉ are shown on a gray background. Genomes are indicated by letters. The cross indicates that *5Bq* is a pseudogene. Numbering of amino acid residues is as in *5AQ*.

Role of *Q/q* in Controlling Morphology and Domestication Traits of Hexaploid Wheat. Correlation between *Q/q* loci and various traits revealed pleiotropic effects of *5AQ* influencing glume toughness, threshability, rachis fragility, spike length and shape, spike emergence time, and plant height (Fig. 6 and Table S3). Deletion of *5AQ* (CS-del143) or replacement with *5Aq* (CS-DIC 5A) had similar effects on all traits compared with CS. CS has soft, papery glumes and free-threshing seed, whereas CS-DIC 5A and CS-del143 have tougher, thicker, and more heavily keeled glumes tightly enveloping the seed that was not free threshing. CS-DIC 5A and CS-del143 plants are taller and have longer, lax, and tapering (speltoid) spikes compared with CS (Fig. 6 and Table S3). Spike emergence is 7 d earlier in CS-del143 and 12 d earlier in CS-DIC 5A than CS, which is consistent with the previously identified role of *5AQ* in delaying heading time (23). CS-DIC 5A has more spikelets and seeds per spike than CS-del143 and CS, but grain yield per plant is similar for all three due to more tillers in CS and CS-del143.

The spikes of CS 5BL-14 (*5Bq* deletion) and CS 5DL-5 (*5Dq* deletion) showed slightly more speltoid phenotypes compared with CS, primarily due to reduced spikelet density in the lower portions of the spikes (Fig. 6). The shape of the upper portion of the spikes of CS 5BL-14 and CS 5DL-5 was intermediate between the speltoid spike shape of CS-del143 and the square-headed spike shape of CS. Glumes of CS 5BL-14 and CS 5DL-5 were tougher and more keeled compared with those of CS, but they were not as tough as the CS-del143 glumes. Disarticulation in CS 5BL-14 and CS 5DL-5 occurred at the base of the glumes, as it does in CS. Compared with CS, the number of spikelets per spike and seed weight were reduced in CS 5BL-14 and CS 5DL-5 (Fig. 6 and Table S3). The number of seeds per spike was reduced by 67–75% due to low fertility. All these changes led to substantially reduced grain yield. These observations suggest that *5Dq* and *5Bq* contribute to threshability and the suppression of the speltoid characters, but to a lesser degree than *5AQ*.

Compared with the parental lines CS-del143 and CS 5DL-5, the spikes of the double deletion line CS-5A/5Ddd, which produces no functional *Q* or *q* protein, were much more speltoid in appearance (Fig. 6) with extremely tough glumes suggesting that the genetic effects of *5AQ* and *5Dq* are additive. The double deletion line had fewer spikelets per spike and lower fertility than its parental lines (Table S3), which resulted in the lowest grain yield among all of the stocks tested. Spikes of CS 5DL-5 and CS-5A/5Ddd emerged more than 30 d later than spikes of CS, most likely caused by the absence of gene(s) other than *q* in the terminal deletion of chromosome 5D.

Discussion

It has long been understood that the *Q* allele on wheat chromosome 5A has had a profound influence on wheat domestication. Here, we characterize the structure, evolution, and function of the *Q/q* loci on chromosomes 5A, 5B (and 5S), and 5D, to understand their contribution to domestication traits.

First, the intergenic sequences flanking the *Q/q* loci, composed primarily of TEs, have profoundly diverged by nested insertion and deletion among the A, B, D, and S genomes. This is a common feature of the intergenic regions of the wheat genome (12, 28–31). This important comparative study showed thus no putative regulating conserved sequence, except the ~2 and ~1 kb at 5'- and 3' ends of the *Q/q* gene, respectively (Fig. 2).

Second, *5AQ/q* is a cryptoparalog of the *q* gene on B, S, and D genomes, a result of gene duplication in the progenitor of the diploid wheats followed by selective loss of one copy of the gene. Functional interactions of paralogs may be substantially different from interactions of less divergent homoeologs.

Third, *5AQ/q*, *5Dq*, and *5Sq* are expected to produce functional *Q/q* proteins, but *5Bq* is a pseudogene encoding truncated proteins without the AP2 DNA-binding domains. It is unknown

whether these truncated proteins serve a function. Both *5Bq* and *5Dq* produce multiple transcript variants. One of the *5Dq* transcripts encodes a full-length *q* protein, whereas the other transcript encodes a truncated protein without the AP2 domains. About 20% of plant genes produce multiple transcripts, but their functions often remain unknown (32).

Fourth, the *5Bq* pseudogene remains transcriptionally active. Plant pseudogenes derived from recently duplicated genes often remain transcriptionally active, presumably due to insufficient time for promoter degradation (33). We showed however that the *5Bq* pseudogene participates in regulation of expression of the *5AQ* and *5Dq* genes at the transcript level, suggesting an active interaction between the homoeoalleles (see below).

Fifth, the *5Bq* pseudogenization occurred after allotetraploidization, possibly upon selection imposed by domestication. This suggests that *5Bq* plays a role in regulating expression of the domestication traits in tetraploid and hexaploid wheat.

Given the current findings, we propose that expression of the free-threshing phenotype and other domestication traits associated with the *Q* allele requires not only the V329I amino acid substitution in the *Q* protein, but also a combination of the divergent *Q/q* loci, with features we have identified, brought together in polyploid wheat. This also provides an explanation for the specificity of the free-threshing character to polyploid wheat and its absence among the diploids (26).

To investigate expression regulation and interactions between different *Q/q* homoeoalleles and their roles in domestication of hexaploid wheat, we analyzed different wheat lines altered for the *Q/q* homoeoalleles. Whereas the deletion of *5AQ* includes a small genomic segment of ~2 Mb (34), deletions of *5Bq* and *5Dq* encompass genomic segments representing 25% and 24% of chromosomes 5B and 5D, respectively (35), suggesting that deletion of other genes could also influence *Q/q* expression or the morphological traits that we evaluated. However, loci affecting these traits have not been reported on the deleted segments of chromosomes 5B or 5D. Good evidence of the appropriateness of the genetic material to study interactions between the different *Q/q* homoeoalleles was provided by the observation that the transcript levels and phenotypes for lines lacking *5Bq* and *5Dq*, which both down-regulate expression of *5AQ*, were nearly the same.

Our transcriptional expression analysis indicated complex interactions among the *Q/q* homoeoalleles. Interpretation was complicated by the fact that *5Bq* is a transcriptionally active pseudogene (no *q* protein is made) and *5AQ/q* is a cryptoparalog of *5Bq* and *5Dq*. Clearly, deletion of *5Dq* or *5Bq* leading to reduced expression of *5AQ* suggests that they indirectly up-regulate expression of *5AQ*, whereas the latter plays a repressor role for both *5Bq* and *5Dq* expression. Our results, consistent with previous research (23), demonstrate that *5AQ* is transcribed at higher levels than *5Aq*. The V329I substitution was shown to be responsible for enhanced homodimer formation in *5AQ*, possibly leading to self up-regulation of *5AQ* transcription (23). Our results show that the expression of *5Bq* and *5Dq* increases when *5AQ* is deleted or replaced by *5Aq*, suggesting that the V329I mutation in *5Aq/Q* is also responsible for the down-regulation of *5Bq* and *5Dq* (comparison of expression in lines CS, CS-del143, and CS-DIC 5A). This result could be explained by dosage balance or complementary effects in regulation among the *Q/q* homoeoalleles (36), but more work is needed to decipher the mechanism of this transcriptional regulation.

It has been recently proposed that interactions between mRNAs, transcribed pseudogenes, and long noncoding RNAs form a large-scale regulatory network across the transcriptome using microRNA response elements (11). A putative miRNA172 binding site in exon 10 of the *Q/q* gene (23) points to a possible role of miRNA in regulation of *Q/q* gene expression. The miRNA172 binding site is also present in the *5Bq* pseudogene transcripts.

Interesting correlations were observed between expression variation of *Q/q* homoeoalleles and variations of domestication and agronomic traits. Our study confirms previous findings that the *5AQ/q* genes influence numerous domestication traits (23, 25, 34, 37, 38), and it shows that along with the *5AQ/q* alleles, the *5Dq* and *5Bq* alleles contribute to speltoid suppression, but their function depends on the allelic state of *5AQ/q*. The contribution of *5Bq* to these traits is indirect because it is a pseudogene, but we demonstrated that its deletion significantly reduced transcription of *5Dq*. This apparent coregulation of the *q* genes largely explains the similar phenotype observed when *5Dq* (CS 5DL-5) or *5Bq* (CS 5BL-14) are deleted. The phenotypic differences correlated with expression variations observed between CS-del143 and CS-5A/5Ddd suggest that *5Dq* plays a more important role in controlling spike shape, spikelet number, and glume toughness when *5AQ* is absent.

Subfunctionalization (evolution of partitioned ancestral functions among alleles or homoeoalleles) and neofunctionalization (evolution of novel functions among alleles or homoeoalleles) were suggested as mechanisms leading to the preservation of the duplicated genes after polyploidization (5). Here, we provide a rare example of the fate of a set of homoeologous genes, associated with plant morphology and domestication in hexaploid

wheat. A recent mutation in *5Aq* led to “hyperfunctionalization” of *5AQ*; *5Bq* underwent pseudogenization, but remains transcriptionally active and influences regulation of expression of the homoeologs; and *5Dq* has been subfunctionalized because it contributes to many of the domestication traits.

Materials and Methods

A full description of the materials and methods used in this work is provided in *SI Materials and Methods*. This information includes BAC library screening, sequencing, and annotation; cDNA amplification and sequencing; transcript and expression analysis; phylogenetic analysis; phenotypic analysis; sequences of primer (Table S4); and descriptions of the wheat genotypes used in this research.

ACKNOWLEDGMENTS. We thank Joseph Jahier (Institut National de la Recherche Agronomique) and Moshe Feldman (Weizmann Institute of Science) for valuable discussions and for providing some of the wheat samples. This project was supported by US Department of Agriculture (USDA), Agricultural Research Service, Current Research Information Systems Project 5442-22000-033-00D and the National Center for Sequencing (Centre National de Séquençage-Génoscope)/APCNS2003 Project: *Triticum* Species Comparative Genome Sequencing in Wheat. PCR-based tracing of retrotransposon insertions was funded by the Agence Nationale pour la Recherche (ANR) Biodiversité Project (ANR-05-BDIV-015) and the ANR-05-Blanc Project impact of transposable elements on gene regulation.

- Adams KL, Wendel JF (2005) Novel patterns of gene expression in polyploid plants. *Trends Genet* 21:539–543.
- Soltis PS, Soltis DE (2009) The role of hybridization in plant speciation. *Annu Rev Plant Biol* 60:561–588.
- Van de Peer Y, Fawcett JA, Proost S, Sterck L, Vandepoele K (2009) The flowering world: A tale of duplications. *Trends Plant Sci* 14:680–688.
- Prince VE, Pickett FB (2002) Splitting pairs: The diverging fates of duplicated genes. *Nat Rev Genet* 3:827–837.
- Lynch M, Force A (2000) The probability of duplicate gene preservation by subfunctionalization. *Genetics* 154:459–473.
- Comai L (2005) The advantages and disadvantages of being polyploid. *Nat Rev Genet* 6:836–846.
- Chen ZJ, Ni Z (2006) Mechanisms of genomic rearrangements and gene expression changes in plant polyploids. *Bioessays* 28:240–252.
- Birchler JA, Auger DL, Riddle NC (2003) In search of the molecular basis of heterosis. *Plant Cell* 15:2236–2239.
- Chaudhary B, et al. (2009) Reciprocal silencing, transcriptional bias and functional divergence of homoeologs in polyploid cotton (gossypium). *Genetics* 182:503–517.
- Hovav R, et al. (2008) Partitioned expression of duplicated genes during development and evolution of a single cell in a polyploid plant. *Proc Natl Acad Sci USA* 105: 6191–6195.
- Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP (2011) A ceRNA hypothesis: The Rosetta Stone of a hidden RNA language? *Cell* 146:353–358.
- Chalupska D, et al. (2008) Acc homoeoloci and the evolution of wheat genomes. *Proc Natl Acad Sci USA* 105:9691–9696.
- Huang S, et al. (2002) Genes encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the *Triticum/Aegilops* complex and the evolutionary history of polyploid wheat. *Proc Natl Acad Sci USA* 99:8133–8138.
- Kihara H (1944) Discovery of the DD-analyser, one of the ancestors of vulgare wheats. *Ag. Hort. (Tokyo)* 19:889–890.
- McFadden ES, Sears ER (1946) The origin of *Triticum spelta* and its free-threshing hexaploid relatives. *J Hered* 37:81–89, 107.
- Nesbitt M, Samuel D (1996) From the staple crop to extinction? The archaeology and history of hulled wheats. *Hulled Wheats. Proceedings of the First International Workshop on Hulled Wheats*, eds Padulosi S, Hammer K, Heller J (International Plant Genetic Resources Institute, Rome), pp 41–100.
- Blake NK, Lehfelddt BR, Lavin M, Talbert LE (1999) Phylogenetic reconstruction based on low copy DNA sequence data in an allopolyploid: The B genome of wheat. *Genome* 42:351–360.
- Dvorák J, Terlizzi P, Zhang H-B, Resta P (1993) The evolution of polyploid wheats: Identification of the A genome donor species. *Genome* 36:21–31.
- Dvorák J, Zhang HB (1990) Variation in repeated nucleotide sequences sheds light on the phylogeny of the wheat B and G genomes. *Proc Natl Acad Sci USA* 87:9640–9644.
- Riley R, Unrau J, Chapman V (1958) Evidence on the origin of the B genome of wheat. *J Hered* 49:91–98.
- Takumi S, Nasuda S, Liu YG, Tsunewaki K (1993) Wheat phylogeny determined by RFLP analysis of nuclear DNA. I. Einkorn wheat. *Jpn J Genet* 68:73–79.
- Talbert LE, Blake NK, Storlie EW, Lavin M (1995) Variability in wheat based on low-copy DNA sequence comparisons. *Genome* 38:951–957.
- Simons KJ, et al. (2006) Molecular characterization of the major wheat domestication gene *Q*. *Genetics* 172:547–555.
- Faris JD, Simons KJ, Zhang Z, Gill BS (2005) The wheat super domestication gene *Q*. *Frontiers of wheat Bioscience. Wheat Information Service* 100(Memorial Issue): 129–148.
- Faris JD, Gill BS (2002) Genomic targeting and high-resolution mapping of the domestication gene *Q* in wheat. *Genome* 45:706–718.
- Sharma HC, Waines JG (1982) Attempted gene transfer from tetraploids to diploids in *Triticum*. *Can J Genet Cytol* 23:639–645.
- Faris JD, Zhang Z, Fellers JP, Gill BS (2008) Micro-colinearity between rice, Brachypodium, and *Triticum monococcum* at the wheat domestication locus *Q*. *Funct Integr Genomics* 8:149–164.
- Salse J, et al. (2008) New insights into the origin of the B genome of hexaploid wheat: Evolutionary relationships at the SPA genomic region with the S genome of the diploid relative *Aegilops speltoides*. *BMC Genomics* 9:555.
- Chantret N, et al. (2005) Molecular basis of evolutionary events that shaped the hardness locus in diploid and polyploid wheat species (*Triticum* and *Aegilops*). *Plant Cell* 17:1033–1045.
- Dvorak J, Akhunov ED, Deal KR, Luo MC (2006) Molecular characterization of a diagnostic DNA marker for domesticated tetraploid wheat provides evidence for gene flow from wild tetraploid wheat to hexaploid wheat. *Mol Biol Evol* 23:1386–1396.
- Isidore E, Scherrer B, Chalhoub B, Feuillet C, Keller B (2005) Ancient haplotypes resulting from extensive molecular rearrangements in the wheat A genome have been maintained in species of three different ploidy levels. *Genome Res* 15:526–536.
- Barbazuk WB, Fu Y, McGinnis KM (2008) Genome-wide analyses of alternative splicing in plants: opportunities and challenges. *Genome Res* 18:1381–1392.
- Zou C, et al. (2009) Evolutionary and expression signatures of pseudogenes in *Ara-bidopsis* and rice. *Plant Physiol* 151:3–15.
- Faris JD, Fellers JP, Brooks SA, Gill BS (2003) A bacterial artificial chromosome contig spanning the major domestication locus *Q* in wheat and identification of a candidate gene. *Genetics* 164:311–321.
- Endo TR, Gill BS (1996) The deletion stocks of common wheat. *J Hered* 87:295–307.
- Birchler JA, Riddle NC, Auger DL, Veitia RA (2005) Dosage balance in gene regulation: Biological implications. *Trends Genet* 21:219–226.
- Kato K, Miura H, Sawada S (1999) QTL mapping of genes controlling ear emergence time and plant height on chromosome 5A of wheat. *Theor Appl Genet* 98:472–477.
- Muramatsu M (1963) Dosage effect of the spelta gene *Q* of hexaploid wheat. *Genetics* 48:469–482.