

A test for pre-adapted phenotypic plasticity in the invasive tree Acer negundo L.

Laurent Lamarque L.J., Annabel A. Porte, Camille Eymeric, Jean-Baptiste Lasnier, Christopher J. Lortie, Sylvain S. Delzon

▶ To cite this version:

Laurent Lamarque L.J., Annabel A. Porte, Camille Eymeric, Jean-Baptiste Lasnier, Christopher J. Lortie, et al.. A test for pre-adapted phenotypic plasticity in the invasive tree Acer negundo L.. PLoS ONE, 2013, 8 (9), 10 p. 10.1371/journal.pone.0074239 . hal-02651023

HAL Id: hal-02651023 https://hal.inrae.fr/hal-02651023v1

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



A Test for Pre-Adapted Phenotypic Plasticity in the Invasive Tree *Acer negundo* L.

Laurent J. Lamarque^{1,2,3*}, Annabel J. Porté^{2,3}, Camille Eymeric^{2,3}, Jean-Baptiste Lasnier^{2,3}, Christopher J. Lortie¹, Sylvain Delzon^{2,3}

1 Department of Biology, York University, Toronto, Ontario, Canada, 2 University of Bordeaux, UMR 1202 BIOGECO, Talence, France, 3 INRA, UMR 1202 BIOGECO, Cestas, France

Abstract

Phenotypic plasticity is a key mechanism associated with the spread of exotic plants and previous studies have found that invasive species are generally more plastic than co-occurring species. Comparatively, the evolution of phenotypic plasticity in plant invasion has received less attention, and in particular, the genetic basis of plasticity is largely unexamined. Native from North America, *Acer negundo* L. is aggressively impacting the riparian forests of southern and eastern Europe thanks to higher plasticity relative to co-occurring native species. We therefore tested here whether invasive populations have evolved increased plasticity since introduction. The performance of 1152 seedlings from 8 native and 8 invasive populations was compared in response to nutrient availability. Irrespective of nutrients, invasive populations had higher growth and greater allocation to above-ground biomass relative to their native conspecifics. More importantly, invasive genotypes did not show increased plasticity in any of the 20 traits examined. This result suggests that the high magnitude of plasticity to nutrient variation of invasive seedlings might be pre-adapted in the native range. Invasiveness of *A. negundo* could be explained by higher mean values of traits due to genetic differentiation rather than by evolution of increased plasticity.

Citation: Lamarque LJ, Porté AJ, Eymeric C, Lasnier J-B, Lortie CJ, et al. (2013) A Test for Pre-Adapted Phenotypic Plasticity in the Invasive Tree Acer negundo L. PLoS ONE 8(9): e74239. doi:10.1371/journal.pone.0074239

Editor: Gabriele Sorci, CNRS, Université de Bourgogne, France

Received April 15, 2013; Accepted July 31, 2013; Published September 9, 2013

Copyright: © 2013 Lamarque et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was supported by NSERC DG and Canada Foundation for Innovation grant to CJL. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Christopher J. Lortie serves as an editor for this journal. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

1

* E-mail: llamarqueab@gmail.com

Introduction

Phenotypic plasticity has been widely recognized as an important feature for plants to cope with environmental changes [1,2]. Numerous studies have shown that plants are plastic for a large array of traits related to structure, development, metabolic activity, morphology, physiology, phenology, and reproduction [3-10]. Phenotypic plasticity has also been classified as a major determinant of the success of invasive species by increasing fitness relative to native species in recipient communities [11,12,13]. Broader distributions of alien species are correlated with higher levels of plasticity in response to increasing resource availability [14], and on artificial gradients, invasive species are also more plastic than co-occurring native or non-invasive species [15–20]. Davidson et al. [21] recently synthesized this work via a metaanalysis of 75 pairs of invasive/native species concluding that invasive species do express greater phenotypic plasticity than native species irrespective of the response traits measured. However, there are instances that did not support this pattern [22-25], and it has been suggested that the success and fitness advantage of invasive species can be mediated by the expression of constant higher mean trait values across different environments and not necessarily by the plasticity of these traits [26-29]. For instance, invasive species frequently possess higher trait values for growth rate [30,31,32], lower leaf mass per area [30,33,34], and advanced leaf unfolding and flowering periods [35,36,37].

Phenotypic plasticity should therefore be considered in combination with mean trait values when testing for plant invasiveness.

Higher plasticity levels of invasive species are generally hypothesized to be related to post-introduction evolution of phenotypic plasticity [13,38,39]. In this regard, intraspecific contrasts across environmental gradients have been analyzed in the following cases: (i) between populations from native and invasive ranges [40,41] and (ii) between populations within the invasive range [12,42,43]. Overall, no general pattern has emerged to date. Invasive populations of Senecio inaequidens were for instance more plastic than their native conspecifics in response to fertilization [44] while no difference was observed between native and invasive populations of Microstegium vimineum across a large array of environments [45]. Variation in light and soil moisture availability induced differences in plasticity for aboveground biomass and leaf mass per area among invasive populations of Microstegium vimineum but not for reproductionrelated traits among invasive populations of Polygonum cespitosum [42,43]. Pre-adapted plasticity might therefore be a common feature of several invasive plants, and it should now be more explicitly tested.

A total of 357 tree species has been reported to be invasive worldwide disrupting major native ecosystem structure and functioning [46,47,48]. Invasive trees are thus appropriate models to evaluate the role of ecological and evolutionary processes in invasion given their large impacts, frequency, and longevity [49–

Table 1. The 16 source populations sampled to examine phenotypic plasticity of invasive tree species Acer negundo L.

Province/Department	Collection site	River	Latitude	Longitude	Distance to the nearest population (km)
Native range					
Ontario	Paris	Grand	43°12′27″N	80°21′58″W	65
Ontario	Fergus	Grand	43°41′53″N	80°22′50″W	65
Ontario	Nicolston	Nottawasaga	44°10′40″N	79°49′02″W	18
Ontario	Angus	Nottawasaga	44°18′59″N	79°53′08″W	18
Ontario	Toronto Home Smith park	Humber	43°39′06″N	79°29′44″W	26
Ontario	Toronto Serena Gundy park	Don	43°43′05″N	79°21′15″W	26
Quebec	Sherbrooke	Saint-François	45°23′44″N	71°52′50″W	24
Quebec	Windsor	Saint-François	45°34′04″N	72°00′23″W	24
Invasive range					
Landes	Saubusse	Adour	43°39′22″N	01°11′13″W	10
Landes	Riviere-Saas-et-Gourby	Adour	43°40′29″N	01°08′06″W	10
Landes	Pontonx-sur-l'Adour	Adour	43°47′03″N	00°55′30″W	35
Gironde	Cestas	Eau Bourde	44°45′20″N	00°40′49″W	30
Gironde	Bruges	Les Jalles	44°54′13″N	00°36′16″W	30
Gironde	Moulon	Dordogne	44°51′30″N	00°13′10″W	19
Gironde	Castillon-la-Bataille	Dordogne	44°51′04″N	00°02′16″W	19
Gironde	St-Denis-de-Pile	Isle	44°59′34″N	00°12′29″W	22

doi:10.1371/journal.pone.0074239.t001

53]. To date, most studies examining the importance of phenotypic plasticity in tree invasion compared invasive vs. native tree species [15,39,54,55]. With the exception of *Melaleuca quinquenervia* and *Triadica sebifera* [39,56], the genetic differences in plasticity between native and invasive populations of exotic trees are unexamined.

Porté et al. [57] recently found that the invasive tree Acer negundo significantly expressed higher magnitude of phenotypic plasticity than its co-occurring native species with increasing environmental resources, and particularly nutrient availability. The purpose of this study was therefore to examine the genetic basis of plasticity in A. negundo, i.e. to determine whether higher plasticity of invasive populations is due to post-introduction evolution or pre-adaptation in the native range. The performance of native and invasive populations of A. negundo was compared across a gradient of nutrient availability. Life-history traits related to growth, physiology, leaf morphology and biomass and known to promote plant invasiveness [31,32] were measured. We hypothesize that invasive populations possess greater plasticity in growth and associated traits relative to populations from the native range. These findings would support the idea that plasticity could have evolved in the introduced range. In contrast, the absence of difference in plasticity between populations from native and invasive ranges would indicate that higher plasticity of invasive populations of A. negundo could be due to pre-adaptation in the native range.

Materials and Methods

Studied Species

Acer negundo L. (Box Elder or Manitoba maple) is a widely distributed mid-successional species native to North America. Its distribution range extends from southern Alberta and central Manitoba to Mexico and Guatemala southward and from central Montana to New England states and central Florida eastward [58,59,60]. This species is frequently found in floodplains and

riparian habitats but can also occur in dry coniferous forests, oak savannas, and grasslands [61,62]. *A. negundo* was intentionally introduced in the Old Continent at the end of the seventeenth century, *i.e.* in 1688 in England, and in France in 1749 [60,63]. Currently, *A. negundo* is highly invasive throughout southern, central and eastern Europe [60,64]. It frequently occurs not only in riparian habitats characterized by high rate of flood disturbance and high soil nutrient level [65,66] but also under drier conditions along roadsides, industrial wastelands, and dry ruderal sites [52,67].

Experimental Design

Seeds of A. negundo were harvested between September and November 2009 from eight native populations sampled in Ontario and Quebec, Canada and from eight invasive populations located in the Landes and Gironde departments of Aquitaine region, Southern France (Table 1). No specific permissions were required for these locations that are not part of protected areas and do not involve endangered species. All native and invasive populations were sampled from riparian forests. Populations in the invasive range were distributed within the Adour-Garonne river basin. Seeds came from 9 to 12 maternal trees in each source populations with maternal trees randomly selected and at least 10 m apart. In February 2010, 30 seeds per maternal tree were subjected to a cold treatment (14 weeks at 5°C in a cold chamber) at the INRA research station of Pierroton, France (44°44′N, 0°46′W). In spring 2010, 27 seeds per maternal tree were sown into 4 L (15×15×17.7 cm) pots filled with a commercial sphagnum peat soil mixture (organic matter 80%, pH=6). We first sowed three seeds per pot until germination and then kept one seedling in each pot thereby generating a total of 90 seedlings per source population. Pots were then placed under a greenhouse that was side-opened to permit wind and insects to enter. We did not




Figure 1. Mean \pm SE of life-history traits for native and invasive seedlings of *Acer negundo*. Differences in growth (A,B), physiology (C,D), leaf morphology (E), biomass (F,G) and biomass allocation (H) were calculated across nutrient levels. n = 576 (height and diameter), 96 (A_{area}) and 144 (N_{area} , LMA, W_t and A_l) per range. See text for definition of terms. * P < 0.05, ** P < 0.01, *** P < 0.001. doi:10.1371/journal.pone.0074239.g001

control light and temperature that approximated ambient conditions. Seedlings were watered twice a week to saturation.

A split-plot design was used with nutrient level as the fixed main effect and range of *A. negundo* populations (native or invasive) as the fixed sub-effect with all native and invasive populations subjected to three nutrient levels. We selected 72 seedlings from 8 to 10

families (*i.e.* maternal trees) per population for a total of 1152 seedlings structured as follows: 6 blocks \times 3 nutrient levels \times 2 ranges \times 8 populations \times 4 individuals. The experiment was initiated on February 17th, 2011 and lasted 147 days. Nutrients were applied on the 25th, 53th, 81st and 109th days of the experiment. The nutrient treatment corresponded to the addition

Table 2. Generalized linear mixed models (GLMM) analyses of traits related to growth, gas exchange and leaf morphology, biomass and biomass allocation in eight native and eight invasive populations of *Acer negundo* along a nutrient gradient.

		Source of variation									
		Nutrient		Range		Nutrie range	nt ×	Popul (range		Nutrie	ent $ imes$ population (range)
Traits	AIC	F	P	F	P	F	Р	LLR	Р	LLR	Р
Growth											
Diameter (mm)	5237.0	55.93	<.0001	5.92	0.0289	0.14	0.8718	12.3	0.0004	2.9	0.0885
Height (cm)	11004.7	39.43	<.0001	39.33	<.0001	2.37	0.1123	5.3	0.0213	4.2	0.0404
Leaf traits											
A_{area} (µmol CO ₂ .m ⁻² .s ⁻¹)	784.7	38.09	<.0001	0.04	0.8392	0.00	0.9980	2.7	0.1003	5.2	0.0226
A_{mass} (µmol CO ₂ .g ⁻¹ .s ⁻¹)	-467.9	31.78	<.0001	0.06	0.8098	0.24	0.7876	0.3	0.5839	0.9	0.3428
N _{area} (g N.m ⁻²)	1.9	116.61	<.0001	28.96	<.0001	2.95	0.0543	0.0	1	0.0	1
N _{mass} (%)	486.0	194.18	<.0001	7.58	0.0087	0.13	0.8772	0.5	0.4795	2.8	0.0943
PNUE (µmol CO ₂ .g ⁻¹ N.s ⁻¹)	929.2	17.91	<.0001	2.86	0.1135	0.03	0.9748	2.8	0.0943	0.6	0.4386
LMA (g.m ⁻²)	2032.0	0.97	0.4069	8.94	0.0098	0.10	0.9087	0.7	0.4028	1.1	0.2943
L _s (cm ²)	2343.6	14.44	<.0001	20.21	0.0005	0.31	0.7393	0.8	0.3711	0.2	0.6547
Biomass											
W_{t} (g)	2590.9	17.06	<.0001	0.02	0.8985	0.23	0.7928	0.0	1	0.9	0.3428
$W_{\rm a}$ (g)	2454.3	18.93	0.0002	1.24	0.2709	0.25	0.7803	0.0	1	1.5	0.2207
W _I (g)	1559.5	37.29	<.0001	0.16	0.6915	0.11	0.8944	0.0	1	1.0	0.3173
W_{s} (g)	2338.2	14.51	0.0006	1.60	0.2132	0.35	0.7099	0.0	1	1.5	0.2207
$W_{\rm r}$ (g)	1886.0	9.47	0.0001	10.74	0.0059	0.24	0.7837	0.0	1	1.1	0.2943
$A_{\rm I}~({\rm m}^2)$	-609.1	14.42	0.0012	5.62	0.0326	1.45	0.2355	1.3	0.2542	0.5	0.4795
Biomass allocation											
RSR $(g.g^{-1})$	-308.2	20.25	<.0001	54.33	<.0001	0.38	0.6846	0.7	0.4028	3.0	0.0833
LWR (g leaf.g ⁻¹ plant)	-1011.7	32.35	<.0001	5.22	0.0277	1.74	0.1893	0.7	0.4028	0.7	0.4028
SWR (g stem.g ⁻¹ plant)	-772.7	0.06	0.9401	66.33	<.0001	2.43	0.1068	1.6	0.2060	4.2	0.0404
RWR (g root.g ⁻¹ plant)	-705.8	17.63	<.0001	53.89	<.0001	0.11	0.8943	1.3	0.2542	3.7	0.0544
LAR (m ² leaf.g ⁻¹ leaf)	-2577.3	29.76	<.0001	9.40	0.0083	0.79	0.4631	0.1	0.7518	2.9	0.0886

F values are given for fixed effects while log likelihood ratios (LLR) are given for random effects. Statistically significant values (P<0.05) are shown in bold and marginally significant values (P<0.1) are shown in italic. The Akaike Information Criterion (AIC) value of the model used is given for each variable. See text for definition of terms. doi:10.1371/journal.pone.0074239:t002

of the complete slow release 16-7-15 (NPK plus micronutrients) fertiliser Floranid Permanent (Compo France SAS, Levallois-Perret, France). In the low nutrient level (N0), seedlings did not receive any additional fertilizer. In the medium and high nutrient levels (N1 and N2, respectively), seedlings received four fertilizer doses equivalent to 0.125 g and 0.500 g N each, for a total of 0.500 g and 2 g N, respectively. The high nutrient level corresponded to the nutrient availability encountered by *A. negundo* populations in soils of the invaded riparian habitats of southern France [68,69]. A previous study conducted *in situ* also showed that invasive individuals of *A. negundo* had a leaf N content averaging 1.17 gN.m⁻² [57]. The N0 and N1 treatments thus represent levels of nutrient that are below the average field conditions in the introduced range.

Gas Exchange

Photosynthetic rate measurements were performed on 192 seedlings. In each treatment, four individuals from different families and blocks were randomly sampled per source population. The measurements were done on sunny days between June 20th and July 7th. Leaf gas exchange measurements were carried out with a portable steady-state flow-through chamber (PLC6)

connected to an infrared gas analyser (CIRAS-2, PP Systems, Hitchin, UK) equipped with temperature, humidity, light and CO₂ control modules. Net gas exchanges were measured within a sealed cuvette of 2.5 cm², with an air CO₂ concentration of 380±3 ppm, a temperature of 22±0.5°C and a relative humidity of $80\pm10\%$ of ambient, controlled by regulating the flow diverted through a desiccant. To obtain the maximum assimilation rate per unit leaf area (A_{area} , μ mol CO_2 .m⁻².s⁻¹) at ambient CO_2 , leaves were illuminated with a red-blue light source attached to the gas exchange system and maintained at saturated light (PPFD = 1500 μ mol PAR.m⁻².s⁻¹). Prior to the measurements, the gas analyser was calibrated in the laboratory using 400 ppm standard gas, while full CO2 and H2O zero and differential calibrations were performed in the field after each set of six measurements. Up to three measurements were carried out on each sampled individual, and data were recorded when assimilation curves remained stable for more than 20 s. All measurements were taken between 8.00 and 11.00 solar time on fully expanded and sun-exposed leaves to avoid midday stomatal closure.




Figure 2. Plasticity of native and invasive seedlings of *Acer negundo* to nutrient availability. Means \pm SE of traits related to growth (A,B), physiology (C,D), leaf morphology (E), biomass (F,G) and biomass allocation (H) are represented. n = 192 (height and diameter), 32 (A_{area}) and 48 (N_{area} , LMA, W_t and A_l) per range and nutrient level. See text for definition of terms. doi:10.1371/journal.pone.0074239.g002

Leaf Morphology and Biochemistry

Leaf nitrogen content and morphological traits were measured on 288 seedlings representing six individuals per population and per treatment (including those used for gas exchange measurements). Leaves were sampled on the same days as the photosynthetic rate measurements. Three to five leaves were collected per sampled individual. Leaf surface area was measured with a planimeter (Light Box model, Gatehouse, Scientific Instruments LTD, Norfolk, UK) and the average leaf size (L_s, cm²) was

calculated. Leaves were then placed in an oven at 65°C until constant dry weight and leaf dry mass was later weighed with an electronic weighing scale (Explorer Pro, EP 114 model, Ohaus Corporation, Pine Brook, NJ, USA). Leaf mass per area index (LMA, g leaf.m $^{-2}$ leaf) was calculated as the ratio of leaf weight by leaf area. Finally, leaf samples were crushed to a powder with a ball mill (MM 200, Fisher Bioblock Scientific, France) and leaf nitrogen content (N_{mass}, %) was determined using an elementary analyser Eager 300 CHNOS (FlashEA 1112, ThermoElectron

Table 3. Relative Distance Plasticity Index (RDPI) along a nutrient gradient for populations of *Acer negundo* from the native and invasive ranges.

	RDPI low-to-mediu	m nutrient levels	RDPI medium-to-high nutrient levels			
Traits	Invasive	Native	Invasive	Native		
Growth						
Height	0.12±0.02	0.10±0.02	0.04±0.01	0.05 ± 0.01		
Diameter	0.08 ± 0.01	$0.07\!\pm\!0.01$	0.04 ± 0.01	0.04 ± 0.01		
Leaf traits						
A _{area}	$0.29\!\pm\!0.04$	$0.28\!\pm\!0.04$	0.19 ± 0.03	0.14 ± 0.04		
A _{mass}	0.24 ± 0.04	0.25±0.06	0.15±0.03	0.11±0.03		
N_{area}	0.24 ± 0.03	0.22 ± 0.02	$0.27\!\pm\!0.02$	0.28 ± 0.04		
N _{mass}	0.20 ± 0.04	0.19±0.03	0.27±0.02	0.25 ± 0.03		
PNUE	0.11 ± 0.04	0.14 ± 0.04	$0.21\!\pm\!0.05$	0.22 ± 0.04		
LMA	0.06 ± 0.02	0.09±0.02	$0.03\!\pm\!0.01$	0.10±0.02*		
L _s	0.11 ± 0.01	$0.07\!\pm\!0.03$	0.11 ± 0.03	0.08 ± 0.02		
Biomass						
W_{t}	$0.23\!\pm\!0.04$	$0.26 \!\pm\! 0.06$	$0.08\!\pm\!0.02$	0.14 ± 0.04		
W _a	0.24 ± 0.04	0.28±0.07	0.09 ± 0.03	0.15±0.05		
W_{I}	$0.28\!\pm\!0.06$	$0.31\!\pm\!0.08$	$0.12 \!\pm\! 0.04$	0.15 ± 0.04		
W_{s}	$0.23\!\pm\!0.04$	0.28±0.07	0.09 ± 0.02	0.16±0.05		
W _r	$0.21\!\pm\!0.05$	$0.20\!\pm\!0.05$	$0.09\!\pm\!0.04$	0.11 ± 0.03		
A _I	0.27±0.06	0.28±0.08	0.11±0.03	0.10±0.02		
Biomass allocation						
RSR	0.15±0.03	0.12±0.04	0.09±0.03	0.10±0.01		
LWR	0.11 ± 0.04	$0.09 \!\pm\! 0.02$	$0.08\!\pm\!0.04$	$0.06 \!\pm\! 0.04$		
SWR	0.04±0.01	0.04±0.01	0.02±0.01	0.03±0.01		
RWR	0.11±0.02	0.09±0.03	$0.07\!\pm\!0.02$	0.07±0.01		
LAR	0.13±0.04	0.13±0.03	0.12±0.03	0.16±0.03		

Comparisons of RDPI using a Generalized Linear Mixed Model with range as a fixed factor and population nested within range as a random factor. Significant difference between ranges (P<0.05) denoted by an asterisk. See text for definition of terms. doi:10.1371/journal.pone.0074239.t003

Corporation, Waltham, MA, USA). The maximum assimilation rate per unit leaf mass ($A_{\rm mass}$, μ mol CO_2 .g $^{-1}$.s $^{-1}$) was calculated as the $A_{\rm area}$ to LMA ratio, the leaf nitrogen content per leaf area ($N_{\rm area}$, g $N.m^{-2}$) as the product of $N_{\rm mass}$ and LMA, and the photosynthetic N-use efficiency (PNUE, μ mol CO_2 .g $^{-1}$ $N.s^{-1}$) as the $A_{\rm area}$ to $N_{\rm area}$ ratio.

Growth and Biomass

A total of seven individuals died during the course of the experiment and therefore, final height and stem collar diameter of 1145 seedlings were recorded on July 4th. A graduated pole to 0.01 m accuracy was used to record heights, and diameters were measured with an electronic calliper to 0.01 mm accuracy. The 288 individuals previously used for morphological measurements were harvested on July 14th after 147 days of growth. Aboveground biomass was separated into stems and leaves, and roots were separated from soil and washed. Biomass was oven-dried at 65°C until constant dry weight and further weighed using an electronic weighing scale (Explorer Pro, EP 114 model, Ohaus Corporation, Pine Brook, NJ, USA). The following traits were calculated: total biomass (W_t , g), above-ground biomass (W_a , g), leaf biomass (W_b , g), stem biomass (W_s , g), root biomass (W_r , g), total leaf area (A_b , m²), root:shoot ratio (RSR, g.g⁻¹), leaf weight ratio (LWR, g leaf.g⁻¹ plant), stem weight ratio (SWR, g stem.g⁻¹

plant), root weight ratio (RWR, g root.g⁻¹ plant) and leaf area ratio (LAR, m² leaf.g⁻¹ leaf).

Statistical Analyses

Differences in traits were tested with a generalized linear mixed model that was fit to a split-plot design (procedure MIXED, REML method in SAS, version 9.2, SAS Institute, Cary, NC, USA) [70]. We used nutrient level, range, and the interaction of nutrient level × range as fixed factors whilst block, block × range, population nested within range, and the interaction of nutrient level x population nested within range were treated as random factors. To account for the influence of plant size on biomass allocation [71], we used total biomass as a covariate when we tested the following traits: A₁, RSR, LWR, SWR, RWR and LAR. Type III sums of squares were used for the calculation of F statistics. Random effects were further evaluated using a log likelihood ratio (LLR) test from the full and reduced models. All factors were identified significant at alpha <0.05. A significant range effect for a given trait indicated an overall genetic differentiation between seedlings from native and invasive populations. Moreover, phenotypic plasticity was examined here at the population-level [13,72]. A significant effect of nutrient level indicated plasticity for a given trait. The difference in plasticity of a given trait between seedlings from native and invasive populations

Table 4. Plasticity Index (PI; Valladares et al. 2000) along a nutrient gradient for populations of *Acer negundo* from the native and invasive ranges.

	PI low-to-medium nu	utrient levels	PI medium-to-high nutrient levels			
Traits	Invasive	Native	Invasive	Native		
Growth						
Height	-0.22 ± 0.03	-0.17 ± 0.03	-0.04 ± 0.03	-0.01 ± 0.04		
Diameter	-0.14 ± 0.02	-0.13 ± 0.02	-0.06 ± 0.03	$-0.07\!\pm\!0.02$		
Leaf traits						
A _{area}	-0.44 ± 0.04	-0.43 ± 0.05	-0.14 ± 0.11	-0.14 ± 0.09		
A _{mass}	-0.38 ± 0.04	-0.38 ± 0.09	-0.13 ± 0.10	-0.07 ± 0.08		
N_{area}	-0.38 ± 0.03	-0.35 ± 0.03	$-0.43\!\pm\!0.02$	-0.43 ± 0.05		
N _{mass}	-0.31 ± 0.07	-0.32 ± 0.05	$-0.43\!\pm\!0.02$	-0.39 ± 0.04		
PNUE	-0.14 ± 0.07	-0.17 ± 0.09	$0.33\!\pm\!0.07$	0.35 ± 0.06		
LMA	-0.06 ± 0.05	-0.02 ± 0.07	0.01 ± 0.03	-0.02 ± 0.08		
L_s	-0.12 ± 0.06	-0.08 ± 0.07	-0.17 ± 0.05	-0.09 ± 0.05		
Biomass						
W_{t}	$-0.30 \!\pm\! 0.09$	-0.35 ± 0.10	$-0.07\!\pm\!0.06$	$-0.07\!\pm\!0.10$		
$W_{\rm a}$	-0.32 ± 0.10	-0.39 ± 0.09	-0.10 ± 0.06	-0.10 ± 0.11		
W_{I}	-0.40 ± 0.08	-0.43 ± 0.08	$-0.21\!\pm\!0.06$	-0.13 ± 0.10		
W_{s}	-0.29 ± 0.10	-0.37 ± 0.10	-0.07 ± 0.06	-0.09 ± 0.12		
$W_{\rm r}$	-0.23 ± 0.11	-0.28 ± 0.09	-0.01 ± 0.08	0.00 ± 0.08		
A_{I}	-0.40 ± 0.08	-0.39 ± 0.09	-0.18 ± 0.06	-0.13 ± 0.05		
Biomass allocation						
RSR	0.14±0.09	0.20±0.06	0.09 ± 0.06	0.11 ± 0.06		
LWR	-0.18 ± 0.06	-0.17 ± 0.03	-0.13 ± 0.04	-0.06 ± 0.04		
SWR	0.00±0.03	-0.05 ± 0.03	0.01 ± 0.02	-0.02 ± 0.03		
RWR	0.12±0.07	0.14 ± 0.05	0.06 ± 0.05	$0.07\!\pm\!0.04$		
LAR	-0.12±0.09	-0.12 ± 0.08	-0.15 ± 0.07	-0.03 ± 0.11		

Comparisons of PI using a Generalized Linear Mixed Model with range as a fixed factor and population nested within range as a random factor. See text for definition of terms.

doi:10.1371/journal.pone.0074239.t004

was reported when the interaction of nutrient level \times range was significant. The variation of trait of native and invasive seedlings was also reported as follows: $(1-(\text{trait}_{\text{env2}}/\text{trait}_{\text{env1}}))*100$. Lastly, we calculated the Relative Distance Plasticity Index (RDPI) [73], and the Plasticity Index (PI) [5] for two experimental nutrient level changes, low-to-medium and medium-to-high, as follows:

$$RDPI = \frac{|mean(env1) - mean(env2)|}{|mean(env1) + mean(env2)|}$$

 $PI = \frac{mean(env1) - mean(env2)}{max(mean(env1), mean(env2))}$

For each trait, the two indexes were calculated for each population using mean values in each treatment (*i.e.* nutrient level). The difference in RDPI and PI between native and invasive ranges was examined using a generalized linear mixed model with range as a fixed factor and population nested within range as a random factor.

Results

Overall Trends

Irrespective of nutrients, individuals of A. negundo from invasive populations expressed significantly greater heights and smaller diameters than their native conspecifics (significant range effect; Table 2; Fig. 1A, B; see Table S1 for means per treatment). There was no significant difference in maximum assimilation rates ($A_{\rm area}$ and A_{mass} ; Fig. 1C), and invasive seedlings had lower leaf nitrogen contents (N $_{\rm area}$ and N $_{\rm mass};$ Fig. 1D) and greater PNUE (Table 2). Invasive seedlings also had lower average leaf size and LMA (Table 2; Fig. 1E). There were no statistical differences in total and aboveground biomass (Table 2; Fig. 1F). Seedlings from invasive populations however allocated more resources to foliage than to roots, displaying greater A_{l} , LWR, SWR and LAR, and lower W_{r} , RSR and RWR compared to seedlings from native populations (Table 2; Fig. 1G, H). Significant genetic variations were found in height among invasive populations (within invasive range: LLR = 5.6, P = 0.018; within native range: LLR = 0.6, P = 0.44) and in diameter among native populations (within native range: LLR = 14.1, P = 0.0002; within invasive range: LLR = 0.5, P = 0.44).

Trait Plasticity

Seedlings of A. negundo responded significantly to increases in nutrients (significant nutrient effect for all traits but LMA and SWR; Table 2; Fig. 2) with increased growth, maximum assimilation rate, total biomass, and above-ground allocation (AGB, TLA, LWR, LAR) and decreased below-ground allocation (RSR, RWR). The change from low-to-medium nutrient conditions had a stronger effect on seedling trait values than the change from medium-to-high nutrient conditions. Individuals of A. negundo respectively showed a 19%, 44% and 35% increase in height, maximum assimilation rate and total biomass from low-to-medium nutrient conditions but a 2%, 18% and 9% increase from medium-to-high nutrient conditions (Fig. 2; see Tables 3 and 4 for trait RDPI and PI values). Across all populations, traits such as SWR and LMA showed low plasticity along the nutrient $RDPI_{SWR} = 0.03,$ gradient (mean $RDPI_{LMA} = 0.07$) while W_l , A_{area} and N_{area} exhibited larger changes (mean $RDPI_{M} = 0.21$, mean $RDPI_{Aarea} = 0.22$, mean $RDPI_{Narea} = 0.25$).

There were no significant differences in plasticity between seedlings from native and invasive populations for any traits (non-significant nutrient \times range effect; Table 2; Fig. 2). There was also no difference in RDPI or PI for any traits but the RDPI_LMA between medium and high nutrient levels did differ (Tables 3 and 4; across the whole gradient, mean trait RDPI = 0.15 and 0.14 and mean trait PI = -0.14 and -0.15 for native and invasive populations, respectively). The magnitude of plasticity differed at the population level for height, maximum assimilation rate, and SWR (significant nutrient \times population effect; Table 2).

Discussion

Higher magnitudes of plasticity relative to native species are common in invasive plants, particularly in invasive trees [15,21,74]. Nevertheless, these differences are not necessarily a product of post-introduction evolution and can also be explained by innate characteristics. This null hypothesis was tested and supported in this study using the highly invasive tree species *Acer negundo*. Although increased nutrient availability is a key component of tree recruitment dynamics [75,76], this artificial gradient tested here did not elicit differences in plasticity between native and invasive seedlings. Pre-adapted plasticity to nutrient availability is thus a reasonable explanation for the successful spread of this species, at least at this early stage of development.

The evolution of plasticity in invasive species is relatively infrequent and no consensus has been reached in the literature so far (Table S2). Variation in resource conditions lead to differences in plasticity between seedlings from native and invasive populations for perennials Centaurea stoebe and Taraxacum officinale and trees Melaleuca quinquenervia and Triadica sebifera [39,56,77,78] but not for the annual grass Microstegium vimineum, the biennnial forb Alliaria petiolata and the perennial shrub Clidenia hirta [40,45,79]. However, a rigorous assessment of the origin and importance of plasticity in plant invasion requires both inter- and intraspecific contrats [39]. In response to nutrient availability, invasive seedlings of A. negundo, which had shown increased plasticity relative to than their cooccurring native species across the same resource gradient [57], expressed here similar response for all life-history traits compared to their native conspecifics. Our results therefore reflect innate characteristics of plasticity that would be pre-adapted in the native range. This supports the outcome observed for Triadica sebifera in response to water availability: invasive seedlings exhibited greater growth than seedlings of native Schizachyrium scoparium but not than their native conspecifics [39]. The only other study that conducted both inter- and intraspecific comparisons across the same resource gradient did not find any difference in plasticity to CO_2 enrichment between native and invasive populations of *Eupatorium adenopherum* and the native congener *Eupatorium japonicum* [80].

Seedlings from native and invasive populations of A. negundo significantly differed in most of their traits across the gradient of nutrient availability. Invasive seedlings consistently exhibited higher values for traits associated with invasiveness, i.e. higher growth rate, lower LMA, and greater allocation to foliage [30,34]. This supports many other studies which posit that geneticallybased advantages in plant size and above-ground biomass for invasive over native genotypes may promote the success of invasive species [81-84]. For instance, invasive individuals of Melaleuca quinquenervia and Triadica sebifera also outperformed native congeners [39,56,85]. Interestingly, invasive seedlings of A. negundo did not achieve greater height growth via physiological advantages but only via a preferential allocation to foliage. Significant lower leaf nitrogen content and similar maximum assimilation rate were found here. This contradicts recent studies on the geneticallybased difference of functional traits in invasive plant species that showed higher values of physiological traits for invasive genotypes [86,87,88]. These divergences might be due to the rapid adaptation of A. negundo in its introduced range reflecting a change in adaptive strategy. Whilst plasticity may not have evolved de novo, it is possible that most of the traits conferring faster growth (such as greater allocation to above-ground biomass) may have done so to provide a competitive advantage over native species of recipient

Multi-species comparisons in the native range of exotic plant species showed that invasive aliens differed in traits but not in plasticity from their non-invasive alien congeners [26,28], and preadaptation of plasticity in invasive plant species might finally be more common than expected. Phenotypic plasticity is a common denominator for invasive plant species but tolerance of invasive genotypes across a broad range of conditions might rely more on a combination of life-history traits rather than on evolved plasticity in the introduced range. This would be the case for A. negundo since the species occupies wide and similar ranges of habitats such as wet-rich and dry-poor nutrient riparian forests both in North America and in Europe [52,61,89]. Furthermore, various mechanisms such as founder effects, multiple introductions, and selective pressures can drive genetic differentiation between native and invasive populations. Molecular analyses using neutral markers over large areas sampled including whole native and invasive ranges would thus be necessary to fully understand the role of these factors [90]. Given that there was no consistent variation in traits amongst populations from the invasive range, genetic data would provide valuable information on the origin of those populations sampled in French riparian areas, e.g. whether they have all undergone similar selective pressures or come from the same pool of native populations which were not sampled in this study (i.e. founder effects).

Conclusions

The origin of increased plasticity in invasive plant species is an important and relatively understudied set of hypotheses. Given the geographical scope of the populations we were able to sample herein, pre-adaptation is a more viable explanation for the high magnitude of plasticity of invasive A. negundo seedlings to variation in nutrient availability. Future studies should however test in the native range the response of native and invasive genotypes sampled at broader scales to a combination of abiotic factors in order to test more effectively both the importance of evolved versus

pre-adapted plasticity and increases in competitive ability of invasive species.

Supporting Information

Table S1 Mean \pm SE for traits related to growth, gas exchange and leaf morphology, biomass and biomass allocation of eight native and eight invasive populations of *Acer negundo* grown along a nutrient gradient. Sample sizes are n=24 for growth traits, n=4 for physiology traits and n=6 for leaf morphology and biomass related traits. See text for definition of terms.

(DOCX)

Table S2 Intraspecific comparisons of phenotypic plasticity in invasive plant species. Summary of studies comparing phenotypic plasticity between native and invasive populations of exotic plant species in response to variation in environmental conditions. Plasticity was reported for various traits

References

- Schlichting CD (1986) The evolution of phenotypic plasticity in plants. Annu Rev Ecol Syst 17: 667–693.
- Pigliucci M (2001) Phenotypic plasticity: beyond nature and nurture. Baltimore: John Hopkins University Press. 328 p.
- Pintado A, Valladares F, Sancho LG (1997) Exploring phenotypic plasticity in the lichen Ramalina capitata: morphology, water relations and chlorophyll content in north- and south-facing populations. Ann Bot 80: 345–353.
- Sultan SE (2000) Phenotypic plasticity for plant development, function and life history. Trends Plant Sci 5: 537–542.
- Valladares F, Wright SJ, Lasso E, Kitajima K, Pearcy RW (2000) Plastic phenotypic response to light of 16 congeneric shrubs from a Panamanian rainforest. Ecology 81: 1925–1936.
- Sultan SE (2001) Phenotypic plasticity for fitness components in *Polygonum* species of contrasting ecological breadth. Ecology 82: 328–343.
- González AV, Gianoli E (2004) Morphological plasticity in response to shading in three Convolvulus species of different ecological breadth. Acta Oecol 26: 185– 100
- Sánchez-Gómez D, Valladares F, Zavala MA (2006) Functional traits and plasticity underlying shade tolerance in seedlings of four Iberian forest tree species. Tree Physiol 26: 1425–1433.
- Portsmuth A, Niinemets U (2007) Structural and physiological plasticity in response to light and nutrients in five temperate deciduous woody species of contrasting shade tolerance. Funct Ecol 21: 61–77.
- Molina-Montenegro MA, Atala C, Gianoli E (2010) Phenotypic plasticity and performance of *Taraxacum officinale* (dandelion) in habitats of contrasting environmental heterogeneity. Biol Invasions 12: 2277–2284.
- Baker HG (1965) Characteristics and modes of origin of weeds. In: Baker HG, Stebbins GL, editors. The genetics of colonizing species. New York: Academic Press. 147–169.
- Sexton JP, McKay JK, Sala A (2002) Plasticity and genetic diversity may allow saltcedar to invade cold climates in North America. Ecol Appl 12: 1652–1660.
- Richards CL, Bossdorf O, Muth NZ, Gurevitch J, Pigliucci M (2006) Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. Ecol Lett 9: 981–993.
- Dawson W, Rohr RP, Van Kleunen M, Fischer M (2012) Alien plant species with a wider global distribution are better able to capitalize on increased resource availability. New Phytol 194: 959–967.
- Pattison RR, Goldstein G, Ares A (1998) Growth, biomass allocation and photosynthesis of invasive and native Hawaiian rainforest species. Oecologia 117: 449–459.
- Niinemets U, Valladares F, Ceulemans R (2003) Leaf-level phenotypic variability and plasticity of invasive Rhododendron ponticum and non-invasive Ilex aquifolium co-occurring at two contrasting European sites. Plant Cell Environ 26: 941–956
- Burns JH, Winn AA (2006) A comparison of plastic responses to competition by invasive and non-invasive congeners in the Commelinaceae. Biol Invasions 8: 797–807.
- Pan XY, Geng YP, Zhang WJ, Li B, Chen JK (2006) The influence of abiotic stress and phenotypic plasticity on the distribution of invasive *Alternanthera* philoxeroides along a riparian zone. Acta Oecol 30: 333–341.
- Zhao YJ, Qing H, Zhao CJ, Zhou CF, Zhang WG, et al. (2010) Phenotypic plasticity of Spartina alterniflora and Phragmites australis in response to nitrogen addition and intraspecific competition. Hydrobiologia 637: 143–155.
- Skalova H, Havlickova V, Pysek P (2012) Seedling traits, plasticity and local differentiation as strategies of invasive species of Impatiens in central Europe. Ann Bot 110: 1429–1438.

related to biomass (B), defense to herbivory (D), growth (G), leaf morphology (M), phenology (Pe), physiology (P) and reproduction (R).

(DOCX)

Acknowledgments

We thank Maurice Aulen and Mathieu Reveillas for their assistance with seed collection as well as Yann Guengant for his assistance with growth measurements. We are also grateful to Jean-Baptiste Lamy for his advice on statistical analyses and to the INRA experimental unit of Cestas-Pierroton, in particular Frederic Bernier and Henri Bignalet, for their logistical support.

Author Contributions

Conceived and designed the experiments: LJL AJP SD. Performed the experiments: LJL CE JBL. Analyzed the data: LJL AJP CJL SD. Wrote the paper: LJL AJP CJL SD.

- Davidson AM, Jennions M, Nicotra AB (2011) Do invasive species show higher phenotypic plasticity than native species and, if so, is it adaptive? A metaanalysis. Ecol Lett 14: 419–431.
- Peperkorn R, Werner C, Beyschlag W (2005) Phenotypic plasticity of an invasive acacia versus two native Mediterranean species. Funct Plant Biol 32: 933–944.
- Funk JL (2008) Differences in plasticity between invasive and native plants from a low resource environment. J Ecol 96: 1162–1173.
- Funk JL, Zachary VA (2010) Physiological responses to short-term water and light stress in native and invasive plant species in southern California. Biol Invasions 12: 1685–1694.
- Werner C, Zumkier U, Beyschlag W, Maguas C (2010) High competitiveness of a resource demanding invasive acacia under low resource supply. Plant Ecol 206: 83–96.
- Schlaepfer DR, Glattli M, Fischer M, van Kleunen M (2010) A multi-species experiment in their native range indicates pre-adaptation of invasive alien plant species. New Phytol 185: 1087–1099.
- Godoy O, Valladares F, Castro-Diez P (2011) Multispecies comparison reveals
 that invasive and native plants differ in their traits but not in their plasticity.
 Funct Ecol 25: 1248–1259.
- van Kleunen M, Schlaepfer DR, Glaettli M, Fischer M (2011) Preadapted for invasiveness: do species traits or their plastic response to shading differ between invasive and non-invasive plant species in their native range? J Biogeogr 38: 1294–1304.
- Godoy O, Valladares F, Castro-Diez P (2012) The relative importance for plant invasiveness of trait means, and their plasticity and integration in a multivariate framework. New Phytol 195: 912–922.
- Grotkopp E, Rejmanek M, Rost TL (2002) Toward a causal explanation of plant invasiveness: Seedling growth and life-history strategies of 29 pine (*Pinus*) species. Am Nat 159: 396–419.
- van Kleunen M, Weber E, Fischer M (2010) A meta-analysis of trait differences between invasive and non-invasive plant species. Ecol Lett 13: 235–245.
- Lamarque LJ, Delzon S, Lortie CJ (2011) Tree invasions: a comparative test of the dominant hypotheses and functional traits. Biol Invasions 13: 1969–1989.
- Grotkopp E, Rejmanek M (2007) High seedling relative growth rate and specific leaf area are traits of invasive species: Phylogenetically independent contrasts of woody angiospernis. Am J Bot 94: 526–532.
- Pysek P, Richardson DM (2007) Traits associated with invasiveness in alien plants: where do we stand? In: Nentwig W, editor. Biological Invasions. Berlin: Springer-Verlag. 97–125.
- Xu CY, Griffin KL, Schuster WSF (2007) Leaf phenology and seasonal variation
 of photosynthesis of invasive Berberis thunbergii (Japanese barberry) and two cooccurring native understory shrubs in a northeastern United States deciduous
 forest. Oecologia 154: 11–21.
- Shustack DP, Rodewald AD, Waite TA (2009) Springtime in the city: exotic shrubs promote earlier greenup in urban forests. Biol Invasions 11: 1357–1371.
- Wolkovich EM, Cleland EE (2011) The phenology of plant invasions: a community ecology perspective. Front Ecol Environ 9: 287–294.
- Bossdorf O, Auge H, Lafuma L, Rogers WE, Siemann E, et al. (2005) Phenotypic and genetic differentiation between native and introduced plant populations. Oecologia 144: 1–11.
- Zou JW, Rogers WE, Siemann E (2009) Plasticity of Sapium sebiferum seedling growth to light and water resources: Inter- and intraspecific comparisons. Basic Appl Ecol 10: 79–88.
- DeWalt SJ, Denslow JS, Hamrick JL (2004) Biomass allocation, growth, and photosynthesis of genotypes from native and introduced ranges of the tropical shrub Clidemia hirta. Oecologia 138: 521–531.

- Chun YJ, Collyer ML, Moloney KA, Nason JD (2007) Phenotypic plasticity of native vs. invasive purple loosestrife: A two-state multivariate approach. Ecology 88: 1499–1512.
- Droste T, Flory SL, Clay K (2010) Variation for phenotypic plasticity among populations of an invasive exotic grass. Plant Ecol 207: 297–306.
- Matesanz S, Horgan-Kobelski T, Sultan SE (2012) Phenotypic plasticity and population differentiation in an ongoing species invasion. PLoS ONE 7(9): e44955. doi:44910.41371/journal.pone.0044955.
- Bossdorf O, Lipowsky A, Prati D (2008) Selection of preadapted populations allowed Senecio inaequidens to invade Central Europe. Divers Distrib 14: 676–685.
- Flory SL, Long FR, Clay K (2011) Invasive Microstegium populations consistently outperform native range populations across diverse environments. Ecology 92: 2248–2257.
- Castro-Diez P, Gonzalez-Munoz N, Alonso A, Gallardo A, Poorter L (2009) Effects of exotic invasive trees on nitrogen cycling: a case study in Central Spain. Biol Invasions 11: 1973–1986.
- Lorenzo P, Gonzalez L, Reigosa MJ (2010) The genus Acacia as invader: the characteristic case of Acacia dealbata Link in Europe. Ann For Sci 67: 101.
- Richardson DM, Rejmanek M (2011) Trees and shrubs as invasive alien species

 a global review. Divers Distrib 17: 788–809.
- Siemann E, Rogers WE (2001) Genetic differences in growth of an invasive tree species. Ecol Lett 4: 514–518.
- Zou J, Rogers WE, Siemann E (2007) Differences in morphological and physiological traits between native and invasive populations of *Sapium sebiferum*. Funct Ecol 21: 721–730.
- Franks SJ, Pratt PD, Dray FA, Simms EL (2008) Selection on herbivory resistance and growth rate in an invasive plant. Am Nat 171: 678–691.
- Erfmeier A, Bohnke M, Bruelheide H (2011) Secondary invasion of Acer negundo: the role of phenotypic responses versus local adaptation. Biol Invasions 13: 1599–1614
- Delmas CEL, Delzon S, Lortie CJ (2011) A meta-analysis of the ecological significance of density in tree invasions. Community Ecol 12: 171–178.
- Schumacher E, Kueffer C, Edwards PJ, Dietz H (2009) Influence of light and nutrient conditions on seedling growth of native and invasive trees in the Seychelles. Biol Invasions 11: 1941–1954.
- Paquette A, Fontaine B, Berninger F, Dubois K, Lechowicz MJ, et al. (2012) Norway maple displays greater seasonal growth and phenotypic plasticity to light than native sugar maple. Tree Physiol 32: 1339–1347.
- Kaufman SR, Smouse PE (2001) Comparing indigenous and introduced populations of Melaleuca quinquenervia (Cav.) Blake: response of seedlings to water and pH levels. Oecologia 127: 487–494.
- Porté AJ, Lamarque LJ, Lortie CJ, Michalet R, Delzon S (2011) Invasive Acer negundo outperforms native species in non-limiting resource environments due to its higher phenotypic plasticity BMC Ecol 11: 28.
- Little EL Jr (1971) Atlas of United States trees: conifers and important hardwoods. Vol. 1. U.S. Department of Agriculture, Forest Service. Washington: Government Printing Office. 202 p.
- Little EL Jr (1979) Checklist of United States Trees (native and naturalized).
 Agriculture Handbook 541. U.S. Department of Agriculture, Forest Service.
 Washington: Government Printing Office. 375 p.
- Mędrzycki P (2007): NOBANIS Invasive Alien Species Fact Sheet Acer negundo – From: Online Database of the North European and Baltic Network on Invasive Alien Species – NOBANIS www.nobanis.org.
- Maeglin RR, Ohmann LF (1973) Boxelder (Acer negundo): A review and commentary. B Torrey Bot Club 100: 357–363.
- Dewine JM, Cooper DJ (2008) Canopy shade and the successional replacement of tamarisk by native box elder. J Appl Ecol 45: 505–514.
- 63. Kowarik I (2003) Biologische Invasionen: Neophyten und Neozoen in Mitteleuropa. Stuttgart: Eugen Ulmer. 380 p.
- 64. Lamarque LJ, Delzon S, Sloan MH, Lortie CJ (2012) Biogeographical contrasts to assess local and regional patterns of invasion: a case study with two reciprocally introduced exotic maple trees. Ecography 35: 803–810.
- Planty-Tabacchi AM, Tabacchi E, Naiman RJ, Deferrari C, Decamps H (1996) Invasibility of species-rich communities in riparian zones. Conserv Biol 10: 598–607.
- Tabacchi E, Planty-Tabacchi AM (2003) Recent changes in riparian vegetation: Possible consequences on dead wood processing along rivers. River Res Appl 19: 251–263.
- Rothmaler W (1984) Exkursionsflora fur die Gebiete der DDR und der BRD -Gefaßpflanzen, 12th edn. Gefaßpflanzen: Kritischer Band. Berlin: Volk und Wissen. 640 p.

- Pinay G, Black VJ, Planty-Tabacchi AM, Gumiero B, Decamps H (2000) Geomorphic control of denitrification in large river floodplain soils. Biogeo-chemistry 50: 163–182.
- Pinay G, Roques L, Fabre A (1993) Spatial and temporal patterns of denitrification in a riparian forest. J Appl Ecol 30: 581–591.
- Littell RC, Milliken GA, Stroup WW, Wolfinger RD, Schabenberger O (2006) SAS® for Mixed Models, Second Edition. Cary: SAS Institute Inc.
- Maestre FT, Quero JL, Valladares F, Reynolds JF (2007) Individual vs. population plastic responses to elevated CO2, nutrient availability, and heterogeneity: a microcosm experiment with co-occurring species. Plant Soil 296: 53–64.
- Williams JL, Auge H, Maron JL (2008) Different gardens, different results: native and introduced populations exhibit contrasting phenotypes across common gardens. Oecologia 157: 239–248.
- Valladares F, Sanchez-Gomez D, Zavala MA (2006) Quantitative estimation of phenotypic plasticity: bridging the gap between the evolutionary concept and its ecological applications. J Ecology 94: 1103–1116.
- Yamashita N, Koike N, Ishida A (2002) Leaf ontogenetic dependence of light acclimation in invasive and native subtropical trees of different successional status. Plant Cell Environ 25: 1341–1356.
- 75. Sayer EJ (2006) Using experimental manipulation to assess the roles of leaf litter in the functioning of forest ecosystems. Biol Rev $81\colon 1-31$.
- Salazar A, Goldstein G, Franco AC, Miralles-Wilhelm F (2012) Differential seedling establishment of woody plants along a tree density gradient in Neotropical savannas. J Ecol 100: 1411–1421.
- He WM, Thelen GC, Ridenour WM, Callaway RM (2010) Is there a risk to living large? Large size correlates with reduced growth when stressed for knapweed populations. Biol Invasions 12: 3591–3598.
- Molina-Montenegro MA, Quiroz CL, Torres-Diaz C, Atala C (2011) Functional differences in response to drought in the invasive *Taraxacum officinale* from native and introduced alpine habitat ranges. Plant Ecol Divers 4: 37–44.
- Hillstrom C, Cipollini D (2011) Variation in phenotypic plasticity among native and invasive populations of Alliaria petiolata. Int J Plant Sci 172: 763–772.
- Lei YB, Feng YL, Zheng YL, Wang RF, Gong HD, et al. (2011) Innate and evolutionarily increased advantages of invasive *Eupatorium adenophorum* over native *E. japonicum* under ambient and doubled atmospheric CO2 concentrations. Biol Invasions 13: 2703–2714.
- Bastlova D, Kvet J (2002) Differences in dry weight partitioning and flowering phenology between native and non-native plants of purple loosestrife (*Lythrum salicaria* L.). Flora 197: 332–340.
- Leger EA, Rice KJ (2003) Invasive California poppies (Eschscholzia californica Cham.) grow larger than native individuals under reduced competition. Ecol Lett 6: 257–264.
- Güsewell S, Jakobs G, Weber E (2006) Native and introduced populations of Solidago gigantea differ in shoot production but not in leaf traits or litter decomposition. Funct Ecol 20: 575–584.
- Lavergne S, Molofsky J (2007) Increased genetic variation and evolutionary potential drive the success of an invasive grass. P Nat Ac S USA 104: 3883— 2000
- Huang W, Siemann E, Wheeler GS, Zou JW, Carrillo J, et al. (2010) Resource allocation to defence and growth are driven by different responses to generalist and specialist herbivory in an invasive plant. J Ecol 98: 1157–1167.
- Feng YL, Lei YB, Wang RF, Callaway RM, Valiente-Banuet A, et al. (2009) Evolutionary tradeoffs for nitrogen allocation to photosynthesis versus cell walls in an invasive plant. P Nat Ac S USA 106: 1853–1856.
- Mozdzer TJ, Żieman JC (2010) Ecophysiological differences between genetic lineages facilitate the invasion of non-native *Phragmites australis* in North American Atlantic coast wetlands. J Ecol 98: 451–458.
- Feng YL, Li YP, Wang RF, Callaway RM, Valiente-Banuet A, et al. (2011) A
 quicker return energy-use strategy by populations of a subtropical invader in the
 non-native range: a potential mechanism for the evolution of increased
 competitive ability. J Ecol 99: 1116–1123.
- Ward JK, Dawson TE, Ehleringer JR (2002) Responses of Acer negundo genders to interannual differences in water availability determined from carbon isotope ratios of tree ring cellulose. Tree Physiol 22: 339–346.
- Dlugosch KM, Parker IM (2008) Founding events in species invasions: genetic variation, adaptive evolution, and the role of multiple introductions. Mol Ecol 17: 431–449.