



**HAL**  
open science

## Cognitive and emotional alterations are related to hippocampal inflammation in a mouse model of metabolic syndrome

Anne-Laure Dinel, Caroline Andre, Agnès Aubert, Guillaume Ferreira, Sophie Layé, Nathalie Castanon

### ► To cite this version:

Anne-Laure Dinel, Caroline Andre, Agnès Aubert, Guillaume Ferreira, Sophie Layé, et al.. Cognitive and emotional alterations are related to hippocampal inflammation in a mouse model of metabolic syndrome. PLoS ONE, 2011, 6 (9), 10 p. 10.1371/journal.pone.0024325 . hal-02651479

**HAL Id: hal-02651479**

**<https://hal.inrae.fr/hal-02651479>**

Submitted on 29 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Cognitive and Emotional Alterations Are Related to Hippocampal Inflammation in a Mouse Model of Metabolic Syndrome

Anne-Laure Dinel<sup>1,2</sup>, Caroline André<sup>1,2</sup>, Agnès Aubert<sup>1,2</sup>, Guillaume Ferreira<sup>1,2</sup>, Sophie Layé<sup>1,2</sup>, Nathalie Castanon<sup>1,2\*</sup>

**1** Nutrition et Neurobiologie Intégrée, INRA UMR 1286, Bordeaux, France, **2** University of Bordeaux, Bordeaux, France

## Abstract

Converging clinical data suggest that peripheral inflammation is likely involved in the pathogenesis of the neuropsychiatric symptoms associated with metabolic syndrome (MetS). However, the question arises as to whether the increased prevalence of behavioral alterations in MetS is also associated with central inflammation, i.e. cytokine activation, in brain areas particularly involved in controlling behavior. To answer this question, we measured in a mouse model of MetS, namely the diabetic and obese *db/db* mice, and in their healthy *db/+* littermates emotional behaviors and memory performances, as well as plasma levels and brain expression (hippocampus; hypothalamus) of inflammatory cytokines. Our results shows that *db/db* mice displayed increased anxiety-like behaviors in the open-field and the elevated plus-maze (i.e. reduced percent of time spent in anxiogenic areas of each device), but not depressive-like behaviors as assessed by immobility time in the forced swim and tail suspension tests. Moreover, *db/db* mice displayed impaired spatial recognition memory (hippocampus-dependent task), but unaltered object recognition memory (hippocampus-independent task). In agreement with the well-established role of the hippocampus in anxiety-like behavior and spatial memory, behavioral alterations of *db/db* mice were associated with increased inflammatory cytokines (interleukin-1 $\beta$ , tumor necrosis factor- $\alpha$  and interleukin-6) and reduced expression of brain-derived neurotrophic factor (BDNF) in the hippocampus but not the hypothalamus. These results strongly point to interactions between cytokines and central processes involving the hippocampus as important contributing factor to the behavioral alterations of *db/db* mice. These findings may prove valuable for introducing novel approaches to treat neuropsychiatric complications associated with MetS.

**Citation:** Dinel A-L, André C, Aubert A, Ferreira G, Layé S, et al. (2011) Cognitive and Emotional Alterations Are Related to Hippocampal Inflammation in a Mouse Model of Metabolic Syndrome. PLoS ONE 6(9): e24325. doi:10.1371/journal.pone.0024325

**Editor:** Michelle L. Block, Virginia Commonwealth University, United States of America

**Received:** May 13, 2011; **Accepted:** August 6, 2011; **Published:** September 16, 2011

**Copyright:** © 2011 Dinel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by Institut National de la Recherche Agronomique and Région Aquitaine (grant number 2008-1301-038; SL). ALD was supported by a doctoral fellowship from the Institut Danone. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: nathalie.castanon@bordeaux.inra.fr

## Introduction

For several decades, the prevalence of the metabolic syndrome (MetS) and related comorbidities continuously increases worldwide at alarming rate. This syndrome is defined as a constellation of interrelated metabolic dysregulations, including abdominal obesity, hypertension, hyperglycemia, insulin-resistance, leptin-resistance and hypercortisolemia [1–3].

Mounting evidence highlights that patients with MetS often experience a higher prevalence of mood symptoms and cognitive dysfunctions than the general age-matched population [4–6]. The overwhelming influence of metabolic and neuropsychiatric disorders considerably impairs the quality of life of MetS patients and results in incremental costs to health care systems around the world [7]. Actually, neuropsychiatric symptoms emerge as significant risk factors for aggravation of MetS and related health outcomes, particularly cardiovascular diseases and type 2 diabetes (T2D) [8–12]. Even so, relatively little is known about the pathophysiological mechanisms contributing to the development of neuropsychiatric symptoms in the context of MetS. Recent evidence suggests that some major biological systems, including

the inflammatory system, may participate in both MetS and neuropsychiatric disorders [13–15].

Inflammation is a key component of MetS, as elevated circulating levels of inflammatory mediators facilitate the development of this condition [16,17]. Severe obesity is associated with an inflammatory profile characterized by increased plasma production of cytokines and resulting in an imbalance of the cytokine network [18]. Similarly, T2D is associated with a shift of the balance between proinflammatory and anti-inflammatory cytokines toward inflammation [19]. Consequently, MetS is presently viewed not only as a metabolic disorder, but also as an inflammatory disease affecting both innate and acquired immune systems [17,20].

Abundant evidence supports immune-to-brain communication, with peripheral cytokines acting on the brain to induce local production of cytokines and to influence pathways involved in the regulation of mood and cognition, including neurotransmitter metabolism, neuroendocrine function and neural plasticity [21,22]. Interestingly, we and others have identified peripheral and central inflammatory factors as important mediators of the neuropsychiatric symptoms observed in many medical illnesses

sharing chronic systemic inflammation as a common denominator [22–26]. We have found in mice that the selective activation of enzymatic pathways involved in brain monoamine metabolism by interferon- $\gamma$  (IFN- $\gamma$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) mediates development of depressive-like behaviors in response to chronic infection [24–26]. The same mechanisms likely contribute to the high incidence of depressive disorders reported in medically ill patients chronically treated with IFN- $\alpha$  [27] or in elderly subjects exhibiting systemic low-grade inflammation as manifested by increased serum levels of C-reactive protein (CRP) and interleukin-6 (IL-6) [28]. Besides, converging animal [29–31] and clinical findings [32–34] support a main role for IL-6 in mood disorders and cognitive decline. Mounting evidence suggests that systemic inflammation, particularly higher peripheral IL-6, is associated with brain inflammation [35,36] that might adversely affect mood, learning and memory through processes related to neurodegeneration and structural remodeling [37–39]. These processes mainly affect the hippocampus [40–42], a key brain area for memory formation and mood [43,44].

A sizeable number of molecules are well-known to influence hippocampal synaptic plasticity contributing to mood and memory processes, but the brain-derived neurotrophic factor (BDNF) is a major candidate in the context of MetS. BDNF has been found critical in a number of memory tasks [45] and represents an essential constituent of central neural circuits involved in regulating energy homeostasis [46]. Intact hippocampal BDNF signaling determines antidepressant efficacy and influences anxiety-like behaviors [47]. Moreover, hippocampal BDNF mRNA expression and signal transduction are negatively regulated by proinflammatory cytokines [48–50], with reported consequences on cytokine-induced behavioral alterations [21,48,49].

There are but few reports concerned with the role of inflammation in the onset of neuropsychiatric symptoms in patients with MetS and/or obesity. Increased brain levels of TNF- $\alpha$  and IL-6 induced by consumption of high-fat diet in mice are associated with reduced BDNF levels and cognitive performances [51]. At the clinical levels, cognitive impairments have been recently reported to be more likely associated with MetS in the presence of elevated circulating levels of CRP [11,52]. Other clinical reports point to elevated circulating levels of IL-6 as important determinant of mood symptoms [53,54] and cognitive decline [32] associated with MetS and/or obesity. In light of the present knowledge on the neurobiological and behavioral consequences of elevated IL-6 concentrations, these data strongly suggest that brain inflammation is likely involved in the pathogenesis of the neuropsychiatric symptoms associated with MetS. However, clinical studies in patients with MetS only assess systemic inflammation and it is still unknown if enhanced cytokine production associated with neuropsychiatric disorders in MetS does exist in brain areas involved in controlling behavior, particularly the hippocampus. In this context, corollary questions arise as to whether all neuropsychiatric symptoms are equally affected in MetS and which inflammatory pathways are potentially activated in the hippocampus.

In the present study, we sought to answer these questions in a mouse model of MetS, the *db/db* mice, which display T2D, obesity, hyperglycemia, insulin-resistance and hyperinsulinemia as a consequence of an inactivating mutation in the leptin receptor [55]. There are but few reports showing that *db/db* mice also display immune changes, particularly altered immune response to infection both in terms of cytokine production [56,57] and sickness behavior [58]. Few studies have dealt with the emotional behaviors and cognitive performances of *db/db* mice [59–62], but there are as yet no report regarding the potential involvement

of inflammatory processes, particularly within the brain, in impaired emotional and cognitive behaviors. Therefore, the present set of experiments was carried out to identify in *db/db* mice potential alterations in depressive-like, anxiety-like and cognitive behaviors together with a detailed assessment of their peripheral and central inflammatory status. According to the role of the hippocampus in emotional and cognitive behaviors [43,44] we hypothesized that the behavioral deficits of *db/db* mice would be specifically associated with activation of cytokine pathways in the hippocampus.

## Results

### 1-*db/db* mice display expected metabolic dysregulations

In accordance with their expected phenotype, *db/db* mice were markedly heavier than *db/+* controls (weight (g):  $25.5 \pm 0.3$  vs.  $40.5 \pm 1.0$  at 10 week-old;  $F_{(1,26)} = 220.0$ ,  $p < 0.0001$ ), ate more (daily food intake (g):  $4.5 \pm 0.2$  vs.  $7.8 \pm 0.5$  during the last week before sacrifice;  $F_{(1,25)} = 43.6$ ,  $p < 0.0001$ ) and showed a marked increase in the proportion of adipose tissue compared to *db/+* mice (data not shown). This was associated with hyperleptinemia ( $F_{(1,10)} = 266.2$ ,  $p < 0.0001$ ), hyperinsulinemia ( $F_{(1,10)} = 21.1$ ,  $p < 0.001$ ), hyperglycemia ( $F_{(1,8)} = 14.3$ ,  $p < 0.005$ ) and hypercortisolemia ( $F_{(1,10)} = 11.4$ ,  $p < 0.01$ ), but reduced plasma concentrations of resistin ( $F_{(1,10)} = 13.4$ ,  $p < 0.005$ ) (Table 1).

### 2-*db/db* mice display increased anxiety-like behaviors and impaired spatial working memory performances

Anxiety-like behaviors were assessed in the open-field (OF) and elevated plus maze (EPM). When exposed to the OF, *db/db* mice clearly spent less time ( $F_{(1,19)} = 4.9$ ,  $p < 0.05$ ) and visited less often ( $F_{(1,19)} = 13.5$ ,  $p < 0.005$ ) the central area than *db/+* mice (Fig. 1A). The detailed time-course analysis of the number of entries performed in this anxiogenic central area revealed a progressive over-time increase in *db/+* mice (time:  $F_{(1,110)} = 4.4$ ,  $p < 0.0001$ ) whereas central entries remained very few in *db/db* mice throughout the test. Similarly, although all mice equally visited the closed arms of the EPM, *db/db* mice displayed less entries ( $F_{(1,20)} = 4.5$ ,  $p < 0.05$ ; Fig. 1B) and reduced percent of time spent into the open arms than *db/+* mice ( $F_{(1,20)} = 6.48$ ,  $p < 0.05$ ; Fig. 1C). Moreover, these behavioral differences were not due to

**Table 1.** Concentrations of metabolic hormones and cytokines in the plasma of *db/+* and *db/db* mice.

Plasma	<i>db/+</i>	<i>db/db</i>
Leptin (ng/ml)	$2.8 \pm 0.7$	$63.7 \pm 3.7^{***}$
Resistin (ng/ml)	$3.4 \pm 0.1$	$1.7 \pm 0.1^{**}$
Insulin (ng/ml)	$1.1 \pm 0.2$	$5.9 \pm 1.0^{***}$
Glucose (mg/dl)	$220.2 \pm 14.7$	$632.8 \pm 135.5^{**}$
Corticosterone (ng/ml)	$26.2 \pm 6.1$	$148.2 \pm 38.61^{**}$
IL-1 $\beta$ (pg/ml)	$17.4 \pm 9.6$	$29.0 \pm 11.5$
TNF- $\alpha$ (pg/ml)	$8.7 \pm 1.1$	$13.9 \pm 5.1$
IL-6 (pg/ml)	$11.7 \pm 5.3$	$136.7 \pm 59.9^*$
MCP-1 (pg/ml)	$36.8 \pm 9.2$	$80.6 \pm 11.9^*$


Data represent means  $\pm$  SEM (n = 6–7/group).

\* $p < .05$ ,

\*\* $p < .01$ ,

\*\*\* $p < .001$  for *db/db* vs. *db/+* mice.

doi:10.1371/journal.pone.0024325.t001




**Figure 1. Anxiety-like behaviors of *db/db* and *db/+* mice.** (A) Percent of time spent in the center area of the open-field and temporal evolution of the number of entries into this area. (B) Number of entries and (C) percent of time spent into the open arms of the elevated plus-maze. Data represent means  $\pm$  SEM (n=6/group). \*  $p < .05$ , \*\*  $p < .01$  for *db/db* vs. *db/+* mice. doi:10.1371/journal.pone.0024325.g001

locomotor impairment since *db/db* mice displayed proportionally more entries into the peripheral squares of the OF than *db/+* mice (63.37% vs. 55.89%;  $F_{(1,19)} = 8.7$ ,  $p < 0.01$ ) and both groups exhibited similar number of total arm entries in the EPM (data not shown). Thus, results obtained from the OF and EPM paradigms converge to demonstrate greater anxiety-like behaviors in *db/db* mice compared to their *db/+* counterparts. In contrast, assessment of depressive-like behaviors in the tail suspension test (TST) and forced swim test (FST) reveals similar duration of immobility in both genotypes, suggesting that *db/db* mice did not display greater depressive-like behavior than *db/+* mice (Fig. 2).

We also measured spatial working memory performances of *db/db* and *db/+* mice using a spatial recognition test designed as a hippocampus-dependent task [63,64]. During the pre-test session of free exploration of two arms out of three in the Y-maze, the number of visits and the time spent in each open arm were similar in both genotypes indicating unaltered spontaneous alternation (data not shown). Spatial memory performances differed across the genotypes when a 30-min retention interval was used, as revealed by main effects of genotype ( $F_{(1,18)} = 5.6$ ,  $p < 0.05$ ) and arm ( $F_{(1,18)} = 7.6$ ,  $p < 0.05$ ) (Fig. 3A). Further analysis indicated that *db/+* mice spent more time exploring the novel arm than the familiar arm ( $p < 0.05$ ), whereas *db/db* mice randomly explored the different arms. Consequently, the index of discrimination calculated as the time spent exploring the novel arm divided by the total time exploring the different arms multiplied by 100, was significantly higher than chance level in *db/+* mice (59%, one sample t-test:  $p < 0.05$ ), but not in *db/db* mice (55%,  $p > 0.05$ ) (data not shown). When tested with a short

2-min ITI that corresponds to a minimal mnemonic demand, both *db/db* and *db/+* mice preferentially explored the novel arm compared to familiar arms ( $F_{(1,46)} = 10.2$ ,  $p < 0.005$ ; 62% and 60% for *db/db* and *db/+* mice, respectively  $p < 0.05$ ) (data not shown). In summary, *db/db* mice displayed normal spatial recognition performances with a short retention interval, but impaired performances once the mnemonic demand increased together with the ITI. These data likely reflect a deficit in hippocampus-dependent working memory rather than motor or motivational disturbances.

To evaluate whether working memory performances were also affected in a non-spatial task, mice were submitted to the novel object recognition (NOR) task used as a hippocampus-independent task [64,65]. After 30-min retention, all mice explored more the novel object than the familiar one ( $F_{(1,36)} = 22.5$ ,  $p < 0.0001$ ) (Fig. 3B), although *db/db* mice spent globally less time exploring the objects than *db/+* mice (genotype:  $F_{(1,36)} = 12.7$ ,  $p < 0.005$ ; object  $\times$  genotype:  $F_{(1,36)} = 4.4$ ,  $p < 0.05$ ). Both *db/+* and *db/db* mice were therefore able to discriminate the novel object from the familiar one, as confirmed by the fact that the index of discrimination was statistically different from chance level (50%) whatever the genotype (70.9% and 68.3% respectively;  $p < 0.001$ ). To evaluate whether longer ITI could reveal an impairment of object recognition in *db/db* mice, other mice were tested with a 1-h ITI. Neither *db/db* nor *db/+* mice preferentially explored the novel object over the familiar one with a 1-h ITI, although *db/+* mice still spent more time exploring the objects than *db/db* mice (genotype:  $F_{(1,34)} = 15.4$ ,  $p < 0.001$ ) (data not shown). In summary, *db/db* mice displayed selective behavioral alterations affecting anxiety-like behavior and spatial working memory.



**Figure 2. Depressive-like behaviors of *db/db* and *db/+* mice.** Immobility time in the (A) tail suspension test and (B) forced swim test. Data represent means  $\pm$  SEM ( $n=7$ /group). doi:10.1371/journal.pone.0024325.g002

### 3–Behavioral alterations are associated with peripheral and brain inflammation in *db/db* mice

Table 1 shows peripheral levels of several well-known markers of inflammation. *db/db* mice displayed significantly higher plasma levels of IL-6 ( $F_{(1,10)}=6.2$ ,  $p<0.05$ ) and the chemokine MCP-1 ( $F_{(1,12)}=3.0$ ,  $p<0.05$ ) than *db/+* mice. Plasma levels of IL-1 $\beta$  and TNF- $\alpha$  were low and similar in both genotypes, whereas no IFN- $\gamma$  was detected whatever the group.

Cytokine mRNA expression was measured in the hippocampus, a key brain areas for the control of spatial memory and anxiety-like behaviors [43,44]. Concomitantly, the hypothalamus was used as a control structure involved in physiological homeostasis, but not in the control of anxiety-like and cognitive behaviors [44,46]. IFN- $\gamma$  and MCP-1 mRNA expression levels were similar regardless the genotype and the area. Hippocampus mRNA expression of IL-1 $\beta$  ( $F_{(1,9)}=6.8$ ,  $p<0.05$ ), TNF- $\alpha$  ( $F_{(1,10)}=7.4$ ,  $p<0.05$ ) and IL-6 ( $F_{(1,8)}=7.1$ ,  $p<0.05$ ) was significantly higher in *db/db* mice than in their *db/+* counterparts (Fig. 4A). In contrast, hypothalamus TNF- $\alpha$  and IL-6 mRNA expression was similar in both genotypes whereas *db/db* mice displayed lower levels of IL-1 $\beta$  mRNA expression than *db/+* mice ( $F_{(1,9)}=7.6$ ,  $p<0.05$ ; Fig. 4B).


To further understand the functional consequences of increased expression of cytokines, we next measured mRNA expression of one of the key element of the IL-6 receptor complex, the glycoprotein 130 (GP130), IL-6 being the only cytokine increased at the periphery and in the hippocampus of *db/db* mice.

Additionally, we measured the mRNA expression of cytokine signaling-3 (SOCS3), a classical indicator of cytokine signaling pathway activation [66]. GP130 mRNA expression was stable across the brain area and the genotype (Fig. 5A, 5C). On the contrary, *db/db* mice displayed increased levels of SOCS3 mRNA expression in the hippocampus ( $F_{(1,9)}=6.7$ ,  $p<0.05$ ; Fig. 5A), together with a reduced expression in the hypothalamus ( $F_{(1,9)}=12.1$ ,  $p<0.01$ ; Fig. 5C). Alterations displayed by *db/db* mice in anxiety-like behaviors and spatial working memory were therefore associated with hippocampal, but not hypothalamic, low-grade inflammation.


Finally, we measured the mRNA expression of BDNF, a potent neuroprotective growth factor regulated by cytokines [48–50] and well-known to participate in mood regulation and memory function [45,47]. BDNF was significantly reduced in the hippocampus of *db/db* mice compared to *db/+* mice ( $F_{(1,9)}=7.8$ ,  $p<0.05$ ; Fig. 5B), whereas no significant differences were observed between both groups in the hypothalamus (Fig. 5D).

## Discussion


Our study allows for the first time to directly relate in *db/db* mice increased anxiety-like behaviors and impaired spatial memory performances with activation of specific inflammatory pathways within the hippocampus, a key brain area for the control of emotional and cognitive behaviors.



**Figure 3. Working memory performances of *db/db* and *db/+* mice.** (A) Spatial recognition in the Y-maze expressed as the time spent exploring the novel and the familiar arms. (B) Time spent exploring the novel and the familiar object in the novel object recognition task. In both tasks, measures were assessed over a 5-min test and after 30-min retention. Data represent means  $\pm$  SEM ( $n=7-10$ /group). \*  $p<0.05$ , \*\*  $p<0.01$  for *db/db* vs. *db/+* mice. doi:10.1371/journal.pone.0024325.g003



**Figure 4. mRNA expression levels of cytokines in the hippocampus and hypothalamus of *db/db* and *db/+* mice.** Relative fold changes in the levels of (A) hippocampal and (B) hypothalamic IL-1β, TNF-α, IL-6, IFN-γ and MCP-1 mRNA expression, as calculated in relation to the averaged value for control saline group. IL-1β, interleukin-1β; TNF-α, tumor necrosis factor-α; IFN-γ, interferon-γ; MCP-1, monocyte chemoattractant protein-1. Data represent means ± SEM (n=6/group). \*  $p < .05$  for *db/db* vs. *db/+* mice. doi:10.1371/journal.pone.0024325.g004



**Figure 5. mRNA expression levels of GP130, SOCS3 and BDNF in the hippocampus and hypothalamus.** Relative fold changes in the levels of (A–B) hippocampal and (C–D) hypothalamic GP130, SOCS3 and BDNF mRNA expression, as calculated in relation to the averaged value for control saline group. GP130, glycoprotein 130; SOCS3, suppressor of cytokine signaling-3; BDNF, brain-derived neurotrophic factor. Data represent means ± SEM (n=6/group). \*  $p < .05$  for *db/db* vs. *db/+* mice. doi:10.1371/journal.pone.0024325.g005

Most clinical studies investigating the link between MetS and mood disorders report a positive association [6,10,15]. Although measuring mood in rodents could appear limiting, the development of consistent and reliable behavioral tests modeling different core symptoms of anxiety and depression rather than the entire syndromes has provided very useful tools to study their respective pathophysiology [67]. Here, data obtained in different experimental paradigms converge to indicate that *db/db* mice display increased anxiety-like behaviors, as previously suggested [62], but not depressive-like behaviors. Increased immobility in the FST has been reported in leptin-deficient (*ob/ob*) mice when experimental conditions much more stressful than ours are used (longer exposure to the FST and for 3 consecutive days) [68]. Of note, depressive-like behaviors in the FST and/or TST mostly increase under challenging conditions such as stress exposure [69] or immune stimulation [24–26,70]. The possibility that *db/db* mice would display greater depressive-like behaviors in such conditions is currently under study. While this may be, the current study clearly shows that *db/db* mice displayed increased anxiety-like, but not depressive-like, behaviors in basal conditions.

Clinical studies have reported that MetS adversely impairs cognition, although not all cognitive domains are equally affected [4,5,11]. Our study further extends these findings by demonstrating task-specific cognitive impairments in *db/db* mice using two working memory paradigms (the Y-maze and NOR). After 30 minutes of retention, *db/db* mice exhibited spatial working memory impairment in the Y-maze while their performances were unaltered in NOR memory task. Moreover, the spatial deficit in *db/db* mice is not due to motor and/or motivational problems as their spatial performance was intact with a shorter retention interval (2-min). Intact performances of *db/db* mice in the NOR task disagree with recently published data showing NOR impairment in these mice with the same 30-min ITI [61]. However, two important differences between their experimental protocol and ours can likely explain this apparent discrepancy. First, they used C57BL/6J mice as controls instead of *db/+* mice (as we used), which have similar genetic background (C57BL/6J×DBA/2J) and perinatal environment as their littermates *db/db* mice. Interestingly, we recently showed differences in the duration of object exploration between C57BL/6J and *db/+* mice (data not shown) that can likely be explained by their different genetic background and/or different perinatal environment. Second and more importantly, they exposed *db/db* mice only once to the testing cage before the NOR training and test, whereas in our protocol mice were extensively habituated to the testing cage (15 min per day during 8 days). According to the fact that the level of habituation to the experimental context influences the emotional arousal and performances during NOR [71,72] and that *db/db* mice show greater anxiety-like behaviors than controls, it is likely that testing *db/db* mice in higher stressful conditions than ours may interfere with their NOR performances.

In parallel to behavioral alterations, we report neurobiological changes in *db/db* mice affecting molecules well-known to participate in emotional and cognitive behaviors, particularly cytokines [21,22]. Expression level of IL-6, TNF $\alpha$  and IL-1 $\beta$  mRNAs was selectively increased by ~80% in the hippocampus, but not the hypothalamus, of *db/db* mice. Although we only measured cytokine transcripts, these changes are likely to be reflected in changes of protein levels contributing to downstream neurobiological and behavioral modulations [73]. This assumption is supported by the concomitant increase of hippocampal SOCS3 mRNA expression that is classically used as an indicator of cytokine signaling pathway activation, mainly IL-6 [66]. In contrast, IL-1 $\beta$  mRNA expression is significantly reduced in the

hypothalamus of *db/db* mice. This can be linked to the increased food intake of these mice as mounting evidence indicates the anorectic effect of enhanced hypothalamic expression of IL-1 [21,74], including in response to leptin [75]. Of note, this result is particularly relevant to better understand the role of brain cytokines in the impairment of food intake in *db/db* mice, although this question has not been directly addressed in the present study. As reported for cytokines, BDNF is also well-known to contribute to mood regulation and memory function, particularly in the hippocampus [45,47] where its expression is negatively regulated by cytokines [48,49]. Accordingly, the increase of proinflammatory cytokine expression we found in the hippocampus of *db/db* mice is accompanied by ~60% decrease of hippocampal, but not hypothalamic, BDNF mRNA expression. This change is likely reflected in changes of functional protein levels since significant correlations have been reported between both BDNF mRNA expression and protein levels [76,77].

These neurobiological changes are observed in the hippocampus, but not in the hypothalamus. This is particularly relevant in light of the behavioral alterations of *db/db* mice. Indeed, we found that *db/db* mice are impaired in spatial, but not in object, recognition tasks for a similar delay and considerable literature shows that spatial memory, but not object memory, requires normal hippocampal functioning [65,78]. This interpretation agrees with data reporting that *db/db* mice are impaired in the hidden-platform version of the Morris water-maze test (a hippocampus-dependent situation), but not under visible-platform conditions (a hippocampus-independent situation) [59–61]. Moreover, the hippocampus is also involved in anxiety-related behaviors [43]. Therefore the higher level of anxiety-like behaviors we found in *db/db* mice (see also Stranahan et al. [62]) could also be linked to the neurobiological changes we found in the hippocampus of these mice, although further studies are needed to measure potential neurobiological changes in other brain areas well-known to participate in controlling emotional behavior, particularly the amygdala complex (for instance see [79]).

All these data support a key role for the hippocampal changes (increased cytokines, decreased BDNF) in the cognitive and emotional alterations displayed by *db/db* mice. Interestingly, chronic overexpression of brain IL-6 or hippocampal IL-1 $\beta$  in transgenic mice impairs learning [37,41]. These findings concur with several reports highlighting the relationship between neuroinflammation, alterations of hippocampal synaptic plasticity and impaired cognitive performances [38,39,51]. Interestingly, different animal studies report that increased levels of hippocampal IL-6 interfere with local long-term potentiation, neurogenesis and synaptic plasticity [40–42], whereas blocking hippocampal IL-6 intracellular pathway prevents cytokine-induced alterations of synaptic activity and spatial learning [80]. In humans, circulating IL-6 concentrations have been shown to covary inversely with hippocampal grey matter volume [36] and cognitive performances [32]. Chronic proinflammatory cytokines have also been shown to produce detrimental effects on mood [22–26]. Consistent clinical data report a significant correlation between elevated circulating levels of inflammatory markers, particularly IL-6, and development of anxiety symptoms [33,81], including in obese patients [53,54]. Likewise, experimental studies using IL-6-deficient mice [30] or rats bred for extremes in anxiety-related behavior [82] support a role of IL-6 in anxiety-like behavior. Concerning BDNF, decreased level in the hippocampus is associated with impaired synaptic plasticity, cognitive performances [45] and mood-related behaviors [47]. Interestingly, IL-1 $\beta$  modulates memory-induced increase in hippocampal BDNF mRNA expression and signal transduction [21,48,50]. Accordingly, elevated cytokine levels

observed in the hippocampus of *db/db* mice may contribute to decrease BDNF level and consequently to impair anxiety-like behaviors and spatial cognitive performances.

Beyond elevated cytokine levels and related neurobiological changes, *db/db* mice also displayed elevated plasma levels of leptin, insulin and corticosterone. In addition to their metabolic properties, leptin and insulin can modulate behavior by acting within the brain [83,84]. Consequently, impaired leptin and/or insulin signaling activation, as observed in *db/db* mice, has been proposed as potential link between behavioral and metabolic disorders. Although this assumption cannot be totally excluded based on the present data, several lines of evidence suggest that leptin does not play a main role in mediating behavioral alterations of *db/db* mice. Indeed, cognitive impairment displayed by *db/db* mice can be improved by normalization of corticosterone levels, without restoring impaired leptin signaling pathway [61,62]. Moreover, caloric restriction and running enhance exploratory behavior in *db/db* mice, without changing serum leptin levels [62]. Similarly, insulin unlikely contributes to the behavioral alterations reported in *db/db* mice. Indeed, brain concentrations of glucose and insulin are similar in both *db/db* and *db/+* mice and remained unchanged after normalization of peripheral hyperinsulinemia [61]. Moreover, caloric restriction and running completely reversed hyperglycemia in *db/db* mice, without reducing anxiety-like behavior in the open-field [62]. Likewise, normalizing hyperglycemia in insulin-deficient rats does not reverse spatial cognitive impairment [85]. These data match those showing that the increased risk of cognitive dysfunction reported in MetS patients is independent from the presence of diabetes [4]. Additionally, it was recently suggested that elevated corticosterone contributes to cognitive alterations in *db/db* mice by acting within the hippocampus [61]. Normalizing plasma corticosterone levels reverses alterations of hippocampal plasticity and improves hippocampus-dependent memory [61]. However, these findings do not preclude the involvement of other factors, namely proinflammatory cytokines, in cognitive and emotional alterations. As both the inflammatory system and the hypothalamic-pituitary-adrenal axis have been found tightly interrelated [86], it would be of interest to assess in *db/db* mice whether normalizing plasma corticosterone also normalizes hippocampal proinflammatory cytokines. Additionally, this study should be broadened to other brain areas, particularly the amygdala complex, since 1) it plays a key role in controlling emotional behavior, particularly anxiety [79] and 2) it can be targeted by both corticosterone and cytokines [87]. Based on all these converging data, it can reasonably be proposed that the behavioral profile displayed by *db/db* mice likely results from the concerted interactions of cytokines and glucocorticoids with central processes involving the hippocampus and controlling cognition and emotionality.

In conclusion, although only correlative and with the limitation inherent to the measurement of neurobehavioral reactivity in rodents, the present results strongly point to hippocampal inflammation as important contributing factors to the pathophysiology of neuropsychiatric complications associated with MetS. These findings may prove valuable for introducing novel approaches to treatment, alongside currently used non-pharmacologic and pharmacologic interventions.

## Materials and Methods

### Animals

All animal experiments were conducted according to the INRA Quality Reference System, and to relevant French (Directive 87/148, Ministère de l'Agriculture et de la Pêche) and international

(Directive 86/609, November 24<sup>th</sup> 1986, European Community) legislation. They adhered to protocols approved by Région Aquitaine Veterinary Services (Direction Départementale de la Protection des Animaux, approval ID: A33-063-920). Every effort was made to minimize suffering and the number of animal used. Male *db/db* (C57BLKS/J-*lepr<sup>db</sup>/lepr<sup>db</sup>*; n=40) and *db/+* (C57BLKS/J-*lepr<sup>db</sup>/+*; n=40) mice obtained from Charles River Laboratories (France) were housed individually under a normal 12-hour light:dark cycle with food and water available *ad libitum*. Mice were handled daily for 1 week before the experiment onset to minimize stress reactions to manipulation. They were 8 to 10 weeks-old at the time of behavioral assessments.

### Behavioral measurements

Experiments were performed in the morning under conditions of dim light and low noise. Behavior was videotaped and scored using "The Observer Basic" software (Noldus, Netherlands). Each mouse was submitted to a maximum of two different behavioral tests, with a 1-week interval between tests. All testing equipment was thoroughly cleaned between each session.

**Open-field (OF).** Mice were exposed to an unknown square area (40×40 cm) from which escape is prevented by surrounding walls (16 cm high). The apparatus was virtually divided into 4 central squares defined as the central area and 12 squares along the walls that are defined as the periphery. Each mouse was placed in the central area and allowed to freely explore the OF for 10 min. Parameters recorded to evaluate anxiety-like behavior were the number of entries and the percent of time spent in the central area [88].

**Elevated Plus Maze (EPM).** The EPM was a plus shaped acryl maze with two opposing open arms (30×8 cm) and two opposing closed arms (30×8×15 cm) connected by a central platform (8×8 cm) and elevated 120 cm above the floor. Each mouse was placed in the center of the maze facing an open arm and the number of arm entries, as well as the percent of time spent in open arms, was assessed during a 5-min period. An entry was scored as such only when the mouse placed all four limbs into any given arm. A reduction of the percent of time spent and number of entries into the open arms is considered as an anxiety-like index, independent of locomotor activity [88].

**Tail Suspension Test (TST).** This standardized test of depressive-like behavior was carried out as previously described [24,70]. Briefly, an adhesive tape was fixed to the mouse tail and hooked to a horizontal ring stand bar placed 30 cm above the floor. The test was conducted for a 6-min period in a visually isolated area. Mice demonstrated several escape attempts interspersed with immobility periods during which they hung passively and completely motionless. Depressive-like behavior was inferred from increased duration of immobility.

**Forced Swim Test (FST).** This standardized test of depressive-like behavior was essentially conducted as previously described [24,70]. Briefly, each mouse was placed individually in a cylinder (16×31 cm) containing warm water (25±1°C) to avoid temperature-related stress response. Mice were tested during a 6-min period. Immobility time was determined by the time a mouse stopped struggling and moved only slowly to remain floating in the water, keeping its head above water. Increased duration of immobility has been proposed to reflect a state of helplessness that is reduced by antidepressants.

**Y-maze.** Spontaneous spatial recognition in the Y-maze was used as a hippocampal-dependent test as previously described [63,64]. The apparatus was a Y-shaped acryl maze with three identical arms (34×8×14 cm). Corncob litter covered the floor and was mixed between each trial in order to remove olfactory



cues. Visual cues were placed in the testing room and kept constant during the whole test. Discrimination of novelty versus familiarity was based on the different aspects of the environment that the mouse can perceive from each arm of the Y-maze. In the first trial of the test (acquisition), one arm was closed with a door and mice were allowed to freely visit the two other arms for 5 min. After a 30-min inter-trial interval (ITI), mice were again placed in the start arm for the second trial (retrieval) and allowed free access to all three arms for 5 min. Start and closed arms were randomly assigned to each mouse. Arm entries were defined as all four paws entering the arm. Preference for novelty was also measured using a short 2-min ITI between acquisition and retrieval in order to control for potential motivational disturbances [63,64]. Analyses were based on the time spent exploring the novel and the familiar arms during the second trial. An index of discrimination between novel and familiar arms was calculated as the ratio of the [time spent in the novel arm/(time spent in the novel + adjacent arms)] $\times$ 100. An index of discrimination significantly higher than chance level (50%) indicates therefore that mice indeed recognize the novel arm.

**Novel object recognition (NOR).** This task is a free exploration paradigm allowing mice to explore objects in a non-threatening and familiar environment. It was used as a hippocampal-independent task [64,65]. Mice were first acclimatized in an opaque acrylic cage (43 $\times$ 28 $\times$ 19 cm) for 15 min per day during the week before training. The floor was covered with corncob litter that was mixed between each trial in order to remove olfactory cues. On the ninth day, mice were placed in the experimental cage with two identical objects during a 10-min training session. Three sets of objects with different shapes and colors were used for discrimination. Each object was chosen on the basis of a preliminary study examining the preferences of a separate group of mice (data not shown) and was heavy enough not to be displaced by a mouse. After a 30-min or 1-hour ITI, one of the familiar objects was replaced by a novel object, with a different shape and color, to test for memory retention. During the 5-min test, exploration of each object was defined as sniffing or touching the object with the nose and/or forepaws. Analyses were based on the time spent exploring the novel and the familiar objects. An index of discrimination was calculated as the ratio of the [time spent exploring the novel object/(total time spent exploring both objects)] $\times$ 100.

### Biochemical measurements

After completion of the behavioral experiments, some *db/db* and *db/+* mice were euthanized by CO<sub>2</sub> inhalation (10–12 weeks-old at the time of sacrifice). Blood samples were immediately collected

via cardiac puncture into EDTA (10%)-coated chilled tubes. After centrifugation (10 min, 3000 *g*, 4°C), aliquots of plasma were stored at –80°C. Mice were perfused with chilled PBS via the ascending aorta to remove all traces of blood from tissues. Brains were rapidly extracted from the skulls and carefully dissected. The hippocampus and the hypothalamus were immediately collected, dried frozen and stored until assaying.

**Hormones and cytokines assay.** As previously described [24,35], plasma leptin, insulin, resistin, monocyte chemoattractant protein-1 (MCP-1), IL-1 $\beta$ , IL-6, TNF- $\alpha$  and IFN- $\gamma$  were measured with the mouse adipokine and cytokine LINCOplex kits (Linco Research, Inc., St. Charles, MO, USA) following the manufacturer's instructions. Plasma corticosterone concentrations were measured using a commercial RIA Kit (Diasorin, Antony, France). Plasma glucose levels were measured using a One Touch Ultra glucometer per the manufacturer's instructions. All samples were run in duplicate.

**Reverse transcription and real-time RT-PCR.** Total RNA was extracted from the hippocampus and hypothalamus using a RNeasy Mini Kit (Qiagen) and reverse-transcribed as previously described [25,26,35,64]. Real-time RT-PCR was performed on an ABI Prism 7700 using Taqman gene expression assays for sequence-specific primers purchased from Applied Biosystems (Foster City, CA). Reactions were performed in duplicate according to manufacturer instructions as previously described [25,26,35]. Relative expression levels were calculated according to the methods of Livak and Schmittgen [89] and plotted as fold change relative to the appropriate control condition.

### Statistical analysis

Results are presented as mean  $\pm$  SEM and were analyzed using a one-way (genotype) or a two-way (genotype $\times$ arm; genotype $\times$ object) ANOVA followed by a *post-hoc* pair wise multiple comparison procedure using the Fischer's LSD method, if the interaction was significant.

### Acknowledgments

The authors thank P. Birac and C. Tridon for tacking care of the mice. Real-time PCR experiments were performed at the Genotyping and Sequencing Facility of Bordeaux.

### Author Contributions

Conceived and designed the experiments: NC A-LD. Performed the experiments: A-LD CA AA NC. Analyzed the data: A-LD CA NC. Wrote the paper: NC A-LD GF SL.

### References

- Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, et al. (2009) Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 120: 1640–1645.
- Anagnostis P, Athyros VG, Tziomalos K, Karagiannis A, Mikhailidis DP (2009) Clinical review: The pathogenetic role of cortisol in the metabolic syndrome: a hypothesis. *J Clin Endocrinol Metab* 94: 2692–2701.
- Gupta A, Gupta V (2010) Metabolic syndrome: what are the risks for humans? *Biosci Trends* 4: 204–212.
- Muller M, van Raamt F, Vissers FL, Kalmijn S, Geerlings MI, et al. (2010) Metabolic syndrome and cognition in patients with manifest atherosclerotic disease: the SMART study. *Neuroepidemiology* 34: 83–89.
- Raffaitin C, Feart C, Le Goff M, Amieva H, Helmer C, et al. (2011) Metabolic syndrome and cognitive decline in French elders: the Three-City Study. *Neurology* 76: 518–525.
- van Reedt Dortland AK, Giltay EJ, van Veen T, Zitman FG, Penninx BW (2010) Metabolic syndrome abnormalities are associated with severity of anxiety and depression and with tricyclic antidepressant use. *Acta Psychiatr Scand* 122: 30–39.
- Padwal RS, Sharma AM (2010) Prevention of cardiovascular disease: obesity, diabetes and the metabolic syndrome. *Can J Cardiol* 26 Suppl C: 18C–20C.
- Blalock EM, Grondin R, Chen KC, Thibault O, Thibault V, et al. (2010) Aging-related gene expression in hippocampus proper compared with dentate gyrus is selectively associated with metabolic syndrome variables in rhesus monkeys. *J Neurosci* 30: 6058–6071.
- Engum A (2007) The role of depression and anxiety in onset of diabetes in a large population-based study. *J Psychosom Res* 62: 31–38.
- Goldbacher EM, Bromberger J, Matthews KA (2009) Lifetime history of major depression predicts the development of the metabolic syndrome in middle-aged women. *Psychosom Med* 71: 266–272.
- Roberts RO, Geda YE, Knopman DS, Cha RH, Boeve BF, et al. (2010) Metabolic syndrome, inflammation, and nonamnestic mild cognitive impairment in older persons: a population-based study. *Alzheimer Dis Assoc Disord* 24: 11–18.
- Zeugmann S, Quante A, Heuser I, Schwarzer R, Anghelescu I (2010) Inflammatory biomarkers in 70 depressed inpatients with and without the metabolic syndrome. *J Clin Psychiatry* 71: 1007–1016.

13. Elenkov IJ (2008) Neurohormonal-cytokine interactions: implications for inflammation, common human diseases and well-being. *Neurochem Int* 52: 40–51.
14. Emery CF, Fondow MD, Schneider CM, Christofi FL, Hunt C, et al. (2007) Gastric bypass surgery is associated with reduced inflammation and less depression: a preliminary investigation. *Obes Surg* 17: 759–763.
15. McIntyre RS, Rasgon NL, Kemp DE, Nguyen HT, Law CW, et al. (2009) Metabolic syndrome and major depressive disorder: co-occurrence and pathophysiologic overlap. *Curr Diab Rep* 9: 51–59.
16. Dandona P, Aljada A, Chaudhuri A, Mohanty P, Garg R (2005) Metabolic syndrome: a comprehensive perspective based on interactions between obesity, diabetes, and inflammation. *Circulation* 111: 1448–1454.
17. Marsland AL, McCaffery JM, Muldoon MF, Manuck SB (2010) Systemic inflammation and the metabolic syndrome among middle-aged community volunteers. *Metabolism* 59: 1801–1808.
18. Cancellor R, Clement K (2006) Is obesity an inflammatory illness? Role of low-grade inflammation and macrophage infiltration in human white adipose tissue. *Bjog* 113: 1141–1147.
19. Donath MY, Boni-Schnetzler M, Ellingsgaard H, Halban PA, Ehlers JA (2010) Cytokine production by islets in health and diabetes: cellular origin, regulation and function. *Trends Endocrinol Metab* 21: 261–267.
20. Schmidt MI, Duncan BB (2003) Diabesity: an inflammatory metabolic condition. *Clin Chem Lab Med* 41: 1120–1130.
21. Anisman H, Merali Z, Hayley S (2008) Neurotransmitter, peptide and cytokine processes in relation to depressive disorder: comorbidity between depression and neurodegenerative disorders. *Prog Neurobiol* 85: 1–74.
22. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW (2008) From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci* 9: 46–56.
23. Evans DL, Charney DS, Lewis L, Golden RN, Gorman JM, et al. (2005) Mood disorders in the medically ill: scientific review and recommendations. *Biol Psychiatry* 58: 175–189.
24. Moreau M, Andre C, O'Connor JC, Dumich SA, Woods JA, et al. (2008) Inoculation of *Bacillus Calmette-Guerin* to mice induces an acute episode of sickness behavior followed by chronic depressive-like behavior. *Brain Behav Immun* 22: 1087–1095.
25. O'Connor JC, Andre C, Wang Y, Lawson MA, Szegedi SS, et al. (2009) Interferon-gamma and tumor necrosis factor-alpha mediate the upregulation of indoleamine 2,3-dioxygenase and the induction of depressive-like behavior in mice in response to *bacillus Calmette-Guerin*. *J Neurosci* 29: 4200–4209.
26. O'Connor JC, Lawson MA, Andre C, Briley EM, Szegedi SS, et al. (2009) Induction of IDO by bacille Calmette-Guerin is responsible for development of murine depressive-like behavior. *J Immunol* 182: 3202–3212.
27. Raison CL, Dantzer R, Kelley KW, Lawson MA, Woolwine BJ, et al. (2010) CSF concentrations of brain tryptophan and kynurenines during immune stimulation with IFN- $\alpha$ : relationship to CNS immune responses and depression. *Mol Psychiatry* 15: 393–403.
28. Capuron L, Schroecksnadel S, Fearf C, Aubert A, Huguere D, et al. (2011) Chronic Low-Grade Inflammation in Elderly Persons Is Associated with Altered Tryptophan and Tyrosine Metabolism: Role in Neuropsychiatric Symptoms. *Biol Psychiatry* 70: 175–82.
29. Braida D, Sacerdote P, Panerai AE, Bianchi M, Aloisi AM, et al. (2004) Cognitive function in young and adult IL (interleukin)-6 deficient mice. *Behav Brain Res* 153: 423–429.
30. Chourbaji S, Urani A, Inta I, Sanchis-Segura C, Brandwein C, et al. (2006) IL-6 knockout mice exhibit resistance to stress-induced development of depression-like behaviors. *Neurobiol Dis* 23: 587–594.
31. Sparkman NL, Buchanan JB, Heyen JR, Chen J, Beverly JL, et al. (2006) Interleukin-6 facilitates lipopolysaccharide-induced disruption in working memory and expression of other proinflammatory cytokines in hippocampal neuronal cell layers. *J Neurosci* 26: 10709–10716.
32. Marsland AL, Petersen KL, Sathanoori R, Muldoon MF, Neumann SA, et al. (2006) Interleukin-6 covaries inversely with cognitive performance among middle-aged community volunteers. *Psychosom Med* 68: 895–903.
33. O'Donovan A, Hughes BM, Slavich GM, Lynch L, Cronin MT, et al. (2010) Clinical anxiety, cortisol and interleukin-6: evidence for specificity in emotion-biology relationships. *Brain Behav Immun* 24: 1074–1077.
34. Weaver JD, Huang MH, Albert M, Harris T, Rowe JW, et al. (2002) Interleukin-6 and risk of cognitive decline: MacArthur studies of successful aging. *Neurology* 59: 371–378.
35. Andre C, O'Connor JC, Kelley KW, Lestage J, Dantzer R, et al. (2008) Spatio-temporal differences in the profile of murine brain expression of proinflammatory cytokines and indoleamine 2,3-dioxygenase in response to peripheral lipopolysaccharide administration. *J Neuroimmunol* 200: 90–99.
36. Marsland AL, Gianaros PJ, Abramowitch SM, Manuck SB, Hariri AR (2008) Interleukin-6 covaries inversely with hippocampal grey matter volume in middle-aged adults. *Biol Psychiatry* 64: 484–490.
37. Hein AM, Stasko MR, Matousek SB, Scott-McKean JJ, Maier SF, et al. (2010) Sustained hippocampal IL-1 $\beta$  overexpression impairs contextual and spatial memory in transgenic mice. *Brain Behav Immun* 24: 243–253.
38. Lee JW, Lee YK, Yuk DY, Choi DY, Ban SB, et al. (2008) Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment through enhancement of beta-amyloid generation. *J Neuroinflammation* 5: 37.
39. Noble F, Rubira E, Boulanouar M, Palmier B, Plotkine M, et al. (2007) Acute systemic inflammation induces central mitochondrial damage and mnesic deficit in adult Swiss mice. *Neurosci Lett* 424: 106–110.
40. Bellinger FP, Madamba SG, Campbell IL, Siggins GR (1995) Reduced long-term potentiation in the dentate gyrus of transgenic mice with cerebral overexpression of interleukin-6. *Neurosci Lett* 198: 95–98.
41. Heyser CJ, Masliah E, Samimi A, Campbell IL, Gold LH (1997) Progressive decline in avoidance learning paralleled by inflammatory neurodegeneration in transgenic mice expressing interleukin 6 in the brain. *Proc Natl Acad Sci U S A* 94: 1500–1505.
42. Monje ML, Toda H, Palmer TD (2003) Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302: 1760–1765.
43. Bannerman DM, Rawlins JN, McHugh SB, Deacon RM, Yee BK, et al. (2004) Regional dissociations within the hippocampus—memory and anxiety. *Neurosci Biobehav Rev* 28: 273–283.
44. Maletic V, Robinson M, Oakes T, Iyengar S, Ball SG, et al. (2007) Neurobiology of depression: an integrated view of key findings. *Int J Clin Pract* 61: 2030–2040.
45. Yamada K, Nabeshima T (2003) Brain-derived neurotrophic factor/TrkB signaling in memory processes. *J Pharmacol Sci* 91: 267–270.
46. Benarroch EE (2010) Neural control of feeding behavior: Overview and clinical correlations. *Neurology* 74: 1643–1650.
47. Martinowich K, Manji H, Lu B (2007) New insights into BDNF function in depression and anxiety. *Nat Neurosci* 10: 1089–1093.
48. Barrientos RM, Sprunger DB, Campeau S, Watkins LR, Rudy JW, et al. (2004) BDNF mRNA expression in rat hippocampus following contextual learning is blocked by intrahippocampal IL-1 $\beta$  administration. *J Neuroimmunol* 155: 119–126.
49. Tanaka S, Ide M, Shibusaki T, Ohtaki H, Numazawa S, et al. (2006) Lipopolysaccharide-induced microglial activation induces learning and memory deficits without neuronal cell death in rats. *J Neurosci Res* 83: 557–566.
50. Tong L, Balazs R, Sojampornkul R, Thangnipon W, Cotman CW (2008) Interleukin-1 $\beta$  impairs brain derived neurotrophic factor-induced signal transduction. *Neurobiol Aging* 29: 1380–1393.
51. Pistell PJ, Morrison CD, Gupta S, Knight AG, Keller JN, et al. (2010) Cognitive impairment following high fat diet consumption is associated with brain inflammation. *J Neuroimmunol* 219: 25–32.
52. Yaffe K (2007) Metabolic syndrome and cognitive decline. *Curr Alzheimer Res* 4: 123–126.
53. Capuron L, Poitou C, Machaux-Tholliez D, Frochot V, Bouilliot JL, et al. (2010) Relationship between adiposity, emotional status and eating behaviour in obese women: role of inflammation. *Psychol Med*. pp 1–12.
54. Capuron L, Su S, Miller AH, Bremner JD, Goldberg J, et al. (2008) Depressive symptoms and metabolic syndrome: is inflammation the underlying link? *Biol Psychiatry* 64: 896–900.
55. Chen H, Charlat O, Tartaglia LA, Woolf EA, Weng X, et al. (1996) Evidence that the diabetes gene encodes the leptin receptor: identification of a mutation in the leptin receptor gene in db/db mice. *Cell* 84: 491–495.
56. Naguib G, Al-Mashat H, Desta T, Graves DT (2004) Diabetes prolongs the inflammatory response to a bacterial stimulus through cytokine dysregulation. *J Invest Dermatol* 123: 87–92.
57. Rummel C, Inoue W, Poole S, Luhsheh GN (2010) Leptin regulates leukocyte recruitment into the brain following systemic LPS-induced inflammation. *Mol Psychiatry* 15: 523–534.
58. O'Connor JC, Satpathy A, Hartman ME, Horvath EM, Kelley KW, et al. (2005) IL-1 $\beta$ -mediated innate immunity is amplified in the db/db mouse model of type 2 diabetes. *J Immunol* 174: 4991–4997.
59. Li XL, Aou S, Oomura Y, Hori N, Fukunaga K, et al. (2002) Impairment of long-term potentiation and spatial memory in leptin receptor-deficient rodents. *Neuroscience* 113: 607–615.
60. Oomura Y, Aou S, Fukunaga K (2010) Prandial increase of leptin in the brain activates spatial learning and memory. *Pathophysiology* 17: 119–127.
61. Stranahan AM, Arumugam TV, Cutler RG, Lee K, Egan JM, et al. (2008) Diabetes impairs hippocampal function through glucocorticoid-mediated effects on new and mature neurons. *Nat Neurosci* 11: 309–317.
62. Stranahan AM, Lee K, Martin B, Maudsley S, Golden E, et al. (2009) Voluntary exercise and caloric restriction enhance hippocampal dendritic spine density and BDNF levels in diabetic mice. *Hippocampus* 19: 951–961.
63. Dellu F, Contarino A, Simon H, Koob GF, Gold LH (2000) Genetic differences in response to novelty and spatial memory using a two-trial recognition task in mice. *Neurobiol Learn Mem* 73: 31–48.
64. Labrousse VF, Costes L, Aubert A, Darnaudey M, Ferreira G, et al. (2009) Impaired interleukin-1 $\beta$  and c-Fos expression in the hippocampus is associated with a spatial memory deficit in P2X(7) receptor-deficient mice. *PLoS One* 4: e6006.
65. Dere E, Huston JP, De Souza Silva MA (2007) The pharmacology, neuroanatomy and neurogenetics of one-trial object recognition in rodents. *Neurosci Biobehav Rev* 31: 673–704.
66. Lebel E, Vallieres L, Rivest S (2000) Selective involvement of interleukin-6 in the transcriptional activation of the suppressor of cytokine signaling-3 in the brain during systemic immune challenges. *Endocrinology* 141: 3749–3763.
67. Cryan JF, Slattery DA (2007) Animal models of mood disorders: Recent developments. *Curr Opin Psychiatry* 20: 1–7.
68. Collin M, Hakansson-Ovesjo ML, Misane I, Ogren SO, Meister B (2000) Decreased 5-HT transporter mRNA in neurons of the dorsal raphe nucleus and

- behavioral depression in the obese leptin-deficient ob/ob mouse. *Brain Res Mol Brain Res* 81: 51–61.
69. Lu A, Steiner MA, Whittle N, Vogl AM, Walser SM, et al. (2008) Conditional mouse mutants highlight mechanisms of corticotropin-releasing hormone effects on stress-coping behavior. *Mol Psychiatry* 13: 1028–1042.
  70. Frenois F, Moreau M, O'Connor J, Lawson M, Micon C, et al. (2007) Lipopolysaccharide induces delayed FosB/DeltaFosB immunostaining within the mouse extended amygdala, hippocampus and hypothalamus, that parallel the expression of depressive-like behavior. *Psychoneuroendocrinology* 32: 516–531.
  71. Maroun M, Akirav I (2008) Arousal and stress effects on consolidation and reconsolidation of recognition memory. *Neuropsychopharmacology* 33: 394–405.
  72. Okuda S, Roozendaal B, McGaugh JL (2004) Glucocorticoid effects on object recognition memory require training-associated emotional arousal. *Proc Natl Acad Sci U S A* 101: 853–858.
  73. van Dam AM, Poole S, Schultzberg M, Zavala F, Tilders FJ (1998) Effects of peripheral administration of LPS on the expression of immunoreactive interleukin-1 alpha, beta, and receptor antagonist in rat brain. *Ann N Y Acad Sci* 840: 128–138.
  74. Laye S, Gheusi G, Cremona S, Combe C, Kelley K, et al. (2000) Endogenous brain IL-1 mediates LPS-induced anorexia and hypothalamic cytokine expression. *Am J Physiol Regul Integr Comp Physiol* 279: R93–98.
  75. Luheshi GN, Gardner JD, Rushforth DA, Loudon AS, Rothwell NJ (1999) Leptin actions on food intake and body temperature are mediated by IL-1. *Proc Natl Acad Sci U S A* 96: 7047–7052.
  76. Wu A, Ying Z, Gomez-Pinilla F (2004) The interplay between oxidative stress and brain-derived neurotrophic factor modulates the outcome of a saturated fat diet on synaptic plasticity and cognition. *Eur J Neurosci* 19: 1699–1707.
  77. Tozuka Y, Kumon M, Wada E, Onodera M, Mochizuki H, et al. (2010) Maternal obesity impairs hippocampal BDNF production and spatial learning performance in young mouse offspring. *Neurochem Int* 57: 235–247.
  78. Broadbent NJ, Squire LR, Clark RE (2004) Spatial memory, recognition memory, and the hippocampus. *Proc Natl Acad Sci U S A* 101: 14515–14520.
  79. Tye KM, Prakash R, Kim SY, Fenno LE, Grosenick L, et al. (2011) Amygdala circuitry mediating reversible and bidirectional control of anxiety. *Nature* 471: 358–362.
  80. Balschun D, Wetzel W, Del Rey A, Pitossi F, Schneider H, et al. (2004) Interleukin-6: a cytokine to forget. *FASEB J* 18: 1788–1790.
  81. Pervanidou P, Margeli A, Lazaropoulou C, Papassotiiriou I, Chrousos GP (2008) The immediate and long-term impact of physical and/or emotional stress from motor vehicle accidents on circulating stress hormones and adipo-cytokines in children and adolescents. *Stress* 11: 438–447.
  82. Salome N, Tasiemski A, Dutriez I, Wigger A, Landgraf R, et al. (2008) Immune challenge induces differential corticosterone and interleukin-6 responsiveness in rats bred for extremes in anxiety-related behavior. *Neuroscience* 151: 1112–1118.
  83. Akanmu MA, Nwabudike NL, Ilesanmi OR (2009) Analgesic, learning and memory and anxiolytic effects of insulin in mice. *Behav Brain Res* 196: 237–241.
  84. Finger BC, Dinan TG, Cryan JF (2010) Leptin-deficient mice retain normal appetitive spatial learning yet exhibit marked increases in anxiety-related behaviours. *Psychopharmacology* 210: 559–568.
  85. Kuhad A, Bishnoi M, Tiwari V, Chopra K (2009) Suppression of NF-kappabeta signaling pathway by tocotrienol can prevent diabetes associated cognitive deficits. *Pharmacol Biochem Behav* 92: 251–259.
  86. Raison CL, Miller AH (2003) When not enough is too much: the role of insufficient glucocorticoid signaling in the pathophysiology of stress-related disorders. *Am J Psychiatry* 160: 1554–1565.
  87. Konsman JP, Veeneman J, Combe C, Poole S, Luheshi GN, et al. (2008) Central nervous action of interleukin-1 mediates activation of limbic structures and behavioural depression in response to peripheral administration of bacterial lipopolysaccharide. *Eur J Neurosci* 28: 2499–2510.
  88. Belzung C, Griebel G (2001) Measuring normal and pathological anxiety-like behaviour in mice: a review. *Behav Brain Res* 125: 141–149.
  89. Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3: 1101–1108.

