



HAL
open science

Acute Schmallenberg virus infections, France, 2012

Corinne Sailleau, Emmanuel Breard, Cyril Viarouge, Alexandra Desprat, Virginie Doceul, Estelle Lara, Jérôme Languille, Damien Vitour, Houssam Attoui, Stéphan Zientara

► **To cite this version:**

Corinne Sailleau, Emmanuel Breard, Cyril Viarouge, Alexandra Desprat, Virginie Doceul, et al.. Acute Schmallenberg virus infections, France, 2012. *Emerging Infectious Diseases*, 2013, 19 (2), pp.321-322. 10.3201/eid1902.121281 . hal-02651880

HAL Id: hal-02651880

<https://hal.inrae.fr/hal-02651880>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Acute Schmallenberg Virus Infections, France, 2012

To the Editor: After unexpected emergence of bluetongue virus serotype 8 in northern Europe in 2006 (1), another arbovirus, Schmallenberg virus (SBV), which is transmitted by *Culicoides* spp. biting midges, emerged in Europe in 2011 and caused disease outbreaks among ruminants (2). Nonspecific clinical signs such as fever, decreased milk production, and diarrhea were associated with acute infection in cattle, and late abortions and birth defects in newborns were associated with infection of pregnant cows, ewes, and goats (2,3).

SBV, which belongs to the family *Bunyaviridae* and genus *Orthobunyavirus*, was detected in Germany, the Netherlands, and Belgium in 2011. This virus was later detected in the United Kingdom, France, Italy, Luxembourg, Spain, Denmark, and Switzerland (4). As of August 1, 2012, a total of 5,701 infected farms were reported in Europe (2,498 sheep farms, 3,124 cattle farms, and 79 goat farms) (www.survepi.org/cerepi/). France has been the country most affected: it had 2,650 SBV-infected farms (5), (i.e., in which ≥ 1 malformed offspring was positive for SBV by real-time reverse transcription PCR [RT-PCR] on 1,128 sheep farms, 1,505 cattle farms, and 17 goat farms).

Abnormalities detected in offspring in 2011 and in early 2012 were caused by infections acquired in 2011 (4). At that time, it was unclear whether SBV could survive the 2011–2012 winter and remain a threat to Europe. We report data suggesting that SBV overwintered or was reintroduced in France.

On May 16, 2012, a herd of 75 dairy cows in southwestern France (Pyrénées-Atlantiques) had hyperthermia and decreased milk

production. Of 18 cows tested by the Agence Nationale de Sécurité Sanitaire (Maisons-Alfort, France), 9 were positive for SBV by PCR (6) (cycle threshold [C_t] range 17–36.5) and negative for SBV by ELISA (IDVet, Montpellier, France), 1 was positive by PCR and ELISA, and 8 were negative by PCR and ELISA (Figure).

On May 23, a week after the first samples were collected, all cows tested were positive by ELISA and only 1/18 cows were positive by RT-PCR. On June 29, all 18 cows tested were negative by RT-PCR. Detection of SBV-neutralizing antibodies by virus neutralization assay in serum samples obtained on May 23 confirmed SBV ELISA results and showed that a commercial IgG ELISA is suitable for detection of acute cases of SBV.

Viremia, as measured by RT-PCR, occurs during the first 5 days after acute infection (2). Antibody response against SBV, as measured by ELISA, is detected during or after the first 10 days after experimental infection (C. Sailleau et al., unpub. data). Accordingly, serologic and

molecular data showed that acute SBV infection occurred in cattle in southwestern France in May 2012, suggesting that SBV overwintered or was reintroduced. Moreover, in July 2012, another case of acute SBV infection was identified in Finistère (Brittany). A cow with hyperthermia and diarrhea was SBV positive by RT-PCR (C_t 31) and negative by ELISA, which indicated a recent SBV infection.

Three blood samples (each 170 μ L) from RT-PCR-positive cows (C_t range 18–13) (1 from the Finistère and 2 from Pyrénées-Atlantiques) were injected into 3 adult IFNAR^{-/-} mice. Seventy microliters of each sample was injected intraperitoneally and 100 μ L was injected into the neck scruff of the same mice. After 4 days, the mice showed no clinical signs and their weights were unchanged. Blood samples were collected and tested for SBV by RT-PCR. All mice were positive for SBV RNA (C_t range 18–31). These data showed that blood samples from cows contained virus RNA and confirmed that SBV reemerged in 2012.




Figure. Number of cows positive for Schmallenberg virus according to reverse transcription PCR (RT-PCR) and ELISA, France, 2012.

On July 25, 2012, SBV infection was identified in a cow in Jura Canton in the northwestern, French-speaking region of Switzerland (Romandie) (7). A serologic study conducted in the United Kingdom showed that several cattle and sheep seroconverted for SBV in 2012 (8). However, our data show that SBV survived the winter, when midge numbers decreased. The precise mechanisms of SBV overwintering are not known and need to be explored.

The consequences of SBV recirculation should be investigated, particularly in pregnant cows, ewes, and goats. The 2 SBV-positive farms described in this report are located in a previously SBV-free area (Finistère-Brittany) or an area in which the infection rate was low (Pyrénées-Atlantiques) in the winter of 2011–2012, during which seroprevalence for most herds was probably weak (C. Sailleau et al., unpub. data). Therefore, reemergence of cases of congenital forms of SBV infection in France and others areas of Europe can be expected.

Acknowledgments

We thank the veterinary services in France and Nicolas Fradin for contributing to this study, the veterinary practitioners for collecting blood samples, and the farmers for permitting their animals to be included in this study.

Corinne Sailleau, Emmanuel Bréard, Cyril Viarouge, Alexandra Desprat, Virginie Doceul, Estelle Lara, Jérôme Languille, Damien Vitour, Houssam Attoui, and Stéphan Zientara

Author affiliations: Agence Nationale de Sécurité Sanitaire, Maisons-Alfort, France (C. Sailleau, E. Bréard, C. Viarouge, A. Desprat, V. Doceul, E. Lara, D. Vitour, S. Zientara); Direction Générale de l'Alimentation, Paris, France (J. Languille); and Institute for Animal Health, Pirbright, UK (H. Attoui)

DOI: <http://dx.doi.org/10.3201/eid1902.121281>

References

1. Toussaint JF, Vandenbussche F, Mast J, De Meester L, Goris N, Van Dessel W, et al. Bluetongue in northern Europe. *Vet Rec.* 2006;159:327. <http://dx.doi.org/10.1136/vr.159.10.327-a>
2. Hoffmann B, Scheuch M, Hoper D, Jungblut R, Holsteg M, Schirmmeier H, et al. Novel orthobunyavirus in cattle, Europe, 2011. *Emerg Infect Dis.* 2012;18:469–72. <http://dx.doi.org/10.3201/eid1803.111905>
3. van den Brom R, Lutikholt SJ, Lievaart-Peterson K, Peperkamp NH, Mars MH, van der Poel WH, et al. Epizootic of ovine congenital malformations associated with Schmallenberg virus infection. *Tijdschr Diergeneeskd.* 2012;137:106–11.
4. European Food Safety Authority. Schmallenberg virus: analysis of the epidemiological data and assessment of impact. European Food Safety Authority. 2012;10: 2768–857. <http://www.efsa.europa.eu/en/efsajournal/doc/2768.pdf>
5. Dominguez M, Hendriks P, Zientara S, Calavas D, Jay M, Touratier A, et al. Preliminary estimate of Schmallenberg virus infection impact in sheep flocks, France. *Vet Rec.* 2012;171:426. <http://dx.doi.org/10.1136/vr.100883>
6. Bilk S, Schulze C, Fischer M, Beer M, Hlinak A, Hoffmann B. Organ distribution of Schmallenberg virus RNA in malformed newborns. *Vet Microbiol.* 2012;159:236–8. <http://dx.doi.org/10.1016/j.vetmic.2012.03.035>
7. Schmallenberg virus - Europe (48), 1st case 2012. *ProMed* 2012 [cited 2012 Jul 22]. www.promedmail.org, archive no. 20120721.1210343
8. Anonymous. Schmallenberg virus 'still circulating' in the UK. *Vet Rec.* 2012;171:140. <http://dx.doi.org/10.1136/vr.e5373>

Address for correspondence: Corinne Sailleau, Unité Mixte de Recherche Virologie 1161, Agence Nationale de Sécurité Sanitaire, 23 Ave du Générale de Gaulle, Maisons-Alfort 94076, France; email: corinne.sailleau@anses.fr

Toscana Virus Isolated from Sandflies, Tunisia

To the Editor: Toscana virus (TOSV; genus *Phlebovirus*, family *Bunyaviridae*) is transmitted by sandflies, mostly the species *Phlebotomus perniciosus* and *P. perfiliewi* (1). Initially discovered in central Italy, TOSV was recently identified in other European countries (i.e., Portugal, Spain, France, Croatia, and Turkey) (2). TOSV is a primary cause of aseptic meningitis during warm months (2). A seroprevalence study suggested that TOSV is present in Tunisia and may cause neuroinvasive infections, but definitive evidence of TOSV circulation has not been possible because it is difficult to distinguish from the antigenically related phlebovirus Punique virus (3,4); both viruses are members of the species *Sandfly fever Naples virus*. We investigated the prevalence of TOSV among sandflies in northern Tunisia.

A total of 5,288 sandflies (3,547 females, 1,740 males) were collected during June–October 2010 by using CDC light traps (John W. Hock Company, Gainesville, FL, USA) at Utique (37°08'N, 7°74'E), a focus for visceral leishmaniasis in northern Tunisia. Sandflies were separated by sex and trapping nights and pooled with ≥ 30 specimens by pool. Pools were processed as described (4) and subjected to PCR detection of phlebovirus RNA targeting 2 genes independently (4–6) and virus isolation onto Vero cells. Of 249 pools processed, 8 strains of phleboviruses were isolated: 2 TOSV, 3 Punique virus, and 3 other phleboviruses currently being characterized.

TOSV strains were obtained from 2 pools of sandflies trapped in September 2010, T152 and T166, consisting of 30 males and 30 females, respectively. These pools were positive for TOSV RNA by sequencing of

