



HAL
open science

Extracellular superoxide dismutase in insects: characterization, function and interspecific variation in parasitoid wasps' venom

Dominique Colinet, Dominique Cazes, Maya Belghazi, Jean-Luc Gatti,
Marylene Poirie

► To cite this version:

Dominique Colinet, Dominique Cazes, Maya Belghazi, Jean-Luc Gatti, Marylene Poirie. Extracellular superoxide dismutase in insects: characterization, function and interspecific variation in parasitoid wasps' venom. *Journal of Biological Chemistry*, 2011, 286, pp.40110-40121. 10.1074/jbc.M111.288845 . hal-02652213

HAL Id: hal-02652213

<https://hal.inrae.fr/hal-02652213v1>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manuscript Title: Extracellular Superoxide Dismutase in Insects: Characterization, Function and Interspecific Variation in Parasitoid Wasps' Venom

Manuscript No: JBC/2011/288845

Manuscript Type: Regular Paper

Date Submitted by the Author: 2 Aug 2011

Complete List of Authors: Dominique Colinet, Dominique Cazes, Maya Belghazi, Jean-Luc Gatti, and Marylene Poirie

Keywords: Drosophila; Insect immunity; Invertebrates; Superoxide dismutase (SOD) ; Virulence factors; Parasitoid wasp; Phenoloxidase

Extracellular Superoxide Dismutase in Insects:
Characterization, Function and Interspecific Variation in Parasitoid Wasps' Venom*

**Dominique Colinet^{1,2,3}, Dominique Cazes^{1,2,3}, Maya Belghazi⁴,
Jean-Luc Gatti^{1,2,3} and Marylène Poirié^{1,2,3}**

¹From the Evolution and Specificity of Multitrophic Interactions (ESIM),
UMR 1301 "Biotic Interactions and Plant Health" (IBSV),
Institut National de la Recherche Agronomique, INRA PACA, Sophia Antipolis, France

²UMR 6243, Centre National de la Recherche Scientifique, CNRS, France

³Université Nice Sophia Antipolis, UFR Sciences, France

⁴Centre d'Analyses Protéomiques de Marseille (CAPM), IFR Jean Roche, Faculté de Médecine -
Secteur Nord, Université de la Méditerranée, Marseille, France

*Running title: *Superoxide dismutases in parasitoids' venom*

To whom correspondence should be addressed: Colinet D., Evolution and Specificity of Multitrophic Interactions (ESIM), UMR 1301 "Biotic Interactions and Plant Health" (IBSV), Institut National de la Recherche Agronomique, INRA PACA, 400 route des Chappes, Sophia Antipolis, 06903, France. Tel.: +33492386409; E-mail: Dominique.Colinet@sophia.inra.fr

Key-words: Superoxide dismutases, phenoloxidase, parasitoid wasp, venom

Background: Endoparasitoid wasps inject venom proteins at oviposition to alter host immunity.

Results: The venom of *Leptopilina boulardi* but not of a related species contains an active extracellular SOD. Recombinant SODs inhibit the *Drosophila* host phenoloxidase activity.

Conclusion: An extracellular SOD is secreted and active in an insect fluid.

Significance: SODs may be used as immune suppressive virulence factors by parasitoid wasps.

SUMMARY

Endoparasitoid wasps inject venom proteins with their eggs to protect them from the host immune response and ensure successful parasitism. Here, we report identification of intracellular (SOD1) and extracellular (SOD3) Cu,Zn superoxide dismutase transcripts in the venom apparatus of two *Leptopilina* species, parasitoids of *Drosophila*. *Leptopilina* SODs show sequence and structure similarity to human SODs, but phylogenetic analyses indicate that the extracellular SODs are more related to cytoplasmic vertebrate SODs than to extracellular ones, a feature shared by predicted insect extracellular SODs. We

demonstrate that *L. boulardi* SOD3 (LbSOD3) is indeed secreted and active as monomeric glycosylated forms in venom. Our results also evidence quantitative variation in SOD3 venom contents between closely related parasitoid species since *sod3* is 100-fold less expressed in *L. heterotoma* venom apparatus and no protein and SOD activity are detected in its venom. *Leptopilina* recombinant SOD3s as well as a mammalian SOD *in vitro* inhibit the *Drosophila* phenoloxidase activity in a dose-dependent manner, demonstrating that SODs may interfere with the *Drosophila* melanization process and therefore with production of cytotoxic compounds. Although the recombinant LbSOD3 quantity needed to observe this effect precludes a systemic effect of the wasp venom SOD3, it is still consistent with a local action at oviposition. This work provides the first demonstration that insect extracellular SODs are indeed secreted and active in an insect fluid and can be used as virulence factors to counteract the host immune response, a strategy largely used by bacterial and fungal pathogens but also protozoan parasites during infection.

Endoparasitoid wasps are insects that lay eggs in arthropods, generally other insects, eventually killing them as the result of their

development. To ensure successful parasitism, they inject with the egg various components, including venom factors that alter the host immunity and/or manipulate its development and physiology (1,2). The typical host immune defense against parasitoids is the formation of a multicellular melanized capsule around the egg, which requires the coordination of cellular and humoral components (3,4). Following egg parasite recognition, host hemocytes proliferate, become adhesive, attach and spread at the surface of the egg (5). In the meantime, a cascade involving several serine proteases is triggered, leading to the cleavage of prophenoloxidase (proPO) into active phenoloxidase (PO) which catalyzes oxidation of phenolic compounds that polymerize to form melanin (4,6). This melanization pathway involves oxidoreduction reactions and generates diverse cytotoxic molecules including reactive oxygen and nitrogen species (ROS/RNS) that have been suggested to play a major role in the death of the parasitoid egg (7-9).

Until now, a limited number of proteins have been identified and characterized from parasitoid venom fluids and only a few of them have been demonstrated to interfere with host immune defenses (10). Known immune suppressive proteins target either host hemocytes or the PO cascade (10), as shown for *Leptopilina boulardi*, a parasitoid of *Drosophila* species of the *melanogaster* subgroup (2). This species is likely the best known parasitoid model regarding venom virulence factors, and it is also an important model for understanding intraspecific variation of virulence with occurrence of two well-characterized strains, ISm and ISy (reviewed in 11). The venom of ISm females, a highly virulent strain against *D. melanogaster*, contains a RhoGAP protein that is required for parasitism success (12). This factor targets *Drosophila* lamellocytes, the main capsule-forming hemocytes, and induces changes in their morphology by inactivating the RhoGTPases Rac1 and Rac2, both required for successful encapsulation (13-17). *L. boulardi* ISy females differ from ISm ones in their ability to suppress immune defenses of both *D. melanogaster* and *D. yakuba* hosts, but depending on the host resistance genotype (11,18). The main identified effect of ISy female venom is the inhibition of the host serine protease cascade leading to the proPO-PO conversion thanks to the action of a serpin, a specific serine protease inhibitor (19).

Although it has been suggested that *L. boulardi* ISm venom might interfere with the oxidation pathways during the melanization process (20), no parasitoid-injected protein had ever been described that could prevent the deleterious effects of the production of ROS or RNS. This is quite surprising since a well-known virulence strategy employed by numerous bacterial and fungal pathogens, including pathogens of insects, as well as protozoan parasites, is the detoxification of reactive species by antioxidant enzymes, mainly superoxide dismutases (SODs) (21-26). SODs are one of the most important cellular enzymatic defenses against the detrimental ROS generated by the aerobic metabolism. These ubiquitous metalloenzymes catalyze the dismutation of the superoxide anion, O_2^- , to hydrogen peroxide H_2O_2 , which in turn is converted into H_2O by catalase and peroxidases (reviewed in 27). In mammals, three distinct SODs have been identified and characterized which have a similar enzymatic function, although they differ in their metal cofactor requirements and/or cellular compartmentalization (reviewed in 28). The cytoplasmic Cu/ZnSOD (SOD1) is primarily localized in cytoplasm and also found in the mitochondrial inter-membrane space (29). The extracellular Cu,ZnSOD (SOD3) is similar in structure to SOD1 but contains a N-terminal signal peptide and a C-terminal region involved in extracellular matrix binding (30). Finally, the MnSOD (SOD2), which is very different in protein sequence and structure from Cu/ZnSODs, is restricted to the inner matrix of mitochondria (31).

While studying candidate virulence proteins in *Leptopilina* wasps' venom, we have identified predicted cytoplasmic and extracellular Cu,ZnSODs from the two strains of *L. boulardi* as well as from *L. heterotoma*, a more generalist species than *L. boulardi*. The extracellular SOD gene is specifically overexpressed in the venom apparatus of *L. boulardi* compared to the remnant body, in contrast to what observed for *L. heterotoma*. The corresponding extracellular SOD protein as well as SOD activity have been detected in *L. boulardi* venom only, and SOD identity was confirmed by mass spectrometry. *In vitro* assays using a recombinant protein show a direct inhibitory effect of the *L. boulardi* extracellular SOD on *D. melanogaster* PO activity.

To our knowledge, this work is the first demonstration that insect predicted extracellular

SODs are indeed secreted. It is also the first report of the presence and specific overproduction of a SOD protein in parasitoid wasps' venom and of its *in vitro* effect on host immune components. We suggest that parasitoid venom SODs likely have a role in parasitism success either directly, by interfering with the local host melanization pathway, or indirectly by protecting other virulence factors from ROS damages during their storage in the wasp's venom reservoir.

EXPERIMENTAL PROCEDURES

Biological material - The origin of *L. bouleardi* ISy (Lby; Gif stock 486) and ISm (Lbm; Gif stock 431) lines has been previously described (32). Briefly, they derive from single females collected respectively in Brazzaville (Congo) and Nasrallah (Tunisia). The *L. heterotoma* (Lh) strain originates from South of France (Gotheron). All parasitoid lines were reared at 25°C on a susceptible *D. melanogaster* strain (Gif stock 1333). The wasp venom apparatus that can be easily isolated consists in a unique long gland connected to a reservoir (2), that can be separated and processed independently.

Cloning and molecular analysis of Leptopilina Cu,ZnSODs - Total RNA was extracted from 100 venom apparatus (venom gland + reservoir) collected from Lbm, Lby or Lh, using TRIzol Reagent (Invitrogen). Three cDNA libraries were constructed using 1 µg of RNA and the Creator SMART cDNA Library Construction Kit (Clontech) as previously described (19). Among the 120 clones sequenced from each library and analyzed by BLAST against the NCBI non-redundant databases, three sequences had very high similarities with Cu,ZnSODs. They were named LbmSOD3, LbySOD3 and LhSOD1 according to the presence or the absence of a putative peptide signal predicted using the CBS prediction servers (<http://www.cbs.dtu.dk/services/>).

To obtain the corresponding sequences LbmSOD1, LbySOD1 and LhSOD3, RNA isolated from venom apparatus of Lbm, Lby or Lh was reverse transcribed using the iScript cDNA Synthesis Kit (BioRad). PCR were performed using the GoTaq DNA Polymerase (Promega) and the primer pairs designed from LhSOD1 and LbmSOD3/ LbySOD3 sequences (Supplemental Table 1). Amplified fragments were cloned into the pCR2.1-TOPO vector (Invitrogen) and sequenced.

Pairwise sequence comparisons were performed using the EMBOSS program Needle at EMBL-EBI (<http://www.ebi.ac.uk/emboss/align/>). Multiple amino acid sequence alignment was performed using MUSCLE (33), and phylogenetic analysis was conducted using PhyML (34) at LIRMM (<http://www.phylogeny.fr/>). Molecular modeling of *Leptopilina* Cu,ZnSODs was performed using the Phyre server (35) (<http://www.sbg.bio.ic.ac.uk/~phyre/>).

Visualization of structures, structural alignments and root-mean-square deviations (RMSD) were done using Pymol (<http://pymol.sourceforge.net/>). N-glycosylation sites were predicted using the CBS prediction servers (<http://www.cbs.dtu.dk/services/>).

Semi-quantitative and quantitative RT-PCR - RNA isolated from dissected Lby, Lbm or Lh venom apparatus or from the remnant female bodies (without venom apparatus) were reverse-transcribed as described above. PCRs were carried out with primers common to LbmSOD1/ LbySOD1, common to LbmSOD3/LbySOD3, specific to LhSOD1 and specific to LhSOD3 (Supplemental Table 1). Semi-qRT-PCR experiments were performed for 28 or 33 cycles. PCR products were then separated and quantified using the GelEval software (FrogDance Software). qRT-PCR experiments were carried out on an Opticon monitor 2 (BioRad) using the Absolute qPCR SYBR MasterMix Plus (Eurogentec). PCR conditions were 50°C for 2 min and 95°C for 10 min, followed by 40 cycles at 95°C for 30 s, 60°C for 30 s and 68°C for 30 s. Each reaction was performed in triplicate and the mean of three independent biological replicates was calculated. All data were normalized using the ITS2 ribosomal sequence (Internal Transcribed Spacer 2; Supplemental Table 1), and analysed by a Student's t test (R software package ; <http://www.r-project.org/>).

Production of recombinant proteins - For production of the thioredoxin (Trx) fusion proteins Trx-LbmSOD3, Trx-LbySOD3 and Trx-LhSOD3, cDNA fragments corresponding to the mature proteins were amplified by RT-PCR from RNA isolated from Lbm, Lby and Lh venom apparatus, respectively. Amplified fragments were cloned into pET-32a(+) vector (Novagen), and recombinant plasmids were transformed in competent Origami (DE3) *Escherichia coli* cells. Production and purification of Trx-LbmSOD3, Trx-LbySOD3, Trx-LhSOD3 and Trx-tag alone

were performed according to manufacturer's instructions.

Immunoblotting experiments - For dot blot experiments, serial 1:3 dilutions of recombinant Trx-LbmSOD3, Trx-LbySOD3, Trx-LhSOD3 and of Trx-tag, starting from 1 µg of protein, were spotted onto a nitrocellulose membrane. Following blocking with TBST-2% milk, the membrane was probed overnight at 4°C with a *H. sapiens* Superoxide Dismutase 1 (SOD1) rabbit polyclonal antibody (1:1000, ab13498, Abcam). After washing and a 2 h incubation at RT with a goat anti-rabbit IgG horseradish peroxidase conjugate (1:25000, Sigma), chemoluminescence signal detection was performed (Immobilon Western, Millipore). Relative spot intensities were quantified using GeneSnap and GeneTools (Syngene).

For all gel and Western blot experiments, venom reservoirs were dissected in insect Ringer and residual tissues were removed by centrifugation 2 min at 500g. For SDS-PAGE comparison, the content of 10 reservoirs was separated on a 12.5 % polyacrylamide gel before transfer. For reducing and non-reducing conditions, 60 reservoirs were dissected in 40 µl of Ringer and 20 µl were mixed with sample buffer and heated either in presence (R) or absence (NR) of 2.5% 2-mercaptoethanol before SDS-PAGE and transfer. For native PAGE comparison, the contents of 20 reservoirs was mixed (50/50, v/v) with sample buffer without SDS and 2-mercaptoethanol, and separated on a 7.5 % native PAGE. The gel was soaked 15 min in SDS-PAGE running buffer before transfer. All membranes were probed as described above.

N-Glycosylation assay - The content of 10 Lbm reservoirs was collected in 50 mM sodium phosphate buffer (pH 7.5) before supplementation with 0.2% SDS and 0.5% 2-mercaptoethanol. The sample was heated at 95°C for 10 min, cooled on ice for 2 min, and incubated with one unit of N-Glycosidase F (Roche) overnight at 37°C. The reaction was stopped at 95°C for 5 min. The extent of carbohydrate removal was determined by Western blot analysis as described above.

Mass spectrometry - Silver stained bands cut from a 6-16% SDS-PAGE gel were destained, reduced and alkylated with iodoacetamide and incubated overnight at 37°C with 12.5 ng/µl trypsin (Sigma). The solution was dried, reconstituted with 0.1% formic acid and sonicated for 10 min. Generated peptides were sequenced by nano-LC-tandem mass

spectrometry (MS/MS) (Q-TOF Ultima with a nano-electrospray ionization source, Waters/Micromass, UK) in data-dependent acquisition (DDA) mode using the five most intense parent ions. Peptides were loaded on a C18 column (XBridge™ BEH130 3,5 µm, 75 µm×150 mm, Waters) and eluted with a 5 to 60% linear gradient at a flow rate of 180 nl/min over 90 min (buffer A: water/acetonitrile (98:2, v/v) and 0.1% formic acid; buffer B: water/acetonitrile (20:80, v/v) and 0.1% formic acid). MS/MS data analysis was performed with Mascot software (www.matrixscience.com) using *Leptopilina* Cu,ZnSOD sequences and the following parameters: trypsin specificity, two missed cleavages, carbamidomethyl cysteine and oxidation of methionine as variable modification, and 0.2 Da mass tolerance on both precursor and fragment ions. To validate SOD identification, only matches with peptides with a score above the identity threshold of 13 ($p < 0,005$), and containing at least 5 consecutive y or b ions were considered.

SOD activity assays - SOD activity was detected by zymography following 7.5% non-denaturing, non-reducing electrophoresis at 100 V at 4°C. For recombinant Trx-LbmSOD3, Trx-LbySOD3, Trx-LhSOD3, and Trx-tag, 3 µg of proteins were loaded. For venom assays, the content of 25 venom reservoirs was loaded. SOD activity was visualized as inhibition of the reduction of NBT according to Beauchamp and Fridovich (36). Upon illumination, achromatic bands indicate zones of SOD activity. SOD spectrophotometric assays were performed using the SOD determination kit (Sigma). Assays were performed using 20 and 200 ng of recombinant protein or Trx as a control, 200 ng of BtSOD, a commercial SOD from bovine erythrocytes (Sigma), and protein extracts from 20 Lbm, Lby and Lh reservoirs. Inhibition assays were performed with 2 mM KCN. P values were generated by ANOVA followed by pairwise t-tests (R software package).

PO activity measurements - Phenoloxidase activity was assayed using a method modified from Colinet *et al.* (19). For each replicate, hemolymph of three *D. melanogaster* 96 h aged-larvae was collected in 10 µl cacodylate buffer (cacodylic acid 10 mM, CaCl₂ 5 mM, pH 7). After 15 min incubation at 21°C during which proPO is converted into PO, 10 µl of cacodylate buffer containing 0.003 to 3 µg of the recombinant protein, 3 µg of BtSOD or 3 to 45 µg of venom proteins (estimated using

Bradford assay; Coo Protein Assay, Interchim) was added. Samples were incubated 15 min at 21°C before addition of the substrate (180 µl of 1 mg/ml dopamine hydrochloride in cacodylate buffer). After 1 h at 21°C, PO activity was estimated spectrophotometrically at 490 nm by detecting the quantity of dopachrome formed. Hemolymph samples in which 10 µl of cacodylate buffer alone or containing 1-phenyl-2-thiourea (PTU, final concentration 0.01%), a PO inhibitor, were added after the initial 15 min incubation period were used as controls. To determine whether recombinant *Leptopilina* Cu,ZnSODs affect the PO activity itself, or the proPO to PO activation by serine proteases, absorbance values were compared between hemolymph samples collected in presence of recombinant proteins or in which the recombinant proteins were added after the 15 min incubation period that follows hemolymph collection. Hemolymph samples collected in PTU (0.01%), in which PTU was added after the 15 min incubation period, or collected in cacodylate buffer alone, were used as controls. Three replicates were made for each treatment. P values were generated by ANOVA followed by pairwise t tests (R software package).

RESULTS

Identification of Cu,ZnSOD sequences from Leptopilina venom apparatus - Among the 120 clones sequenced from each venom apparatus cDNA library of *L. bouleari* ISm (Lbm), *L. bouleari* ISy (Lby) and *L. heterotoma* (Lh), full length sequences with high similarity to Cu,ZnSODs were identified. Sequence comparisons and prediction of a peptide signal in *L. bouleari* ISm and ISy Cu,ZnSOD transcripts strongly suggested that they were both extracellular SODs. They were thus named LbmSOD3 and LbySOD3, according to the human nomenclature. The *L. heterotoma* sequence had no predicted peptide signal, suggesting that it was a cytoplasmic Cu,ZnSOD, and it was named LhSOD1. The sequences of *L. bouleari* LbmSOD1 and LbySOD1 and of *L. heterotoma* LhSOD3 were further obtained by RT-PCR, using specific primers designed from the homolog sequences (supplemental Table 1).

The predicted *Leptopilina* Cu,ZnSODs are proteins of 151 amino acids for SOD1 and 173 to 176 amino acids for SOD3 (Table 1 and Fig. 1A). The estimated molecular mass of the mature proteins is 15.5 kDa for SOD1 and 16.4 to 16.6 kDa for SOD3, but two and three N-

glycosylation sites are predicted for *Leptopilina* SOD3 and SOD1 respectively (Table 1). Cu,ZnSODs sequences are nearly identical between *L. bouleari* strains and very similar between *L. bouleari* and *L. heterotoma*, with LhSOD3 and LhSOD1 closely related to extracellular and cytoplasmic *L. bouleari* Cu,ZnSODs, respectively (Table 2 and Fig. 1A). As a whole, extracellular SODs are clearly more divergent from one another than are cytoplasmic SODs (Table 2). Multiple sequence alignment with the fully characterized *H. sapiens* SOD1 and SOD3 indicates that *Leptopilina* SODs contain all the conserved residues essential for dismutation, metal-binding and stability (28), suggesting that they are active Cu,ZnSODs (Fig. 1A). In addition, a strong conservation of key structural elements is shown by superposition of the predicted 3D models for *Leptopilina* proteins with the solved *H. sapiens* SOD structures (37) (see Fig. 1B, illustrated for LbmSOD3 only).

Leptopilina SOD3s are closely related to human SOD1 - Interestingly, sequence comparisons of *Leptopilina* extracellular SODs with human ones clearly show that they are more related to the human intracellular SOD1 than to the human extracellular SOD3 (Table 2 and Fig. 1A). More specifically, the *Leptopilina* SOD3 signal peptide differs from that of human SOD3 and *Leptopilina* SOD3 lack the heparin-binding C-terminal part of human SOD3 involved in the binding to the cellular matrix (30). In addition, *Leptopilina* and human SOD3 differ in the number of disulfide bridges. The human SOD contains two intrasubunit disulfide bonds : Cys107-Cys189, also found in human SOD1 and predicted for *Leptopilina* SOD1s, and Cys45-Cys190 unique to human SOD3 (38,39). The specific C-terminal heparin-binding region of human SOD3 also contains a cystein (Cys219) implicated in an additional intersubunit disulfide bridge, necessary for the formation of the human SOD3 tetramer (38,39). Only the disulfide bridge geometrically similar to Cys107-Cys189 could be predicted for *Leptopilina* SOD3 since a Val and a Gly are found at positions corresponding to Cys45 and Cys190 (Figure 1A), and the absence of the C-terminal region precludes the formation of the intersubunit disulfide bridge. Finally, comparisons of solved human and *Leptopilina* predicted SOD1 and SOD3 structures confirmed that *Leptopilina* extracellular SODs are more different from human SOD3 (with RMSD ranging from 0.839 to 0.851 Å for 116 to 127 α -carbon atoms), than from human SOD1 (with

RMSD ranging from 0.448 to 0.449 Å for 125 α -carbon atoms; Table 3). Among key structural elements, the Greek key loop 1 (GK1, Figure 1B), involved in SOD stability (39), is longer in human SOD3 only.

Phylogenetic analysis with selected vertebrate and insect Cu,ZnSOD sequences (Fig. 2A), and with available hymenopteran Cu,ZnSOD sequences (Fig. 2B), confirms that *Leptopilina* SOD3 clusters with predicted extracellular insect Cu,ZnSODs while *Leptopilina* SOD1 groups with cytoplasmic Cu,ZnSODs (Fig. 2A and B). In agreement with sequence and structure comparisons, extracellular vertebrate and insect Cu,ZnSOD sequences form largely separated clusters, insect extracellular Cu,ZnSODs being more closely related to cytoplasmic vertebrate Cu,ZnSODs than to extracellular ones (Fig. 2A).

The extracellular Cu,ZnSOD gene is overexpressed in L. bouleari venom apparatus - The relative abundance of Cu,ZnSOD transcripts in venom apparatus of *Leptopilina* wasps was estimated by semi-qRT-PCR using ITS2 as a reference gene. After 28 and 33 cycles, *L. bouleari* *lbmsod3* and *lbysod3* genes were found 20 and 30 times more expressed in venom apparatus than *lbmsod1* and *lbysod1*, while in *L. heterotoma*, *lhsod1* was 10 times more expressed in venom tissues than *lhsod3* (Fig. 3A, shown for 28 cycles). Overall, the expression level of *sod3* in venom apparatus was approximately 100 times higher in *L. bouleari* than in *L. heterotoma*, while expression of *sod1* was only twice higher. Using a similar approach, *lbmsod3* and *lbysod3* were found more expressed in venom-producing tissues than in the remnant body in contrast to *lhsod3* and all cytoplasmic Cu,ZnSOD genes (Fig. 3B, shown for 28 cycles). qRT-PCR results confirmed that extracellular Cu,ZnSODs gene expression in *L. bouleari* venom apparatus is 7 to 9 times higher than in the remnant body (Fig. 3C), with no significant difference between *lbmsod3* and *lbysod3* (Student t-test: $t = -0.6117$, $df = 3$, $p = 0.584$). All these results demonstrate that only *L. bouleari* wasps specifically overexpress extracellular Cu,ZnSOD genes in venom-producing tissues.

SOD3 proteins are secreted in different amounts in L. bouleari and L. heterotoma venom - To confirm that predicted extracellular Cu,ZnSODs are secreted in venom, we performed immunoblotting using a polyclonal antibody against *H. sapiens* SOD1, which

recognizes the recombinant Trx-fusion proteins LbmSOD3, LbySOD3 and LhSOD3 with the same efficiency (Fig. 4A). On a Western blot with venom from 10 female reservoirs, four bands of different intensity, ranging from 19 kDa to 25 kDa were detected for Lbm and Lby, whereas only a very faint band at 19 kDa was obtained for Lh (Fig. 4B). By comparing signal intensities between Lbm and Lby venom extracts and dilutions of recombinant LbmSOD3s, we estimated the total quantity of SOD as approximately 10 ng per reservoir for both *L. bouleari* strains (results not shown).

To ascertain the identity of the proteins detected in Lbm and Lby venom, we performed mass spectrometry on gel bands cut at the corresponding position on an SDS-PAGE gel of venom proteins. For both strains, we found peptides specifically matching with the extracellular Cu,ZnSOD (Table 4) demonstrating that this protein is indeed secreted in *L. bouleari* venom. By contrast, no peptides corresponding to SOD3 were found in *L. heterotoma* venom. Some peptides were also found that matched SOD1 in *L. bouleari* and *L. heterotoma*, possibly because of the presence of contaminants originating from cells leakage during venom collection. All these results show that the SOD antibody signal in *L. bouleari* venom essentially corresponds to detection of SOD3.

In mammals, Cu,ZnSODs are classically found in SDS-PAGE as dimers or tetramers even in presence of a reducing agent (28). In contrast, in *L. bouleari*, Cu,ZnSOD migration profiles were similar between reducing and non-reducing conditions (Fig. 4D), which suggests that LbSOD3 is mainly under the monomeric form in venom. Besides, treatment of *L. bouleari* ISm venom extracts with N-Glycosidase F induced a shift in size for the strongest reactive bands, suggesting the presence of N-glycosylations, although deglycosylation might be incomplete (Fig. 4C). The 19 kDa to 25 kDa bands thus likely correspond to SOD3 monomers with post-translational modifications (Fig. 4B).

Cu,ZnSODs are active in the venom of L. bouleari. SOD zymography and spectrophotometric assays clearly demonstrated that the recombinant Trx-LbmSOD3, Trx-LbySOD3 and Trx-LhSOD3 proteins were all active (Fig. 5A). The level of activity was similar between these proteins and with that of BtSOD, a commercial SOD from bovine erythrocytes (Fig. 5B; ANOVA : $df = 9$, $F = 67.238$, $p > 0.9$). All Trx-SODs and BtSOD activities significantly

decreased with the concentration of the protein (Fig. 5B; ANOVA : $df = 9$, $F = 67.238$, $p < 0.001$) and there was no activity for the Trx tag alone. In contrast to that of BtSOD, the activity of Trx-LbmSOD3 decreased at pH 10 and above (Fig. 5C; ANOVA : $df = 11$, $F = 11.954$, $p < 0.05$).

Using zymographic assays, a SOD activity was detected in venom of both *L. boulandi* strains (Fig 6A, and 6C), at a position corresponding to the signal observed on Western blot under native conditions (Fig. 6A, not shown for Lby). In contrast, no activity was observed in *L. heterotoma* venom (Fig. 6A). Spectrophotometric assays confirmed that the venom of *L. boulandi* strains has a similar SOD activity (Fig. 6B ; ANOVA : $df = 3$, $F = 113.94$, $p = 1$). Besides, venom samples, which contain 200 ng of SOD, had a SOD activity equivalent to that of 200 ng of recombinant LbmSOD3, suggesting that recombinant and venom SOD3 have a similar specific activity (Fig. 6B ; ANOVA : $df = 3$, $F = 113.94$, $p = 1$). Finally, a complete inhibition was observed when adding KCN, both in zymography (Fig. 6C) and spectrophotometric assays (Fig. 6D, not shown for Lby), indicating that all venom activity is due to a Cu/ZnSOD activity.

Cu,ZnSODs inhibit PO activity in D. melanogaster. To test whether *Leptopilina* extracellular Cu,ZnSODs may interfere with the host PO cascade, PO assays were conducted on *Drosophila* hemolymph samples in presence of recombinant Trx-LbmSOD3 or Trx tag alone as a control (Fig. 7A). The PO activity was decreased by 70 % using 3 μg of recombinant protein and by 40 % using 0.3 μg (Fig. 6A ; ANOVA : $df = 6$, $F = 49.334$, $p < 0.01$), while addition of lower concentrations (0.03 μg and 0.003 μg) had no inhibitory effect, as Trx tag or buffer alone (Fig. 7A, $p > 0.5$). The same inhibition level was observed with 3 μg of Trx-LbySOD3, of Trx-LhSOD3 and, interestingly, of BtSOD (Fig. 7B ; ANOVA : $df = 5$; $F = 19.3058$, $p < 0.01$). As expected, PO activity was totally inhibited by PTU.

To investigate whether the reduction of absorbance reflected the inhibition of proPO to PO conversion through action on the serine protease cascade or the inhibition of the PO activity itself, we performed parallel experiments in which Trx-LbmSOD3 protein, Trx tag alone, or PTU, were added either before hemolymph collection or after proPO to PO conversion (Figure 7C). The inhibitory effect of PTU and of

Trx-LbmSOD3 did not differ whether they were added before or after PO activation (Figure 7C ; ANOVA : $df = 6$, $F = 18.4594$, $p = 1$), thus demonstrating that recombinant SODs do not affect the PO activation cascade but PO activity itself.

Because of the presence of a serpin in *L. boulandi* ISy venom which inhibits proPO to PO conversion (19), PO assays with *L. boulandi* ISm and ISy venom were performed by adding venom in conditions where the proPO was already converted into PO. Assays were performed using venom quantities containing from 0.04 μg to 0.6 μg of SOD. However, no significant effect was detected, even when using an amount of venom estimated to contain twice the minimal amount of recombinant SOD3 (0.3 μg) that induced a detectable effect on PO activity (data not shown).

DISCUSSION

In contrast to mammals, insects have long been thought to produce only the cytoplasmic and mitochondrial SODs. The presence of extracellular SODs was questioned only recently with the description of two highly divergent Cu/ZnSOD sequences in the ant *Lasius niger*, one of them containing a signal peptide (40). Subsequent database searches and phylogenetic analyses then revealed that insect genomes, including that of *D. melanogaster*, contain sequences corresponding to the two distinct Cu/ZnSOD families related to mammalian SOD1 and SOD3 respectively (40,41). However, although predicted by *in silico* analyses, occurrence of extracellular Cu/ZnSOD in insects had never been proven *in vivo*.

In this work, we demonstrate that insects produce and secrete extracellular Cu,ZnSODs. Indeed, having identified Cu,ZnSOD transcripts containing a signal peptide from venom apparatus of two *Leptopilina* wasps, parasitoids of *Drosophila*, we confirmed the presence of the corresponding secreted SOD3 protein in *L. boulandi* venom using Western blot and mass spectrometry identification. In accordance with the conservation of predicted essential residues and key structural elements, this SOD3 protein is active and its activity is specific of Cu,ZnSODs.

Interestingly, *Leptopilina* SOD3s are more similar in sequence and structure to human SOD1 than to human SOD3, and they lack in particular the human SOD3 motif involved in binding of the protein to the cellular matrix (30). Moreover, our results suggest that *Leptopilina*

venom SOD3s are active as monomers, in agreement with the absence of the intersubunit disulfide bridge and of one of the two intrasubunit disulfide bridges found in human SOD3. Phylogenetic analyses show that not only the *Leptopilina* but all the predicted insect SOD3s are more closely related to vertebrate SOD1 than SOD3 sequences, thus suggesting a rapid divergence of the extracellular vertebrate Cu,ZnSOD sequences.

The characterization of *Leptopilina* SOD3 is the first example of secretion of an active extracellular SOD in hymenopteran venom despite recent proteomic analyses from different species, including parasitoids wasps (42-45). This questions the potential role of SOD3 in *L. bouleardi* which, as most endoparasitoids, inject at oviposition venom toxins that among other effects counteract the host immune response. Production of ROS is part of this response, as evidenced for *D. melanogaster* (7-9), and the detoxification of ROS is known to be part of virulence mechanisms. SODs, for instance, are used as immune suppressive toxins by a large number of pathogens, including eukaryotic protozoan parasites (21-26).

A likely role of *L. bouleardi* venom SOD3 in parasitism success is supported by the specific overexpression of the *sod3* gene in the venom apparatus compared to other tissues, as shown for most parasitoid virulence factors (43-52), including two major venom toxins of *L. bouleardi* (17,19). In *L. heterotoma*, the SOD3 protein is hardly detected in venom, and the venom contains no measurable Cu,ZnSOD-specific activity. This suggests that SOD3 is not used as a virulence factor by *L. heterotoma* as it is by *L. bouleardi*, thus evidencing interspecific variation in venom toxins in *Leptopilina* parasitoids despite phylogenetic proximity and the use of similar hosts. We recently showed that quantitative difference in the LbGAP venom toxin was at the origin of the intraspecific variation of virulence between *L. bouleardi* ISm and ISy lines (12). No such variation was found for SOD3 proteins, neither in sequence, nor in venom quantity and activity, which seems to preclude a role of SODs in intraspecific variation of *L. bouleardi* virulence.

Interestingly, the recombinant *L. bouleardi* SOD3, but also a purified mammalian SOD, have inhibitory effects on *D. melanogaster* PO activity *in vitro*, in a dose-dependent manner. We thus hypothesized that *L. bouleardi* injection

of the SOD3 protein in *Drosophila* could result in a modification of enzymatic equilibria through dismutation of the superoxide anion $\cdot\text{O}_2^-$ that could disturb the host melanization pathway. However, no significant effect of *L. bouleardi* ISm nor ISy venom on PO activity was observed *in vitro* that could correspond to the SOD effect. This was not due to quantitative differences since the venom sample used in the assays contained a SOD3 quantity higher than the SOD3 recombinant protein amount that yielded 40% inhibition of PO activity. However, the very high quantity of venom used (corresponding to the content of 60 reservoirs) concentrated in the small volume of the assay, might have disturbed the PO assay that is very sensitive to salt and/or pH modifications (53). In any case, the SOD3 quantity in one unique female reservoir was estimated to be thirty times smaller than the minimal amount of recombinant SOD3 shown to inhibit *Drosophila* hemolymph PO activity *in vitro*.

The SOD3 effect could also be through direct detoxification of $\cdot\text{O}_2^-$ which is produced at high levels by the host during the melanization process (7). Although $\cdot\text{O}_2^-$ is not cytotoxic against the parasitoid (7), it likely plays a critical role in *Drosophila* cellular immunity by initiating the formation of $\cdot\text{OH}$, the most reactive free radical in biological systems, through the reduction of some metal ions (7,9).

If a systemic effect of injection of SOD3 in the host, whether directly on the superoxide anion or on PO activity, is rather unlikely due to its low quantity in venom, this does not preclude a local effect nearby the deposited parasitoid egg. Following recognition, the egg is indeed rapidly covered by host hemocytes that likely produce killing oxygen radicals (54). Besides, lamellocytes, the main forming-capsule hemocytes, specifically express the PO3 gene (55,56), and a deposit of melanin, indicating PO activity, is observed at the surface of the parasitoid egg (3,57,58).

Finally, a possible function of *Leptopilina* venom SODs might be the protection from oxidative stress of venom apparatus tissues and/or venom proteins during storage. Protection of venom gland tissues by the SOD1 protein was suggested for instance in the honeybee (42) and can be hypothesized as well in *Leptopilina* parasitoids. However, as the SOD3 protein is only secreted in venom and in a high amount in *L. bouleardi*, its role in protecting venom

components would suggest a specific need of this species that remains to be evidenced.

In summary, our results demonstrate that insects indeed produce and secrete extracellular SODs, and that venom-secreted SODs are likely used as virulence factors by parasitoid wasps, as

they are by pathogens of mammals and insects. The exact role of the venom SOD in parasitism success nevertheless remains an open question which might be explored in the next future through development of promising methods, such as RNAi, for parasitoids (59).

REFERENCES

- Godfray H. (1994) *Parasitoids : Behavioral and Evolutionary Ecology*. Princeton University Press, Princeton, NJ.
- Poirié, M., Carton, Y., and Dubuffet, A. (2009) *C. R. Biol.* **332**(2-3), 311–320
- Carton, Y., Poirié, M., and Nappi, A. J. (2008) *Insect Sci.* **15**, 67–87
- Nappi, A. J. (2010) *Invert. Surv. J.* **7**, 198–210
- Strand, M. (2008) *Insect Sci.* **15**, 1–14
- Cerenius, L., Lee, B. L., and Soderhall, K. (2008) *Trends Immunol.* **29**(6), 263–271
- Nappi, A. J., Vass, E., Frey, F., and Carton, Y. (1995) *Eur. J. Cell Biol.* **68**(4), 450–456
- Nappi, A. J. and Christensen, B. M. (2005) *Insect Biochem. Mol. Biol.* **35**(5), 443–459
- Nappi, A. J., Poirié, M., and Carton, Y. (2009) *Adv. Parasitol.* **70**, 99–121
- Asgari, S. and Rivers, D. B. (2010) *Annu. Rev. Entomol.* **56**, 313–335
- Dubuffet, A., Colinet, D., Anselme, C., Dupas, S., Carton, Y., and Poirié, M. (2009) *Adv. Parasitol.* **70**, 147–88
- Colinet, D., Schmitz, A., Cazes, D., Gatti, J.-L., and Poirié, M. (2010) *PLoS Pathog.* **6**(11), e1001206
- Labrosse, C., Eslin, P., Doury, G., Drezen, J. M., and Poirié, M. (2005a) *J. Insect Physiol.* **51**(2), 161–170
- Labrosse, C., Stasiak, K., Lesobre, J., Grangeia, A., Huguet, E., Drezen, J. M., and Poirié, M. (2005b) *Insect Biochem. Mol. Biol.* **35**(2), 93–103
- Williams, M. J., Ando, I., and Hultmark, D. (2005) *Genes to Cells* **10**(8), 813–823
- Williams, M. J., Wiklund, M.-L., Wikman, S., and Hultmark, D. (2006) *J. Cell Sci.* **119**(10), 2015–2024
- Colinet, D., Schmitz, A., Depoix, D., Crochard, D., and Poirié, M. (2007) *PLoS Pathog.* **3**(12), e203
- Dupas, S., Carton, Y., and Poirié, M. (2003) *Heredity* **90**(1), 84–89
- Colinet, D., Dubuffet, A., Cazes, D., Moreau, S., Drezen, J., Poirié, M. (2009) *Dev. Comp. Immunol.* **33**(5), 681–689
- Kohler, L. J., Carton, Y., Mastore, M., and Nappi, A. J. (2007) *Arch. Insect Biochem. Physiol.* **66**(2), 64–75
- Lynch, M. and Kuramitsu, H. (2000) *Microbes Infect.* **2**(10), 1245–55
- Hwang, C.-S., Rhie, G.-E., Oh, J.-H., Huh, W.-K., Yim, H.-S., and Kang, S.-O. (2002) *Microbiology* **148**(11), 3705–3713
- Ghosh, S., Goswami, S., and Adhya, S. (2003) *Biochem. J.* **369**, 447–452
- Narasipura, S., Ault, J. G., Behr, M. J., Chaturvedi, V., and Chaturvedi, S. (2003) *Mol. Microbiol.* **47**(6), 1681–1694
- Ammendola, S., Pasquali, P., Pacello, F., Rotilio, G., Castor, M., Libby, S. J., Figueroa-Bossi, N., Bossi, L., Fang, F. C., and Battistoni, A. (2008) *J. Biol. Chem.* **283**(20), 13688–13699
- Xie, X.-Q., Wang, J., Huang, B.-F., Ying, S.-H., and Feng, M.-F. (2010) *Appl. Microb. Cell Physiol.* **86**(5), 1543–1553
- Fridovich, I. (1995) *Annu. Rev. Biochem.* **64**, 97–112
- Perry, J.J.P., Shin, D.S., Getzoff, E.D., Tainer, J.A. (2010) *Biochem. Biophys. Acta* **1804**(2), 245–262
- Okado-Matsumoto, A., Fridovich, I. (2001) *J. Biol. Chem.* **276**(42), 38388–93
- Stenlund, P., Lindberg, M. J., and Tibell, A. E. (2002) *Biochemistry* **41**(9), 3168–3175
- Weisiger, R. A., and Fridovich, I. (1973) *J. Biol. Chem.* **248**(13), 4793–4796
- Dupas, S., Frey, F., and Carton, Y. (1998) *J. Hered.* **89**(4), 306–311
- Edgar, R. (2004) *Nucl. Acids Res.* **32**, 1792–1797

34. Guindon, S., Gascuel, O. (2003) *Syst. Biol.* **52**(5), 696-704
35. Kelley, L.A., Sternberg, M.J.E. (2009) *Nat. Protoc* **4**(3), 363-371
36. Beauchamp, C., Fridovich, I. (1971) *Anal. Biochem.* **44**(1), 276-287
37. DiDonato, M., Craig, L., Huff, M.E., Thayer, M.M., Cardoso, R.M., Kassmann, C.J., Lo, T.P., Bruns, C.K., Powers, E.T., Kelly, J.W., Getzoff, E.D., and Tainer, J.A. (2003) *J Mol. Biol.* **332**(3), 601-615
38. Petersen, S. V., Oury, T. D., Valnickova, Z., Thogersen, I. B., Hojrup, P., Crapo, J. D., and Enghild, J. J. (2003) *Proc. Natl. Acad. Sci. U S A* **100**(24), 13845-13880
39. Antonyuk, S. V., Strange, R. W., Marklund, S. L., and Hasnain, S. S. (2009) *J Mol. Biol.* **388**(2), 310-326
40. Parker, J. D., Parker, K. M., and Keller, L. (2004) *Insect Mol. Biol.* **13**(6), 587-594
41. Corona, M. and Robinson, G. E. (2006) *Insect Mol. Biol.* **15**(5), 687-701
42. Peiren, N., de Graaf, D. C., Vanrobaeys, F., Danneels, E. L., Devreese, B., Van Beeumen, J., and Jacobs, F. J. (2008) *Toxicon* **52**(1), 72-83
43. de Graaf, D. C., Aerts, M., Brunain, M., Desjardins, C. A., Jacobs, F. J., Werren, J. H., and Devreese, B. (2010) *Insect Mol. Biol.* **19**, 11-26
44. Vincent, B., Kaeslin, M., Roth, T., Heller, M., Poulain, J., Cousserans, F., Schaller, J., Poirié, M., Lanzrein, B., Drezen, J.-M., Moreau, S.J.M. (2010) *BMC Genomics* **11**, 693.
45. Zhu, J.-Y., Fang, Q., Wang, L., Hu, C., and Ye, G. Y. (2010) *Arch. Insect Biochem. Physiol.* **75**(1), 28-44
46. Asgari, S., Reineke, A., Beck, M., and Schmidt, O. (2002) *Insect Mol. Biol.* **11**(5), 477-485
47. Crawford, A. M., Brauning, R., Smolenski, G., Ferguson, C., Barton, D., Wheeler, T. T., and McCulloch, A. (2008) *Insect Mol. Biol.* **17**(3), 313-324
48. Parkinson, N., Smith, I., Weaver, R., and Edwards, J. P. (2001) *Insect Biochem. Mol. Biol.* **31**(1), 57-63
49. Price, D., Bell, H., Hinchliffe, G., Fitches, E., Weaver, R., and Gatehouse, J. (2009) *Insect Mol. Biol.* **18**(2), 195-202
50. Vinchon, S., Moreau, S. J. M., Drezen, J. M., Prévost, G., and Cherqui, A. (2010) *Insect Biochem. Mol. Biol.* **40**(1), 38-48
51. Zhang, G., Schmidt, O., and Asgari, S. (2006) *Dev. Comp. Immunol.* **30**(9), 756-764
52. Zhu, J., Ye, G., and Hu, C. (2008) *Toxicon* **51**(8), 1391-1399
53. Zufelato, M. S., Lourenço, A. P., Simoes, Z. L. P., Jorge, J. A., and Bitondi, M. M. G. (2004) *Insect Biochem. Mol. Biol.* **34**(12), 1257-68
54. Nappi, A. J. and Vass, E. (1998) *J. Parasitol.* **84**(6), 1150-1157
55. Irving, P., Ubeda, J.-M., Doucet, D., Troxler, L., Lagueux, M., Zachary, D., Hoffmann, J. A., Hetru, C., and Meister, M. (2005) *Cell. Microbiol.* **7**(3), 335-350
56. Nam, H., Jang, I., Asano, T., and Lee, W. (2008) *Mol. Cells* **26**(6)
57. Cerenius, L., Kawabata, S.-I., Lee, B. L., Nonaka, M., and Söderhäll, K. (2010) *Trends Biochem. Sci.* **35**(10), 575-583
58. Reumer, A., Van Loy, T., and Schoofs, L. (2010) *Invert. Surv. J.* **7**, 32-44
59. Werren, J. H., Loehlin, D. W., and Giebel, J. D. (2009) *CSH Protoc.* **2009**(10), pdb.prot5311

Acknowledgments- We are grateful to Christian Rebuf and Didier Crochard for technical assistance, to Didier Herouart for help with SOD zymography assays and to Hugo Mathé-Hubert for help with statistical analyses.

FOOTNOTES

* This work was supported by grants CLIMEVOL (ANR-08-BLAN-0231) and PARATOXOSE (ANR-09-BLAN-0243-01) from the French National Agency for Research, and grants from the department "Plant Health and Environment" from the French National Institute for Agricultural Research (INRA).

¹The LbmSOD1, LbmSOD3, LbySOD1, LbySOD3, LhSOD1 and LhSOD3 nucleotide sequences have been submitted to the GenBank database with accession numbers JF770428, JF770429, JF770430, JF770431, JF770432 and JF770433 respectively.

²The abbreviations used are: Lbm, *Leptopilina bouvardi* ISm; Lby, *Leptopilina bouvardi* ISy; Lh, *Leptopilina heterotoma*; PNGase, N-Glycosidase F; PO, phenoloxidase; ROS, reactive oxygen species; SOD, superoxide dismutase; Trx, Thioredoxin.

FIGURE LEGENDS

FIGURE 1. Sequence and structure conservation of Cu,ZnSOD sequences. (A) Multiple sequence alignment of *Leptopilina* and *Homo sapiens* (SOD1, P00441 and SOD3, P08294) Cu,ZnSODs. Residues identical to HsSOD1 sequence are illustrated by a dot. The signal peptide is indicated by a blue box. Functional residues are highlighted in red (28). Red stars : positions of HsSOD3 Cys107-Cys189 involved in the Cu,ZnSODs conserved intrasubunit disulfide bridge ; Blue stars : positions of the Cys45-Cys190 involved in the HsSOD3 intrasubunit disulfide bridge. Green star : position of the the HsSOD3 Cys219 residue involved in the intersubunit disulfide bridge ; Green box : heparin-binding region. (B) Comparison of the predicted tertiary structure of LbmSOD3 with the solved HsSOD1 structure 1PU0. β -strands are displayed as flat arrows and helices are displayed as ribbons. Metal-liganding residues are shown as sticks. Key structural elements are color-coded : E-Loop, Electrostatic loop ; GK1 and GK2, Greek key loops 1 and 2; S-S, Disulfide region ; V-loop, Variable loop; Zn Region, Zinc binding region (28).

FIGURE 2. Maximum-likelihood phylogenetic trees. (A) Using vertebrate and insect Cu,ZnSOD sequences and (B) using Hymenopteran Cu,ZnSOD sequences (SOD3, extracellular SOD ; SOD1, cytoplasmic SOD). The dashed line delineates Hymenopteran extracellular Cu,ZnSODs. Numbers at corresponding nodes are bootstrap support values (500 bootstrap replicates). Numbers followed with * are bootstrap values for *Leptopilina* SOD1 nodes. Numbers followed with # are bootstrap values for *Leptopilina* SOD3 nodes. Only bootstrap support values $\geq 75\%$ are shown. Am, *Apis mellifera* ; Bi, *Bombus ignitus* ; Lb, *Leptopilina bouvardi* ; Lh, *Leptopilina heterotoma* ; Ln, *Lasius niger* ; Ny, *Nasonia vitripennis*.

FIGURE 3. Expression levels of *Leptopilina* Cu,ZnSOD genes. (A) Relative expression ratio of extracellular to cytoplasmic Cu,ZnSOD genes (logarithmic scale) in Lbm, Lby and Lh venom apparatus after 28 cycles of semi-qRT-PCR. (B) Relative expression ratio of *Leptopilina* Cu,ZnSOD genes (logarithmic scale) in venom apparatus compared to the remnant bodies after 28 cycles of semi-qRT-PCR. (C) Relative expression level (qRT-PCR) in venom-producing tissues compared to the remnant bodies for *lbmsod3* and *lbylod3* genes.

FIGURE 4. Immunoblotting experiments using a *H. sapiens* SOD1 polyclonal antibody. (A) Dot blot experiment on serial dilutions starting with 1 μ g of the recombinant proteins Trx-LbmSOD3, Trx-LbySOD3, Trx-LhSOD3 and Trx tag alone. (B) 12.5% SDS-PAGE (left) and equivalent Western blot (right) on protein extracts from 10 reservoirs of Lbm, Lby and Lh. (C) Western blot on protein extracts from 30 Lbm venom reservoirs in non-reducing (NR) and reducing (R) conditions. (D) Glycosylation profile analysis of *Leptopilina* Cu,ZnSOD using Western blot on extracts from 10 *L. bouvardi* ISm reservoirs either untreated (-) or treated (+) with N-Glycosidase F (PNGase, lane 2). MW in kDa.

FIGURE 5. SOD activity assays with recombinant proteins. (A) SOD zymography assay using 3 μ g of recombinant proteins Trx-LbmSOD3, Trx-LbySOD3 and Trx-LhSOD3. (B) SOD spectrophotometric assay using 200 and 20 ng of BtSOD and of Trx-LbmSOD3, Trx-LbySOD3 and Trx-LhSOD3. Trx alone is used as a control. (C) Effect of pH on activity performed with 200 ng of BtSOD and LbmSOD3. Mean values (\pm SE) are given for each category (three replicates). Asterisk indicates statistically significant differences (* : $p < 0.05$; *** : $p < 0.001$).

FIGURE 6. SOD activity assays with venom. (A) SOD zymography assay and Western blot experiment in native conditions with extracts of 25 Lbm and Lh venom reservoirs. (B) SOD spectrophotometric assay using 200 ng of Trx-LbmSOD3 and protein extracts of 20 Lbm, Lby and Lh venom reservoirs (20R). (C) SOD zymography assay using extracts of 25 Lbm and Lby venom reservoirs with (KCN+) and without KCN (KCN-) (D) SOD spectrophotometric assay using 200 ng of

Trx-LbmSOD3 and extracts of 20 Lbm reservoirs (20R), with (KCN+) and without KCN (KCN-). Trx tag alone was used as a control.

FIGURE 7. Effect of *Leptopilina* recombinant SOD3 and venom on *Drosophila* PO activity. PO activity measured as absorbance at 490 nm for (A) *D. melanogaster* larval hemolymph in presence of buffer alone, Trx-LbmSOD3 (3 to 0.003 μ g), Trx-tag alone (3 μ g) or PO inhibitor (PTU). (B) *D. melanogaster* larval hemolymph in presence of buffer alone, 3 μ g of Trx-LbmSOD3, Trx-LhSOD3, BtSOD or Trx-tag alone, or PTU. (C) *D. melanogaster* larval hemolymph in presence of Trx-LhSOD3 (3 μ g), Trx-tag alone (3 μ g) or PTU, added either before or after PO activation. Mean values (\pm SE) are given for each category (three replicates).

	LbmSOD1	LbySOD1	LhSOD1	LbmSOD3	LbySOD3	LhSOD3
ORF (bp)	456	456	456	522	525	531
Protein (aa)	151	151	151	173	174	176
Signal peptide	No	No	No	Yes	Yes	Yes
MW (kDa)	15.5	15.5	15.5	16.4	16.4	16.6
N-glycosylation	2	2	2	3	3	3

Table 1. Main characteristics of Cu,ZnSOD sequences identified in *L. boulardi* ISm, *L. boulardi* ISy and *L. heterotoma* venom apparatus.

	LbmSOD3	LbySOD3	LhSOD3	LbmSOD1	LbySOD1	LhSOD1	HsSOD1
LbySOD3	98.9/98.9						
LhSOD3	73.3/88.1	73.9/88.6					
LbmSOD1	44.6/56.0	44.3/55.7	45.5/55.1				
LbySOD1	44.6/56.0	44.3/55.7	45.5/55.1	100/100			
LhSOD1	44.0/54.9	43.8/54.5	43.8/53.9	95.4/98.7	95.4/98.7		
HsSOD1	47.7/58.5	47.5/58.2	47.5/56.5	65.8/74.2	65.8/74.2	64.5/74.8	
HsSOD3	25.5/37.7	25.0/35.9	24.1/37.3	23.5/30.8	23.5/30.8	24.0/31.1	25.1/34.4

Table 2. Sequences identity (%)/similarity (%) between *Leptopilina* and *Homo sapiens* (SOD1, NP_000445 ; SOD3, P08294) Cu,ZnSODs (amino acid sequence alignment using the EMBOSS program Needle).

	LbmSOD3	LbySOD3	LhSOD3	LbmSOD1	LbySOD1	LhSOD1	HsSOD1
HsSOD1	0.449 (125)	0.449 (125)	0.448 (125)	0.420 (132)	0.420 (132)	0.440 (133)	
HsSOD3	0.839 (127)	0.839 (127)	0.851 (116)	0.881 (108)	0.881 (108)	1.090 (100)	0.615 (110)

Table 3. Structural alignments between *Leptopilina* and *Homo sapiens* (SOD1, 1PU0; SOD3, 2JLP) Cu,ZnSODs based on the root-mean-square deviations (RMSD, Angstrom) using Pymol. Values in brackets correspond to number of aligned α -carbon atoms.

Superoxide dismutases in parasitoids' venom

Organism	Protein	Peptides
Lbm	LbmSOD1	LACGVIGITKN
	LbmSOD3	GDISKGCISTGK HVGDLGNVIANK AFVVHEK MVPNNVQK
Lby	LbySOD1	GTVFFEESGDSVK LACGVIGITKN
	LbySOD3	HVGDLGNVIANK (2 bands) GNTSLSLETGDAGDR
Lh	LhSOD1	GNNNIIGR LACGVIGITK TLVVHGDPPDLGK GTVFFEEAGDSVK

Table 4. Cu,ZnSOD peptides identified by mass spectrometry in *L. boulardi* and *L. heterotoma* venom.

Figure 1

A

HsSOD1	1	-----MATKAVCVL
LbmSOD1	1	-----I.....
LbySOD1	1	-----I.....
LhSOD1	1	-----I.....
LbmSOD3	1	-----MMKIFPI-F--VLVAVSYGQELR..VKM
LbySOD3	1	-----MMKIFPIVF--VLVAVSYGQELR..VKM
LhSOD3	1	-----MKKVFSIVFIPLLIAASCGQEL...VK.
HsSOD3	1	<u>MLALLCSCLLLAAGASDAW</u> TGEDSAEPNSDSAEWIRDMYAKVTEIWQEVQRDDDG.LH.A.QV*
HsSOD1	10	KGD-----GPVQGIINFEQKESNGPVKVGSIKGLT----EGLHGF ^H VHEFGDNTAG ^C TSAGP
LbmSOD1	10	Q.E-----K.TVF..ESGDS---.T.EVT..K---K.....I.....N.....
LbySOD1	10	Q.E-----K.TVF..ESGDS---.T.EVT..K---K.....I.....N.....
LhSOD1	10	Q.E-----K.TVF..EAGDS---.T.EVT..K---K.....I.....N.....
LbmSOD3	26	VPN---NVQKNN.T.NLLIV.DGK...TIT...Y..S---P.S...K..ISK..I.T.K
LbySOD3	27	VPN---NVQKNN.T.NLLIV.DGK...IIT...Y..S---P.S...K..ISK..I.T.K
LhSOD3	29	VPN---NVAKLN.T.LLIS.SVK...TIT.T.Y.IP---P.....K..M.K..I.T.K
HsSOD3	66	QPSATLDAAQPR.T.VVL.R.LAPRAKLDAFFALE.FPTEPNSSRAI...Q...LSQ..E.T..*
HsSOD1	64	^H FNPLSRKHGGPKDEER ^H VG ^D LGNVTADKDGADVSIEDSVISLSGDHCCIIGRTL ^H VVEKADDLG
LbmSOD1	60	...SGMD...T.SV...E.SG...K.N.T.KQ.Q.K.NNN...GDP...
LbySOD1	60	...SGMD...T.SV...E.SG...K.N.T.KQ.Q.K.NNN...GDP...
LhSOD1	60	...SGVE...T.SV...I...E.GS...K.N.S.KQ.Q.K.NNN...GDP...
LbmSOD3	84	...EKVN..A.D.KV...I.N.E.E.VIN.T..I...PNNVL..AF...E...
LbySOD3	85	...EKVN..A.D.KV...I.N.E.E.VIN.T..I...PNNVL..AF...E...
LhSOD3	87	...ERVN..A.N.RV...LN.SE.WT.K.D.T.TM...PNS...AF...T...
HsSOD3	131	Y...AVP.-----PQ ^R P ^C .F..F..VR..SLWR ^R YRAGLAA..A.P.S.V..AV...AGE....*
HsSOD1	129	KGGNEESTKTGNAGS ^R LA ^C GVIGIAQ-----154
LbmSOD1	125	...H.L.KT....A.....TKN-----151
LbySOD1	125	...H.L.KT....A.....TKN-----151
LhSOD1	125	...H.L.KT....A.....TKN-----151
LbmSOD3	149	..NTSL.LE..D..D...IV..Q-----173
LbySOD3	150	..NTSL.LE..D..D...IV..Q-----174
LhSOD3	152	..NSTL.LE..D..D..I..IV..Q-----176
HsSOD3	189	R...QA.VEN...R...C.V.VCGPGLWERQAREHSE ^R RKKRRRESECKAA 240*

B





Figure 2

A



B






Figure 3

A



B



C




Figure 4







A**B****C****D**

Figure 5

A



B



C




Figure 6




Figure 7

