



**HAL**  
open science

## Effect of oral exposure to polycyclic aromatic hydrocarbons on goat's milk contamination

Nathalie Grova, Guido Rychen, Fabrice Monteau, Bruno Le Bizec, Cyril Feidt

### ► To cite this version:

Nathalie Grova, Guido Rychen, Fabrice Monteau, Bruno Le Bizec, Cyril Feidt. Effect of oral exposure to polycyclic aromatic hydrocarbons on goat's milk contamination. *Agronomy for Sustainable Development*, 2006, 26 (3), pp.195-199. 10.1051/agro:2006019 . hal-02654144

**HAL Id: hal-02654144**

**<https://hal.inrae.fr/hal-02654144>**

Submitted on 29 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

## Effect of oral exposure to polycyclic aromatic hydrocarbons on goat's milk contamination

Nathalie GROVA<sup>a\*</sup>, Guido RYCHEN<sup>a</sup>, Fabrice MONTEAU<sup>b</sup>, Bruno LE BIZEC<sup>b</sup>, Cyril FEIDT<sup>a</sup>

<sup>a</sup> URAPA, INRA-INPL-UHP, BP 172, 2 avenue de la Forêt de Haye, 54505 Vandœuvre-lès-Nancy Cedex, France

<sup>b</sup> LABERCA, ENVN, route de Gachet, 44307 Nantes Cedex 3, France

(Accepted 24 July 2006)

**Abstract** – The impact of chronic exposure to polycyclic aromatic hydrocarbons (PAHs) on milk contamination was evaluated by oral administration of a mixture of fluorene, phenanthrene, anthracene, fluoranthene, pyrene, chrysene, benzo(k)fluorene, benzo(a)pyrene and benzo(g,h,i)perylene at 0.02 mg/kg to lactating goats for 28 days. We analysed PAHs and their major metabolites in milk by gas chromatography-mass spectrometry. The results evidence several major points: (1) benzo(k)fluorene, benzo(a)pyrene and benzo(g,h,i)perylene were not detected in the milk; (2) unexpectedly, the concentration of fluorene, phenanthrene, anthracene, fluoranthene, pyrene and chrysene did not change with time; (3) monohydroxylated PAH metabolites (-OH), namely 2-OH-fluorene, 3-OH-phenanthrene and 1-OH-pyrene were detected shortly after administration. The concentrations of 2-OH-fluorene and 3-OH-phenanthrene reached, respectively, maxima of 0.41 and 0.22 ng/mL during the first exposure week, whereas the concentration of 1-OH pyrene increased to reach a maximum of 0.97 ng/mL on day 14, then slightly decreased during the last two exposure weeks. Those findings suggest a lack of activation of a metabolism that could lead to an excretion of PAHs into milk under native forms. However, a slight increase in concentration could induce the metabolism, which should lead to an increase in the excretion of metabolites into the milk. In spite of the absence of a significant transfer of parent PAHs to milk, the appearance of metabolites in milk raises questions of their impact on human health.

### Chronic exposure / PAHs / metabolites / milk contamination

#### 1. INTRODUCTION

Polycyclic aromatic hydrocarbons (PAHs) are potentially mutagenic compounds widely occurring in natural media such as soils, atmosphere, sediments and plants (Wild et al., 1992; Bryselbout et al., 2000; Juhasz et Naidu, 2000, Amellal et al., 2006). The potential carcinogenicity of PAHs is well known, and the ingestion of contaminated food is considered to be the main source of human exposure (IARC, 1983). Consumers might be exposed to PAHs by eating grilled or charred meats, contaminated cereals, flour, bread and vegetables. New rules established in 2005 set the maximum levels for benzo(a)pyrene intakes at 5000 ng/kg in fish and meat products and at 1000 ng/kg in children's foods. Currently, milk and dairy products represent more than 406 kg/eq/year and there is no legislation regulating PAHs and benzo(a)pyrene in particular. Several studies achieved in environmental conditions show that PAHs can be excreted in the milk of ruminants (Husain et al., 1997; Grova et al., 2000, 2002a; Feidt et al., 2005).

Studies to investigate the transfer of these pollutants in lactating ruminants have previously been carried out according to various modalities. West and Horton (1976) evaluated the transfer factor of <sup>14</sup>C benzo(a)pyrene at 0.01% in milk after

chronic oral administration (1 mg/kg per day) in sheep. A similar study was performed in lactating goats with a single oral administration of <sup>14</sup>C-phenanthrene, <sup>14</sup>C-pyrene and <sup>14</sup>C benzo(a)pyrene at levels of 0.005 mg/kg. It showed transfer factors to the milk of 1.5%, 1.9% and 0.2%, respectively (Grova et al., 2002b). A recent study focused on the transfer and metabolism of phenanthrene in lactating goats allowed the detection and identification of the native compound and its metabolites in various compartments such as plasma, excretion products, and particularly in milk (Grova et al., 2005). In terms of food safety, knowledge of PAH metabolites' levels in food is of some interest, as they appear to be more reactive than parent compounds in terms of mutagenesis and carcinogenesis (Denison and Whitlock, 1995; Uno et al., 2001) or endocrine disruption (Fertuck et al., 2001; van de Wiele et al., 2005; van Lipzig et al., 2005). Unfortunately, these results do not correlate to the potential chronic exposure of animals by ingestion of contaminated ground and fodders (Crépineau et al., 2003). This study performed on transfer and metabolism of PAHs in lactating goats will contribute to the evaluation of the effect of chronic exposure to PAH mixture on milk contamination and give new insight into the form (parent or metabolised molecule) transferred to the milk.

\* Corresponding author: nathalie.grova@lns.etat.lu

Current address: Laboratoire National de Santé, Institute of Immunology, PO Box 1102, 1011 Luxembourg.

## 2. MATERIALS AND METHODS

### 2.1. Animals

Three Alpine goats (50 ± 5 kg; second and third lactations, second month post-partum) from the herd of the "Domaine Expérimental de la Bouzule" (Champenoux, France) were used for this experiment as a model of the lactating ruminant. They received a seven-day adaptation period within individual boxes. Ambient temperature was close to 22 °C and natural light conditions prevailed. The animals were mechanically milked twice a day, and the average milk production during the seven days before the experiment was 2.5 ± 0.8 L/day. The goats were fed with meadow hay, water and mineral salt ad libitum and received a concentrate ration twice a day (at 8 am and 5 pm) consisting of dehydrated fodder beet (1100 g), crushing maize (400 g), soya meal (200 g) and dehydrated alfalfa (800 g). The diet composition was established to meet the maintenance and milk production (about 3 L/day) needs of the lactating ruminants (INRA, 1988). Measurements of the production (mL), protein level (g/L) and fat level (g/L) were performed every week in order to verify the quality of the milk.

### 2.2. Experimental design

The animal protocol was in accordance with the general guidelines of the Council of European Communities (1986, No. 86/609/CEE) and the French Animal Care Guidelines. A control sample of milk was collected from each goat three days before starting the daily administration of PAHs. After morning milking, each animal received 2 mL of vegetal oil (ISIO 4, Lesieur, Neuilly-sur-Seine, France) containing 1 mg of each PAH (0.02 mg/kg of body weight) for 28 consecutive days. PAHs were directly administered into the mouth of the animal with a syringe. The syringe was then flushed twice with 2 mL of vegetal oil and the contaminated oil was given to the animals. The total of 6 mL oil ingested by each animal is not expected to affect rumen activity (Murphy and Morgan, 1983). The PAH mixture was composed of 1 mg of the following compounds: Fluorene (Fluo), Phenanthrene (Phe), Anthracene (Ant), Fluoranthene (Fluot), Pyrene (Pyr), Chrysene (Chrys), Benzo(k)Fluoranthene (B(k)F), Benzo(a)pyrene (B(a)P) and Benzo(g,h,i)perylene (B(g,h,i)Per). Milk samples were collected on days 7, 14, 21 and 28 of the trial. Both evening and the following morning milking were pooled, weighed and stored at -20 °C. The levels of PAHs in water and ration samples were quantified by gas chromatography coupled to mass spectrometry in the single ion monitoring mode. The target PAHs in the diet (Fluo 1.8 ng/g DM, Phe 9.3 ng/g DM, Pyr 0.9 ng/g DM and B[a]P not detected) and water (Fluo and B[a]P not detected, Phe 24 ng/L, Pyr 16 ng/L) were analysed and the concentrations revealed a total daily supply of less than 1% of the intake.

### 2.3. Milk analyses

The analytical procedure for the extraction and purification of PAHs and their metabolites has been previously described (Grova et al., 2005; Lutz et al., 2006). Milk concentrations of the following PAHs: Fluo, Phe, Ant, Fluot, Pyr, Chrys, B(k)F, B(a)P and B(g,h,i)Per (purchased from Sigma Aldrich, St

Quentin Fallavier, France) and major monohydroxylated metabolites: 2-hydroxyfluorene (2-OH Fluo), 3-hydroxyphenanthrene (3-OH Phe) and 1-hydroxypyrene (1-OH Pyr) from Chiron (Trondheim, Norway), were determined as follows. The 3-OH B(a)P is considered as the most abundant metabolite of B(a)P, which is the reference of PAH toxicity. The 1-OH Pyr was selected as the usual indicator of environmental contamination (Jacob and Seidel, 2002). The 3-OH Phe is known to be the major metabolite excreted in goat's milk (Grova et al., 2005). Moreover, the choice of the 2-OH Fluo enables us to consider a decreasing number of benzene cycles.

Prior to extraction, 10 mL of milk fortified with several internal standards (d10-Phe, d10-Pyr and d12-perylene, Interchim, Montluçon, France) was adjusted to pH 5.7 with 100 µL of glacial acetic. Deconjugation was performed with 3750 units of purified *Helix pomatia* juice (Biosepra, Villeneuve-la-Garenne, France) and milk samples were incubated for 16 hours at 37 °C to induce hydrolysis of glucuronide and sulphate conjugates of hydroxymetabolites. Samples were then shaken with 20 mL cyclohexane/ethyl acetate (50:50, v/v; SDS, Peypin, France) for 30 minutes. After centrifugation (15 min at 1000 g) the supernatant was evaporated. Residues were dissolved in 3 mL cyclohexane and applied onto an Envi-Chrom P SPE (Styrene-divinylbenzene copolymer resin, Envi Chrom P: 0.5 g) column previously conditioned with water, methanol and cyclohexane (SDS, Peypin, France). After rinsing with 3 mL cyclohexane, PAHs were eluted with 12 mL cyclohexane/ethyl acetate (50:50, v/v). After evaporation to dryness, 2 mL cyclohexane and 2 mL methanol/water (80:20, v/v) were added, mixed for 30 seconds, and the cyclohexane phase was extracted after centrifugation (1000 g, 5 min). The methanol/water phase was washed again with 2 mL cyclohexane, centrifuged (1000 g, 5 min) and the cyclohexane layer was added to the first cyclohexane phase. At this point, the methanol layer containing hydroxylated metabolites was set aside for later analysis. The supernatant containing the PAHs in cyclohexane was evaporated; saponification was achieved with 5 mL 10% KOH for 80 minutes at 90 °C to prevent any fatty residues from being present in the final product. 3 mL water and 5 mL cyclohexane were then added and the mixture was shaken prior to centrifugation (1000 g, 5 min). The supernatant was harvested, an external standard (d12-chrysene) was added, and the cyclohexane was evaporated. The residue was dissolved in 20 µL toluene. The previously separated methanol fraction, as described above, was evaporated and extracted with 4 mL water/ethyl acetate (50:50; v/v). After vortexing and centrifugation (1000 g, 5 min), the supernatant was evaporated, supplemented with an external standard (1-OH chrysene, Interchim, Montluçon, France), and hydroxymetabolites were derivatised with 20 µL N-methyl-N-(trimethylsilyl) trifluoroacetamide (MSTFA, Fluka, Buchs, Switzerland).

### 2.4. Chromatography conditions

The quadrupole MS (HP-5973) system used was coupled to a HP-6890 gas chromatograph, both from Hewlett-Packard (Palo Alto, CA, USA). The split/splitless injector was maintained at 250 °C, time of splitless mode was set at 1.5 minutes, and the injected volume was 2 µL. The column used for the separation of phenanthrene and its metabolites was an OV-1



**Table III.** Concentration of monohydroxylated polycyclic aromatic hydrocarbons (PAHs) in goat's milk during the exposure period. 2-OH Fluo: 2-OH Fluorene, 3-OH Phe: 3-OH Phenanthrene, 1-OH Pyr: 1-OH Pyrene, 3-OH BaP: 3-OH Benzo(a)Pyrene, ND: not detected, SD: standard deviation, M: mean, RMSE: root mean square error. \*  $P < 0.05$ . \*\*  $P < 0.01$  difference statistically significant versus control group (Tukey's t-test).

Concentration ng/mL milk (n = 3)	Day 0		Day 7		Day 14		Day 21		Day 28		Time Effect	RMSE
	M	SD	M	SD	M	SD	M	SD	M	SD		
2-OH Fluo	0.00	0.00	0.41	0.25	0.51	0.34	0.67	0.34	0.47	0.21	*	0.057
3-OH Phe	0.00	0.00	0.22	0.09	0.37	0.17	0.30	0.12	0.13	0.03	**	0.009
1-OH Pyr	0.00	0.00	0.86	0.53	0.97	0.27	0.53	0.41	0.32	0.15	*	0.048
3-OH BaP	ND		ND		ND		ND		ND		–	–

However, the statistical analysis ( $P > 0.05$ ) did not reveal an increase in these molecules during the 4 weeks of exposure (Tab. II). This result is in accordance with Lutz et al. (2006), who indicated that a chronic administration of contaminated soil by cow's ruminal canula did not increase the level of PAHs in milk. Lactating ruminants exposed daily to PAHs via the ingestion of contaminated matrices (at an environmental level of 0.02 mg/kg/day) did not present a significant increase in PAH concentration in milk.

Of the 4 monohydroxy-metabolites investigated, only 3 of them were detected: the 2-OH Fluo, the 3-OH Phe and the 1-OH Pyr (Tab. III). Three different profiles were observed: firstly, the concentrations of 2-OH Fluo and 3-OH Phe seemed to plateau out at day 7 ( $0.41 \pm 0.25$  and  $0.22 \pm 0.09$  ng/mL, respectively). The highest daily excreted quantities were 1582 ng ( $\pm 330$ ) and 930 ng ( $\pm 360$ ), respectively, for 2-OH Fluo and 3-OH Phe. Secondly, the variance analysis revealed an increase in 1-OH Pyr during the first 2 weeks of exposure ( $P < 0.05$ ,  $0.97 \pm 0.27$  ng/mL on day 14, corresponding to  $2382 \pm 141$  ng excreted), followed by a slight decrease over the two last weeks. Finally, this study did not allow the detection of the presence of 3-OH B(a)P in milk before or after the treatment (Tab. III).

The difference between the three compounds may be explained by the reduced number of metabolites for Pyr compared with the numerous metabolites of Phe and Fluo. Moreover, the absence of 3-OH B(a)P could be partially explained by the very poor absorption of B(a)P and its strong metabolism at the enterocyte levels (Cavret and Feidt, 2005). The same concentration profiles of metabolites in milk were also described in cow's milk after chronic exposure to contaminated soil (Lutz et al., 2006).

Contrary to the native compounds, target monohydroxylated metabolites were not detected in control samples. However, the 2-OH Fluo, the 3-OH Phe and the 1-OH Pyr were present in the milk at day seven, and their levels increased significantly compared with controls ( $P < 0.05$ ) during the rest of the exposure period (Tab. III). The absence of metabolites in control milk may suggest a lack of activation of the metabolism at an environmental level, which could lead to an excretion of native forms of PAHs into milk. According to Grova et al. (2005), this experiment highlighted a metabolism induction in goats chronically exposed to a slight increase in PAH concentration (0.02 mg/kg/day), revealed by the presence of metabolites in milk. This metabolism induction contributes to the increase in excretion of metabolites into urine and milk (e.g., 40% of the

radioactivity associated with Pyr was excreted via the urine, Grova et al., 2002b) and partially explains the significant increase in metabolite levels observed during the experiment.

The impact of metabolites in milk on human health is not currently understood; little information is available. Analytical analysis showed that 100% of metabolites present in milk are under conjugated forms (Grova et al., 2005). These sulphate and glucuronide groups increase the molecular weight and the solubility of PAH metabolites. This increase in solubility could change their availability and their absorption in the intestinal tract, where they could again be metabolised under reactive forms.

#### 4. CONCLUSION

The daily oral exposure of lactating goats to a mixture of 9 PAHs for 28 days resulted in a significant increase in monohydroxylated metabolites in milk. The constant level of native PAH forms and the significant appearance of metabolites confirm that in terms of risk assessment, exposure of lactating ruminants to PAHs remains problematic. Considering that their toxicity and bioavailability are not known, it is surprising that the proven documented transfer of metabolites is largely ignored. The authors believe that it would be useful to characterise these two parameters in order to evaluate the potential risk of transfer within the food chain.

**Acknowledgments:** We would like to thank the A.D.E.M.E (Agence De l'Environnement et de la Maîtrise de l'Énergie), the « Région Lorraine », and the M.E.N.R.T. (Ministère de l'Éducation Nationale de la Recherche et de la Technologie) for their financial support.

#### REFERENCES

- Amellal S., Boivin A., Perrin-Ganier C., Schiavon M. (2006) High sorption of phenanthrene in agricultural soils, *Agron. Sustain. Dev.* 26, 99–106. DOI: 10.1051/agro:2006002.
- Bryselbout C., Henner P., Carsignol J., Lichtfouse E. (2000) Polycyclic aromatic hydrocarbons in highway plants and soils, Evidence for a local distillation effect, *Analisis* 28, 32–35.
- Buesen R., Mock M., Nau H., Seidel A., Jacob J., Lampen A. (2003) Human intestinal Caco-2 cells display active transport of benzo(a)pyrene metabolites, *Chem.-Biol. Inter.* 142, 201–221.

- Cavret S., Feidt C. (2005) Intestinal metabolism of PAH: in vitro demonstration and study of its impact on PAH transfer through the intestinal epithelium, *Environ. Res.* 98, 22–32.
- Cavret S., Feidt C., Laurent F. (2005) Differential transfer of organic micropollutants through intestinal barrier using Caco-2 cell line, *J. Agr. Food Chem.* 53, 2773–2777.
- Crépineau C., Rychen G., Feidt C., Leroux Y., Lichtfouse E., Laurent F. (2003) Contamination of pastures by polycyclic aromatic hydrocarbons (PAHs) in the vicinity of a highway, *J. Agr. Food Chem.* 51, 4841–4845.
- Denison M.S., Whitlock J.P. Jr. (1995) Xenobiotic inducible transcription of cytochrome P450 genes, *J. Biol. Chem.* 270, 18175–18178.
- Feidt C., Cavret S., Grova N., Laurent C., Rychen G. (2005) Evaluation of the risk of PAHs and Dioxins transfer to humans via the dairy ruminant, in: Lichtfouse E., Schwarzbauer J., Robert D. (Eds.), *Environmental chemistry, Green chemistry and pollutants in ecosystems*, Springer, Berlin, pp. 419–430.
- Fertuck K.C., Kumar S., Sikka H.C., Matthews J.B., Zacharewski T.R. (2001) Interaction of PAH-related compounds with alpha and beta isoforms of the estrogen receptor, *Toxicol. Lett.* 121, 167–177.
- Grova N., Laurent C., Feidt C., Rychen G., Laurent F. (2000) Gas chromatography-mass spectrometry of polycyclic aromatic hydrocarbons in grass and milk from urban and rural farms, *Eur. J. Mass Spectrom.* 6, 457–460.
- Grova N., Feidt C., Crépineau C., Laurent C., Lafargue P.E., Hachimi A., Rychen G. (2002a) Influence of potential PAHs emission sources on milk contamination, *J. Food Agr. Chem.* 50, 4640–4642.
- Grova N., Feidt C., Laurent C., Rychen G. (2002b) [<sup>14</sup>C] Milk, urine and faeces excretion kinetics in lactating goats after an oral administration of [<sup>14</sup>C]polycyclic aromatic hydrocarbons, *Int. Dairy J.* 12, 1025–1031.
- Grova N., Monteau F., Le Bizec B., Feidt C., Andre F., Rychen G. (2005) Determination of Phenanthrene and Hydroxyphenanthrenes in Various Biological Matrices at Trace Levels using Gas Chromatography-Mass Spectrometry, *J. Anal. Toxicol.* 29, 175–181.
- Husain A., Naeemi E., Dahsti B., Al-Omirah H., Al-Zenki S. (1997) Polycyclic aromatic hydrocarbons in food products originating from locally reared animals in Kuwait, *Food Addit. Contam.* 14, 295–299.
- IARC (International Agency for Research on Cancer) (1983) IARC monographs on the evaluation of carcinogenic risks of chemicals to humans. Polynuclear aromatic compounds. Part 1: Chemical, environmental and experimental data 32, International Agency for Research on Cancer, Lyon, France.
- INRA (1988) Tables de l'alimentation des bovins, ovins et caprins, Services des publications de l'INRA, Versailles.
- Jacob J., Seidel A. (2002) Biomonitoring of polycyclic aromatic hydrocarbons in human urine, *J. Chromatogr. B* 778, 31–47.
- Juhász A.L., Naidu R. (2000) Bioremediation of high molecular weight polycyclic aromatic hydrocarbons: a review of the microbial degradation of benzo(a)pyrene, *Int. Biodegr. Biodegr.* 45, 57–88.
- Littell C.R., Milliken G.A., Stroup W.W., Wolfinger R.D. (1996) SAS system for Mixed Models, Chap. 3: Analysis of Repeated Measures Data, Cary, NC, SAS Institute Inc., 633 p.
- Lutz S., Feidt C., Monteau F., Rychen G., Le Bizec B., Jurjanz S. (2006) Effect of exposure to soil-bound polycyclic aromatic hydrocarbons on milk contaminations of parent compounds and their monohydroxylated metabolites, *J. Food Agr. Chem.* 54, 263–268.
- Murphy J.J., Morgan D.J. (1983) Effect of inclusion of protected and unprotected tallow in the supplement on the performance of lactating dairy cows, *Anim. Prod.* 37, 203–210.
- Uno S., Dalton T.P., Shertzer H.G., Gener M.B., Warshawsky D., Talaska G., Nebert D.W. (2001) Benzo[a]pyrene induced toxicity: paradoxical protection in Cyp1a1(–/–) knockout mice having increased hepatic BaP-DNA adduct levels, *Biochem. Biophys. Res. Co.* 289, 1049–1056.
- van de Wiele T., Vanhaecke L., Bockaert C., Peru K., Headley J., Verstraete W., Siciliano S. (2005) Human colon microbia transform polycyclic aromatic hydrocarbons to estrogenic metabolites, *Environ. Health Persp.* 113, 6–10.
- van Lipzig M.M.H., Vermeulen N.P.E., Gusinu R., Legler J., Frank H., Seidel A., Meerman J.H.N. (2005) Formation of estrogenic metabolites of benzo[a]pyrene and chrysene by cytochrome P450 activity and their combined and supra-maximal estrogenic activity, *Environ. Toxicol. Phar.* 19, 41–55.
- West C.E., Horton B.J. (1976) Transfer of polycyclic aromatic hydrocarbons from diet to milk in rats, rabbits and sheep, *Life Sci.* 19, 543–551.
- Wild S.R., Jones K.C., Johnston A.E. (1992) The polycyclic aromatic hydrocarbon (PAH) content of herbage from a long-term grassland experiment, *Atmos. Environ.* 26A, 1299–1307.