



A stochastic simulation study on using different models for prediction of breeding values while changing the breeding goal

J. Lassen, M.K. Sorensen, P. Madsen, Vincent Ducrocq

► To cite this version:

J. Lassen, M.K. Sorensen, P. Madsen, Vincent Ducrocq. A stochastic simulation study on using different models for prediction of breeding values while changing the breeding goal. *Animal*, 2007, 1, pp.631-636. hal-02655015

HAL Id: hal-02655015

<https://hal.inrae.fr/hal-02655015>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A stochastic simulation study on using different models for prediction of breeding values while changing the breeding goal

J. Lassen^{1,2†}, M.K. Sørensen¹, P. Madsen¹ and V. Ducrocq³

¹Department of Genetics and Biotechnology, Danish Institute of Agricultural Sciences, PO Box 50 DK-8830 Tjele, Denmark; ²Department of Large Animal Sciences, The Royal Veterinary and Agricultural University, Ridebanevej 12, DK-1870 Frederiksberg C Denmark; ³Station de Génétique Quantitative et Appliquée, Institut National de la Recherche Agronomique, 78352 Jouy-en-Josas, France

(Received 13 October 2006; Accepted 21 February 2007)

In a stochastic simulation study the effect of simultaneously changing the model for prediction of breeding values and changing the breeding goal was studied. A population of 100 000 cows with registrations on seven traits was simulated in two steps. In the first step of 15 years the population was selected for production and mastitis occurrence using a univariate model for prediction of breeding values for production and a trivariate model using information on mastitis treatments, udder depth and somatic cell score for prediction of breeding values for mastitis occurrence. In the second step six different scenarios were set up and simulated for 15 years combining two different breeding goals and three different models for prediction of breeding values in 20 replicates. Breeding goal 1 had relative economic value per genetic standard deviation on production (19.4) and mastitis occurrence (–50) whereas breeding goal 2 had a economic value on production (19.4), udder depth (4.2), mastitis occurrence (–50), non return rate (13.0) and days open (–16.75). Model 1 was a model similar to the one used in the first 15 years. Model 2 was an approximate multitrait model where solutions for fixed effects from a model corresponding to model 1 were subtracted from the phenotypes and a multitrait model with an overall mean, a year effect, an additive genetic and a residual effect were applied. Model 3 was a full multitrait model. Average genetic trends for total merit and each individual trait over 20 replicates were compared for each scenario. With the number of replicates the genetic responses using model 2 and 3 were not significant different. With a broad breeding goal using, model 2 or model 3 gave a significantly higher response in total merit than using model 1. Using a narrow breeding goal there was no significant difference between models used for prediction of breeding values. Results showed that with a breeding goal with a lot of emphasis on low heritable traits with a high economic value using a multitrait methodology for prediction of breeding values will redistribute the genetic progress in the total merit index. More gain will come from the low heritable traits in the breeding goal and less from traits with higher heritability. With a broad breeding goal and exploiting the available information in the data the inbreeding coefficient increased though not significantly.

Keywords: breeding goal, dairy cattle, multitrait model, selection, stochastic simulation

Introduction

Over the last years focus in breeding dairy cattle has moved from mainly production to a situation where cost-reducing traits such as udder health, reproduction and calving ease receive increasing emphasis. This has been done both to increase the income of the farmer and also to improve animal welfare and sustainability in production (Olesen *et al.*, 2000).

Common for most of the cost-reducing traits is a low heritability and unfavourable genetic correlation to production traits and to some conformation traits. The low

heritabilities lead to lower accuracies for the predicted breeding value (EBV) for these traits. One possible way to increase the accuracy of EBVs is to perform multitrait evaluations using direct and indirect information to take advantage of the genetic relationship between the traits (Van der Werf *et al.*, 1992; Ducrocq, 1994). Complete multitrait evaluation in national breeding programmes with dairy cattle and properly also with other species are in most cases not computationally feasible due to the amount of data and number of traits. Therefore, approximations have been proposed (Ducrocq *et al.*, 2001)

Multitrait evaluations using direct and indirect information have been shown to increase accuracy on longevity

[†] E-mail: jan.lassen@agrsci.dk

by more than 15% for newly proven sires (Ducrocq *et al.*, 2001). Another benefit of multitrait evaluation is that it accounts for the fact that the main selection has been on production, and thereby selection bias is taken into account.

In a stochastic simulation study of a nucleus breeding scheme Sørensen *et al.* (1999) showed that a multitrait breeding goal gave a higher genetic gain than only selecting for milk production. However the selection for the multitrait breeding goal was based on breeding values from primarily single-trait models. Genetic gain for production was smaller, selecting for total merit, but due to a considerable increase in genetic gain for cost-reducing traits total economic gain increased. This has also been shown in deterministic simulation studies (Christensen, 1998; Willam *et al.*, 2002).

Ducrocq *et al.* (2001) compared the selection differential for the traits in the breeding goal for the 10% best bulls when using an approximate multitrait model for breeding value estimation instead of ranking according to single-trait EBVs. This showed that changing the model modified the genetic profile of the best animals, since genetic merit increased for functional longevity, somatic cell score (SCS) and female fertility and decreased for milk yield, overall type and overall udder. However, at the same time as this approximate multitrait model, a new breeding goal was introduced, so it is hard to distinguish whether it was the effect of the model or the effect of the breeding goal which changed the new selection differentials. Comparing the use of single-trait models with an approximate multitrait model similar to the one used by Ducrocq *et al.* (2001) in laying hens, Besbes *et al.* (2002) found an improvement for all traits in the breeding goal.

The aim of this study is to quantify the effect of at the same time changing the methods used for prediction of breeding values and the breeding goal into a more sustainable breeding programme including functional traits. This is compared with the methods and breeding goal used today. The hypothesis is that the benefit from using multitrait evaluation is bigger when selection is on a broader

breeding goal. This is tested in a stochastic simulation study of a dairy cattle population.

Material and methods

Population structure

A population of approximately 100 000 cows was simulated for a 30-year period. For all animals true breeding values for seven traits were simulated, and for cows phenotypes for these traits were simulated (Table 1). Genetic parameters presented in this table come from an intensive literature study. All traits were simulated as normally distributed traits with a phenotypic variance of 1 and for all traits there was exactly one observation per cow except for dairy form and udder depth where only 40 females from each progeny group were measured. These 40 cows were randomly chosen. None of the traits was measured on the bulls. Mastitis occurrence and non-return rate were converged to binary traits with thresholds of -1 and 0 respectively, indicating a frequency of 0.18 and 0.5 in the starting population. All traits were realised and included in the breeding value estimation when cows were 35 months old. The average herd size was 100 cows. Progeny group size was 100 and the usage of semen from young sires was 35%. Selection of proven bulls was based on EBVs and from each proven bull 20 000 doses of semen could be used per year. Cows older than 6 years and bulls older than 8 years were culled and furthermore 5% of all animals were culled every year at random for other reasons. Cow replacement was done according to the EBVs, so cows with the highest EBV for total merit were kept to produce the next generation of offspring following a truncation selection strategy. This was done yearly after each round of breeding value estimation, and in each round of selection the same selection intensity was used due to restrictions on number of doses of semen per bull and population size of females. Sire-daughter and half- and full-sib matings were not allowed. Two different breeding goals were constructed (BG1 and BG2) using the economic

Table 1 Name and type of traits, economic values in Danish Kroner per phenotypic standard unit in the two breeding goals (BG1 and BG2), heritability (diagonal), genetic (below diagonal) and residual correlations (above diagonal) used in the simulation to generate records

Trait	Production	Udder depth	Mastitis occurrence	Non-return Rate	Dairy form	Days open	Somatic cell score
Type	Linear	Linear	Binary	Binary	Linear	Linear	Linear
Weight in BG1 (Narrow)	19.4	0	−50	0	0	0	0
Weight in BG2 (Broad)	19.4	4.2	−50	13	0	−16.75	0
Production	0.30	−0.20	0	−0.10	0.25	0.20	−0.15
Udder depth	−0.35	0.30	−0.05	0	0	−0.10	−0.05
Mastitis	0.35	−0.60	0.04	0	0	0	0.20
Non-return rate	−0.35	0.30	−0.10	0.03	0	0.05	0
Dairy form	0.45	−0.10	0.25	0	0.25	0.10	0
Days open	0.55	−0.10	0.05	−0.10	0.45	0.04	0
Somatic cell score	0.15	−0.30	0.75	−0.20	0.25	−0.25	0.10

values in Danish Kroner per genetic standard deviation corresponding to the values used in the Holstein population in Denmark in 2005 (Danish Agricultural Advisory Centre, 2006). BG1 were constructed to be narrow and therefore only had economic value on production and mastitis occurrence, whereas BG2 was constructed to mimic a broader breeding goal having economic values on production, udder depth, mastitis occurrence, non-return rate and days open (Table 1).

The seven traits were chosen to represent the traits that have been in focus over the last years and likely to be in focus in the near future in dairy cattle breeding. The most important trait is production. Two conformation traits were chosen – dairy form and udder depth. These two traits have relatively high genetic correlations to the other traits and have received some attention in the breeding goal so far. Mastitis occurrence and SCS were chosen to represent health traits and days open and non-return rate at day 56 were chosen to represent reproduction traits. Longevity could also be included when using this approximate method (Besbes *et al.*, 2002; Tarrés *et al.*, 2006) but was not considered here.

Simulation

The simulation was conducted using a modified version of the DAIRYSIM program. (Sørensen *et al.*, 1999). This program simulates all animals in the population.

The simulation was in two steps. The first step was to simulate a population comparable with a real population in which selection has been going on for several years. This first step covered 15 years, and selection was on BG1. EBVs were obtained using single trait animal model (AM) Best linear unbiased prediction (BLUP) for production and by a trivariate AM-BLUP for mastitis treatment with information from mastitis, SCS and udder depth. After this initial step, six different scenarios were simulated in order to quantify the effects of changing breeding goal and/or changing method for prediction of breeding values. The selection in each scenario was on BLUP EBVs obtained from animal models. In scenarios N1, N2 and N3 (N for 'Narrow'), BG1 was used as selection criterion, while in scenarios B1, B2 and B3 (B for 'Broad'), BG2 was used as selection criterion (Table 1). In scenarios N1 and B1, a multivariate model was used for the evaluation of mastitis, udder depth and SCS, and a univariate model was used for the other four traits. In scenario N2 and B2, an approximate multitrait model was used and in scenario N3 and B3 a complete multivariate model was applied. All six scenarios were started with estimates of (co)-variance components coming from analysis using the same model as later used for prediction of breeding values. These parameters were later used in the breeding value estimation whereas the ones used to simulate the data stayed the same. For each scenario 20 replicates were simulated.

Since genetic superiority depends on the reliability of the index of the individual traits, genetic progress is a way to describe the effect of changing the method for breeding

value estimation (Falconer and Mackay, 1996). Therefore, genetic progress in total merit was used as the main criterion for comparison of the six scenarios.

Model for prediction of breeding values

The following linear lactation model was used:

$$y_i = hys_i + a_i + e_i \quad (1)$$

where y_i is the observations for trait i , hys_i is the herd-year-season effects on trait i , a_i is the genetic effects for trait i and e_i is the residuals for trait i . Traditional effects such as parity and age at calving are assumed not to be statistically significant in the simulation. The models for prediction of EBVs in scenario N1 and B1 (model 1) were single trait AM for all traits, except for mastitis occurrence where a trivariate AM model for mastitis treatments, udder depth and SCS. This corresponds to the way the breeding values are calculated today in Denmark (Danish Agricultural Advisory Centre, 2006). In the approximate multitrait (MT) model scenarios (N2 and B2), a two-step analysis was performed. Data were first analysed based on models corresponding to the models used in scenario N1 and B1. Then, fixed effects estimates for (hys) were used to pre-adjust the data and on these pre-adjusted records, a full multitrait model (model 2) only containing an overall mean, a year effect, a genetic and a residual term was applied (Ducrocq *et al.*, 2003):

$$\text{model for the first step: } y_{i,m} = hys_i + a_{i,m} + e_{i,m} \quad (2a)$$

$$\begin{aligned} \text{model for the second step: } \hat{y}_{i,m} &= y_{i,m} - \hat{h}ys_i \\ &= \mu_i + \text{year} + \hat{a}_{i,m} + \hat{e}_{i,m} \end{aligned} \quad (2b)$$

where m is individual animals. The year effect was included in order to account for potential selection biases in the models used to generate the preadjusted data (Ducrocq *et al.*, 2003; Lassen *et al.*, 2007). The multivariate model was a complete multitrait model based on model 1.

(Co)-variance components for the prediction models were obtained by averaging estimates from three subsets of data, each including records from around 10 000 cows originating from 20 randomly chosen herds from the simulated data. Estimates were obtained using the AI-REML algorithm implemented in the DMU package (Madsen and Jensen, 2000).

The different scenarios were compared by differences in the achieved genetic trends computed as regressions of true genetic values on year over the last 13 years and for the different traits for the different scenarios as averages over replicates. From that point on, all females with realised phenotypes in the population were offspring from animals selected with respect to the changes in breeding goal or model for evaluation. For total merit, the regressions of both true and predicted genetic values on year were calculated using the economic values from BG2.

Results

Genetic gain

Scenario B3 led to the highest progress in true genetic total merit (Table 2). Scenario B3 was not significantly better than scenario B2 – at least with the number of replicates used to compute observed standard deviations, but these two scenarios were both significantly better than scenario B1. Similar patterns were observed for scenarios N3, N2 and N1, respectively. Using a multitrait or an approximate multitrait setting for prediction of breeding values led to a higher genetic total merit, especially when selection was done on BG2. The relative increase in true genetic trend from scenario B1 to B3 was 15.0% and from scenario B1 to B2, the increase was 11.8%. The increase between scenario N1 and N3 was 14.6% and between N1 and N2 it was 8.8%. In general the predicted breeding values for total merit were lower than the true breeding values.

Scenario BG1 gave a higher genetic merit for production than scenario BG2. For all the other traits, BG2 gave a more favourable genetic merit than BG1 (Tables 3 and 4). For non-return rate and days open, there was very little progress if any from changing to a multitrait setting. In scenarios N1, N2 and N3 where there was no selection for non-return rate and days open, reproduction traits declined substantially.

In general, using a method for prediction of breeding values where more of the available information is taken into account led to a more desirable genetic progress for the individual traits whatever the breeding goal. These differences were not significant, but in total they led to a higher total genetic merit.

In Table 5, the total merit is calculated using the economic values from breeding goals 1 and 2, respectively. Also the contribution of the different traits to the total merit is shown. Calculating the total merit using the weights from breeding goal 1 gives the highest merit as well as the lowest variation between scenarios. The results using weights from breeding goal 1 and breeding goal 2 cannot be compared since they are calculated using different economic expressions. In all scenarios, there is an improvement of production and mastitis and an unfavourable trend in the group of other traits containing the reproduction traits. This pattern is less pronounced when a multitrait model is

Table 2 Mean regression coefficients of 20 replicates in relative economic units of true and predicted genetic trends on year for total merit in simulation using the economic values from breeding goal 2 with standard errors

Breeding goal	Breeding value	Model 1	Model 2	Model 3
BG1	True	1.716 ± 0.060	1.867 ± 0.051	1.968 ± 0.060
	Predicted	†	1.753 ± 0.232	1.879 ± 0.212
BG2	True	2.672 ± 0.060	2.989 ± 0.044	3.073 ± 0.069
	Predicted	2.346 ± 0.212	2.989 ± 0.188	2.996 ± 0.208

† Estimated breeding values were not calculated for non-return rate, days open and dairy form.

Table 3 Mean (with s.e.) of true average genetic trends from scenarios using breeding goal 1 for the seven traits on year in simulation over 20 replicates (the desired direction of selection is indicated)

Trait	Desired direction	N1		N2		N3	
		Mean	s.e.	Mean	s.e.	Mean	s.e.
Production	+	0.139	0.005	0.145	0.007	0.142	0.011
Udder depth	+	−0.014	0.010	−0.031	0.005	−0.031	0.012
Mastitis occurrence	−	−0.002	0.005	−0.002	0.004	−0.002	0.008
Non-return rate	+	−0.028	0.002	−0.021	0.002	−0.020	0.007
Dairy form	0	0.050	0.005	0.055	0.005	0.048	0.005
Days open	−	0.033	0.004	0.035	0.002	0.033	0.004
Somatic cell score	−	−0.004	0.006	−0.006	0.005	−0.008	0.005

used to predict breeding values. Using a more advanced model for prediction of EBVs for production did not increase genetic merit for the trait. Despite selecting for udder depth, non-return rate and days open, there was a combined decline for these traits in all scenarios.

Inbreeding

The yearly increase in inbreeding in scenario N1 was high (0.5%). In Table 6, the average increase in inbreeding for cows at year 30 in each scenario is compared with the increase in scenario N1 which is set to 100. There is a higher but non-significant increase in inbreeding when comparing BG1 with BG2 whatever the model. In this study there was no significant effect of using a more advanced model on increase in inbreeding.

Generation interval

The change in accuracy of selection influenced the generation interval for both sexes but mainly for bulls (Table 7). When using the available information in the breeding value estimation, animals earlier showed their potential as candidates for selection and therefore were used to produce the next generation of offspring. Therefore generation interval decreased in scenarios where the available information in the data was fully used. This decrease in generation interval also contributed to the increase in genetic progress shown in the scenarios.

Table 4 Mean (with s.e.) of true average genetic trends from scenarios using breeding goal 2 of for the seven traits on year in simulation over 20 replicates (the desired direction of selection is indicated)

Trait	Desired direction	B1		B2		B3	
		Mean	s.e.	Mean	s.e.	Mean	s.e.
Production	+	0.109	0.009	0.108	0.011	0.107	0.007
Udder depth	+	0.030	0.005	0.024	0.008	0.030	0.008
Mastitis occurrence	−	−0.015	0.006	−0.022	0.006	−0.023	0.004
Non-return rate	+	−0.003	0.004	−0.002	0.002	0.001	0.003
Dairy form	0	0.039	0.008	0.037	0.007	0.030	0.007
Days open	−	0.020	0.004	0.018	0.002	0.018	0.005
Somatic cell score	−	−0.012	0.006	−0.017	0.006	−0.018	0.006

Table 5 Contribution to total merit index in relative economic units from production (A), mastitis (B) and other traits (C) using economic values from BG1 (A + B) and BG2 (A + B + C)

Scenario	Total merit using BG 1 (Narrow) A + B	Total merit using BG 2 (Broad) A + B + C	Production A	Mastitis B	Other traits [†] C
N1	2.80	1.72	2.70	0.10	−1.06
B1	2.86	2.67	2.11	0.75	−0.25
N2	2.91	1.87	2.81	0.10	−0.96
B2	3.23	2.99	2.11	1.12	−0.24
N3	2.85	1.97	2.75	0.10	−0.95
B3	3.25	3.07	2.08	1.17	−0.17

[†] Udder depth, non-return rate and days open.

Discussion

With the increasing number of traits with low heritability included in the breeding goals in dairy cattle breeding around the world (Miglior *et al.*, 2005) the full potential of the traits is not exploited when correlations to other traits are ignored. A multitrait selection would be improved in terms of genetic gain if it followed a multitrait evaluation but this is often computationally impossible for large data sets and many correlated traits described by very different (non) linear models. Here the approximate MT model appeared to work well and led to genetic gains similar to the more computationally demanding full linear multitrait model. The approximate multitrait model was found to be significantly better than a model corresponding to the one used today when a broad breeding goal was applied. This effect is a combination of an increase in accuracy of the breeding values in the selection criterion which directly leads to a higher genetic gain. As the simulation program is constructed this effect also leads to a lower generation interval as animals at an earlier age become interesting to produce the next generation of offspring. The lower generation interval also contributes to a higher genetic progress for total merit. The predicted breeding values for total merit were somewhat lower than the true breeding values, although in no scenarios were they significantly lower. There seems to be no reason for the predicted breeding values to be lower than the true breeding values.

Table 6 Average inbreeding coefficient (delta F) and standard error over 20 replicates for cows at year 30 for the six scenarios relative to the increase in approach N1 which is set to 100

Scenario	Delta F	s.e.
N1	100	10.5
B1	109.9	11.1
N2	103.6	9.6
B2	114.7	9.2
N3	103.4	7.4
B3	115.8	6.6

Table 7 Mean generation interval over 20 replicates for cows and bulls in the six different scenarios with standard errors

Scenario	Cows		Bulls	
	Mean	s.e.	Mean	s.e.
N1	3.43	0.009	5.44	0.12
N2	3.36	0.011	5.36	0.09
N3	3.35	0.012	5.21	0.09
B1	3.25	0.010	5.16	0.10
B2	3.16	0.009	4.88	0.12
B3	3.12	0.008	4.78	0.12

Production was more affected by a radical change of breeding goal than by a change in method for prediction of breeding values. In fact there was a small decline in genetic trend for production when a more advanced model was applied together with a broad breeding goal. So the additional information obtained by multitrait evaluation for production from other traits did not compensate for the genetic decline for the trait related to the relatively lower weight put on production. For all traits with low heritability, genetic trends improved in the desired direction. However, for reproduction, it was not possible to obtain a positive genetic trend. This was certainly due to too low an economic value for the reproduction traits in the breeding goal chosen here. In general, few theoretical or real breeding schemes have exhibited an improvement for fertility (Christensen, 1998; Sørensen *et al.*, 1999). Ducrocq *et al.* (2001) showed that the decline in fertility could be stopped and the selection differential of the 10% best bulls for total merit would have a mean genetic merit of 0.00 genetic standard deviation for fertility when using an approximate multitrait model compared with a mean of −0.49 when a single-trait model was used. When changing model for prediction of breeding values to an approximate multitrait model as described in this study, Ducrocq *et al.* (2001) also found a decline for production. But this was also a result of changing breeding goal at the same time. Besbes *et al.* (2002) used similar methods and combined linear traits, a survival trait and a categorical trait into a total merit index and found more favourable selection differentials for all traits in the breeding goal in laying hens.

Direct comparison of the effect of the six different selection criteria in the two first columns in Table 5 is not valid since they are results of differently expressed breeding goals and economic values. However, comparison of N1, N2 and N3 with B1, B2 and B3, respectively within BG1 and BG2 seems to be valid. These results suggest that selecting for a narrow breeding goal when the true economic criterion was in fact broad was worse than selecting for a broad breeding goal when the true economic criterion was in fact narrow. This effect was more pronounced when the prediction of breeding values was done with a more advanced model. What was gained from production when using a narrow breeding goal as in scenario N1, N2, and N3 did not match the lower gain in mastitis and therefore less

progress in total merit was obtained. This effect was even more pronounced when the selection criterion was broad.

Method for prediction of breeding values had no significant effect on inbreeding in this study, whereas a broad breeding goal with many traits with a low heritability gave an increase in inbreeding of approximately 10%. The reason for that was that BG2 included more traits with low heritability than BG1 which means that selection to a higher extent relies more on pedigree information and on related animals than when selecting mainly for production. This corresponds to the effect found by Sørensen *et al.* (1999). The choice of model had little or no effect on inbreeding, which might be surprising. One would expect a higher rate of inbreeding with a more advanced model due to a more extensive use of the pedigree but this effect was not significant in this study. With a higher number of replicates this might also be the case for the set up in this study. The level of inbreeding was relatively high. This was mainly due to the mating strategy, which only did not allow for matings between full- and half-sibs. All other matings were allowed. In this study any inbreeding depression was not simulated. Neither was inbreeding taken account for when predicting breeding values. Such an inclusion might have given a clearer image of the effect of increased inbreeding when applying a multitrait model and a multitrait selection criterion.

This study has a dairy-cattle specific approach. In general, more and more emphasis has been put on low heritability traits when defining breeding goals for dairy cattle over recent years world-wide (Miglior *et al.*, 2005). Also with the progeny testing system that most dairy cattle populations practise the generation interval is relatively long, which makes decisions on bulls to use as proven sires crucial for long-term genetic progress. Multitrait breeding evaluation where the available information is fuller used is appealing to improve accuracy of selection and thereby be able to make better decisions in the breeding scheme. This is not just relevant for dairy cattle breeding schemes but in any species where low heritability traits play a major role in the breeding objective. Consumers have more and more focus on animal health and welfare. Traits related to this often show low heritability independent of species. Multitrait breeding value estimation could be one way to improve animal wellbeing through improved selection.

The approximate multitrait model as described in this study using pre-adjusted data in a two-step procedure has been further developed to include more complex traits, described by non-linear models (Besbes *et al.*, 2002; Tarrés *et al.*, 2006). In this study, the main focus was on the effect of using different models for prediction of breeding values and at the same time using different breeding goals more than on model development. Therefore all models were linear. Linear models are also still widely used for breeding value estimation of binary traits which justifies the use of such models in this study.

Conclusion

Using an approximate multitrait model as described for prediction of breeding values will lead to higher genetic progress for total merit than using a univariate approach. This effect is more pronounced when using a broad breeding goal than when using a narrow breeding goal. Given a selection criterion where several traits with low heritability are a major part of the breeding goal and most traits have mutually high or moderate genetic correlations a broad breeding goal will lead to a higher genetic progress in total merit. Another result can be a redistribution of the genetic merit to get a relatively lower progress in high heritability traits such as production and a relatively higher progress for low heritability traits such as cost-reducing traits. This study shows that multitrait selection should follow multitrait evaluation.

References

- Besbes B, Ducrocq V and Protais M 2002. An approximate total merit index combining linear traits, a survival trait and a categorical trait in laying hens. Proceedings of the seventh world congress on genetics applied to livestock production. CD-ROM communication no 20–05.
- Christensen LG 1998. Possibilities for genetic improvement of disease resistance, functional traits and animal welfare. Acta Agriculturae Scandinavica Section A, Animal Science 29, (suppl.) 77–99.
- Danish Agricultural Advisory Centre 2006. <http://www.lr.dk/kvaeg/diverse/principles.pdf> (viewed 13 October 2006).
- Ducrocq V 1994. Multi trait prediction: principles and problems. Proceedings of the fifth world congress on genetics applied to livestock production, August 6–11, Guelph, Canada, 18, 455–462.
- Ducrocq V, Boichard D, Barbat A and Larroque H 2001. Implementation of an approximate multitrait BLUP to combine production traits and functional traits into a total merit index. Proceedings of the 52nd annual meeting of the European Association for Animal Production, Budapest, Hungary.
- Ducrocq V, Delaunay I, Boichard D and Mattalia S 2003. A general approach for international genetic evaluations robust to inconsistencies of genetic trends in national evaluations. Interbull technical workshop, Beltsville, Maryland, USA, 2–3. March, Interbull Bulletin 30, 101–111.
- Falconer DS and Mackay TFC 1996. An introduction to quantitative genetics, 4th ed. Longman, Harlow, UK.
- Lassen J, Sørensen MK, Madsen P and Ducrocq V 2007. Robustness of an approximate multitrait model and correction for selection bias. Genetics, Selection, Evolution (in press).
- Madsen P and Jensen J 2000. DMU: a user's guide. A package for analysing multivariate mixed models. Version 6, release 4. DJF, Foulum, Denmark.
- Miglior F, Muir BL and Van Doormaal BJ 2005. Selection indices in Holstein cattle of various countries. Journal of Dairy Science 88, 1255–1263.
- Olesen I, Groen AF and Gjerde B 2000. Definition of animal breeding goals for sustainable production systems. Journal of Animal Science 78, 570–582.
- Sørensen MK, Berg P, Jensen J and Christensen LG 1999. Stochastic simulation of breeding schemes for total merit in dairy cattle. GIFT seminar on genetic improvement of functional traits in cattle, Wageningen, The Netherlands, 7–9. November. Interbull Bulletin 23, 183–192.
- Tarrés J, Piedrafitá J and Ducrocq V 2006. Validation of an approximate approach to compute genetic correlations between longevity and linear traits. Genetics, Selection, Evolution 38, 65–85.
- Van der Werf JHJ, Van Arendonk JAM and De Vries AG 1992. Improving selection of pigs using correlated characters. Proceedings of the 43rd annual meeting of the European Association for Animal Production, Madrid, Spain.
- Willam A, Egger Danner C, Sölkner J and Gierzinger E 2002. Optimization of progeny testing schemes when functional traits play an important role in the total merit index. Livestock Production Science 77, 217–225.