Kathleen C. Horner, Yves Cazals, Régis Guieu, Marc Lenoir and Nicole Sauze

Am J Physiol Endocrinol Metab 293:1224-1232, 2007. First published Aug 21, 2007;

doi:10.1152/ajpendo.00279.2007

You might find this additional information useful...

This article cites 96 articles, 22 of which you can access free at:

<http://ajpendo.physiology.org/cgi/content/full/293/5/E1224#BIBL>

Updated information and services including high-resolution figures, can be found at:

<http://ajpendo.physiology.org/cgi/content/full/293/5/E1224>

Additional material and information about *AJP - Endocrinology and Metabolism* can be found at:

<http://www.the-aps.org/publications/ajpendo>

This information is current as of September 2, 2010 .

Experimental estrogen-induced hyperprolactinemia results in bone-related hearing loss in the guinea pig

Kathleen C. Horner,¹ Yves Cazals,¹ Régis Guieu,² Marc Lenoir,³ and Nicole Sauze²

¹Centre National de la Recherche Scientifique UMR 6153, Institut National de la Recherche Agronomique 1147, Université Paul Cézanne; ²Centre National de la Recherche Scientifique FRE 2738, Université de la Méditerranée, Marseille; and

³Institut National de la Santé et de la Recherche Médicale, Unité 583, Montpellier, France

Submitted 3 May 2007; accepted in final form 10 August 2007

Horner KC, Cazals Y, Guieu R, Lenoir M, Sauze N. Experimental estrogen-induced hyperprolactinemia results in bone-related hearing loss in the guinea pig. *Am J Physiol Endocrinol Metab* 293: E1224–E1232, 2007. First published August 21, 2007; doi:10.1152/ajpendo.00279.2007.—Our group (Horner KC, Guieu R, Magnan J, Chays A, Cazals Y. *Neuropsychopharmacology* 26: 135–138, 2002) has earlier described hyperprolactinemia in some patients presenting inner ear dysfunction. However, in that study, it was not possible to determine whether hyperprolactinemia was a cause or an effect of the symptoms. To investigate the effect of hyperprolactinemia on inner ear function, we first developed a model of hyperprolactinemia in estrogen-primed Fischer 344 rats and then performed functional studies on pigmented guinea pigs. Hyperprolactinemia induced, after 2 mo, a hearing loss of ~30–40 dB across all frequencies, as indicated by the compound action potential audiogram. During the 3rd mo, the hearing loss continued to deteriorate. The threshold shifts were more substantial in males than in females. Observations under a dissection microscope revealed bone dysmorphology of the bulla and the cochlea. Light microscopy observations of cryostat sections confirmed bone-related pathology of the bony cochlear bulla and the cochlear wall and revealed morphopathology of the stria vascularis and spiral ligament. Scanning electron microscopy revealed loss of hair cells and stereocilia damage, in particular in the upper three cochlear turns and the two outermost hair cell rows. The data provide the first evidence of otic capsule and hair cell pathology associated with estrogen-induced prolonged hyperprolactinemia and suggest that conditions such as pregnancy, anti-psychotic drug treatment, aging, and/or stress might lead to similar ear dysfunctions.

hormones; deafness; prolactin; estrogen; osteoprotegerin

PROLACTIN IS A POLYPEPTIDE hormone, originally believed to be uniquely secreted by the lactotroph cells of the anterior pituitary and was shown to have a major role in inducing lactation. Now, it is recognized to be synthesized in a wide range of tissues, including the brain (33). Receptor binding sites have been identified in a wide range of tissues, including the brain and peripheral organs, which include the otic capsule (78). Indeed, the biological action of prolactin is so widespread that it is now known to have at least 300 different functions (10, 33). Furthermore, prolactin is now recognized, together with other hormones such as cortisol, growth hormone, and endorphins, as a major stress-related hormone and is known to participate in the psychoneuroendocrinology of depression (67).

Address for reprint requests and other correspondence: K. C. Horner, CNRS UMR 6153-1147 INRA, Université Paul Cézanne, Faculté des Sciences et Techniques, Campus St Jérôme, Cases 351 352, Ave. Escadrille Normandie Nièmen, 13397 Marseille Cedex 20, France (e-mail: kathleen.horner@univ-cezanne.fr).

Some ear pathologies, such as tinnitus, are known to be affected by situations of stress (45). Stress is widely believed to trigger episodes of Ménière disease, characterized by tinnitus, fluctuant hearing loss, and episodic vertigo. Recently, our group (49) investigated stress hormones in patients with Ménière disease and, unexpectedly, revealed several cases of Ménière disease with marked hyperprolactinemia and subsequent MRI-confirmed pituitary adenoma. A further study in Sweden confirmed the association of hyperprolactinemia in several patients with Ménière disease, and one had a pituitary adenoma (30). The data are particularly intriguing because hyperprolactinemia has never before been described in this pathology; furthermore, clinical diagnostic schema for hyperprolactinemia do not include any otological pathological features (6). On the other hand, giant pituitary masses can be associated with hearing loss (9, 16, 21, 65), tinnitus (27), and vertigo (82).

To shed some light on the effect of hyperprolactinemia on inner ear function, in the present study, we first reproduced the well-established rat model of estrogen-induced hyperprolactinemia (19, 66). With this model, we could demonstrate that estrogen implants did create hyperprolactinemia. We then went on to provoke hyperprolactinemia in the pigmented guinea pig for which our group has long experience with audiometry. We hypothesized that experimentally induced hyperprolactinemia would modify cochlear function.

MATERIALS AND METHODS

Experimental procedures were reviewed and approved by the committee of the French Ministry of Agriculture and carried out under license (to K. C. Horner). The experimental protocols followed the guidelines of the Centre National de la Recherche Scientifique and the French and European laws on laboratory animal care and use.

Fischer 344 (F344) male rats (total of 30 rats) weighing ~200 g at the start of the study were supplied by Charles River (Lyon, France). Pigmented guinea pigs (total of 30 males and 30 females) weighing ~250 g at the start of the study were supplied by Centre de Production Animale (Olivet, France).

Protocol chronology. This study was carried out over a period of 3 yr. Most of the first year was devoted to investigations on F344 rats, which is a long-standing model used to induce hyperprolactinemia but which proved not to be a good model for the present study of auditory function (see RESULTS and DISCUSSION). The investigation was continued on pigmented guinea pigs. Guinea pigs were first chronically implanted with electrodes on the cochlear round window. Some animals were implanted with bilateral electrodes, and other animals

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

received only one electrode. The choice of unilateral or bilateral electrodes depended on time and material availability. The eighth nerve audiograms were tested several times over a period of 2 wk to ensure the stability of the recordings from each animal. The animals then underwent operations to introduce implants subcutaneously. Thereafter, the audiograms were monitored weekly to detect short-term and long-term effects of the hormonal changes. At the end of the observation period, the animals were anesthetized and perfused for histological investigations.

Implantation of the round window electrodes in guinea pigs. Guinea pigs were anesthetized with a mixture of 10 mg/kg xylazine (Rompun from Bayer) and 50 mg/kg ketamine (Imalgène from Merial). A Teflon-coated platinum electrode was surgically implanted on the round window of the cochlea. The electrode was led, under the skin, to a connector that was fixed by four stainless steel screws to the skull. The connector was secured with dental cement. All materials were purchased from Phymep (Paris, France).

Round-window compound action potential audiograms. Guinea pigs were restrained in a holding tube in which head movements were prevented by the use of neck and nose rings. Recordings were carried out in a sound-proof booth (Amplifon, Milan, Italy). After several sessions, the animals became familiarized with the protocol and remained calm during testing.

The compound action potential (CAP) audiogram was constructed from the visual detection of averaged evoked CAP responses. The CAP was filtered (30 Hz to 10 kHz) and averaged (~200 responses) via a Cambridge Electronic Design computer/averager. Visual threshold detection was determined in response to tone pips (2-ms rise/fall, no plateau) at different frequencies (500 Hz to 32 kHz) presented at a rhythm of 30 per second. The step size for CAP thresholds was 10 dB down toward thresholds, and thereafter it was 3 dB. The reproducibility of threshold determination was close to 3 dB across different recording sessions and across experienced raters. The tone pips were presented in free field via an earphone (Sennheiser HD 424) at a distance of 1 cm from the ear pinna. The acoustic levels at the entrance of the acoustic meatus were determined by placing a 0.5-in. condenser microphone (Bruel and Kjaer 4191) at the position of the animal's pinna. Acoustic levels were calibrated and expressed in dB sound pressure level (SPL).

Only ears presenting normal CAP audiograms, at the end of the 2-wk observation period, were included in the study. According to our criteria for normality of audiogram, threshold at 8 kHz was <10 dB SPL; threshold at 16 kHz, 4 kHz, and 2 kHz was below 15 dB SPL; and threshold at 32 kHz and 1 kHz was below 30 dB SPL. These thresholds were similar to the best estimates from behavioral or other electrophysiological studies (31). We therefore eliminated animals that manifested signs of pathology such as otitis media. In addition, we did not include data from animals that were not followed regularly every week and those that had lost their electronic connector or electrode before the end of the observation period.

Estrogen implants. Silastic medical tubing (1.98-mm inner diameter, 3.18-mm outer diameter; Dow-Corning) was cut into 3-cm lengths. One end was sealed with silicon adhesive. The tubes were incubated overnight at 38°C. Then, 25 mg of estradiol (Sigma-Aldrich, Lyon, France) were introduced into the test tubes, and some control tubes were left empty. The open end was sealed with silicon adhesive, and the tubes were again incubated overnight at 38°C. Under Rompun/Ketalar anesthesia, an incision was made in the scapula region of the skin of the animal, and one implant, filled or empty for controls, per 250 g body wt was introduced. The incision was sutured.

Blood sampling. At the end of the observation period, animal groups (treated: 10 males and 10 females; and controls: 2 males and 2 females) were decapitated under anesthesia. Whole trunk blood was collected in dry tubes, allowed to clot at 4°C, and then centrifuged at 2,000 rpm. The serum was then stored at -20°C until assay.

Serum prolactin determination. The serum prolactin level was determined with the use of a colorimetric enzyme immunoassay kit (Biotrak, RPN 2563) supplied by Amersham Life Sciences. The protocol supplied with the kit was followed, and serum samples were diluted 1:5 in assay buffer (supplied). Briefly, samples, as well as the blanks and standards, were run simultaneously on the same assay plate. All incubations were carried out at room temperature. All samples were added to the coated wells and incubated with antiserum for 3 h. Prolactin conjugate was added and incubated for 30 min. The plate was decanted and washed four times with wash buffer. Amplex amplification reagent (horseradish peroxidase coupled) was added and again incubated for 30 min. After samples were washed four times with wash buffer, substrate was added and incubated for a further 30 min. The coloration was arrested by adding 1 M sulfuric acid to the wells. The plate was read on a spectrophotometer plate reader at 450 nm. A calibration curve was constructed from working standards supplied (0.82–200 ng/ml), and the experimental values were read directly from the curve. The assay kit had an analytic sensitivity of 0.44 ng/ml.

Serum estrogen determination. Serum estrogen was determined by the double antibody estradiol ¹²⁵I radioimmunoassay kit supplied by DPC France. The protocol supplied with the kit was followed, and serum samples were diluted 1:10 in the zero calibrator (supplied). Incubations were carried out at room temperature. Briefly, the serum samples were pipetted, antiserum was added, and samples were vortexed, followed by incubation for 2 h. ¹²⁵I-labeled estradiol was added, vortexed, and incubated for 1 h. Cold precipitating solution was added, vortexed, and incubated for 10 min. After centrifugation for 15 min at 3,000 g, the tubes were decanted and inverted on absorbent paper for 10 min. The tubes were then passed in a gamma counter, and counting was carried out over 1 min. A calibration curve was constructed from supplied calibrators (5–500 pg/ml), and the experimental values were read off directly from the curve. The assay kit had an analytic sensitivity of 1.4 pg/ml.

Light microscopy. The animals were anesthetized and then underwent intracardiac perfusion with ice-cold saline followed by paraformaldehyde (4% in phosphate buffer 0.1 M). The cochleae were removed and immersed in fixative for at least 2 h or up to overnight. They were then rinsed in phosphate buffer (0.1 M). The cochleae were then examined under the dissection microscope for detection of indexes of pathology. They were then decalcified in EDTA for up to 3 wk. The cochleae were then rinsed in phosphate buffer and immersed in sucrose (30%) overnight. The specimens were then bathed in cryoblock and rapidly frozen to -20°C in methyl butane. The cochleae were sectioned (15 µm) on a cryostat and colored with hematoxylin-eosin. Observations were carried out on a Nikon Eclipse E600 light microscope, and photography was carried out via a camera (Nikon) linked to ACT-1 software.

Scanning electron microscopy. Two alternative fixation procedures were employed. Some animals underwent intracardiac perfusion with ice-cold saline followed by paraformaldehyde (4% in phosphate buffer, 0.1 M). The cochleae were removed and perfused with phosphate buffer followed by glutaraldehyde (2.5% in phosphate buffer, 0.1 M). After 2 h, the cochleae were perfused and rinsed three times (each 15 min) in phosphate buffer (0.1 M).

For animals that had been decapitated for trunk blood sampling, some cochleae were rapidly removed and perfused via the round and oval windows with glutaraldehyde (2.5% in phosphate buffer). After 2 h, the cochleae were perfused and rinsed three times (each 15 min) in phosphate buffer (0.1 M).

In all cases, the cochleae were then perfused with osmic acid (1% in phosphate buffer) and further fixed for 1 h. This was followed by perfusion and rinsing three times (each 15 min) in phosphate buffer. Dehydration was carried out to up to 70% alcohol (each 15 min). Dissection was performed to remove the bony wall of the cochlea and to expose the organ of Corti. The specimens were then prepared for

scanning electron microscopy (SEM) after critical point drying and coating with gold-palladium.

Statistical analysis. Statistical analysis was carried out with the SigmaStat 3.1 program (Systat Erkrath). Data at 1, 2, and 3 mo were considered for statistics. For analysis of acoustic sensitivity data, a three-way (sex or laterality, time, frequency) ANOVA was performed followed by examination of all pair-wise comparisons using the Holm-Sidak method.

RESULTS

Attempts to follow auditory function in F344 rats, by brain stem recordings, were abandoned for two reasons. We did find some hearing sensitivity loss (~ 20 dB after 1 mo) in the F344 rats. However, several recent publications, which appeared after the start of our study, showed that hearing function in untreated F344 rats deteriorates with aging faster than in other strains of rats (74, 75). In addition, we encountered technical complications of instability of long-term recordings associated with rejection of the screws from the skull after ~ 1 mo. Because this did not happen in control rats and because we used the same type of screws and cement as for guinea pigs, we suggest that one plausible explanation would be a modified immune response related to hyperprolactinemia in these rats (33).

Pigmented guinea pigs. The outward sign of the changing hormonal status in the guinea pigs was a break in the body weight growth curve, which could be detected 1 mo after insertion of the estrogen implants (data not shown). At the end of 3-mo treatment, some animals started to lose their hair.

Hormone assays. Standard control prolactin values are between 10 and 50 ng/ml in normal young rat. In the absence of published data, to our knowledge, the standard control values for guinea pigs were assumed to be similar. Control values for one male and one female were below the lower end of the calibration curve (ends at 10 ng/ml) calculated to be ~ 10 ng/ml. In treated animals, prolactin had minimum-median-maximum values of 78–143–438 ng/ml for males ($n = 10$) and 23–83–625 ng/ml for females ($n = 10$). There was no significant difference between males and females according to a Mann-Whitney rank sum test ($P = 0.121$; Fig. 1, left).

The experimental estrogen levels increased above standard control values of 5–20 pg/ml (reference in normal young rat). Control values for one male and one female were below the lower end of the calibration curve (ends at 50 pg/ml) but calculated to be ~ 10 and ~ 15 pg/ml, respectively. For estrogen, the minimum-median-maximum values were 138–861–1,688 pg/ml for males ($n = 10$) and 175–751–8,900 pg/ml for females ($n = 10$). There was no significant difference between males and females according to a Mann-Whitney rank sum test ($P = 0.91$; Fig. 1, right).

At the time of sampling, after 3 mo of treatment, we found no significant relationship, according to Pearson product-moment correlation, between prolactin and estrogen for males ($P = 0.19$), females ($P = 0.33$), or males and females together ($P = 0.15$).

Audiogram changes. Twelve test males (4 bilateral and 8 unilateral; total ears: $n = 16$) and 13 test females (7 bilateral and 6 unilateral; total ears: $n = 20$) presented good audiograms at the onset of the study and could be followed over 3 mo. Control animals with empty tubes (5 males and 5 females) showed no change in the audiogram over the observation

Fig. 1. Immunoassay for determination of guinea pig prolactin and estrogen. Blood serum from estrogen-primed guinea pigs was employed for determination of prolactin (ng/ml) by enzyme immunoassay (left) and for estrogen (pg/ml) by radioimmunoassay (right). ●, Males; ○, females. In each case, the calibration curve was supplied and adjusted here to take into account the respective dilutions (black plot with fixed reference values represented as black stars). On the y-axes, %B/B₀ represents the percentage of immunobinding for the calibration values from which the experimental values were calculated according to the kit instructions and plotted (gray plot).

period. In test animals with estrogen implants, some threshold fluctuations could be observed over several weeks (not shown here). These fluctuations were considered particularly interesting but not detailed for the purpose of the present long-term study. No consistent deterioration of the audiogram could be detected before several weeks. After 2 mo, definite threshold shifts could be detected, and, after 3 mo, the threshold shifts progressed, in general, in males and in females. At this stage, some threshold shifts from males ($n = 7$) were more than 60 dB and in one case above 80 dB at the limit of the calibration curve of the acoustic setup (Fig. 2).

For all animals (males or females) with bilateral data, ANOVA (3 way: laterality, months, frequency) showed no statistical difference between left and right ears at any month or frequency. Thereafter, all results from all ears were considered in further statistical analysis. A three-way ANOVA (sex, months, frequency) indicated significant differences between months ($P < 0.001$), between sexes ($P < 0.001$), and between frequencies ($P < 0.001$), with significant interactions between months and sex ($P < 0.019$) and frequency and sex ($P < 0.01$). All pair-wise multiple comparisons were then made with the Holm-Sidak method, and a significance level of 0.05 was chosen. All results cannot be listed here; however, it appeared, as can also be seen in Fig. 2, that threshold shifts increased and differed significantly at all 3 mo for males and for females. Males differed significantly from females at all 3 mo.

Light microscopy observations. All cochleae were visualized under the dissection microscope during the perfusion for fixation. At dissection, the bulla wall was perceived to be thicker and harder than in control specimens. When the bulla was opened, the middle ear was clearly visible, and the absence of otitis media was confirmed in all cases (Fig. 3). At this stage, the presence of some type of bone remodeling could be clearly identified within the cochlear bulla (arrow 1) and the cochlea (arrow 3) compared with the normal specimens. In addition, the dark band corresponding to the position of stria vascularis was less easily identified than normal (arrow 2). Within the

Fig. 2. Audiogram changes in estrogen-primed guinea pigs over 3 mo. Auditory nerve compound action potential (CAP) threshold shift (dB) is plotted as a function of frequency (kHz). There was no significant difference between right and left ears for males or females; therefore, data from both ears are presented without distinction. *Left:* females ($n = 20$). *Right:* males ($n = 16$). Progressive threshold shifts observed throughout 3 mo demonstrated more loss for males than for females.

limits of the macro-observations under the dissection microscope, the middle ear ossicles appeared normal in the majority of cases, but clear bone dysmorphology of the middle ear ossicles could clearly be identified in two specimens (*arrow 4*).

Light microscopy was carried out on cochleae from treated females (4 unilateral and 3 bilateral; total ears: $n = 10$) and males (4 unilateral and 2 bilateral; total ears: $n = 8$) and control cochleae (3 unilateral female and 3 unilateral male; $n = 6$). The cochlea images presented in Fig. 4, *D–F*, were from a male animal and are representative of the cochlear pathology observed. There was no evidence for otitis media. The middle ear ossicles could be clearly identified at the base of the cochlea (Fig. 4*D*). The most obvious pathological feature was the bone dysmorphology throughout the cochlear bone, which could represent otospongiotic-like foci (Fig. 4*D*). This was particu-

larly evident immediately behind the stria vascularis/spiral ligament (Fig. 4*E*). In addition, the spiral ligament showed signs of deterioration and the collagen fibers were less easily identified than in control cochleae (Fig. 4*F*). The stria vascularis also appeared degenerated (Fig. 4*F*). No control cochlea presented such anomalies.

One specimen from a female presented normal thresholds in both ears for frequencies between 16 and 32 kHz together with a sensitivity loss of more than 40 dB for all frequencies below. Because the two ears had similar audiograms, one cochlea was observed in light microscope and the other in SEM. Light microscopy did not reveal changes to either bone or the stria vascularis. The other cochlea was observed in SEM, where pathological changes could be detected at hair cell level, as detailed below.

Fig. 3. Macroscopic view of cochleae illustrating the bone dysmorphology. Dissection microscope view of a control cochlea (*left*) and a cochlea from a treated animal (*right*). Obvious pathological features could be observed, including thickening of the bulla wall (arrow 1) and cochlear wall dysmorphology (arrow 3). Stria vascularis was less visible through the bony wall of the cochlea (arrow 2); however, middle ear bones were exceptionally calcified (2 cases), as presented here (arrow 4).

Scanning microscopy. SEM observations were carried out on cochleae from male (2 unilateral, $n = 2$) and female (1 bilateral and 1 unilateral, $n = 3$) guinea pigs. The cochleae were observed throughout their length.

The two cochleae from the male guinea pigs presented at least 30-dB losses across the audiogram, and SEM observa-

tions confirmed the loss of hair cells and stereocilia pathology throughout the cochlea with a slightly better preservation in the upper basal turn.

One female guinea pig presented a substantial sensitivity loss across the audiogram (>75 dB) for which SEM confirmed the degeneration of stereocilia bundles and loss of hair cells

Fig. 4. Light microscopy showing pathology of the bony cochlea. Cryostat sections (15 μ m, hematoxylin-eosin stained) from a control cochlea (A–C) are compared with those from a treated animal (D–F). For the experimental cochleae, there was complete absence of otitis media, and the middle ear ossicles could be clearly identified (arrow in D). The most striking pathology was the apparent osteoporosis of the bony wall of the cochlea. At higher magnification (rectangle in D presented in E), the bone appeared particularly fragile close to the spiral ligament. In addition, the stria vascularis (arrow in E presented in F) showed thinning. The spiral ligament appeared fragile, and collagen fibers were less easily identified.

throughout the cochlea. The other ear, on the other hand, presented very little sensitivity loss, and indeed the SEM observations confirmed the excellent condition of the stereocilia and hair cells. Indeed, they were comparable to those seen in the organ of Corti of a control guinea pig (Fig. 5, A–C).

For the female guinea pig presenting very similar audiograms in the two ears, one cochlea was observed in light microscopy and the other in SEM. Light microscope observations did not reveal modifications. In contrast, SEM revealed stereocilia pathology and hair cell loss above the basal turn of the cochlea. The damage essentially involved loss of hair cells from the two more external rows. The stereociliary bundles of the remaining hair cells of rows 2 and 3 appeared limp or folded, in particular for row 3 (Fig. 5, D–F).

DISCUSSION

Estrogen treatment in pigmented guinea pigs led to body-weight loss, in agreement with first studies on this animal model (19, 66) and in accordance with its anorectic property described more recently (34, 35). In the longer term, estrogen induced hyperprolactinemia, for the large part via inhibition of dopamine synthesis in tuberinfundibular dopaminergic neurons (3, 5). The treatment resulted in body hair loss in accordance with prolactin modulation of hair growth in humans (32) and in mice (24). Beyond these expected results, our study revealed two major new findings concerning the ear. Long-term estrogen treatment led to hearing loss, which was more substantial

in males than in females. There was a close link between threshold shifts and bone dysmorphology of the cochlear bulla and cochlea.

Limitations of the present data. Elevated experimentally induced levels of estrogen and prolactin were confirmed. We were not able to correlate hormone levels with threshold shifts because different experimental groups of animals were involved. On the other hand, when substantial threshold shifts were recorded, otic bone dysmorphology was systematically observed. At early stages of the pathology, hearing loss was fluctuant (not documented). In one case, bone dysmorphology was not detected by light microscopy, but SEM confirmed hair cell stereocilia disarray in the three upper cochlear turns, which followed a gradient from the third outer hair cell row to first row, similar to that described earlier for experimental endolymphatic hydrops in the guinea pig (50, 79, 80). Actin depolymerization of the stereocilia was hypothesized in that model to account for the fluctuant hearing losses (48). Interestingly, a causal link between F-actin disassembly and increased prolactin has been established for pituitary cells in culture (13). It would be interesting to pursue this hypothesis, to determine the relationship between stereocilia pathology and early threshold fluctuations.

We found that males were more sensitive than females, although at the end of 3-mo observation period the median levels of prolactin and estrogen in males and females were not significantly different. The sex difference might have been due

Fig. 5. Cochlear hair cells as seen in scanning electron microscopy. In cases where the hearing loss across the audiogram was moderate, the stereociliary bundles appeared, as shown for a control guinea pig (A–C). A more profound low hearing loss was associated with essentially normal basal turn of the cochlea (D) and substantial damage, including loss of hair cells, from the 2 more external rows in the middle turn (E) and apical turn (F). The stereociliary bundles of the remaining hair cells of rows 2 and 3 appeared limp and folded, particularly for row 3 (E and F). I, inner hair cells; O, outer hair cells; iP, pillar supporting cells.

to dynamic interactions between estrogen and prolactin or the interplay of other hormones such as cortisol, which can protect against acoustic trauma (12) and can be sex-difference sensitive in some patients with inner ear pathology (47).

Estrogen and the cochlea. Estrogen receptors α and β have been identified within the inner ear of rats and mice in various cell types, including hair cells, spiral ganglion cells, stria vascularis, and spiral ligament (91). Observations on the isolated stria vascularis from gerbils have shown that estrogen rapidly inhibited the slow component of the potassium current channel in the apical membrane of the marginal cells, suggesting that estrogen could influence K^+ secretion in vivo via a nongenomic mechanism (56). Data from human studies are equally scarce. Estrogen receptor α -containing cells have been located selectively in the spiral ganglion and β -containing cells in the stria vascularis (90).

Estrogen can positively affect hearing. Estrogen does affect hearing, but the data are often contradictory (51). Aging or ovariectomy in rats resulted in increased latency of the acoustically evoked brain stem responses. Estrogen treatment was not found to change these observations in older rats (22) but did reverse the latency changes after ovariectomy in younger rats (20). Hearing in mice was best preserved in females relative to males until menopause, beyond which the advantage was lost (40). Indeed, when estrogen was suppressed in female mice, by administration of tamoxifen, reduced medial olivocochlear feedback was observed (93). Sex differences in age-related human hearing loss is well documented (37). Hearing sensitivity declines rapidly above the age of 20–30 yr in men and above the age of 50 yr in women (71). Better hearing has been reported to be correlated with higher serum estradiol levels in postmenopausal women (53).

Estradiol can negatively affect hearing. Several reports have indeed suggested that oral contraceptives may increase risk of hearing loss and in particular otosclerosis, although no clear picture has emerged (96). Otosclerosis is estimated to have a prevalence of 0.3% (36) according to clinical manifestations and up to 3% (26) or more according to histological examination of temporal bones postmortem. It is an age-related ear pathology that becomes manifest, in general, between ages 20 and 40 yr, affects twice as many women as men, and is usually bilateral (15, 25, 58). Hearing loss related to cochlear otosclerosis worsens during periods of intense hormonal activity (83). Otosclerosis often begins during a first pregnancy and is seriously aggravated during successive pregnancies (38).

Reversible (44) and irreversible hearing losses have also been described in women after the use of contraceptives (68), as well as with hormone replacement therapy (92). Elevated serum prolactin levels in women can occur with high-dose estrogen contraceptive pills (52). A recent report has revealed that hormone replacement therapy, using estrogen and progesterin together, negatively affects hearing in aged women, whereas estrogen alone does not (39). Hyperprolactinemia can be induced by an estrogen-progesterone synergy (72, 97). Interestingly, treatment with estrogen together with progesterone for 30 days in guinea pigs resulted in minor changes in latency and decreased amplitude of brain stem-evoked responses, as well as an inflammatory pattern within the cochlea (7).

Hyperprolactinemia. Hyperprolactinemia prevalence in an unselected normal population has been reported to be 0.4% but

can be as high as 17% in a population of women with reproductive problems (6). Hyperprolactinemia is associated with aging in rats (29, 89) and in women and men (4, 8, 77, 81, 95). Prolactin levels rise during pregnancy and in particular during the last trimester (73) as well as during lactation (94). In addition, schizophrenia is supposed to be associated with hyperactivity of dopamine, such that neuroleptic treatment has its therapeutic effect through inhibition of central dopamine functions (14). Hyperprolactinemia is a well recognized side effect of antipsychotic drug treatment (28) and in schizophrenia (41, 42).

Hyperprolactinemia can lead to osteoporosis and possibly otosclerosis. Prolactin influences calcium mobilization in rats (55, 69). One major side effect of hyperprolactinemia in humans is decreased bone mineral density (54, 64) and osteoporosis (76). Hyperprolactinemia is associated with decreased bone mass density, which occurs during pregnancy (2), lactation (70), and schizophrenia (1, 43, 62, 63).

Strangely, hyperprolactinemia has not previously been linked to otosclerosis. On the other hand, decreased bone mineral density has been found to be moderately associated with hearing loss in older men but not in women (46). In the presence of osteoporosis, the chances of having otosclerosis are increased by a factor of two (17) or four (18). Furthermore, abnormal *COL1A1* collagen gene expression might be implicated in the two pathologies (18, 61). At this point, it is particularly interesting to recall the earlier reports of middle ear bone remodeling in the gerbil after chemical sympathectomy by guanethidine sulfate (86) or hydroxydopamine (85). Although these neurotoxins do eliminate sympathetic activity, there are, in parallel, major central consequences. In particular, both treatments reduce hypothalamic dopamine, which leads to an increase in serum prolactin levels (57, 59). In light of the present data, the possibility of a prolactin involvement in chemical sympathectomy should certainly be considered.

Prolactin is known to have a direct inhibitory effect on osteoblast activity (23); in addition, prolactin regulates the expression of receptor activator of NF- κ B ligand (RANKL), at least in mammary gland (88). RANKL, expressed in osteoblasts, is highly involved in bone remodeling and activates its receptor RANK on osteoclasts (60). Osteoprotegerin, which is an antagonist of RANKL, has been found to be produced within the cochlea in the spiral ligament (98) from where it is likely to be secreted into the perilymph and can hence diffuse into surrounding otic capsule bone (60). Estrogen-induced hyperprolactinemia, described in the present study, may lead to bone remodeling of the otic capsule by interfering with the fine balance between osteoprotegerin-RANKL-RANK within the otic capsule.

Conclusions. Although experimental chronic estrogen-induced hyperprolactinemia in the guinea pig may not represent all categories of hyperprolactinemia in humans, the data allow us to speculate that hyperprolactinemia observed by us earlier, in some patients with Ménière disease (30, 49), might have played some causative role in the symptoms. In fact, otosclerosis and Ménière disease can coexist, although the causal relationship remains controversial (84, 87). Because both can present low-frequency hearing loss, tinnitus, and vestibular dysfunction, otosclerosis can mimic the symptoms of Ménière disease (11), and the diagnosis of otosclerosis can even be missed at first (87).

The unexpected novel finding of prolactin-related otic capsule bone dysmorphology fits with recent concepts regarding the molecular biology of otosclerosis, and our data suggest that prolactin is acting via a osteoprotegerin-RANKL-RANK mechanism.

ACKNOWLEDGMENTS

The authors are grateful for the technical aid for the preliminary part of this study on rats by C. Lopez and M. Lucciano and technical advice for the final part of the study by J. Pio. We sincerely thank A. Jean for providing laboratory space. We are grateful for helpful comments of S. D. G. Stevens and S. Rydmarker and to anonymous reviewers who improved the final version of the manuscript.

REFERENCES

1. Abraham G, Halbreich U, Friedman RH, Josiassen RC. Bone mineral density and prolactin associations in patients with chronic schizophrenia. *Schizophr Res* 59: 17–18, 2003.
2. Akesson A, Vahter M, Berglund M, Eklof T, Bremme K, Bjellerup P. Bone turnover from early pregnancy to postweaning. *Acta Obstet Gynecol Scand* 83: 1049–1055, 2004.
3. Arita J, Kimura F. Direct inhibitory effect of long term estradiol treatment on dopamine synthesis in tuberoinfundibular dopaminergic neurons: in vitro studies using hypothalamic slices. *Endocrinology* 121: 692–698, 1987.
4. Baranowska B, Wolinska-Witort E, Bik W, Baranowska-Bik A, Martynska L, Chmielowska M. Evaluation of neuroendocrine status in longevity. *Neurobiol Aging* 28: 774–783, 2007.
5. Ben-Jonathan N, Hnasko R. Dopamine as a prolactin (PRL) inhibitor. *Endocr Rev* 22: 724–763, 2001.
6. Biller BM, Luciano A, Crosignani PG, Molitch M, Olive D, Rebar R, Sanfilippo J, Webster J, Zacur H. Guidelines for the diagnosis and treatment of hyperprolactinemia. *J Reprod Med* 44: 1075–1084, 1999.
7. Bittar RS, Cruz OL, Lorenzi MC, Marone SA, Miniti A. Morphological and functional study of the cochlea after administration of estrogen and progesterone in the guinea pig. *Int Tinnitus J* 7: 41–45, 2001.
8. Blackman MR, Kowatch MA, Wehmann RE, Harman SM. Basal serum prolactin levels and prolactin responses to constant infusions of thyrotropin releasing hormone in healthy aging men. *J Gerontol* 41: 699–705, 1986.
9. Blagoveshchenskaia NS, Leushkina LN. [Otoneurological symptomatology in tumors of the hypophysis with marked extrasellar growth]. *Zh Vopr Neirokhir Im N N Burdenko*: 24–28, 1988.
10. Bole-Feysot C, Goffin V, Edery M, Binart N, Kelly PA. Prolactin (PRL) and its receptor: actions, signal transduction pathways and phenotypes observed in PRL receptor knockout mice. *Endocr Rev* 19: 225–268, 1998.
11. Brookler KH, Glenn MB. Meniere's syndrome: an approach to therapy. *Ear Nose Throat J* 74: 534–538, 540, 542, 1995.
12. Canlon B, Meltser I, Johansson P, Tahera Y. Glucocorticoid receptors modulate auditory sensitivity to acoustic trauma. *Hear Res* 226: 61–69, 2007.
13. Carbajal ME, Vitale ML. The cortical actin cytoskeleton of lactotropes as an intracellular target for the control of prolactin secretion. *Endocrinology* 138: 5374–5384, 1997.
14. Carlsson A. Antipsychotic drugs, neurotransmitters, schizophrenia. *Am J Psychiatry* 135: 165–173, 1978.
15. Chole RA, McKenna M. Pathophysiology of otosclerosis. *Otol Neurotol* 22: 249–257, 2001.
16. Ciftci E, Erden I, Akay S. Brucellosis of the pituitary region. *MRI Neuroradiol* 40: 383–384, 1998.
17. Clark K, Sowers MR, Wallace RB, Jannausch ML, Lemke J, Anderson CV. Age-related hearing loss and bone mass in a population of rural women aged 60 to 85 years. *Ann Epidemiol* 5: 8–14, 1995.
18. Clayton AE, Mikulec AA, Mikulec KH, Merchant SN, McKenna MJ. Association between osteoporosis and otosclerosis in women. *J Laryngol Otol* 118: 617–621, 2004.
19. Clifton KH, Meyer RK. Mechanism of anterior pituitary tumor induction by estrogen. *Anat Rec* 125: 65–81, 1956.
20. Coleman JR, Campbell D, Cooper WA, Welsh MG, Moyer J. Auditory brainstem responses after ovariectomy and estrogen replacement in rat. *Hear Res* 80: 209–215, 1994.
21. Connolly ES Jr, Winfree CJ, Carmel PW. Giant posterior fossa cystic craniopharyngiomas presenting with hearing loss. Report of three cases and review of the literature. *Surg Neurol* 47: 291–299, 1997.
22. Cooper WA, Ross KC, Coleman JR. Estrogen treatment and age effects on auditory brainstem responses in the post-breeding Long-Evans rat. *Audiology* 38: 7–12, 1999.
23. Coss D, Yang L, Kuo CB, Xu X, Luben RA, Walker AM. Effects of prolactin on osteoblast alkaline phosphatase and bone formation in the developing rat. *Am J Physiol Endocrinol Metab* 279: E1216–E1225, 2000.
24. Craven AJ, Nixon AJ, Ashby MG, Ormandy CJ, Blazek K, Wilkins RJ, Pearson AJ. Prolactin delays hair regrowth in mice. *J Endocrinol* 191: 415–425, 2006.
25. Cross NA, Hillman LS, Allen SH, Krause GF, Vieira NE. Calcium homeostasis and bone metabolism during pregnancy, lactation, and post-weaning: a longitudinal study. *Am J Clin Nutr* 61: 514–523, 1995.
26. Declau F, Van Spaendonck M, Timmermans JP, Michaels L, Liang J, Qiu JP, Van de Heyning P. Prevalence of otosclerosis in an unselected series of temporal bones. *Otol Neurotol* 22: 596–602, 2001.
27. Dominguez Ugidos LJ, Martinez Subias J, Urpugui Garcia A, Sancho Serrano E, Alfonso Collado JL, Valles Varela H. [Invasive giant prolactinoma.] *Acta Otorrinolaringol Esp* 49: 156–158, 1998.
28. Emiliano AB, Fudge JL. From galactorrhea to osteopenia: rethinking serotonin-prolactin interactions. *Neuropsychopharmacology* 29: 833–846, 2004.
29. Esquifino AI, Cano P, Jimenez V, Reyes Toso CF, Cardinali DP. Changes of prolactin regulatory mechanisms in aging: 24-h rhythms of serum prolactin and median eminence and adenohypophyseal concentration of dopamine, serotonin, gamma-aminobutyric acid, taurine and somatostatin in young and aged rats. *Exp Gerontol* 39: 45–52, 2004.
30. Falkenius-Schmidt K, Rydmarker S, Horner KC. Hyperprolactinemia in some Meniere patients even in the absence of incapacitating vertigo. *Hear Res* 203: 154–158, 2005.
31. Fay RR. Comparative psychoacoustics. *Hear Res* 34: 295–305, 1988.
32. Foitzik K, Krause K, Conrad F, Nakamura M, Funk W, Paus R. Human scalp hair follicles are both a target and a source of prolactin, which serves as an autocrine and/or paracrine promoter of apoptosis-driven hair follicle regression. *Am J Pathol* 168: 748–756, 2006.
33. Freeman ME, Kanyicska B, Lerant A, Nagy G. Prolactin: structure, function, and regulation of secretion. *Physiol Rev* 80: 1523–1631, 2000.
34. Gao Q, Mezei G, Nie Y, Rao Y, Choi CS, Bechmann I, Leranth C, Toran-Allerand D, Priest CA, Roberts JL, Gao XB, Mobbs C, Shulman GI, Diano S, Horvath TL. Anorectic estrogen mimics leptin's effect on the rewiring of melanocortin cells and Stat3 signaling in obese animals. *Nat Med* 13: 89–94, 2007.
35. Geary N. Estradiol, CCK and satiation. *Peptides* 22: 1251–1263, 2001.
36. Gordon MA. The genetics of otosclerosis: a review. *Am J Otol* 10: 426–438, 1989.
37. Gordon-Salant S. Hearing loss and aging: new research findings and clinical implications. *J Rehabil Res Dev* 42: 9–24, 2005.
38. Gristwood RE, Venables WN. Pregnancy and otosclerosis. *Clin Otolaryngol* 8: 205–210, 1983.
39. Guimaraes P, Frisina ST, Mapes F, Tadros SF, Frisina DR, Frisina RD. Progesterone negatively affects hearing in aged women. *Proc Natl Acad Sci USA* 103: 14246–14249, 2006.
40. Guimaraes P, Zhu X, Cannon T, Kim S, Frisina RD. Sex differences in distortion product otoacoustic emissions as a function of age in CBA mice. *Hear Res* 192: 83–89, 2004.
41. Halbreich U, Kahn LS. Hyperprolactinemia and schizophrenia: mechanisms and clinical aspects. *J Psychiatr Pract* 9: 344–353, 2003.
42. Halbreich U, Kinon BJ, Gilmore JA, Kahn LS. Elevated prolactin levels in patients with schizophrenia: mechanisms and related adverse effects. *Psychoneuroendocrinology* 28, Suppl 1: 53–67, 2003.
43. Halbreich U, Rojansky N, Palter S, Hreshchysyn M, Kreeger J, Bakhai Y, Rosan R. Decreased bone mineral density in medicated psychiatric patients. *Psychosom Med* 57: 485–491, 1995.
44. Hanna GS. Sudden deafness and the contraceptive pill. *J Laryngol Otol* 100: 701–706, 1986.
45. Hebert S, Païement P, Lupien SJ. A physiological correlate for the intolerance to both internal and external sounds. *Hear Res* 190: 1–9, 2004.
46. Helzner EP, Cauley JA, Pratt SR, Wisniewski SR, Talbott EO, Zmuda JM, Harris TB, Rubin SM, Taaffe DR, Tykavsky FA, Newman AB. Hearing sensitivity and bone mineral density in older adults: the Health, Aging and Body Composition Study. *Osteoporos Int* 16: 1675–1682, 2005.

47. Horner KC, Cazals Y. Stress hormones in Meniere's disease and acoustic neuroma. *Brain Res Bull* 66: 1–8, 2005.
48. Horner KC, Cazals Y, Guilhaume A. Round window cochlear microphonics and atrophy of short and middle stereocilia on outer hair cells in hydropic cochleas in the guinea pig. In: *Cochlear Mechanisms*, edited by Wilson JP and Kemp DT. London: Plenum, 1989, p. 207–215.
49. Horner KC, Guieu R, Magnan J, Chays A, Cazals Y. Prolactinoma in some Meniere's patients—is stress involved? *Neuropsychopharmacology* 26: 135–138, 2002.
50. Horner KC, Guilhaume A, Cazals Y. Atrophy of middle and short stereocilia on outer hair cells of guinea pig cochleas with experimentally induced hydrops. *Hear Res* 32: 41–48, 1988.
51. Hultcrantz M, Simonoska R, Stenberg AE. Estrogen and hearing: a summary of recent investigations. *Acta Otolaryngol (Stockh)* 126: 10–14, 2006.
52. Ismail MS, Serour GI, Torsten U, Weitzel H, Berlien HP. Elevated serum prolactin level with high-dose estrogen contraceptive pills. *Eur J Contracept Reprod Health Care* 3: 45–50, 1998.
53. Kim SH, Kang BM, Chae HD, Kim CH. The association between serum estradiol level and hearing sensitivity in postmenopausal women. *Obstet Gynecol* 99: 726–730, 2002.
54. Klibanski A, Neer RM, Beitins IZ, Ridgway EC, Zervas NT, McArthur JW. Decreased bone density in hyperprolactinemic women. *N Engl J Med* 303: 1511–1514, 1980.
55. Krishnamra N, Seemoung J. Effects of acute and long-term administration of prolactin on bone ⁴⁵Ca uptake, calcium deposit, and calcium resorption in weaned, young, and mature rats. *Can J Physiol Pharmacol* 74: 1157–1165, 1996.
56. Lee JH, Marcus DC. Estrogen acutely inhibits ion transport by isolated stria vascularis. *Hear Res* 158: 123–130, 2001.
57. Lin JY, Mai LM, Pan JT. Effects of systemic administration of 6-hydroxydopamine, 6-hydroxydopa and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) on tuberoinfundibular dopaminergic neurons in the rat. *Brain Res* 624: 126–130, 1993.
58. Lopez JM, Gonzalez G, Reyes V, Campino C, Diaz S. Bone turnover and density in healthy women during breastfeeding and after weaning. *Osteoporos Int* 6: 153–159, 1996.
59. Machado LJ, Reis AM, Coimbra CC. Evidence of angiotensin II involvement in prolactin secretion in response to hemorrhage in adrenalectomized and guanethidine-treated rats. *Eur J Endocrinol* 146: 439–446, 2002.
60. McKenna MJ, Kristiansen AG. Molecular biology of otosclerosis. *Adv Otorhinolaryngol* 65: 68–74, 2007.
61. McKenna MJ, Nguyen-Huynh AT, Kristiansen AG. Association of otosclerosis with Sp1 binding site polymorphism in COL1A1 gene: evidence for a shared genetic etiology with osteoporosis. *Otol Neurotol* 25: 447–450, 2004.
62. Meaney AM, Smith S, Howes OD, O'Brien M, Murray RM, O'Keane V. Effects of long-term prolactin-raising antipsychotic medication on bone mineral density in patients with schizophrenia. *Br J Psychiatry* 184: 503–508, 2004.
63. Meltzer HY, Busch DA, Fang VS. Serum neuroleptic and prolactin levels in schizophrenic patients and clinical response. *Psychiatry Res* 9: 271–283, 1983.
64. Naidoo U, Goff DC, Klibanski A. Hyperprolactinemia and bone mineral density: the potential impact of antipsychotic agents. *Psychoneuroendocrinology* 28, Suppl 2: 97–108, 2003.
65. Negishi K, Suzuki T, Masuda Y, Masugi Y, Teramoto A, Ohama E. [Two autopsy cases of primary pituitary carcinoma.] *No To Shinkei* 40: 491–496, 1988.
66. Nelson W. Some effects of hypophyseal tumors induced by treatment with estrogens. *Cancer Res* 2: 728, 1942.
67. Nicholas L, Dawkins K, Golden RN. Psychoneuroendocrinology of depression. *Prolactin Psychiatr Clin North Am* 21: 341–358, 1998.
68. Okulicz G. [Inner ear disturbance following long-term usage of hormonal contraceptives.] *Hno* 26: 330–334, 1978.
69. Pahuja DN, DeLuca HF. Stimulation of intestinal calcium transport and bone calcium mobilization by prolactin in vitamin D-deficient rats. *Science* 214: 1038–1039, 1981.
70. Pearson D, Kaur M, San P, Lawson N, Baker P, Hosking D. Recovery of pregnancy mediated bone loss during lactation. *Bone* 34: 570–578, 2004.
71. Pearson JD, Morrell CH, Gordon-Salant S, Brant LJ, Metter EJ, Klein LL, Fozard JL. Gender differences in a longitudinal study of age-associated hearing loss. *J Acoust Soc Am* 97: 1196–1205, 1995.
72. Pecins-Thompson M, Bethea CL. RU 486 blocks and fluoxetine augments progesterone-induced prolactin secretion in monkeys. *Neuroendocrinology* 65: 335–343, 1997.
73. Pitkin RM. Calcium metabolism in pregnancy: a review. *Am J Obstet Gynecol* 121: 724–737, 1975.
74. Popelar J, Groh D, Mazelova J, Syka J. Cochlear function in young and adult Fischer 344 rats. *Hear Res* 186: 75–84, 2003.
75. Popelar J, Groh D, Pelanova J, Canlon B, Syka J. Age-related changes in cochlear and brainstem auditory functions in Fischer 344 rats. *Neurobiol Aging* 27: 490–500, 2006.
76. Riggs BL, Melton LJ 3rd. Involutional osteoporosis. *N Engl J Med* 314: 1676–1686, 1986.
77. Rolandi E, Magnani G, Sannia A, Barreca T. Evaluation of Prl secretion in elderly subjects. *Acta Endocrinol* 100: 351–357, 1982.
78. Royster M, Driscoll P, Kelly PA, Freemarm M. The prolactin receptor in the fetal rat: cellular localization of messenger ribonucleic acid, immunoreactive protein, and ligand-binding activity and induction of expression in late gestation. *Endocrinology* 136: 3892–3900, 1995.
79. Rydmarker S, Horner KC. Atrophy of outer hair cell stereocilia and hearing loss in hydropic cochleae. *Hear Res* 53: 113–122, 1991.
80. Rydmarker S, Horner KC. Morphological changes of hair cell stereocilia and tectorial membrane in guinea pigs with experimentally induced hydrops. *Scanning Microsc* 4: 705–713, 1990.
81. Sawin CT, Carlson HE, Geller A, Castelli WP, Bacharach P. Serum prolactin and aging: basal values and changes with estrogen use and hypothyroidism. *J Gerontol* 44: M131–M135, 1989.
82. Schaller B, Kirsch E, Tolnay M, Mindermann T. Symptomatic granular cell tumor of the pituitary gland: case report and review of the literature. *Neurosurgery* 42: 166–170, 1998.
83. Sellari-Franceschini S, Ravecca F, De Vito A, Berrettini S. [Progressive sensorineural hearing loss in cochlear otosclerosis.] *Acta Otorhinolaryngol Ital* 18: 59–65, 1998.
84. Shea JJ Jr, Ge X, Orchik DJ. Endolymphatic hydrops associated with otosclerosis. *Am J Otol* 15: 348–357, 1994.
85. Sherman BE, Chole RA. Effect of chemical sympathectomy with 6-hydroxydopamine on osteoclast activity in the gerbilline middle ear bulla. *Otol Neurotol* 22: 237–241, 2001.
86. Sherman BE, Chole RA. Effect of pharmacological sympathectomy on osteoclastic activity in the gerbilline auditory bulla in vivo. *Ann Otol Rhinol Laryngol* 108: 1078–1087, 1999.
87. Sismanis A, Hughes GB, Abedi E. Coexisting otosclerosis and Meniere's disease: a diagnostic and therapeutic dilemma. *Laryngoscope* 96: 9–13, 1986.
88. Srivastava S, Matsuda M, Hou Z, Bailey JP, Kitazawa R, Herbst MP, Horseman ND. Receptor activator of NF- κ B ligand induction via Jak2 and Stat5a in mammary epithelial cells. *J Biol Chem* 278: 46171–46178, 2003.
89. Steger RW. Age related changes in the control of prolactin secretion in the female rat. *Neurobiol Aging* 2: 119–123, 1981.
90. Stenberg AE, Wang H, Fish J, Schrott-Fischer A 3rd, Sahlin L, Hultcrantz M. Estrogen receptors in the normal adult and developing human inner ear and in Turner's syndrome. *Hear Res* 157: 87–92, 2001.
91. Stenberg AE, Wang H, Sahlin L, Hultcrantz M. Mapping of estrogen receptors alpha and beta in the inner ear of mouse and rat. *Hear Res* 136: 29–34, 1999.
92. Strachan D. Sudden sensorineural deafness and hormone replacement therapy. *J Laryngol Otol* 110: 1148–1150, 1996.
93. Thompson SK, Zhu X, Frisina RD. Estrogen blockade reduces auditory feedback in CBA mice. *Otolaryngol Head Neck Surg* 135: 100–105, 2006.
94. Toverud SU, Harper C, Munson PL. Calcium metabolism during lactation: enhanced effects of thyrocalcitonin. *Endocrinology* 99: 371–378, 1976.
95. Vekemans M, Robyn C. Influence of age on serum prolactin levels in women and men. *Br Med J* 4: 738–739, 1975.
96. Vessey M, Painter R. Oral contraception and ear disease: findings in a large cohort study. *Contraception* 63: 61–63, 2001.
97. Williams RF, Gianfortoni JG, Hodgen GD. Hyperprolactinemia induced by an estrogen-progesterone synergy: quantitative and temporal effects of estrogen priming in monkeys. *J Clin Endocrinol Metab* 60: 126–132, 1985.
98. Zehnder AF, Kristiansen AG, Adams JC, Merchant SN, McKenna MJ. Osteoprotegerin in the inner ear may inhibit bone remodeling in the otic capsule. *Laryngoscope* 115: 172–177, 2005.