



**HAL**  
open science

## Interactive effects of high irradiance and moderate heat on photosynthesis, pigments, and tocopherol in the tree-fern *Dicksonia antarctica*

Liubov Volkova, Michael Tausz, Lauren T. Bennett, Erwin Dreyer

### ► To cite this version:

Liubov Volkova, Michael Tausz, Lauren T. Bennett, Erwin Dreyer. Interactive effects of high irradiance and moderate heat on photosynthesis, pigments, and tocopherol in the tree-fern *Dicksonia antarctica*. *Functional Plant Biology*, 2009, 36 (12), pp.1046-1056. 10.1071/FP09098 . hal-02658657

**HAL Id: hal-02658657**

**<https://hal.inrae.fr/hal-02658657>**

Submitted on 30 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Interactive effects of high irradiance and moderate heat on photosynthesis, pigments, and tocopherol in the tree-fern *Dicksonia antarctica*

Liubov Volkova<sup>A,C</sup>, Michael Tausz<sup>A</sup>, Lauren T. Bennett<sup>A</sup> and Erwin Dreyer<sup>A,B</sup>

<sup>A</sup>Department of Forest and Ecosystem Science, Melbourne School of Land and Environment, The University of Melbourne, Water Street, Creswick, Vic. 3363, Australia.

<sup>B</sup>INRA, Nancy-Université, UMR 1137 'Ecologie et Ecophysiologie Forestières', F-54280 Champenoux, France.

<sup>C</sup>Corresponding author. Email: l.volkova@pgrad.unimelb.edu.au

**Abstract.** Effects of high irradiance and moderate heat on photosynthesis of the tree-fern *Dicksonia antarctica* (Labill., Dicksoniaceae) were examined in a climate chamber under two contrasting irradiance regimes (900 and 170  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) and three sequential temperature treatments (15°C; 35°C; back to 15°C). High irradiance led to decline in predawn quantum yield of photochemistry,  $F_v/F_m$  (0.73), maximal Rubisco activity ( $V_{\text{cmax}}$ ; from 37 to 29  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ), and electron transport capacity ( $J_{\text{max}}$ ; from 115 to 67  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). Temperature increase to 35°C resulted in further decreases in  $F_v/F_m$  (0.45) and in chlorophyll bleaching of high irradiance plants, while  $V_{\text{cmax}}$  and  $J_{\text{max}}$  were not affected. Critical temperature for thylakoid stability ( $T_c$ ) of *D. antarctica* was comparable with other higher plants (c. 47°C), and increases of  $T_c$  with air temperature were greater in high irradiance plants. Increased  $T_c$  was not associated with accumulation of osmotica or zeaxanthin formation. High irradiance increased the xanthophyll cycle pigment pool (V+A+Z, 91 v. 48  $\text{mmol mol}^{-1}$  chlorophyll<sup>-1</sup>), de-epoxidation state (56% v. 4%), and  $\alpha$ -tocopherol. Temperature increase to 35°C had no effect on V+A+Z and de-epoxidation state in both light regimes, while lutein,  $\beta$ -carotene and  $\alpha$ -tocopherols increased, potentially contributing to increased membrane stability under high irradiance.

**Additional keywords:**  $\alpha$ -tocopherols,  $\beta$ -carotene, critical temperature, de-epoxidation state,  $J_{\text{max}}$ , light regime, lutein, osmolality, photoinhibition,  $V_{\text{cmax}}$ , temperature, xanthophyll pigments.

## Introduction

While understorey species of evergreen forests often experience high intensity sunflecks, they are not usually exposed to prolonged periods of high irradiance (Lovelock *et al.* 1998; Tausz *et al.* 2005). A protective canopy usually creates a favourable microclimate with more moderate temperature fluctuations and greater air humidity than in the above-canopy atmosphere. However, during the course of forest ecosystem dynamics including gap formation, bushfires, or anthropogenic management such as forest harvesting, understorey species may be suddenly exposed to full sunlight and high temperatures, stress factors that can contribute to temporary decline of these species. According to climate-change projections, these factors are likely to become even more significant in the future, as temperatures are predicted to increase, and disturbances in forest canopies may become more frequent (Hennessy *et al.* 2007).

In the short-term, exposure of shade-acclimated plants to high levels of irradiance often leads to photoinhibition and photo-oxidative stress. Photoinhibition alone is rarely responsible for plant mortality and the plant may recover and become fully acclimated. Photo-oxidative stress is caused by the toxic effects of reactive oxygen species (ROS) produced in the photosynthetic apparatus under high irradiance when carbon

assimilation is light-saturated (Niyogi 2000). Many plants can, to a certain extent, acclimate to increased irradiance through enhanced dissipation of absorbed light energy in the thylakoids, a process related to the conversion of the light harvesting xanthophyll violaxanthin to the energy quenching zeaxanthin (Demmig-Adams and Adams 2006). Protection against high irradiance can also involve the accumulation of tocopherol (Munné-Bosch 2005), an antioxidant that scavenges toxic ROS and contributes to thylakoid membrane stability.

When, in addition to high irradiance, leaves are exposed to other environmental stress factors such as high temperature, there can be sustained reductions in the efficiency of photosynthetic energy conversion and inhibition of repairs to photodamaged photosystem II (PSII; Murata *et al.* 2007). Photosynthesis is particularly sensitive to inhibition by heat stress due to labile components in the photosynthetic apparatus (Salvucci and Crafts-Brandner 2004). The thylakoid membrane is one of the main temperature stress targets and changes during acclimation occur at that level (Ducruet *et al.* 2007). The degree of thermostability of the thylakoids can be estimated by the critical temperature  $T_c$  – the temperature threshold above which irreversible damage occurs to PSII (Schreiber and Berry 1977).  $T_c$  changes with growing conditions, reflecting thermal acclimation of the

photosynthetic apparatus (Ducruet *et al.* 2007). Chlorophyll fluorescence yield is a sensitive indicator of the state of thylakoids, and can be used to assess  $T_c$  in plants as the temperature threshold above which ground level fluorescence ( $F_o$ ) increases (e.g. Froux *et al.* 2004). The mechanisms underlying acclimatory changes in  $T_c$  are still poorly understood, although some results point towards stabilising effects of protective compounds on thylakoids. For example, the xanthophyll zeaxanthin (Havaux and Gruszecki 1993; Havaux and Tardy 1996), as well as increased soluble sugar concentration (Hüve *et al.* 2006), are believed to have a stabilising effect and shift  $T_c$  towards higher temperatures.

Our model understorey species, the tree-fern *Dicksonia antarctica* (Labill., Dicksoniaceae), is known to decline after clear-fell logging in Victoria, Australia (Ough and Murphy 2004). These tree-ferns are iconic and ecologically significant understorey species in many humid forest types in the southern hemisphere, including Australian temperate rain forests and wet sclerophyll (eucalypt) forests (Large and Braggins 2004). They support a large epiphytic diversity on their trunks and provide nursery sites for many tree and shrub species as well as nesting and feeding sites for marsupials, insects and birds (Magrath and Lill 1983; Lindenmayer *et al.* 1994; Roberts *et al.* 2005). Decline in *D. antarctica* numbers is expected to negatively impact on many dependent species and thus maintenance of tree-ferns is often an objective of forest management plans (Department of Natural Resources and Environment 2002).

The reasons for poor survival and ongoing decline of *D. antarctica* after logging remain uncertain (Ough and Murphy 2004), but exposure to high irradiance, combined with increased air and frond temperatures, could be contributing factors. Periodically disturbed by wildfires in their natural habitat, *D. antarctica* are exposed to a broad range of irradiance during their lifetime (Hunt *et al.* 2002), which suggests that this species is able to at least partly acclimate to different levels of irradiance. Certainly, other studies indicate potential for fern acclimation to different light regimes. For example, New Zealand ferns from contrasting habitats displayed contrasting characteristics in terms of photosynthetic light compensation point, which were tightly correlated with specific frond area (Bannister and Wildish 1982). Frond characteristics (frond surface area, epidermis thickness, palisade/spongy mesophyll ratio, blade size, petiole length) of a South American *Cyathea* species (another important tree fern genus) were also correlated with its local irradiance (Arens 1997). However, other studies suggest limited capacity of shade-acclimated tree-ferns to efficiently adjust to increased irradiance (Durand and Goldstein 2001). To our knowledge, only a few studies have examined effects of high temperature, either alone or with high irradiance on the physiological performance of tree-ferns: Tingey *et al.* (1987) found that photosynthesis of *D. antarctica* was particularly susceptible to inhibition with increasing temperature and high light; and Nobel *et al.* (1984) also mentioned negative effects of high temperature on gas exchange of ferns. Natural distribution of *D. antarctica* in Australia is limited to the temperate zone (McCarthy 1998), characterised by cool to warm conditions (Köppen classification; Australian Bureau of Meteorology, [http://www.bom.gov.au/iwk/climate\\_zones](http://www.bom.gov.au/iwk/climate_zones), verified 13 August

2009), perhaps indicating that the species has limited potential for acclimation to temperature increases (such as after clear-fell logging, but potentially also due to climate change), making it susceptible to ongoing decline.

In this study, we investigated the responses of *D. antarctica* to high irradiance, moderately high temperature (+35°C), and a combination of both under fully controlled climate chamber conditions. Measuring  $T_c$  together with several variables related to photosynthesis, chlorophyll fluorescence, and chloroplast pigments, we addressed the following specific questions:

- (1) Are photosynthetic parameters of *D. antarctica* adversely affected by (a) high irradiance; (b) high temperatures; and (c) their interactions?
- (2) Does membrane stability (measured via critical temperature,  $T_c$ ) increase in *D. antarctica* fronds with increased temperature (indicative of an acclimation to high temperature), and if yes, are  $T_c$  changes associated with accumulation of osmotica or zeaxanthin formation?
- (3) Do other potentially protective thylakoid compounds, such as carotenoids and tocopherol, change in relation to high irradiance and high temperature?
- (4) Are effects of high temperature on the above parameters reversible?

## Materials and methods

### Plant material

One-year-old sporophytes of *Dicksonia antarctica* Labill (HSK Gardening and Leisure Avon Dassett, UK) were transplanted into 10-L pots, containing a mixture of sand and peat (50/50, v/v) and 40 g slow release fertiliser (Nutricote 100, Chisso-Asahi Fertilizer Co. Ltd, Tokyo, Japan; N/P/K, %, 13/13/13) per pot. The plants were grown under uniform sunlit conditions in a naturally illuminated glasshouse at INRA, Champenoux, France (48°44'N, 6°14'E) for 2 months in spring 2007. At the end of this period, the plants were transferred to a climate chamber (Chambre Phytotronique STRADER, Angers, France) at Champenoux.

### Climate chamber conditions and experimental design

Irradiance in the climate chamber was provided by two types of 400 W lamps (HQI Philips (mercury halide) and SONT Philips (sodium halide), Koninklijke Philips Electronics N.V., Eindhoven, The Netherlands) and resulted in a photosynthetic photon flux density (PPFD) of 900  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  at plant height (PAR range, 400–700 nm, measured with a Li-Cor quantum sensor; Li-Cor, Lincoln, Nebraska, USA). Relative humidity was 70–80%, air temperature was controlled to  $\pm 0.5^\circ\text{C}$  (see temperature treatments below), and photoperiod was 16 h  $\text{day}^{-1}$ .

The tree-ferns were randomly assigned to two experimental groups ( $n = 7$  in each). One group was shielded by a wavelength-neutral shade mesh (17% light transmission, PPFD: 170  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) – ‘shade’, the other exposed to full light (900  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) – ‘high irradiance’. Such levels are representative of irradiance conditions in the open on overcast winter and clear summer days respectively, at typical *D. antarctica* field sites in mountain-ash forests of Victoria,

Australia. Plants were kept well watered at all times and were rotated daily at random within their designated irradiance regime.

A sequence of temperature treatments was applied as follows:

- (1) 10 days at 15°C day and night;
  - (1a) 3 days at 25°C day and night;
- (2) 12 days at 35°C/25°C day/night (typical hot summer days for *D. antarctica* in the field);
- (3) 10 days at 15°C day and night (to check reversibility of temperature effects).

Chlorophyll *a* fluorescence and critical frond temperature were measured for each individual every 1 to 2 days throughout the experiment. Photosynthesis was recorded from net CO<sub>2</sub> uptake (*A*) v. intercellular CO<sub>2</sub> concentration (*A*-*C<sub>i</sub>* curves) on the first day of the experiment and at the end of each temperature treatment. Samples for nitrogen (N) content, xanthophyll analyses, and osmolality were taken at the end of each of the three temperature treatments. All measurements were made on the mid-third of the youngest fully-expanded fronds that were of healthy appearance (i.e. not discoloured).

Total chlorophyll content was measured for each individual every 1 to 2 days throughout the experiment; measurements were made over the entire plant irrespective of frond condition.

#### *Frond temperature (T<sub>frond</sub>)*

Frond temperature (as 10 random points across the entire plant) was measured twice during the 35°C temperature treatment (beginning/end) at predawn and midday, and once at the end of the experiment (during the second 15°C temperature treatment), using an IR laser thermometer (Raynger PM, Raytech Inc., Santa Cruz, CA, USA).

#### *Maximal quantum yield of photochemistry (F<sub>v</sub>/F<sub>m</sub>)*

F<sub>v</sub>/F<sub>m</sub> was derived from chlorophyll *a* fluorescence measured on dark-acclimated fronds (at the end of the 'night' period) with a modulated fluorometer (PAM 2000, Heinz Walz GmbH, Effeltrich, Germany). Maximum quantum yield of PSII was estimated as F<sub>v</sub>/F<sub>m</sub> = (F<sub>m</sub> - F<sub>o</sub>)/F<sub>m</sub>, after Maxwell and Johnson (2000).

#### *Gas exchange measurements*

Gas exchange was measured with a Li-Cor 6400 portable photosynthesis measurement device, equipped with a 2 × 3 cm broadleaf chamber with red-blue LEDs (Li-Cor). All gas exchange measurements were conducted at the reference frond temperature of 25°C. For each plant, an *A*-*C<sub>i</sub>* curve was generated at PPFD 1000 μmol m<sup>-2</sup> s<sup>-1</sup>, frond temperature 25°C, air flow rate 400 μmol air s<sup>-1</sup>, and relative humidity (RH) >60%. Photosynthesis was induced at a CO<sub>2</sub> mole fraction of 50 μmol mol<sup>-1</sup> for 15–20 min before measurements to ensure maximal stomatal opening and maximal activity of Calvin cycle enzymes. CO<sub>2</sub> mole fraction was then increased in 13 successive steps to 2200 μmol mol<sup>-1</sup> with two measurements at each step. After finishing the *A*-*C<sub>i</sub>* curve, illumination in the leaf chamber was turned off, CO<sub>2</sub> mole fraction was decreased to 400 μmol mol<sup>-1</sup> and respiration rate was recorded after 5 min in the dark. The frond area enclosed in the Li-Cor chamber was

marked, photographed and calculated using imaging software (UTHSCSA Image Tool Version 3, University of Texas, USA). Values for *A*<sub>max</sub> and *g<sub>s</sub>* at ambient CO<sub>2</sub> (400 μmol mol<sup>-1</sup>) were derived from these curves.

Using a biochemical photosynthetic model (Farquhar *et al.* 1980), apparent (i.e. assuming that mesophyll conductance to CO<sub>2</sub>, *g<sub>i</sub>*, is infinite) maximum carboxylation rate (apparent *V<sub>cmax</sub>*<sup>25</sup>), and the maximum apparent rate of electron transport (apparent *J<sub>max</sub>*<sup>25</sup>) were estimated by fitting the *A*-*C<sub>i</sub>* curves to the model as described in Montpied *et al.* (2009). Triose phosphate use (TPU) limitation was not included in the model; points with decreasing *A* at high CO<sub>2</sub> mole fractions were disregarded. A set of primary parameters of Rubisco kinetic properties used herein, *K<sub>c</sub>* = 327 μmol mol<sup>-1</sup>, *K<sub>o</sub>* = 282 600 μmol mol<sup>-1</sup>, *Γ\** = 43.7 μmol mol<sup>-1</sup>, were taken from von Caemmerer *et al.* (1994).

Values of *g<sub>i</sub>* were then estimated with the curve fitting approach introduced by Ethier and Livingston (2004) and described by Montpied *et al.* (2009) and real *V<sub>cmax</sub>*<sup>25</sup> and *J<sub>max</sub>*<sup>25</sup> were computed based on the *A*-*C<sub>c</sub>* (chloroplastic CO<sub>2</sub> mole fraction); therefore, only corrected *V<sub>cmax</sub>* and *J<sub>max</sub>* (i.e. under the hypothesis of finite *g<sub>i</sub>*) are given and discussed in this study.

#### *Frond nitrogen and chlorophyll content*

Frond samples were analysed for nitrogen (N) content using an elemental analyser (NCS 2500, CE instrument Thermo Quest, Milano, Italy). Samples were dried at 60°C for 48 h (to determine dry weight) and then ground to a fine powder. Frond area of fresh samples was scanned and calculated using Scion Image software (Scion Corporation 2000–2001, USA), and these data used to calculate N content on a frond area basis.

Frond chlorophyll content was estimated from transmittance values measured with the Minolta SPAD-502 chlorophyll meter (Minolta, Illinois, USA; hereafter 'SPAD'). Values were the mean of two to three separate pinnules per plant (randomly selected irrespective of frond colour).

Total chlorophyll (*a* + *b*) was also measured by HPLC (see Pigments and tocopherol determination).

#### *Critical temperature (T<sub>c</sub>)*

Critical temperature was estimated *in vivo* from the sharp rise of basal chlorophyll *a* fluorescence under increasing temperature (Schreiber and Berry 1977). Disks of tree-fern pinnules were placed into a temperature-controlled aluminium body, with the fibre-optics of the fluorometer (PAM 2000) pointing at the sample. Ground fluorescence (*F<sub>o</sub>*) was induced with a red diode at low PPFD of ~1 μmol m<sup>-2</sup> s<sup>-1</sup>. Temperature of the aluminium body was increased gradually (1°C min<sup>-1</sup>) from 20°C to 60°C. *F<sub>o</sub>* was continuously recorded and critical temperature (*T<sub>c</sub>*) was estimated graphically at the beginning of the heat-induced fluorescence rise (Froux *et al.* 2004).

#### *Total tissue osmolality*

Total tissue osmolality was measured using freeze-point depression from hot water extracts of dried frond tissue (Callister *et al.* 2006). Approximately 60 mg of dried ground frond tissue (as prepared for N analysis) was placed in a 2-mL polypropylene vial to which 1.6 mL of deionised water was

added. The samples were placed in a water bath at 90°C for 60 min. The samples were left to cool to room temperature, centrifuged at 10 000g for 2 min and 1 mL of the supernatant was transferred to a 1.7-mL polypropylene vial. Osmolality of the solution was measured using an OSMOMAT 030 cryoscopic osmometer (Gonotec, Berlin, Germany).

#### Pigments and tocopherol determination

Froned discs (3.75 mm diameter) were collected at midday at the end of each of the three temperature treatments and immediately frozen in liquid nitrogen. Samples were freeze-dried, sealed with silicagel in airtight plastic bags, and kept at -20°C until analysis.

Four discs per plant were ground in a Matrix Mill (Retsch MM301, Germany) at the temperature of liquid nitrogen. To avoid the presence of traces of acid in the acetone used for the extraction, 0.5 g L<sup>-1</sup> of calcium carbonate were added to samples before grinding (García-Plazaola and Becerril 1999). The resulting powder was extracted with 0.5 mL of ice-cold acetone, homogenised and centrifuged at 4°C for 1 min at 15 000g. The pellets were re-extracted as described above to a combined sample volume of 1 mL. Extracts were stored in sealed vials at -20°C until analysis. Prior to injection, samples were centrifuged at 4°C for 20 min at 15 000 g and the clean supernatant was transferred into HPLC vials.

HPLC separation of chloroplast pigments and tocopherols was according to the methods given in Tausz *et al.* (2003). Chromatographic conditions were:

- Pigments: 25 × 4.6 mm Spherisorb ODS 25 µm column. gradient: solvent A: acetonitrile: methanol: water = 100 : 5 : 10 (v/v/v), solvent B: ethylacetate: acetone = 1 : 2 (v/v), 10% B to 70% B in 17 min, hold at 70% B for 5 min, return to 10% B in 5 min. Flow rate 1 mL min<sup>-1</sup>. The injection volume was 20 µL, photometric detection at 440 nm.
- α-Tocopherol: 25 × 4.6 mm Spherisorb ODS 25 µm column. Solvent 100% methanol isocratic. Flow rate 1 mL min<sup>-1</sup>. Injection volume was 20 µL. Fluorescence detection excitation 295 nm, emission 325 nm.
- Acetone, acetonitrile, methanol and ethyl acetate were of HPLC grade and water was deionised. A standard of α-tocopherol was purchased from Sigma (Sigma-Aldrich, Castle Hill, NSW, Australia); standards for carotenoids and chlorophyll *a* and *b* were prepared as follows: several generic extracts were prepared in 100% acetone and measured at

three wavelengths in the spectrophotometer (Varian UV/V 300, USA). Using the equations of Lichtenthaler (1987), chlorophyll *a*, *b* and total carotenoid concentrations were calculated at a spectrophotometer resolution range of 1–4 nm. The same extracts were then re-run in the HPLC and conversion factors for chlorophyll *a*, *b* and total carotenoids were calculated, disregarding the minor differences in carotenoid absorption coefficients at the wavelength in question.

#### Statistical analysis

Repeated-measures models of SPSS 15 (SPSS Inc., Chicago, USA) were used for statistical analysis, with irradiance as the between-subject factor and temperature as the within-subject factor (both fixed). Effects of irradiance (high irradiance, shade), and 3 levels of temperature (15°C, 35°C, back to 15°C), and irradiance by temperature interactions on each dependent variable were analysed. Data for statistical analyses were the values per individual plant at the end of each temperature treatment. Photosynthetic parameters, measured before the start of the experiment (i.e.  $A_{max}$ ,  $g_s$ ,  $V_{cmax}$  and  $J_{max}$ ) were not used in the model.

#### Results

##### Froned temperature ( $T_{frond}$ )

$T_{frond}$  did not differ among plants at the beginning of the experiment (data not shown). After the temperature increased to 35°C,  $T_{frond}$  was similar in shaded and high irradiance plants at predawn (below 25°C) but differed on average by 1.5°C at midday (around 35°C v. 33.5°C; Table 1). By the end of the treatment at 35°C,  $T_{frond}$  of shaded plants was on average 0.7°C cooler at predawn, and 3.3°C cooler at midday (Table 1).

##### Maximum quantum yield of PS II ( $F_v/F_m$ ) and photosynthetic capacity parameters


Predawn  $F_v/F_m$  remained close to the optimum value of 0.83 in shaded plants across all temperature treatments. In contrast,  $F_v/F_m$  declined after the first day of exposure to high irradiance (Fig. 1a). The 35°C treatment resulted in further decreases of  $F_v/F_m$  (Fig. 1a). After return to 15°C, a partial recovery of  $F_v/F_m$  was detected.

Photosynthetic parameters were comparable among all plants at the start of the experiment (Table 2, 'before'). Ten days of high irradiance (at 15°C) resulted in decreases in both light-saturated

**Table 1.** Temperature of *Dicksonia antarctica* fronds ( $T_{frond}$ ) exposed to high irradiance and under shade

Values are means ± s.e. ( $n = 7$  plants, 10 measurements per plant) at predawn and midday on the first (day 1) and last (day 12) days of two temperature treatments (35°C and back to 15°C). *P*-values indicate significance of difference between high irradiance and shaded plants within temperature treatments (Student's *t*-test)

	35°C			<i>P</i> -value	Back to 15°C			<i>P</i> -value
	High irradiance	Shaded	Difference		High irradiance	Shaded	Difference	
	<i>Predawn</i>							
Day 1	24.3 ± 0.1	24.2 ± 0.1	0.1	0.30	15.3 ± 0.0	16.0 ± 0.0	-0.7	<0.001
Day 12	23.0 ± 0.2	22.3 ± 0.2	0.7	0.02				
	<i>Midday</i>							
Day 1	34.9 ± 0.6	33.4 ± 0.3	1.5	0.02	19.7 ± 0.4	17.2 ± 0.4	2.5	<0.001
Day 12	34.6 ± 0.2	31.3 ± 0.5	3.3	0.002				



**Fig. 1.** Time course of (a) maximum quantum efficiency of PSII ( $F_v/F_m$ ) recorded after 8 h of darkness (predawn); and (b) chlorophyll content in SPAD units of high irradiance (open symbols) and shaded (closed symbols) *Dicksonia antarctica* during three successive temperature treatments (delineated by dotted lines). Values are means of  $n=7$  (error bars indicate s.e.).  $P$ -values indicate significance of effects of irradiance (I), temperature (T), and irradiance by temperature interaction (I  $\times$  T).

rate of net photosynthesis at a reference temperature of 25°C,  $A_{\max}$ , and corresponding stomatal conductance  $g_s$  (Table 2; Fig. 2). Increasing the temperature to 35°C induced stomatal opening in all plants (Table 2). However, while  $A_{\max}$  of all shaded plants also increased (Table 2; Fig. 2),  $A_{\max}$  was less responsive in high irradiance plants (Fig. 2). Nonetheless, all changes in  $g_s$  and  $A_{\max}$  were fully reversible upon return to 15°C (Table 2).

When measured at a reference temperature of 25°C, maximal carboxylation rate,  $V_{\max}^{25}$ , and maximal light-driven electron flux  $J_{\max}^{25}$  decreased in response to increased irradiance. Both were insensitive to temperature treatments regardless of irradiance regime (Table 2). Mesophyll conductance to CO<sub>2</sub>,  $g_i$ , was highly variable, probably owing to the low accuracy of the fitting procedure used. As a result, no significant effects of irradiance and temperature could be detected (Table 2).

Nitrogen content per frond area (g m<sup>-2</sup>) was not affected by either irradiance or temperature (Table 2). Photosynthetic nitrogen use efficiency (PNUE), calculated as  $V_{\max}/N$ , was lower under high irradiance than in shade and remained such until the end of the experiment. Changes in temperature did not affect PNUE (Table 2).

Total chlorophyll content (in SPAD units) decreased at 35°C under high irradiance and remained low thereafter (Fig. 1b). A similar albeit not significant effect was detected for chlorophyll  $a + b$  in HPLC extracts (Table 2). It should be noted that visually damaged fronds were avoided for HPLC analyses, while SPAD measurements were made across entire fronds. Chlorophyll  $a/b$  ratios were similar between irradiance regimes during the 15°C temperature treatment. With temperature increase to 35°C, chl  $a/b$  ratios decreased in high irradiance plants in contrast to a significant increase in shaded plants (Table 2). With temperature return to 15°C, chl  $a/b$  ratios of high irradiance plants recovered to the initial values.

#### Critical temperature ( $T_c$ )

During the 15°C treatment, critical temperature for photochemistry ( $T_c$ ) was similar under shade and high irradiance (means of 47.5°C and 47.2°C, respectively; Fig. 3). Increase in temperature resulted in significant rises of  $T_c$  that were greatest under high irradiance.  $T_c$  started to decrease after return to 15°C, although pretreatment values were not reached by the end of the experiment (Fig. 3).

#### Total tissue osmolality

Total tissue osmolality was not affected by irradiance but decreased significantly during the 35°C treatment (Table 3). It increased to close to the original values after return to 15°C.

#### Carotenoids and $\alpha$ -tocopherol

Neoxanthin and lutein contents (per mol total chlorophyll) were significantly greater under high irradiance, whereas  $\alpha$ - and  $\beta$ -carotene contents remained comparable to shaded plants (Table 3). Temperature increase to 35°C led to significant increase in  $\beta$ -carotene and lutein in high irradiance plants. These pigments tended to remain high on return to 15°C, although  $\alpha$ -carotene significantly decreased under high irradiance (Table 3).


The xanthophyll cycle pigment pool (i.e. violaxanthin, antheraxanthin, and zeaxanthin, V+A+Z) was significantly greater under high irradiance for the whole experiment. Changes in temperature did not affect V+A+Z under either irradiance regime (Table 3). The de-epoxidation state of xanthophylls (expressed as  $(0.5A+Z)/(V+A+Z)$ ) was greater under high irradiance than shade (on average 44% v. 3%; Table 3). It decreased by c. 32% under high irradiance upon return to 15°C, but was insensitive to temperature under shade. Small amounts of lutein-epoxide were detected (Table 3), but they were not affected by light, temperature and their interactions.

Content of  $\alpha$ -tocopherol was significantly greater under high irradiance than shade (Fig. 4). Whereas  $\alpha$ -tocopherol was invariant to temperature changes in shaded plants,  $\alpha$ -tocopherol increased in high irradiance plants with temperature increase to 35°C, and remained high after the temperature returned to 15°C; due to the lack of a high light-low temperature control, we cannot clearly separate the high irradiance and high temperature effects on these changes.

**Table 2. Photosynthesis and frond traits of *Dicksonia antarctica* exposed to high irradiance and under shade before and during three successive temperature treatments**

Values are means  $\pm$  s.e. ( $n=7$ ) of  $A_{\max}^{25}$  – light-saturated net  $\text{CO}_2$  assimilation rate at  $25^\circ\text{C}$ ;  $g_s$ , stomatal conductance under saturating irradiance at  $25^\circ\text{C}$ ;  $N_A$ , nitrogen content on a frond area basis; PNUE, photosynthetic nitrogen use efficiency ( $V_{\text{cmax}}/N_A$ ) at  $25^\circ\text{C}$ ; Chl total, total chlorophyll content in SPAD units and on a frond area basis; Chl  $a/b$ , chlorophyll  $a/b$  ratio;  $V_{\text{cmax}}^{25}$ , maximal carboxylation rate of Rubisco at  $25^\circ\text{C}$ ;  $J_{\max}^{25}$ , maximal light driven electron flux at  $25^\circ\text{C}$ ;  $g_i$ , mesophyll conductance to  $\text{CO}_2$  measured at  $25^\circ\text{C}$ ; Abbreviations: I, Irradiance; T, temperature;  $I \times T$ , irradiance by temperature interaction; n.s., not-significant; n.d., no data

Parameter	Irradiance regime	Temperature treatments				Significance of effects ( $P$ )		
		Before	$15^\circ\text{C}$	$35^\circ\text{C}$	Back to $15^\circ\text{C}$	I	T	$I \times T$
$A_{\max}^{25}$ ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ )	High irradiance	$6.0 \pm 0.5$	$3.7 \pm 0.5$	$5.1 \pm 0.6$	$3.9 \pm 0.4$	<0.001	0.02	n.s. (0.3)
	Shade	$6.4 \pm 0.4$	$6.4 \pm 0.8$	$7.6 \pm 0.3$	$5.6 \pm 0.5$			
$g_s$ at $A_{\max}^{25}$ ( $\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$ )	High irradiance	$82 \pm 11$	$57 \pm 6$	$108 \pm 7$	$50 \pm 6$	0.02	<0.001	n.s. (0.3)
	Shade	$80 \pm 7$	$80 \pm 14$	$152 \pm 12$	$70 \pm 5$			
$V_{\text{cmax}}^{25}$ ( $\mu\text{mol m}^{-2} \text{ s}^{-1}$ )	High irradiance	$37.0 \pm 2$	$29.3 \pm 2.7$	$23.2 \pm 2.8$	$29.1 \pm 3.3$	0.001	n.s. (0.7)	
	Shade	$36.0 \pm 1$	$37.1 \pm 2.9$	$40.6 \pm 3.8$	$34.5 \pm 4.2$			
$J_{\max}^{25}$ ( $\mu\text{mol m}^{-2} \text{ s}^{-1}$ )	High irradiance	$115 \pm 6$	$67.8 \pm 8.7$	$50.2 \pm 7.3$	$84.7 \pm 10.8$	0.01	n.s. (0.1)	
	Shade	$105 \pm 6$	$84.6 \pm 10.2$	$85.6 \pm 11.7$	$97.7 \pm 9.8$			
$g_i$ ( $\text{mmol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ )	High irradiance	$155 \pm 64$	$141 \pm 53$	$143 \pm 115$	$115 \pm 25$	n.s. (0.2)	n.s. (0.8)	
	Shade	$115 \pm 35$	$222 \pm 63$	$294 \pm 195$	$259 \pm 112$			
$N_A$ ( $\text{g m}^{-2}$ )	High irradiance	n.d.	$16.9 \pm 1.0$	$15.9 \pm 0.6$	$17.4 \pm 0.7$	n.s. (0.9)	n.s. (0.4)	
	Shade	n.d.	$16.5 \pm 0.7$	$16.0 \pm 0.8$	$15.5 \pm 2.1$			
PNUE ( $V_{\text{cmax}}/N_A$ ) ( $\mu\text{mol mol}^{-1} \text{ N}^{-1} \text{ m}^{-2}$ )	High irradiance	n.d.	$204 \pm 21$	$201 \pm 24$	$175 \pm 13$	0.01	n.s. (0.4)	
	Shade	n.d.	$278 \pm 18$	$311 \pm 30$	$300 \pm 41$			
Chl total (SPAD units)	High irradiance	n.d.	$49.1 \pm 1.7$	$39.3 \pm 2.3$	$34.1 \pm 2.1$	0.002	<0.001	0.001
	Shade	n.d.	$48.9 \pm 0.4$	$47.1 \pm 0.5$	$47.3 \pm 1.0$			
Chl total ( $a + b$ ) ( $\mu\text{mol m}^{-2}$ )	High irradiance	n.d.	$782 \pm 142$	$721 \pm 111$	$675 \pm 89.6$	n.s. (0.2)	n.s. (0.8)	
	Shade	n.d.	$810 \pm 62.7$	$914 \pm 127$	$843 \pm 81.1$			
Chl $a/b$	High irradiance	n.d.	$2.36 \pm 0.06$	$2.29 \pm 0.05$	$2.41 \pm 0.09$	0.002	0.005	<0.001
	Shade	n.d.	$2.46 \pm 0.04$	$2.71 \pm 0.05$	$2.66 \pm 0.03$			



**Fig. 2.** Stomatal conductance ( $g_s$ ) at  $A_{\max}$  v. light-saturated rate of net photosynthesis ( $A_{\max}$ ) for high irradiance and shaded *Dicksonia antarctica* measured at a standardised temperature of  $25^\circ\text{C}$  during three temperature treatments:  $15^\circ\text{C}$  (open circle),  $35^\circ\text{C}$  (closed diamond) and back to  $15^\circ\text{C}$  (open triangle). Values are means of  $n=6$  (high irradiance) and  $n=7$  (shaded).


#### Correlations between $T_c$ and biochemical parameters

No significant correlation was detected between  $T_c$  and osmolality, tocopherol, and  $\alpha$ - and  $\beta$ -carotenes (correlations not shown). Correlations between  $T_c$  and zeaxanthin, presented as  $Z/(Z+V)$  (after Havaux and Gruszecki 1993) were also not significant, either for each irradiance treatment separately or for the combined data (Fig. 5).

#### Discussion

##### Effect of high irradiance, high temperature and their interaction on photosynthetic capacity parameters of *D. antarctica*

Photosynthetic capacity of *Dicksonia antarctica* in this study was within ranges reported in the literature. Maximum light saturated rates of net photosynthesis ( $A_{\max}^{25}$ ) were comparable with values



**Fig. 3.** Time course of critical temperature ( $T_c$ ) of high irradiance (open symbols) and shaded (closed symbols) *Dicksonia antarctica* across the experiment. Values are means of  $n=7$  ( $\pm$ s.e.).  $P$ -values indicate significance of effects of irradiance (I), temperature (T), and irradiance by temperature interaction (I  $\times$  T).

reported for the same species:  $6 - 10.8 \mu\text{mol m}^{-2} \text{s}^{-1}$  (Nobel *et al.* 1984) and  $8.3 \mu\text{mol m}^{-2} \text{s}^{-1}$  (Hunt *et al.* 2002). Maximum carboxylation rates,  $V_{\text{cmax}}^{25}$  (the maximal *in vivo* Rubisco activity), and the maximum rate of electron transport,  $J_{\text{max}}$ , at a reference temperature of  $25^\circ\text{C}$  were comparable to some shade tolerant tree species, such as silver fir (*Abies alba* Mill; Robakowski *et al.* 2002), and mesophyll conductance,  $g_i$ , corresponded to typical values of evergreen trees among the species reviewed by Ethier and Livingston (2004).

Photosynthetic capacity of *D. antarctica* was adversely affected by high irradiance.  $J_{\text{max}}^{25}$  decreased with respect to shaded plants, and this decrease was paralleled by a decrease

of maximum quantum yield of PSII,  $F_v/F_m$ , indicating a moderate but chronic photoinhibition (Table 2; Tallon and Quiles 2007). High irradiance also led to decreases in  $A_{\text{max}}^{25}$ , due to both reduced stomatal conductance  $g_s$  and decreased photosynthetic capacity, i.e. Rubisco activity and  $J_{\text{max}}^{25}$ . Several processes may result in deactivation of Rubisco – for example, remobilisation and export of nitrogen from the leaves, interruption in the electron transport chain, or the presence of reactive oxygen species (ROS). In our case, deactivation of Rubisco in high irradiance plants was not related to a remobilisation of frond nitrogen – the N content per frond area was not affected by irradiance. This can be seen also through the decline of photosynthetic nitrogen use efficiency (PNUE). However, high irradiance alone did not affect total chlorophyll content and chlorophyll *a/b* ratios, indicating that photoprotective mechanisms were efficient enough to avoid chlorophyll degradation.

High temperature had no negative effects on photosynthetic capacity of *D. antarctica* under shade. Increasing temperature to  $35^\circ\text{C}$  even stimulated  $A_{\text{max}}^{25}$ . Increases in  $A_{\text{max}}^{25}$  were in this case solely due to increases in  $g_s$ , as we found no effects of temperature on photosynthetic capacity ( $V_{\text{cmax}}^{25}$ ,  $J_{\text{max}}^{25}$ ). The increase in chlorophyll *a/b* ratio in shaded plants with increasing temperature perhaps indicates temperature-stimulated resynthesis of photosynthetic reaction centres relative to light-harvesting antenna complexes, which commonly coincides with other changes in pigment composition, e.g. lutein,  $\beta$ -carotene (Haldimann 1999).


The interactive effect of high irradiance and high temperature led to severe photoinhibition in agreement with earlier findings (Berry and Björkman 1980). Temperature increase stimulated  $g_s$  yet without commensurate increases in  $A_{\text{max}}^{25}$ , indicating that metabolic limitations (e.g. Rubisco activity) governed  $A_{\text{max}}^{25}$  (consistent with the findings of e.g. Law and Crafts-Brandner 1999; on Rubisco activation). Many studies have shown a

**Table 3.** Pigment content and osmolality of *Dicksonia antarctica* fronds exposed to high irradiance and under shade during three successive temperature treatments


Values are means  $\pm$  s.e. ( $n=7$ ) of: osmolality; carotenoids (on a chlorophyll basis): lutein, neoxanthin and xanthophyll pool (i.e. Violanxanthin, Antheraxanthin, Zeaxanthin, V+A+Z),  $\alpha$ - and  $\beta$ -carotene, lutein-epoxide and the de-epoxidation state of violanxanthin =  $(0.5A+Z)/(V+A+Z)$ . Abbreviations: I, Irradiance; T, temperature; I  $\times$  T, irradiance by temperature interaction; n.s., not significant

	Irradiance regime	Temperature treatments			Significance of effects ( $P$ )		
		15°C	35°C	Back to 15°C	I	T	I $\times$ T
Osmolality (mosmol g d.w. <sup>-1</sup> )	High irradiance	1.56 $\pm$ 0.06	1.33 $\pm$ 0.04	1.45 $\pm$ 0.05	n.s. (0.4)	<0.001	
	Shade	1.55 $\pm$ 0.02	1.40 $\pm$ 0.03	1.51 $\pm$ 0.01			
V+A+Z (mmol mol <sup>-1</sup> chl <sup>-1</sup> )	High irradiance	91.9 $\pm$ 9.2	118.1 $\pm$ 15.0	113.7 $\pm$ 15.9	<0.001	n.s. (0.2)	
	Shade	47.8 $\pm$ 1.7	45.8 $\pm$ 1.2	45.5 $\pm$ 1.2			
De-epoxidation (%)	High irradiance	56.6 $\pm$ 3.4	53.1 $\pm$ 7.7	21.4 $\pm$ 4.7	<0.001	<0.001	0.001
	Shade	4.4 $\pm$ 1.1	2.3 $\pm$ 0.8	1.6 $\pm$ 0.6			
Lutein (mmol mol <sup>-1</sup> chl <sup>-1</sup> )	High irradiance	263 $\pm$ 9.0	339 $\pm$ 25.4	388 $\pm$ 28.5	<0.001	0.002	0.008
	Shade	185 $\pm$ 3.9	186 $\pm$ 3.3	198 $\pm$ 5.9			
Neoxanthin (mmol mol <sup>-1</sup> chl <sup>-1</sup> )	High irradiance	61.3 $\pm$ 2.6	65.9 $\pm$ 3.8	69.3 $\pm$ 3.6	<0.001	n.s. (0.06)	
	Shade	45.2 $\pm$ 1.9	46.2 $\pm$ 1.1	48.4 $\pm$ 2.0			
$\alpha$ -carotene (mmol mol <sup>-1</sup> chl <sup>-1</sup> )	High irradiance	11.1 $\pm$ 2.3	24.2 $\pm$ 3.4	5.8 $\pm$ 1.5	0.03	<0.001	0.003
	Shade	16.1 $\pm$ 2.6	24.1 $\pm$ 2.4	20.9 $\pm$ 1.8			
$\beta$ -carotene (mmol mol <sup>-1</sup> chl <sup>-1</sup> )	High irradiance	62.2 $\pm$ 7.8	94.6 $\pm$ 8.7	89.3 $\pm$ 7.0	0.001	0.005	n.s. (0.4)
	Shade	44.7 $\pm$ 5.7	59.4 $\pm$ 5.3	59.3 $\pm$ 3.3			
Lutein-epoxide (mmol mol <sup>-1</sup> chl <sup>-1</sup> )	High irradiance	5.2 $\pm$ 1.7	6.3 $\pm$ 1.4	6.2 $\pm$ 2.2	n.s. (0.1)	n.s. (0.2)	
	Shade	3.4 $\pm$ 0.4	3.9 $\pm$ 0.4	4.3 $\pm$ 0.3			





**Fig. 4.**  $\alpha$ -Tocopherol content relative to total chlorophylls of high irradiance (open bars) and shaded (closed bars) *Dicksonia antarctica* under three temperature treatments. Values are means of  $n=7$  ( $\pm$ s.e.).  $P$ -values indicate significance of effects of irradiance (I), temperature (T), and irradiance by temperature interaction ( $I \times T$ ).



**Fig. 5.** Critical temperature ( $T_c$ ) v. xanthophyll zeaxanthin (expressed as  $Z/(Z+V)$ ; after Havaux and Gruszecki 1993) of high irradiance (open symbols) and shaded (closed symbols) *Dicksonia antarctica* during three temperature treatments: 15°C (circle), 35°C (diamond) and back to 15°C (triangle). Each point represents an individual measurement. Relationships were also non-significant when data were separated by irradiance treatment. Z, zeaxanthin; V, violaxanthin.

negative effect of moderate heat and photoinhibition on the activation of Rubisco mediated by an activase (for details, see Salvucci and Crafts-Brandner 2004). In our study, we did not measure the activity of Rubisco-activase, but found no increase in photosynthetic capacity to increased temperatures. Prolonged photoinhibition and heat interaction resulted in decreases in chlorophylls as in numerous other studies (e.g. Lambers *et al.* 2008). Decreases in chlorophyll  $a/b$  ratio indicated that under prolonged light stress, chlorophylls were destabilised with Chl  $a$  being more sensitive than Chl  $b$  (Yamamoto *et al.* 2008).

A return of temperature to 15°C induced stomatal closure thus reversing  $A_{\max}^{25}$  to the initial values at 15°C, as found by Ghoul *et al.* (2003). The only partial recovery of  $F_v/F_m$  in high irradiance plants demonstrated detrimental interactive effects of high irradiance and temperature on  $F_v/F_m$ , suggesting that degradation processes (i.e. bleaching of chlorophylls or photodegradation of thylakoid complexes) did not allow rapid recovery of  $F_v/F_m$  (Ottander *et al.* 1995). The fact that  $V_{\max}$  and  $J_{\max}$  remained non-responsive possibly underlined the high sensitivity of Rubisco to high irradiance in *D. antarctica*.

In summary, photosynthetic capacity and photosynthetic nitrogen use efficiency were rapidly affected by exposure to high irradiance under 15°C, while chlorophylls and predawn  $F_v/F_m$  declined further under the combination of high irradiance and high temperature.

#### Membrane stability of *D. antarctica* measured via critical temperature

Critical temperature,  $T_c$ , recorded in *D. antarctica* was *c.* 47°C, comparable with a range of overstorey species, such as *Quercus petraea* Matt. Liebl. (46.7°C; Dreyer *et al.* 2001). Comparable data for other tree-ferns species are currently lacking.

An increase in air temperature induced an increase in  $T_c$ , as found by many authors (e.g. Dreyer *et al.* 2001). This increase in  $T_c$  was larger under high irradiance than shade, coinciding with a 3.3°C higher midday frond temperature under high irradiance. Previously published data suggest that an increase in  $T_c$  by 1°C (as found in our study) requires an increase in ambient temperatures of  $\sim 10^\circ\text{C}$  (Froux *et al.* 2004). We therefore believe that the larger rise in  $T_c$  in high irradiance plants was not solely caused by the difference in frond temperature, but directly related to high irradiance effects, which resulted in enhanced thermostability of thylakoid membranes.

Contrary to findings by Hüve *et al.* (2006), increased thermostability of the thylakoid membranes was not associated with the accumulation of osmotically active substances. In our study, osmotically active solutes even decreased when temperature increased. It may be argued that the hot water extract method results in artefacts, because some cell wall or other material can be brought into solution as a result of the grinding and extraction procedures (Callister *et al.* 2006). Yet this method is widely used (e.g. Merchant *et al.* 2006) and moreover, a tight correlation was found among methods even though the absolute results were different (Callister *et al.* 2006). Our observation therefore seems reliable.

With subsequent temperature decrease to 15°C, we found a tendency for  $T_c$  to return to the initial values, even though this return was not complete. The observed lag confirms the hysteresis found by Froux *et al.* (2004), when the increase in  $T_c$  with increasing temperature is faster than relaxation from this effect after temperature decrease.

#### Xanthophyll cycle carotenoids, pigments and $\alpha$ -tocopherol

Exposure of *D. antarctica* to high irradiance alone resulted in almost 2-fold increase in xanthophyll cycle carotenoids ( $V+A+Z$ ). A high de-epoxidation state was also recorded

(50% v. 2–4% in the shade). Values for V+A+Z were within the range of values presented for other species (Thayer and Björkman 1990). The maximum de-epoxidation state of the xanthophyll cycle was lower than maximum values measured in epiphytic ferns (Tausz *et al.* 2001). An increase was also recorded in pools of other carotenoids such as neoxanthin and lutein in response to high irradiance. Values were within the range reported for other ferns, e.g. a range of epiphytic species (Tausz *et al.* 2001) or the tree-fern *Cyathea microdonta* (Desv.) Domin (Matsubara *et al.* 2009). These carotenoids (e.g. neoxanthin) may preserve PSII from photo-inactivation and protect membrane lipids from photo-oxidation by ROS (North *et al.* 2007). The observed increase in lutein (located primarily in both the proximal and distal light harvesting centres of PSI and PSII) is probably associated with the acclimation of antennae to increasing irradiance (Senger *et al.* 1993). As antenna size is usually reduced in response to increasing irradiance, it is also likely that an increasing fraction of these carotenoids is not bound to antenna proteins. High irradiance also induced increases in  $\alpha$ -tocopherols, known for their protective function in thylakoids, consistent with findings elsewhere (e.g. García-Plazaola and Becerril 1999; Munné-Bosch 2005).

Increased temperature stimulated an increase in  $T_c$  in shaded plants but without simultaneous accumulation of zeaxanthin, which is contrary to observations by Havaux and Gruszecki (1993) and Havaux and Tardy (1996). Irrespective of an increased  $T_c$ , high temperature had no effect on concentration of  $\alpha$ -tocopherol in shaded plants. According to our results, it is therefore unlikely that increased membrane stability as measured by  $T_c$  is directly and generally dependent on zeaxanthin or  $\alpha$ -tocopherol. There was some coincidence of increased  $T_c$  with increases in  $\alpha$ - and  $\beta$ -carotenes in shaded plants, which may have reflected changes in carotenoid synthesis rates. Carotenes may also contribute to an overall increase in membrane stability, as proposed by Tausz *et al.* (2001).

Interactions between irradiance and temperature had no additional effect on the xanthophyll cycle pool and de-epoxidation state of high irradiance plants. In contrast to V+A+Z, the combination of high irradiance with high temperature led to further increases in the amount of lutein and  $\beta$ -carotene, which remained greater until the end of experiment. It is not clear from our experiment whether sustained concentrations of carotenoids indicated their role as a last resort in membrane photoprotection under extreme stress, or simply reflected their superior stability under such conditions. In contrast to shaded plants, temperature increase stimulated an almost 2-fold increase in  $\alpha$ -tocopherol in high irradiance plants. Although these results for high irradiance plants appear to support findings of Llusà *et al.* (2005) – who suggested that increased tolerance to high temperatures might be at least partly due to an increase in  $\alpha$ -tocopherol – our results for shaded plants indicate that changes in  $T_c$  can occur independently of changes in  $\alpha$ -tocopherol.

Temperature return to 15°C did not affect total V+A+Z pool of high irradiance plants, but the de-epoxidation state decreased by 30% to remain significantly higher than in shaded plants. De-epoxidation of the xanthophyll cycle is driven by an acidic thylakoid lumen, which can be the consequence of an imbalance between electron transport and electron consumption (Demmig-

Adams and Adams 2006). It seems that heat related changes in the photosynthetic apparatus lead to a relaxation of the pH gradient upon temperature decrease, despite the continuation of high irradiance. This may be related to a sustained decrease in light use efficiency as suggested by persistently low  $F_v/F_m$  values, and other, as yet unexplained, changes in the photosynthetic membrane. Such further changes were also expressed in the observed change in  $T_c$ , possibly in combination with an increased electron consumption rate upon relaxation of the high temperature. Temperature decrease did not affect concentration of carotenoids except for  $\alpha$ -carotene. Significant decreases in  $\alpha$ -carotene in high irradiance plants can be explained in terms of its ease of oxidation to lutein under conditions of oxidative stress (Senger *et al.* 1993), or its conversion to  $\beta$ -carotene to increase scavenging of free radicals in core complexes under conditions of stress (Kirchgeßner *et al.* 2003).

In summary, increased thylakoid stability in *D. antarctica* observed during our experiment could not be explained by any of the measured changes in pigments or  $\alpha$ -tocopherol, although they may all play partial roles. Alternative or additional explanations may involve presence of certain heat shock proteins or changes in the composition of membrane lipids, as suggested by Sinsawat *et al.* (2004). Discrepancies with earlier literature may be related to the fact that pigment changes are fast responses to temperature increases (in the order of 1 day; e.g. as in Havaux and Tardy 1996), while we investigated longer term acclimation (12 days) to high temperature. Longer term acclimation of plants to rising temperatures is also related to the appearance of polar lipids with saturated fatty acids causing a decrease in membrane fluidity (Zsófi *et al.* 2009). In parallel with an increased threshold temperature for thermal inactivation of PSII (Downton *et al.* 1984), this can increase thylakoid stability. The difference between the rate of acclimation and de-acclimation supports this hypothesis: the fast rise may be initially due to rapid changes in pigments that are later completed by slower changes in lipid composition. The reversal of these changes may be slower, which would be the cause for a slow return to initial levels of stability. However, our data do not allow confirmation of this speculation.

## Conclusion

High irradiance caused chronic photoinhibition (measured as sustained decrease in maximum PSII quantum efficiency), and decreases in all photosynthetic capacity parameters of *D. antarctica*. Whilst we observed some acclimation in terms of increases in protective carotenoids, which may have sufficed to avoid chlorophyll degradation, similar or even decreasing chlorophyll *a/b* ratios indicated limited short-term acclimation potential of *D. antarctica* fronds to high irradiance. Temperature alone appeared to have no negative effect on photosynthesis of *D. antarctica*, possibly suggesting that this species can thrive under moderately high temperature as long as it is shaded. However, there was ample evidence of severe damage to the photosynthetic apparatus when high irradiance was combined with moderately high temperatures. Therefore we can speculate that future scenarios predicting higher temperatures and more frequent disturbances may decrease the competitiveness of *D. antarctica*.

## Acknowledgements

Liubov Volkova acknowledges support by a Melbourne Research Scholarship and by an S.F. Pond Trust Travelling Scholarship for 2006. The work was in part supported by a research contract with the Victorian Department of Sustainability and Environment and by an internal grant of the University of Melbourne to M Tausz. Thanks to Andrew Merchant for useful hints on the experimental design, to all the kind and helpful staff of INRA Nancy, in particular to Christophe Bailly for ensuring smooth operation of the climate chamber during the experiment, Jean Marie Gioria for growing the tree-ferns, and to Jacqueline Marchand for N quantification in the fronds. Damien L. Callahan, School of Botany, University of Melbourne, is personally thanked for help with operating the HPLC.

## References

- Arens NC (1997) Responses of leaf anatomy to light environment in the tree fern *Cyathea caracasana* (Cyatheaceae) and its application to some ancient seed ferns. *Palaios* **12**, 84–94. doi: 10.2307/3515296
- Bannister P, Wildish KL (1982) Light compensation points and specific leaf areas in some New-Zealand ferns. *New Zealand Journal of Botany* **20**, 421–424.
- Berry J, Björkman O (1980) Photosynthetic response and adaptation to temperature in higher plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **31**, 491–543.
- Callister AN, Arndt SK, Adams MA (2006) Comparison of four methods for measuring osmotic potential of tree leaves. *Physiologia Plantarum* **127**, 383–392. doi: 10.1111/j.1399-3054.2006.00652.x
- Demmig-Adams B, Adams WW III (2006) Photoprotection in an ecological context: the remarkable complexity of thermal energy dissipation. *New Phytologist* **172**, 11–21. doi: 10.1111/j.1469-8137.2006.01835.x
- Department of Natural Resources and Environment (2002) ‘Victorian tree fern management plan.’ (Department of Natural Resources and Environment: East Melbourne, Australia)
- Downton WJS, Berry JA, Seemann JR (1984) Tolerance of photosynthesis to high temperature in desert plants. *Plant Physiology* **74**, 786–790. doi: 10.1104/pp.74.4.786
- Dreyer E, Le Roux X, Montpied P, Daudet FA, Masson F (2001) Temperature response of leaf photosynthetic capacity in seedlings from seven temperate tree species. *Tree Physiology* **21**, 223–232.
- Ducruet JM, Peeva V, Havaux M (2007) Chlorophyll thermofluorescence and thermoluminescence as complementary tools for the study of temperature stress in plants. *Photosynthesis Research* **93**, 159–171. doi: 10.1007/s11120-007-9132-x
- Durand LZ, Goldstein G (2001) Photosynthesis, photoinhibition, and nitrogen use efficiency in native and invasive tree ferns in Hawaii. *Oecologia* **126**, 345–354. doi: 10.1007/s004420000535
- Ethier GJ, Livingston NJ (2004) On the need to incorporate sensitivity to CO<sub>2</sub> transfer conductance into the Farquhar–von Caemmerer–Berry leaf photosynthesis model. *Plant, Cell & Environment* **27**, 137–153. doi: 10.1111/j.1365-3040.2004.01140.x
- Farquhar GD, von Caemmerer S, Berry JA (1980) A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species. *Planta* **149**, 78–90. doi: 10.1007/BF00386231
- Froux F, Ducrey M, Epron D, Dreyer E (2004) Seasonal variations and acclimation potential of the thermostability of photochemistry in four Mediterranean conifers. *Annals of Forest Science* **61**, 235–241. doi: 10.1051/forest:2004016
- García-Plazaola JI, Becerril JM (1999) A rapid high performance liquid chromatography method to measure lipophilic antioxidants in stressed plants: simultaneous determination of carotenoids and tocopherols. *Phytochemical Analysis* **10**, 307–313. doi: 10.1002/(SICI)1099-1565(199911/12)10:6<307::AID-PCA477>3.0.CO;2-L
- Ghoul H, Montpied P, Epron D, Ksontini M, Hanchi B, Dreyer E (2003) Thermal optima of photosynthetic functions and thermostability of photochemistry in cork oak seedlings. *Tree Physiology* **23**, 1031–1039.
- Haldimann P (1999) How do changes in temperature during growth affect leaf pigment composition and photosynthesis in *Zea mays* genotypes differing in sensitivity to low temperature? *Journal of Experimental Botany* **50**, 543–550. doi: 10.1093/jexbot/50.333.543
- Havaux M, Gruszecki WI (1993) Heat-induced and light-induced chlorophyll a fluorescence changes in potato leaves containing high or low levels of the carotenoid zeaxanthin – indications of a regulatory effect of zeaxanthin on thylakoid membrane fluidity. *Photochemistry and Photobiology* **58**, 607–614. doi: 10.1111/j.1751-1097.1993.tb04940.x
- Havaux M, Tardy F (1996) Temperature-dependent adjustment of the thermal stability of photosystem II *in vivo*: Possible involvement of xanthophyll cycle pigments. *Planta* **198**, 324–333. doi: 10.1007/BF00620047
- Hennessy KJ, Fitzharris B, Bates BC, Harvey N, Howden SM, Hughes L, Salinger J, Warrick R (2007) Australia and New Zealand. In ‘Climate change 2007: impacts, adaptation and vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change’. (Eds ML Parry, OF Canziani, JP Palutikof, PJ van der Linden, CE Hanson) pp. 507–540. (Cambridge University Press: Cambridge, UK)
- Hunt MA, Davidson NJ, Unwin GL, Close DC (2002) Ecophysiology of the soft tree fern *Dicksonia antarctica* Labill. *Austral Ecology* **27**, 360–368. doi: 10.1046/j.1442-9993.2002.01190.x
- Hüve K, Bichele I, Tobias M, Niinemets Ü (2006) Heat sensitivity of photosynthetic electron transport varies during the day due to changes in sugars and osmotic potential. *Plant, Cell & Environment* **29**, 212–228. doi: 10.1111/j.1365-3040.2005.01414.x
- Kirchgeßner HD, Reichert K, Hauff K, Steinbrecher R, Schnitzler JP, Pfündel EE (2003) Light and temperature, but not UV radiation, affect chlorophylls and carotenoids in Norway spruce needles (*Picea abies* (L.) Karst.). *Plant, Cell & Environment* **26**, 1169–1179. doi: 10.1046/j.1365-3040.2003.01043.x
- Lambers H, Chapin FS, Pons TL (2008) ‘Plant physiological ecology.’ (Springer: New York)
- Large MF, Braggins JE (2004) ‘Tree ferns.’ (CSIRO Publishing: Collingwood, Vic.)
- Law RD, Crafts-Brandner SJ (1999) Inhibition and acclimation of photosynthesis to heat stress is closely correlated with activation of ribulose-1,5-bisphosphate carboxylase/oxygenase. *Plant Physiology* **120**, 173–182. doi: 10.1104/pp.120.1.173
- Lichtenthaler HK (1987) Chlorophylls and carotenoids – pigments of photosynthetic biomembranes. *Methods in Enzymology* **148**, 350–382. doi: 10.1016/0076-6879(87)48036-1
- Lindenmayer DB, Cunningham RB, Donnelly CF, Triggs BJ, Belvedere M (1994) The conservation of arboreal marsupials in the montane ash forests of the central highlands of Victoria, South-Eastern Australia. 5. Patterns of use and the microhabitat requirements of the mountain brushtail possum *Trichosurus caninus* ogilby in retained linear habitats (wildlife corridors). *Biological Conservation* **68**, 43–51. doi: 10.1016/0006-3207(94)90545-2
- Llusià J, Peñuelas J, Munné-Bosch S (2005) Sustained accumulation of methyl salicylate alters antioxidant protection and reduces tolerance of holm oak to heat stress. *Physiologia Plantarum* **124**, 353–361. doi: 10.1111/j.1399-3054.2005.00519.x
- Lovelock CE, Kursar TA, Skillman JB, Winter K (1998) Photoinhibition in tropical forest understorey species with short- and long-lived leaves. *Functional Ecology* **12**, 553–560. doi: 10.1046/j.1365-2435.1998.00235.x
- Magrath R, Lill A (1983) The use of time and energy by the crimson rosella in a temperate wet forest in winter. *Australian Journal of Zoology* **31**, 903–912. doi: 10.1071/ZO9830903

- Matsubara S, Krause GH, Aranda J, Virgo A, Beisel KG, Jahns P, Winter K (2009) Sun-shade patterns of leaf carotenoid composition in 86 species of neotropical forest plants. *Functional Plant Biology* **36**, 20–36. doi: 10.1071/FP08214
- Maxwell K, Johnson GN (2000) Chlorophyll fluorescence – a practical guide. *Journal of Experimental Botany* **51**, 659–668. doi: 10.1093/jexbot/51.345.659
- McCarthy PM (1998) 'Ferns, gymnosperms and allied groups.' (CSIRO Publishing: Melbourne, Australia)
- Merchant A, Tausz M, Arndt SK, Adams MA (2006) Cyclitols and carbohydrates in leaves and roots of 13 *Eucalyptus* species suggest contrasting physiological responses to water deficit. *Plant, Cell & Environment* **29**, 2017–2029. doi: 10.1111/j.1365-3040.2006.01577.x
- Montpied P, Granier A, Dreyer E (2009) Seasonal time-course of gradients of photosynthetic capacity and mesophyll conductance to CO<sub>2</sub> across a beech (*Fagus sylvatica* L.) canopy. *Journal of Experimental Botany* **60**, 2407–2418. doi: 10.1093/jxb/erp093
- Munné-Bosch S (2005) The role of  $\alpha$ -tocopherol in plant stress tolerance. *Journal of Plant Physiology* **162**, 743–748. doi: 10.1016/j.jplph.2005.04.022
- Murata N, Takahashi S, Nishiyama Y, Allakhverdiev SI (2007) Photoinhibition of photosystem II under environmental stress. *Biochimica Et Biophysica Acta – Bioenergetics* **1767**, 414–421. doi: 10.1016/j.bbabi.2006.11.019
- Niyogi KK (2000) Safety valves for photosynthesis. *Current Opinion in Plant Biology* **3**, 455–460. doi: 10.1016/S1369-5266(00)00113-8
- Nobel PS, Calkin HW, Gibson AC (1984) Influences of PAR, temperature and water vapor concentration on gas exchange by ferns. *Physiologia Plantarum* **62**, 527–534. doi: 10.1111/j.1399-3054.1984.tb02794.x
- North HM, De Almeida A, Boutin JP, Frey A, To A, Botran L, Sotta B, Marion-Poll A (2007) The *Arabidopsis* ABA-deficient mutant *aba4* demonstrates that the major route for stress-induced ABA accumulation is via neoxanthin isomers. *The Plant Journal* **50**, 810–824. doi: 10.1111/j.1365-313X.2007.03094.x
- Ottander C, Campbell D, Öquist G (1995) Seasonal changes in photosystem II organization and pigment composition in *Pinus sylvestris*. *Planta* **197**, 176–183. doi: 10.1007/BF00239954
- Ough K, Murphy A (2004) Decline in tree-fern abundance after clearfell harvesting. *Forest Ecology and Management* **199**, 153–163.
- Robakowski P, Montpied P, Dreyer E (2002) Temperature response of photosynthesis of silver fir (*Abies alba* Mill.) seedlings. *Annals of Forest Science* **59**, 163–170. doi: 10.1051/forest:2002003
- Roberts NR, Dalton PJ, Jordan GJ (2005) Epiphytic ferns and bryophytes of Tasmanian tree-ferns: a comparison of diversity and composition between two host species. *Austral Ecology* **30**, 146–154. doi: 10.1111/j.1442-9993.2005.01440.x
- Salvucci ME, Crafts-Brandner SJ (2004) Mechanism for deactivation of Rubisco under moderate heat stress. *Physiologia Plantarum* **122**, 513–519. doi: 10.1111/j.1399-3054.2004.00419.x
- Schreiber U, Berry JA (1977) Heat-induced changes of chlorophyll fluorescence in intact leaves correlated with damage of photosynthetic apparatus. *Planta* **136**, 233–238. doi: 10.1007/BF00385990
- Senger H, Wagner C, Hermsmeier D, Hohl N, Urbig T, Bishop NI (1993) The influence of light intensity and wavelength on the contents of  $\alpha$ -carotene and  $\beta$ -carotene and their xanthophylls in green algae. *Journal of Photochemistry and Photobiology. B, Biology* **18**, 273–279. doi: 10.1016/1011-1344(93)80075-K
- Sinsawat V, Leipner J, Stamp P, Fracheboud Y (2004) Effect of heat stress on the photosynthetic apparatus in maize (*Zea mays* L.) grown at control or high temperature. *Environmental and Experimental Botany* **52**, 123–129. doi: 10.1016/j.envexpbot.2004.01.010
- Tallon C, Quiles MJ (2007) Acclimation to heat and high light intensity during the development of oat leaves increases the NADH DH complex and PTOX levels in chloroplasts. *Plant Science* **173**, 438–445. doi: 10.1016/j.plantsci.2007.07.001
- Tausz M, Hietz P, Briones O (2001) The significance of carotenoids and tocopherols in photoprotection of seven epiphytic fern species of a Mexican cloud forest. *Australian Journal of Plant Physiology* **28**, 775–783. doi: 10.1071/PP01068
- Tausz M, Wonsch A, Grill D, Morales D, Jimenez MS (2003) Measuring antioxidants in tree species in the natural environment: from sampling to data evaluation. *Journal of Experimental Botany* **54**, 1505–1510. doi: 10.1093/jxb/erg175
- Tausz M, Warren CR, Adams MA (2005) Dynamic light use and protection from excess light in upper canopy and coppice leaves of *Nothofagus cunninghamii* in an old growth, cool temperate rainforest in Victoria, Australia. *New Phytologist* **165**, 143–156. doi: 10.1111/j.1469-8137.2004.01232.x
- Thayer SS, Björkman O (1990) Leaf xanthophyll content and composition in sun and shade determined by HPLC. *Photosynthesis Research* **23**, 331–343. doi: 10.1007/BF00034864
- Tingey DT, Evans RC, Bates EH, Gumpertz ML (1987) Isoprene emissions and photosynthesis in 3 ferns – the influence of light and temperature. *Physiologia Plantarum* **69**, 609–616. doi: 10.1111/j.1399-3054.1987.tb01974.x
- von Caemmerer S, Evans JR, Hudson GS, Andrews TJ (1994) The kinetics of ribulose-1,5-bisphosphate carboxylase/oxygenase *in vivo* inferred from measurements of photosynthesis in leaves of transgenic tobacco. *Planta* **195**, 88–97.
- Yamamoto Y, Aminaka R, Yoshioka M, Khatoon M, Komayama K, *et al.* (2008) Quality control of photosystem II: impact of light and heat stresses. *Photosynthesis Research* **98**, 589–608. doi: 10.1007/s11120-008-9372-4
- Zsófi Z, Váradi G, Bálo B, Marschall M, Nagy Z, Dulai S (2009) Heat acclimation of grapevine leaf photosynthesis: mezo- and macroclimatic aspects. *Functional Plant Biology* **36**, 310–322. doi: 10.1071/FP08200

Manuscript received 5 May 2009, accepted 31 July 2009