



**HAL**  
open science

## SNP discovery and linkage map construction in cultivated tomato

Kenta Shirasawa, Sachiko Isobe, Hideki Hirakawa, Erika Asamizu, Hiroyuki Fukuoka, Daniel Just, Christophe Rothan, Shigemi Sasamoto, Tsunakazu Fujishiro, Yoshie Kishida, et al.

► **To cite this version:**

Kenta Shirasawa, Sachiko Isobe, Hideki Hirakawa, Erika Asamizu, Hiroyuki Fukuoka, et al.. SNP discovery and linkage map construction in cultivated tomato. *DNA Research*, 2010, 17 (6), pp.381-91. 10.1093/dnares/dsq024 . hal-02659276

**HAL Id: hal-02659276**

**<https://hal.inrae.fr/hal-02659276>**

Submitted on 30 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## SNP Discovery and Linkage Map Construction in Cultivated Tomato

KENTA Shirasawa<sup>1,\*</sup>, SACHIKO Isobe<sup>1</sup>, HIDEKI Hirakawa<sup>1</sup>, ERIKA Asamizu<sup>1,†</sup>, HIROYUKI Fukuoka<sup>2</sup>, DANIEL Just<sup>3</sup>, CHRISTOPHE Rothan<sup>3</sup>, SHIGEMI Sasamoto<sup>1</sup>, TSUNAKAZU Fujishiro<sup>1</sup>, YOSHIE Kishida<sup>1</sup>, MITSUYO Kohara<sup>1</sup>, HISANO Tsuruoka<sup>1</sup>, TSUYUKO Wada<sup>1</sup>, YASUKAZU Nakamura<sup>1,‡</sup>, SHUSEI Sato<sup>1</sup>, and SATOSHI Tabata<sup>1</sup>

Kazusa DNA Research Institute, 2-6-7 Kazusa-Kamatari, Kisarazu, Chiba 292-0818, Japan<sup>1</sup>; National Institute of Vegetable and Tea Science (NIVTS), National Agriculture and Food Research Organization, 360 Kusawa, Ano, Tsu, Mie 514-2392, Japan<sup>2</sup> and Unité Mixte de Recherche 619, Biologie du Fruit, Institut National de la Recherche Agronomique (INRA Bordeaux), Universités de Bordeaux, 33883 Villenave d'Ornon cedex, France<sup>3</sup>

\*To whom correspondence should be addressed. Tel. +81 438-52-3935. Fax. +81 438-52-3934.  
Email: shirasaw@kazusa.or.jp

Edited by Doil Choi

(Received 14 July 2010; accepted 21 September 2010; published online 2 November 2010)

### Abstract

**Few intraspecific genetic linkage maps have been reported for cultivated tomato, mainly because genetic diversity within *Solanum lycopersicum* is much less than that between tomato species. Single nucleotide polymorphisms (SNPs), the most abundant source of genomic variation, are the most promising source of polymorphisms for the construction of linkage maps for closely related intraspecific lines. In this study, we developed SNP markers based on expressed sequence tags for the construction of intraspecific linkage maps in tomato. Out of the 5607 SNP positions detected through *in silico* analysis, 1536 were selected for high-throughput genotyping of two mapping populations derived from crosses between 'Micro-Tom' and either 'Ailsa Craig' or 'M82'. A total of 1137 markers, including 793 out of the 1338 successfully genotyped SNPs, along with 344 simple sequence repeat and intronic polymorphism markers, were mapped onto two linkage maps, which covered 1467.8 and 1422.7 cM, respectively. The SNP markers developed were then screened against cultivated tomato lines in order to estimate the transferability of these SNPs to other breeding materials. The molecular markers and linkage maps represent a milestone in the genomics and genetics, and are the first step toward molecular breeding of cultivated tomato. Information on the DNA markers, linkage maps, and SNP genotypes for these tomato lines is available at <http://www.kazusa.or.jp/tomato/>.**

**Keywords:** DNA marker; linkage map; single nucleotide polymorphism; *Solanum lycopersicum*; tomato

### 1. Introduction

Genetics in tomato (*Solanum lycopersicum*) and its wild relatives, including *S. chilense*, *S. habrochaites*, *S. pimpinellifolium*, and *S. pennellii*, have been greatly

advanced since molecular markers have become available.<sup>1</sup> During the past two decades, several genetic maps in tomato have been reported, with a total of more than 2000 loci detected by restriction fragment length polymorphism (RFLP), amplified fragment length polymorphism (AFLP), cleaved amplified polymorphic sequence (CAPS), and simple sequence repeat (SSR) markers based on the mapping of populations derived from crosses between tomato and related wild species.<sup>2–6</sup> Recently, 1282 novel SSR markers and 151 intronic polymorphic markers were mapped onto an

† Present address: Gene Research Center, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.

‡ Present address: Center for Information Biology and DNA Data Bank of Japan, National Institute of Genetics, Research Organization for Information and Systems, Yata, Mishima, Shizuoka 411-8510, Japan.

interspecific map, 'Tomato-EXPEN 2000' derived from a cross between *S. lycopersicum* and *S. pennellii*.<sup>7</sup> Such efforts have resulted in the identification of a number of quantitative trait loci (QTLs) and genes for fruit morphology,<sup>8–11</sup> disease resistance,<sup>12–15</sup> and other agronomical traits.<sup>16</sup> The identified genes, e.g. *Cf-4*, *Tm-2*, and *Sw-5*, have already been used for tomato breeding through advanced-backcross and introgression-line strategies using molecular markers.<sup>1</sup> Though significant advances in molecular genetics and breeding have been reported in tomato, most of them were based on interspecific crosses because genetic diversity in the cultivated tomato is lower than in its wild relatives.<sup>17</sup> Meanwhile, intraspecific maps are required to identify QTLs for agronomically important traits, which are the targets of practical breeding programs. However, only one intraspecific map, based on AFLP, RFLP, and random amplified polymorphic DNA (RAPD) markers, has been reported for *S. lycopersicum*.<sup>18</sup>

Single nucleotide polymorphisms (SNPs) are the most abundant source of variation in the genome for both intragenic and intergenic regions. They therefore represent a valuable basis for the development of molecular markers for identification of polymorphisms among closely related lines. Previous studies have suggested that DNA markers developed from intergenic regions tend to cluster in heterochromatic portions of chromosomes, while those derived from genic regions disperse along entire chromosomes.<sup>7,19–22</sup> Therefore, SNPs, especially those located in intragenic regions, are expected to distribute randomly along the whole genome. In addition, novel techniques based on the DNA microarray method allow high-throughput SNP genotyping.<sup>23</sup> For these reasons, SNP markers derived from intragenic regions are the most informative markers for genome-wide genetic analysis in intraspecific tomato populations. By comparing expressed sequence tags (ESTs) in tomato and related wild species, approximately 40 000 candidate SNPs have been identified.<sup>24–27</sup> Since then, the number of ESTs derived from several tomato cultivars has increased to approximately 300 000, all of which are available in the public DNA databases, e.g. DNA Data Bank of Japan (DDBJ: <http://www.ddbj.nig.ac.jp/>), Sol Genomics Network (SGN: <http://solgenomics.net/>), and MiBASE (<http://www.pgb.kazusa.or.jp/mibase/>).

The tomato is regarded as a model plant not only for the *Solanaceae* but also for other fruiting plants.<sup>28</sup> A miniature dwarf cultivar, 'Micro-Tom', originally bred for home gardening purposes,<sup>29</sup> has drawn attention as a model tomato line because of its small plant size, short life cycle, easy transformation, and availability of transposon-tagging systems for use in reverse genetics.<sup>30</sup> Various genomic and genetic

resources have been developed for 'Micro-Tom'. These include mutagenized lines,<sup>31,32</sup> effective transformation systems,<sup>33,34</sup> metabolite annotations,<sup>35</sup> full-length cDNAs,<sup>36</sup> and BAC-end sequences (Asamizu *et al.*, released in the public DNA database with accession numbers: FT227487–FT321168). 'Micro-Tom' seeds are available through two seed stock centers: the Tomato Genetics Resource Center at the University of California, Davis (USA, accession no. LA3911) and the National Bio-Resource Project at the University of Tsukuba (Japan, accession no. TOMJPF00001).

In this study, we developed SNP markers using publicly available ESTs from several tomato cultivars and designed an SNP-genotyping platform using the GoldenGate<sup>®</sup> assay (Illumina, San Diego, CA, USA) in order to accelerate genetic studies and molecular breeding in tomato. SNP markers, along with SSR markers and intronic polymorphic markers, which were developed and mapped onto the interspecific map Tomato-EXPEN 2000 by Shirasawa *et al.*,<sup>7</sup> were applied to create linkage maps using two mapping populations derived from crosses between 'Micro-Tom' and 'Ailsa Craig', a greenhouse-type tomato, and between 'Micro-Tom' and 'M82', a processing tomato. In addition, the polymorphism of the SNP markers was investigated in cultivated tomato lines in order to estimate the transferability of the SNPs to breeding materials.

## 2. Materials and methods

### 2.1. Plant materials

Two F<sub>2</sub> mapping populations, AMF2 and MMF2, each derived by crossing two *S. lycopersicum* lines, were used for the construction of the linkage maps. AMF2 ( $n = 120$ ) was derived from a cross between the 'Ailsa Craig' and 'Micro-Tom' lines, while MMF2 ( $n = 135$ ) was derived from a cross between the 'M82' and 'Micro-Tom' lines. AMF2 and MMF2 were generated in the National Institute of Vegetable and Tea Science, Japan, and in the Institut National de la Recherche Agronomique, France, respectively (Table 1). To address potential residual heterozygosity in the parental 'Micro-Tom' lines used to create AMF2 and MMF2, they are distinguished in this study by the designations 'Micro-Tom\_AM' and 'Micro-Tom\_MM', respectively. Along with the four parental lines of the mapping populations, 22 lines, including 16 inbred and 6 hybrid tomato lines, and an *S. pennellii* line ('LA716') were used for polymorphic analysis of SNPs (Table 1). Total DNA for each line was extracted using the DNeasy Plant Mini kit (Qiagen, Hilden, Germany).

**Table 1.** Description of plant materials

Line name	Note	Source <sup>a</sup>	Accession number	SNP validation <sup>b</sup>
Parental lines of mapping populations				
Micro-Tom_AM	Inbred line	NIVTS		Tested
Ailsa Craig	Inbred line	NBRP	TOMJPF00004	Tested
Micro-Tom_MM	Inbred line	INRA		
M82	Inbred line	INRA		Tested
Tomato lines for SNP typing				
Aichi First	Inbred line	NBRP	TOMJPF00003	
Best of All	Inbred line	NIVTS	LS3908	
Earliana	Inbred line	TGRC	LA3238	Tested
Fruit	Inbred line	NIVTS	LS1100	
Furikoma	Inbred line	NIVTS	LS3903	
Heinz 1706-BG	Inbred line	NIVTS	LS461	Tested
LA925	Inbred line	Cornell University		Tested
Marglobe	Inbred line	TGRC	LA0502	Tested
Money Maker	Inbred line	TGRC	LA2706	Tested
Ponderosa	Inbred line	NIVTS	LS1728	
Rio Grande	Inbred line	TGRC	LA3343	Tested
Rutgers	Inbred line	TGRC	LA1090	Tested
San Marzano	Inbred line	NIVTS	LS4956	Tested
Tomato Chuukanbohon Nou 9	Inbred line	NIVTS		
Tomato Chuukanbohon Nou 11	Inbred line	NIVTS		
Geronimo	F <sub>1</sub> hybrid	De Ruiter Seeds Co.		Tested
Labell	F <sub>1</sub> hybrid	De Ruiter Seeds Co.		Tested
Matrix	F <sub>1</sub> hybrid	De Ruiter Seeds Co.		Tested
Momotaro 8	F <sub>1</sub> hybrid	Takii Seeds Co.		Tested
Reika	F <sub>1</sub> hybrid	Sakata Seeds Co.		Tested
Regina	Inbred line, cherry type	Sakata Seeds Co.		
Sweet100	F <sub>1</sub> hybrid, cherry type	Vilmorin Seeds Co.		Tested
LA716	Inbred line, <i>S. pennellii</i>	Cornell University		

<sup>a</sup>NBRP: University of Tsukuba in National Bio-Resource Project of MEXT, Japan; INRA: National Institute for Agricultural Research, France; NIVTS: National Institute of Vegetable and Tea Science, Japan; TGRC: Tomato Genetics Resource Center, University of California, Davis, USA.

<sup>b</sup>Lines that used for validation of 82 eSNPs prior to design a SNP genotyping platform using Illumina GoldenGate<sup>®</sup> assay.

## 2.2. Development of SNP markers and polymorphic analysis

A total of 229 086 EST sequences from *S. lycopersicum*, retrieved from two public databases, SGN (<http://solgenomics.net/>) and MiBASE (<http://www.pgb.kazusa.or.jp/mibase/>), were used for identification of eSNPs, i.e. SNPs discovered *in silico*. The ESTs registered in MiBASE were derived only from 'Micro-Tom', while those registered in SGN were developed from 19 tomato lines including 'Micro-Tom'. The retrieved EST sequences were assembled using the MIRA program.<sup>37</sup> The eSNPs were then selected according to the following three criteria: (i) only nucleotides with Phred scores of 15 or

more were considered candidates for eSNPs, (ii) a nucleotide at an eSNP site should be identical among multiple sequences within a given line, and (iii) no other SNP candidates should be detected on the flanking sequences 10 bp upstream and downstream of a given candidate.

In order to validate the credibility of the identified eSNP, nucleotide sequences of PCR products containing the eSNP regions were determined by direct sequencing using a DNA sequencer (ABI-3730x1, Applied Biosystems, Foster City, CA, USA). A total of 82 primer pairs were designed in flanking regions of the randomly selected target eSNPs using the Primer3 program.<sup>38</sup> PCR was performed for 17

tomato lines listed in Table 1 in a 5- $\mu$ l reaction mixture containing 0.5 ng genomic DNA, 1 $\times$  PCR buffer (Bioline, London, UK), 3 mM MgCl<sub>2</sub>, 0.04 U BIOTAQ™ DNA polymerase (Bioline), 0.2 mM dNTPs, and 0.8  $\mu$ M of each of the primers. The modified 'touchdown PCR' protocol was used as described previously.<sup>39</sup>

After validation of the 82 eSNPs, a total of 1536 eSNPs were subjected to polymorphic analysis for the two mapping populations and the 23 tomato lines described above using the GoldenGate® assay system (Illumina). Allele- and locus-specific oligonucleotides were designed from the flanking sequences of the 1536 SNP sites using the iCom website (<https://icom.illumina.com/>). Polymorphic analysis of the SNPs was performed according to the standard protocol of the GoldenGate® assay, and the data analysis was performed using GenomeStudio Data Analysis software (Illumina).

SNPs in *DWARF (D)* and *SELF-PRUNING (SP)* were analyzed using the dCAPS and CAPS methods, respectively. PCR was performed under the same conditions as described above. The primer sequences are shown in Supplementary Table S1. The PCR products from the *D* and *SP* genes were digested with *Pst*I and *Mva*I, respectively, and were subjected to electrophoresis on native 10% polyacrylamide gels in 1 $\times$  TBE buffer. The resulting DNA bands were then stained with ethidium bromide.

### 2.3. Mapping of SSR and intronic polymorphic markers on AMF2

A total of 3510 tomato genomic SSR (TGS), 2047 tomato EST-SSR (TES), and 166 tomato EST-derived intronic polymorphic (TEI) markers, developed by Shirasawa *et al.*,<sup>7</sup> were used for segregation analysis of the AMF2 population (Supplementary Table S1). The polymorphic analyses of the markers were performed as described previously.<sup>7</sup> Primer information for the tested markers is available at <http://www.kazusa.or.jp/tomato/>.

### 2.4. Linkage analysis

Linkage analysis was performed using the JoinMap® program, version 4.<sup>40</sup> The segregated data were classified into 12 linkage groups, which corresponded to the Tomato-EXPEN 2000 map,<sup>7</sup> using the grouping module of JoinMap® with LOD scores of 4.0–10.0. The marker order and relative genetic distances were calculated by the regression-mapping algorithm with the following parameters: Haldane's mapping function, recombination frequency  $\leq 0.35$ , and LOD score  $\geq 2.0$ .

## 3. Results

### 3.1. In silico SNP mining and validation

A total of 170 586 and 58 500 EST sequences available in SGN and MiBASE, respectively, along with data on their quality, were used for *in silico* SNP mining. The name of the original tomato line for each EST was obtained from the DDBJ database (<http://www.ddbj.nig.ac.jp/>). In total, 229 086 ESTs derived from 20 tomato cultivars, the average length of which was 497 bp, were used for assembly (Table 2).

Assembly was performed using nucleotides with Phred scores  $\geq 15$ . As a result, a total of 20 274 contigs, the average length of which was 775 bp, and 29 698 singletons were generated. From initial alignment data from all 20 274 contigs, a total of 5607 eSNP sites were identified in 2634 of these contigs (Supplementary Tables S2 and S3). We gave an SNP code to each eSNP according to the following rule: contig name and position of the eSNP on the contig, linked with an underscore, e.g. the 112th position on contig 2758 was given the following SNP code: 2758\_112.

Before designing the SNP genotyping platform (using the Illumina GoldenGate® assay), 82

**Table 2.** Number of ESTs and their original sources used for assembling

Line name <sup>a</sup>	No. of ESTs
TA496	106 142
Micro-Tom	101 157
Rio Grande PtoR	8803
R11-13	5031
R11-12	4925
TA492	2120
West Virginia 106	861
Money Maker	11
Ailsa Craig	7
VF36	7
Momotaro	4
Zhongshu 4	4
Betterboy	3
Vendor	3
House Odoriko	2
Rutgers	2
M82	1
Pera	1
Rio Grande	1
UC82B	1
Total	229 086

<sup>a</sup>Names of tomato lines used for EST generation.

randomly selected eSNPs were tested in 17 tomato lines (Table 1) by direct sequencing of fragments amplified by PCR. As a result, 55 (67%) out of the 82 examined eSNP candidates were experimentally confirmed as SNPs at the predicted positions, indicating that approximately 67% of the 5607 eSNPs detected *in silico* represent true SNPs in the tomato lines used in the present study. In addition, 40 (49%) and 50 (61%) of the 82 eSNPs segregated between the two mapping parents for AMF2 and MMF2, respectively.

For SNP genotyping, a total of 1536 SNPs were selected from the 5607 eSNPs, as follows: (i) one eSNP was selected from each contig and the Selected-BAC-Mixture contig released from the Kazusa Tomato SBM & Marker Database (<http://www.kazusa.or.jp/tomato/>); (ii) an SNP score of more than 0.6, as determined by the iCom website of Illumina (<https://icom.illumina.com/>), was required for each of these eSNPs. As reported by the GoldenGate<sup>®</sup> assay, 1338 (87%) out of the 1536 SNPs could be properly genotyped in the 279 plants. These included the two mapping populations (AMF2 and MMF2) and 23 other tomato lines. The remaining 198 (13%) eSNPs failed to be genotyped because fluorescent signals for these eSNPs did not form clusters pursuant to the criteria required by the GenomeStudio Data Analysis software (Illumina).

### 3.2. Mapping of SNP, SSR, and intronic markers

In the AMF2 population, 648 of the 1338 available SNPs (48.4%) generated segregation data, a similar ratio to that determined in the validation of the 82 eSNPs. Two SNP markers designed in the *D* and *SP* genes, for which 'Micro-Tom' has mutant alleles,<sup>41</sup> showed polymorphism between 'Ailsa Craig' and 'Micro-Tom'. Along with the SNP markers, a total of 5723 previously reported markers, including 2047 EST-SSR (TES), 3510 genomic-SSR (TGS), and 166 intronic (TEI) markers, were used for the polymorphic analysis. As a result, 96 TES (4.7%), 223 TGS (6.3%), and 28 TEI (16.8%) markers exhibited polymorphism between the parental lines. In total, 997 markers were used to construct the AMF2 linkage map.

In the MMF2 population, 640 of the 1338 available SNPs (47.9%) segregated. This ratio was over 10% less than that determined in the validation of the 82 eSNPs, suggesting that the result of the eSNP validation was overestimated. The SNP on the *D* gene showed polymorphism in the MMF2 mapping population, while two parental lines detected the mutated *sp* allele for the *SP* gene. In total, 641 segregated markers were used to construct the MMF2 map.

### 3.3. Construction of linkage maps

For AMF2, a total of 989 of the 997 segregated loci (99.2%) formed 12 linkage groups (LGs), while 637 of the 641 segregated loci (99.4%) formed 13 linkage groups for MMF2. The total sizes of the LGs of the AMF2 and MMF2 maps were 1467.8 and 1422.7 cM, respectively (Table 3, Fig. 1, Supplementary Table S1). Combining the two maps yielded a total of 1137 markers, including 793 SNP, 221 TGS, 93 TES, 28 TEI, and 2 gene markers, located on the intraspecific map. Among these, 488 SNP markers were commonly located on both linkage maps, while 157 and 148 marker loci were specific to the AMF2 map and the MMF2 map, respectively. Chromosome 7 (Chr07) of MMF2 divided into two linkage groups, Chr07p and Chr07q, which were located at the upper and the lower portions, respectively, of Chr07 of Tomato-EXPEN 2000. The average lengths of the intervals between two loci on the AMF2 and the MMF2 maps were calculated to be 1.5 and 2.2 cM, respectively.

Segregation distortions were observed in the two maps. In the AMF2 map, 9.8% of the marker loci showed segregation distortions, ranging from 0.0% for Chr01, Chr08, and Chr12, to 55.4% for Chr11 (Table 3). In the MMF2 map, 5.3% of the marker loci were distorted, ranging from 0.0% for Chr05 and Chr08, to 17.0% for Chr09 (Table 3). The linkage groups harboring severe segregation distortions were different between the two mapping populations, especially between Chr11 of AMF2 (55.4%) and that of MMF2 (2.3%), suggesting Chr11 of 'Ailsa Craig' might have transmission ratio distorters.


### 3.4. Polymorphic analysis of the SNP markers in tomato cultivars and *S. pennellii*

A total of 916 (68.5%) out of the 1338 SNP markers showed polymorphisms in at least one line among the 27 tomato lines listed in Table 1 (Supplementary Table S4). The polymorphic ratio was similar to the ratio determined during the PCR-based validation of the 82 eSNPs. In 'LA719' (*S. pennellii*) and 'Sweet 100', no data were obtained for 229 (17.1%) and one SNP markers, respectively. The polymorphic ratios differed according to the combination of tomato lines (Fig. 2), and the number of segregated SNPs between any two lines among the 27 lines was 255.0 (19.1%) on average. A total of 608.2 SNPs (45.5%) were identified between 'Micro-Tom' and the other inbred lines, on average, while only 80.8 SNPs (6.0%) were identified among the 17 inbred tomato lines. Within the 17 inbred tomato lines, 'M82' showed the highest number of polymorphisms: 176.3 SNPs (13.2%) on average, which was twice as high as that of the other lines.

**Table 3.** Length, number of mapped loci, and segregation distortion of the AMF2 and MMF2 maps

Chromosome	AMF2							Segregation distortion ratio (%)	MMF2				Segregation distortion ratio (%)
	Length (cM)	Number of loci							Length (cM)	Number of loci			
		TGS	TES	TEI	SNP	Gene	Total		SNP	Gene	Total		
Chr01	187.0	18	8	5	62	0	93	0.0	158.4	57	0	57	3.5
Chr02	141.9	24	6	2	39	1	72	2.8	105.1	60	1	61	9.8
Chr03	143.2	29	6	3	70	0	108	11.1	130.4	100	0	100	4.0
Chr04	132.0	15	15	4	114	0	148	3.4	125.3	97	0	97	4.1
Chr05	54.2	45	8	1	51	0	105	2.9	69.2	70	0	70	0.0
Chr06	95.5	3	0	1	9	1	14	7.1	95.1	18	0	18	16.7
Chr07	117.9	8	17	1	91	0	117	3.4	29.0 <sup>a</sup>	54 <sup>a</sup>	0	54	1.9
Chr08	124.0	5	5	0	22	0	32	0.0	92.8	23	0	23	0.0
Chr09	121.7	13	7	7	53	0	80	15.0	118.2	53	0	53	17.0
Chr10	97.3	6	3	0	22	0	31	6.5	87.2	26	0	26	11.5
Chr11	118.4	25	11	3	62	0	101	55.4	115.3	43	0	43	2.3
Chr12	134.7	30	7	1	50	0	88	0.0	104.0	35	0	35	2.9
Total	1467.8	221	93	28	645	2	989	9.8	1422.7	636	1	637	5.3

<sup>a</sup>Chr07 of the MMF2 were divided into two linkage groups, Chr07p and Chr07q. The numbers indicate total value of two linkage groups.



**Figure 1.** Intrasppecific linkage maps for tomato. The middle and right bars of each linkage group represent maps of the AMF2 and MMF2 lines, respectively. The left bar represents a linkage map of Tomato-EXPEN 2000, which is an interspecific map constructed in our previous study.<sup>7</sup> The lines on each linkage group show SNP (green), TGS (red), TES (blue), TEI (yellow), and the other available (black) markers. Common markers between two maps are connected with black lines. Boxes on the Tomato-EXPEN 2000 map indicate heterochromatic regions. Detailed information on the markers, including marker positions, is shown in Supplementary Table S1, and is also available at <http://www.kazusa.or.jp/tomato/>.

SNPs between the F<sub>1</sub> hybrid cultivars and the inbred lines were found at 190.6 loci (14.2%) on average. The two cherry-type tomato cultivars showed higher polymorphisms than the inbred tomato lines, with 310.1 (23.2%) SNPs on average. When 26

*S. lycopersicum* lines were compared with *S. pennellii*, on average, 618.5 out of the 1338 loci (46.2%) were polymorphic. Heterozygosity was observed at multiple SNP sites in all six F<sub>1</sub> hybrid cultivars, ranging from 69 (5.2%) in 'Matrix' to 229 (17.1%)

	Micro-Tom_MM	136	
Inbred lines	Aichi First	647 543	
	Alisa Craig	648 544	
	Best Of All	639 535 52 37	
	Earliana	644 546 45 56 41	
	Fruit	660 556 43 64 69 68	
	Furikoma	651 549 60 87 70 73 65	
	Heinz 1706-BG	669 577 58 69 78 53 63 84	
	LA925	679 573 80 69 76 83 73 116 92	
	M82	738 640 177 182 193 190 172 167 161 163	
	Marglobe	657 553 69 41 58 65 63 85 76 82 190	
	Money Maker	648 544 73 22 39 60 66 79 69 77 194 43	
	Ponderosa	650 544 6 68 51 48 43 57 82 82 177 69 74	
	Rio Grande	683 581 69 55 70 76 69 88 79 91 180 55 66 69	
	Rutgers	640 538 41 64 45 46 82 71 71 91 188 72 64 41 71	
	San Marzano	649 547 46 47 44 49 53 76 72 82 181 69 65 49 64 43	
	Tomato Chuukanbohon Nou 9	647 545 62 61 74 73 63 56 70 98 169 73 61 64 66 77 72	
	Tomato Chuukanbohon Nou 11	659 555 86 89 92 99 81 84 98 116 137 94 85 84 95 99 104 82	
	F <sub>1</sub> hybrid	Geronimo	756 658 205 194 193 206 206 192 212 223 286 203 189 203 206 204 206 187 198
		Labell	749 647 199 180 187 201 197 199 204 210 263 196 175 199 197 207 200 185 163 100
		Matrix	762 662 191 180 185 194 189 187 195 214 269 185 178 193 184 202 195 175 178 77 101
Momotaro 8		760 659 158 163 168 173 146 157 165 190 235 164 159 158 162 166 172 149 135 183 181 171	
Reika		747 646 180 183 187 193 166 175 187 201 242 183 174 180 177 186 190 167 148 186 180 190 98	
Inbred lines, cherry type	Regina	560 490 286 243 264 277 283 280 276 290 281 263 253 287 283 281 276 284 278 374 361 386 350 374	
	Sweet100	689 600 359 327 336 350 346 347 353 353 366 321 329 359 336 358 351 350 330 413 396 398 382 381 322	
F <sub>1</sub> hybrid, cherry type	LAT16	597 606 612 610 604 604 618 605 611 621 625 614 609 608 623 613 616 613 611 635 638 640 660 667 550 672	
<i>S. pennellii</i>	Micro-Tom_MM		
	Aichi First		
	Alisa Craig		
	Best Of All		
	Earliana		
	Fruit		
	Furikoma		
	Heinz 1706-BG		
	LA925		
	M82		
	Marglobe		
	Money Maker		
Ponderosa			
Rio Grande			
Rutgers			
San Marzano			
Tomato Chuukanbohon Nou 9			
Tomato Chuukanbohon Nou 11			
Geronimo			
Labell			
Matrix			
Momotaro 8			
Reika			
Regina			
Sweet100			
LAT16			
Micro-Tom_MM			
Aichi First			
Alisa Craig			
Best Of All			
Earliana			
Fruit			
Furikoma			
Heinz 1706-BG			
LA925			
M82			
Marglobe			
Money Maker			
Ponderosa			
Rio Grande			
Rutgers			
San Marzano			
Tomato Chuukanbohon Nou 9			
Tomato Chuukanbohon Nou 11			
Geronimo			
Labell			
Matrix			
Momotaro 8			
Reika			
Regina			
Sweet100			
LAT16			
Micro-Tom_MM			
Aichi First			
Alisa Craig			
Best Of All			
Earliana			
Fruit			
Furikoma			
Heinz 1706-BG			
LA925			
M82			
Marglobe			
Money Maker			
Ponderosa			
Rio Grande			
Rutgers			
San Marzano			
Tomato Chuukanbohon Nou 9			
Tomato Chuukanbohon Nou 11			
Geronimo			
Labell			
Matrix			
Momotaro 8			
Reika			
Regina			
Sweet100			
LAT16			

**Figure 2.** Number of SNP markers identified by the GoldenGate<sup>®</sup> assay among 27 tomato and *S. pennellii* lines. The numbers show the loci of the 1338 SNPs genotyped in this study. The colors in each cell represent a continuum of polymorphic ratios: lower ratios are represented by green color, higher ratios by red color, and middle ratios by yellow color.



in 'Sweet 100'. In the inbred line 'Rio Grande', 25 (1.9%) heterozygous SNPs were identified.

It is noteworthy that 136 SNPs (10.2%) were identified between 'Micro-Tom\_AM' and 'Micro-Tom\_MM', the parental lines of AMF2 and MMF2, respectively (Fig. 2). Out of these 136 SNP loci, 134 mapped onto the AMF2 and/or the MMF2 maps, mainly on Chr04 (44 loci), Chr07 (38 loci), and Chr12 (36 loci) (Supplementary Table S1). 'Micro-Tom\_AM' had a higher number of polymorphisms, in comparison with the other 25 examined lines, than 'Micro-Tom\_MM' (Fig. 2). For example, 738 loci (55.2%) showed polymorphisms between 'Micro-Tom\_AM' and 'M82', while only 640 SNPs (47.8%) were found between 'Micro-Tom\_MM' and 'M82'. It is likely that this difference resulted in an overestimation of the number of segregated loci between 'Micro-Tom' and 'M82' in the 82-eSNP PCR-based validation.

#### 4. Discussion

To our knowledge, the two genetic linkage maps presented here are the first intraspecific maps for *S. lycopersicum* with SNPs and other PCR-based co-dominant markers. The AMF2 and MMF2 genetic linkage maps comprise a total of 989 and 637 DNA marker loci, respectively, including SNP, SSR, and intron polymorphic and gene markers. Because the SNP markers developed in this study showed a higher degree of polymorphism among the tomato cultivars than SSR markers, SNP information is greatly important to be utilized for genetic analyses in cultivated tomato, including gene mapping, QTL analysis, population genetics, and marker-assisted breeding. In addition, the genomic tools developed in this study will be valuable for exploiting the extensive artificially induced genetic variability created by ethyl methane sulfonate (EMS) mutagenesis in 'Micro-Tom' mutant collections. For example, they could allow, by forward genetic approaches, the identification of the causal mutations for remarkable fruit and plant phenotypes.

In the SNP genotyping by the GoldenGate<sup>®</sup> assay, 1338 (87%) of the 1536 SNPs could be successfully genotyped. In other crops, successful ratio of SNP genotyping by the GoldenGate<sup>®</sup> assay is reported to be ranging from 79 to 92%,<sup>42-45</sup> which fits to the result of the present study. In order to improve the ratio, we suggest additional three criteria to select SNPs for genotyping. The first is elimination of SNP positioned near junction site of intron and exon, because intron inhibits hybridization of allele- and locus-specific oligonucleotide to the target sequence based on EST. It can be achieved by comparing the sequences of EST with those of genome, if available.

Next is avoidance designing SNP markers on multi-copy genes, which disrupts the fluorescent-signal clusters on the GenomeStudio Data Analysis software. Selection of accurate SNP site, e.g. with high-quality value and/or with highly coverage of sequence fragments, is also important. Large scale of genome analysis by massively parallel DNA sequencers would be convenient to overcome these matters.

In interspecific linkage maps of tomato and its relatives, markers derived from ESTs tend to distribute randomly along the genome, while markers derived from random genomic regions, e.g. RAPD, AFLP, and genomic SSRs, tend to form clusters in heterochromatic regions.<sup>7,19-22</sup> In this study, however, the marker loci did not disperse along the two linkage maps derived from AMF2 and MMF2, despite the fact that most markers were developed from ESTs. Comparison between the maps of Tomato-EXPEN 2000 and the two intraspecific mapping populations did not indicate any evidence of an obvious relationship between the marker clusters on the maps and chromosome structures, i.e. the heterochromatic and euchromatic regions (Fig. 1). It can be assumed that the marker clusters correspond to probable integration regions originating from '*Lycopersicon minutum*', an ancestral line of 'Micro-Tom',<sup>29</sup> which belongs to the *S. chmielewskii* and *S. neorickii* complex.<sup>46</sup> Such regions are expected to show higher frequencies of polymorphism than the other regions, which originate from cultivated lines.

Out of the 989 SNP markers on the AMF2 map, 489 were located on the two intraspecific maps generated in this study and 155 had already been located on the interspecific map generated from Tomato-EXPEN 2000.<sup>7</sup> The common markers for the three maps allow the alignment and connection of these maps as shown in Fig. 1. Significant translocation and inversion of chromosome were not observed between intra- and interspecific maps, meaning the order of genes would be conserved in two species, and the genome sequence of tomato (*S. lycopersicum*) could be used as a reference genome of *S. pennellii*. In addition, it is likely that the AMF2 and MMF2 maps cover the whole tomato genome except for the middle part of Chr07 on the MMF2 map, and that the marker order is mostly conserved in the three maps. One possible approach to connecting the two linkage groups of Chr07 on the MMF2 map would be to develop a novel mapping population between 'M82' and 'Micro-Tom\_AM' instead of 'Micro-Tom\_MM' because SNPs on Chr07 segregated more frequently between 'M82' and 'Micro-Tom\_AM'.

Indeed, we found that 136 of the 1338 tested SNP markers (10.2%) showed polymorphisms between 'Micro-Tom\_AM' and 'Micro-Tom\_MM' (Fig. 2), indicating possible residual heterosis in 'Micro-Tom'. It is

assumed that these loci had not been fixed at the time of 'Micro-Tom' being released, although 'Micro-Tom' seeds are propagated and distributed in the F<sub>12</sub> generation after a crossing.<sup>29</sup> Theoretically, the heterozygosity of genome in the F<sub>12</sub> generation is calculated to be 0.05% [ $=(1/2)^{12-1}$ ] in self-pollinating plants, which means that most of the genomic regions are expected to be homozygous. Most of the polymorphic markers between the two 'Micro-Tom' lines were mapped on Chr04, Chr07, and Chr12. This result suggested that 'Micro-Tom' might have been bred under natural and/or artificial selection pressure from the regions under the influence of heterosis or crossing incompatibilities between *S. lycopersicum* and *L. minutum*. Alternatively, multiple lines might have been selected as 'Micro-Tom' from the breeding population before the complete fixation of the genotypes of each plant.

Though 'Micro-Tom' itself, bred as an ornamental plant, has little agricultural value, its genes may be of great value to agriculture. 'Micro-Tom' has resistance to several diseases, caused by *Alternaria alternata*, *Corynespora cassiicola*, *Fusarium oxysporum*, and *Pseudomonas syringae*.<sup>47</sup> Moreover, a large number of mutant lines have been developed using 'Micro-Tom'.<sup>31,32</sup> The markers and maps developed in this study may therefore be useful for introgression breeding for disease resistance or targeted genes identified in 'Micro-Tom' or its mutant lines. Indeed, 'Micro-Tom' mutant lines carry mutated alleles that may confer high agricultural value to tomato, e.g. alleles causing large variations in fruit color, shape, size, and composition. Mutants may also help to decipher the mechanisms controlling specific traits in tomato.

In this study, we demonstrated the validity of the strategy of combining large-scale eSNP discovery with high-throughput SNP genotyping assays. Comparison of sequence data from tomato cultivars has been reported as an efficient strategy for developing a large number of SNP markers for tomato cultivars.<sup>24,48,49</sup> Today, extensive amounts of sequence data from crop genomes can be easily collected using massively parallel DNA sequencers.<sup>50</sup> In addition, genomic sequences from The International Tomato Genome Sequencing Consortium of SGN will soon become available.<sup>51</sup> The accumulating genome sequences can be used to develop custom SNP markers within cultivated tomato. The molecular markers and genetic linkage maps developed in the present study represent one of the initial milestones in the fusion of genomics, genetics, and molecular breeding in cultivated tomato.

**Acknowledgements:** We are grateful to Dr K. Aoki (Kazusa DNA Research Institute, Japan) for providing the EST data for Micro-Tom. Plant materials were

provided by Dr S. D. Tanksley (Cornell University, USA), Dr T. Ariizumi (University of Tsukuba through the National Bio-Resource Project of the Ministry of Education, Culture, Sports, Science and Technology, Japan), Dr T. Saito (National Institute of Vegetable and Tea Science, Japan), and Dr S. M. Tam (Tomato Genomic Resource Center, University of California, Davis, USA).

**Supplementary data:** Supplementary Data are available at [www.dnaresearch.oxfordjournals.org](http://www.dnaresearch.oxfordjournals.org).

### Availability

Information on the SNP and SSR markers, the AMF2 and MMF2 linkage maps, and the SNP genotypes for the tomato lines investigated in the current study are available at <http://www.kazusa.or.jp/tomato/>.

### Funding

This work was supported by the Kazusa DNA Research Institute Foundation and the Ministry of Agriculture, Forestry, and Fisheries of Japan with the cooperation of the Genomics for Agricultural Innovation Foundation (DD-4010).

### References

1. Labate, J.A., Grandillo, S., Fulton, T., et al. 2007, Tomato, In: Cole, C. (ed.), *Genome Mapping and Molecular Breeding in Plants*, vol. 5. Springer: New York, pp. 1–125.
2. Paterson, A.H., Lander, E.S., Hewitt, J.D., Peterson, S., Lincoln, S.E. and Tanksley, S.D. 1988, Resolution of quantitative traits into Mendelian factors by using a complete linkage map of restriction fragment length polymorphisms, *Nature*, **335**, 721–726.
3. Tanksley, S.D., Ganai, M.W., Prince, J.P., et al. 1992, High density molecular linkage maps of the tomato and potato genomes, *Genetics*, **132**, 1141–1160.
4. de Vicente, M.C. and Tanksley, S.D. 1993, QTL analysis of transgressive segregation in an interspecific tomato cross, *Genetics*, **134**, 585–596.
5. Paran, I., Goldman, I., Tanksley, S.D. and Zamir, D. 1995, Recombinant inbred lines for genetic mapping in tomato, *Theor. Appl. Genet.*, **90**, 542–548.
6. Fulton, T.M., Nelson, J.C. and Tanksley, S.D. 1997, Introgression and DNA marker analysis of *Lycopersicon peruvianum*, a wild relative of the cultivated tomato, into *Lycopersicon esculentum*, followed through three successive backcross generations, *Theor. Appl. Genet.*, **95**, 895–902.
7. Shirasawa, K., Asamizu, E., Fukuoka, H., et al. 2010, An interspecific linkage map of SSR and intronic polymorphism markers in tomato, *Theor. Appl. Genet.*, **121**, 731–739.

8. Frary, A., Nesbitt, T.C., Grandillo, S., et al. 2000, *fw2.2*: a quantitative trait locus key to the evolution of tomato fruit size, *Science*, **289**, 85–88.
9. Liu, J., Van Eck, J., Cong, B. and Tanksley, S.D. 2002, A new class of regulatory genes underlying the cause of pear-shaped tomato fruit, *Proc. Natl Acad. Sci. USA*, **99**, 13302–13306.
10. Cong, B., Barrero, L.S. and Tanksley, S.D. 2008, Regulatory change in YABBY-like transcription factor led to evolution of extreme fruit size during tomato domestication, *Nat. Genet.*, **40**, 800–804.
11. Xiao, H., Jiang, N., Schaffner, E., Stockinger, E.J. and van der Knaap, E. 2008, A retrotransposon-mediated gene duplication underlies morphological variation of tomato fruit, *Science*, **319**, 1527–1530.
12. Dixon, M.S., Jones, D.A., Keddie, J.S., Thomas, C.M., Harrison, K. and Jones, J.D. 1996, The tomato *Cf-2* disease resistance locus comprises two functional genes encoding leucine-rich repeat proteins, *Cell*, **84**, 451–459.
13. Martin, G.B., Brommonschenkel, S.H., Chunwongse, J., et al. 1993, Map-based cloning of a protein kinase gene conferring disease resistance in tomato, *Science*, **262**, 1432–1436.
14. Jones, D.A., Thomas, C.M., Hammond-Kosack, K.E., Balint-Kurti, P.J. and Jones, J.D. 1994, Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging, *Science*, **266**, 789–793.
15. Thomas, C.M., Jones, D.A., Parniske, M., et al. 1997, Characterization of the tomato *Cf-4* gene for resistance to *Cladosporium fulvum* identifies sequences that determine recognition specificity in *Cf-4* and *Cf-9*, *Plant. Cell*, **9**, 2209–2224.
16. Fridman, E., Pleban, T. and Zamir, D. 2000, A recombination hotspot delimits a wild-species quantitative trait locus for tomato sugar content to 484 bp within an invertase gene, *Proc. Natl. Acad. Sci. USA*, **97**, 4718–4723.
17. Miller, J.C. and Tanksley, S.D. 1990, RFLP analysis of phylogenetic relationships and genetic variation in the genus *Lycopersicon*, *Theor. Appl. Genet.*, **80**, 437–448.
18. Saliba-Colombani, V., Causse, M., Gervais, L. and Philouze, J. 2000, Efficiency of RFLP, RAPD, and AFLP markers for the construction of an intraspecific map of the tomato genome, *Genome*, **43**, 29–40.
19. Frary, A., Xu, Y., Liu, J., Mitchell, S., Tedeschi, E. and Tanksley, S. 2005, Development of a set of PCR-based anchor markers encompassing the tomato genome and evaluation of their usefulness for genetics and breeding experiments, *Theor. Appl. Genet.*, **111**, 291–312.
20. Ohshima, A., Asamizu, E., Negoro, S., et al. 2009, Characterization of tomato SSR markers developed using BAC-end and cDNA sequences from genome databases, *Mol. Breed.*, **23**, 685–691.
21. Tang, X., Szinay, D., Lang, C., et al. 2008, Cross-species bacterial artificial chromosome-fluorescence *in situ* hybridization painting of the tomato and potato chromosome 6 reveals undescribed chromosomal rearrangements, *Genetics*, **180**, 1319–1328.
22. Wang, Y., Tang, X., Cheng, Z., Mueller, L., Giovannoni, J. and Tanksley, S.D. 2006, Euchromatin and pericentromeric heterochromatin: comparative composition in the tomato genome, *Genetics*, **172**, 2529–2540.
23. Gupta, P.K., Rustgi, S. and Mir, R.R. 2008, Array-based high-throughput DNA markers for crop improvement, *Heredity*, **101**, 5–18.
24. Jimenez-Gomez, J.M. and Maloof, J.N. 2009, Sequence diversity in three tomato species: SNPs, markers, and molecular evolution, *BMC Plant Biol.*, **9**, 85.
25. Labate, J.A. and Baldo, A.M. 2005, Tomato SNP discovery by EST mining and resequencing, *Mol. Breed.*, **16**, 343–349.
26. Yang, W., Bai, X., Kabelka, E., et al. 2004, Discovery of single nucleotide polymorphisms in *Lycopersicon esculentum* by computer aided analysis of expressed sequence tags, *Mol. Breed.*, **14**, 21–34.
27. Yamamoto, N., Tsugane, T., Watanabe, M., et al. 2005, Expressed sequence tags from the laboratory-grown miniature tomato (*Lycopersicon esculentum*) cultivar Micro-Tom and mining for single nucleotide polymorphisms and insertions/deletions in tomato cultivars, *Gene*, **15**, 127–134.
28. Ezura, H. 2009, Tomato is a next-generation model plant for research and development, *J. Jpn Soc. Hort. Sci.*, **78**, 1–2.
29. Scott, J.W. and Harbaugh, B.K. 1989, Micro-Tom—a miniature dwarf tomato, *Florida Agr. Expt. Sta. Circ.*, **370**, 1–6.
30. Meissner, R., Jacobson, Y., Melamed, S., et al. 1997, A new model system for tomato genetics, *Plant J.*, **12**, 1465–1472.
31. Matsukura, C., Yamaguchi, I., Inamura, M., et al. 2007, Generation of gamma irradiation-induced mutant lines of the miniature tomato (*Solanum lycopersicum* L.) cultivar ‘Micro-Tom’, *Plant Biotechnol.*, **24**, 39–44.
32. Watanabe, S., Mizoguchi, T., Aoki, K., et al. 2007, Ethylmethanesulfonate (EMS) mutagenesis of *Solanum lycopersicum* cv. Micro-Tom for large-scale mutant screens, *Plant Biotechnol.*, **24**, 33–38.
33. Dan, Y., Yan, H., Muniyikwa, T., Dong, J., Zhang, Y. and Armstrong, C.L. 2006, MicroTom—a high-throughput model transformation system for functional genomics, *Plant Cell Rep.*, **25**, 432–441.
34. Sun, H.J., Uchii, S., Watanabe, S. and Ezura, H. 2006, A highly efficient transformation protocol for Micro-Tom, a model cultivar for tomato functional genomics, *Plant Cell Physiol.*, **47**, 426–431.
35. Iijima, Y., Nakamura, Y., Ogata, Y., et al. 2008, Metabolite annotations based on the integration of mass spectral information, *Plant J.*, **54**, 949–962.
36. Aoki, K., Yano, K., Suzuki, A., et al. 2010, Large-scale analysis of full-length cDNAs from the tomato (*Solanum lycopersicum*) cultivar Micro-Tom, a reference system for the Solanaceae genomics, *BMC Genomics*, **11**, 210.
37. Chevreux, B., Pfisterer, T., Drescher, B., et al. 2004, Using the miraEST assembler for reliable and automated mRNA transcript assembly and SNP detection in sequenced ESTs, *Genome Res.*, **14**, 1147–1159.

38. Rozen, S. and Skaletsky, H. 2000, Primer3 on the WWW for general users and for biologist programmers, *Methods Mol. Biol.*, **132**, 365–386.
39. Sato, S., Isobe, S., Asamizu, E., et al. 2005, Comprehensive structural analysis of the genome of red clover (*Trifolium pratense* L.), *DNA Res.*, **12**, 301–364.
40. Van Ooijen, J.W. 2006, *JoinMap<sup>®</sup>4, Software for the Calculation of Genetic Linkage Maps in Experimental Populations*, Kyazma. BV, Wageningen, Netherlands.
41. Marti, E., Gisbert, C., Bishop, G.J., Dixon, M.S. and Garcia-Martinez, J.L. 2006, Genetic and physiological characterization of tomato cv. Micro-Tom, *J. Exp. Bot.*, **57**, 2037–2047.
42. Hyten, D.L., Song, Q., Choi, I.-K., et al. 2008, High-throughput genotyping with the GoldenGate assay in the complex genome of soybean, *Theor. Appl. Genet.*, **116**, 945–952.
43. Muchero, W., Diop, N.N., Bhat, P.R., et al. 2009, A consensus genetic map of cowpea [*Vigna unguiculata* (L) Walp.] and synteny based on EST-derived SNPs, *Proc. Natl Acad. Sci. USA*, **106**, 18159–18164.
44. Hyten, D.L., Song, Q., Fickus, E.W., et al. 2010, High-throughput SNP discovery and assay development in common bean, *BMC Genomics*, **11**, 475.
45. Deulvot, C., Charrel, H., Marty, A., et al. 2010, Highly-multiplexed SNP genotyping for genetic mapping and germplasm diversity studies in pea, *BMC Genomics*, **11**, 468.
46. Rick, C.M., Kesicki, E., Fobes, J.F. and Holle, M. 1976, Genetic and biosystematic studies on two new sibling species of *Lycopersicon* from interandean Peru, *Theor. Appl. Genet.*, **47**, 55–68.
47. Takahashi, H., Shimizu, A., Arie, T., et al. 2005, Catalog of Micro-Tom tomato responses to common fungal, bacterial, and viral pathogens, *J. Gen. Plant Pathol.*, **71**, 8–22.
48. Labate, J.A., Robertson, L.D., Wu, F., Tanksley, S.D. and Baldo, A.M. 2009, EST, COSII, and arbitrary gene markers give similar estimates of nucleotide diversity in cultivated tomato (*Solanum lycopersicum* L.), *Theor. Appl. Genet.*, **118**, 1005–1014.
49. Van Deynze, A., Stoffel, K., Buell, C.R., et al. 2007, Diversity in conserved genes in tomato, *BMC Genomics*, **8**, 465.
50. Varshney, R.K., Hoisington, D.A. and Tyagi, A.K. 2006, Advances in cereal genomics and applications in crop breeding, *Trends. Biotechnol.*, **24**, 490–499.
51. Mueller, L.A., Lankhorst, R.K., Tanksley, S.D., et al. 2009, A snapshot of the emerging tomato genome sequence, *Plant Genome*, **2**, 78–92.