



HAL
open science

Application of SSCP-PCR fingerprinting to profile the yeast community in raw milk Salers cheeses

Cecile C. Callon, Céline C. Delbes, Frédérique Duthoit, Marie-Christine M.-C. Montel

► To cite this version:

Cecile C. Callon, Céline C. Delbes, Frédérique Duthoit, Marie-Christine M.-C. Montel. Application of SSCP-PCR fingerprinting to profile the yeast community in raw milk Salers cheeses. *Systematic and Applied Microbiology*, 2006, 29 (2), pp.172-180. 10.1016/j.syapm.2005.07.008 . hal-02659601

HAL Id: hal-02659601

<https://hal.inrae.fr/hal-02659601>

Submitted on 30 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Application of SSCP–PCR fingerprinting to profile the yeast community in raw milk Salers cheeses

Cécile Callon*, Céline Delbès, Frédérique Duthoit, Marie-Christine Montel

Unité de Recherches Fromagères, INRA, 36 rue de Salers, F-15000 Aurillac, France

Received 23 June 2005

Abstract

Bacteria and yeasts are important sensory factors of raw-milk cheeses as they contribute to the sensory richness and diversity of these products. The diversity and succession of yeast populations in three traditional Registered Designation of Origin (R.D.O.) Salers cheeses have been determined by using phenotypic diagnoses and Single-Strand Conformation Polymorphism (SSCP) analysis. Isolates were identified by phenotypic tests and the sequencing of the D1–D2 domains of the 26S rRNA gene. Ninety-two percent of the isolates were identified as the same species in both tests. Yeast-specific primers were designed to amplify the V4 region of the 18S rRNA gene for SSCP analysis. The yeast species most frequently encountered in the three cheeses were *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Saccharomyces cerevisiae*, *Candida zeylanoides* and *Debaryomyces hansenii*. Detection of less common species, including *Candida parapsilosis*, *Candida silvae*, *Candida intermedia*, *Candida rugosa*, *Saccharomyces unisporus*, and *Pichia guilliermondii* was more efficient with the conventional method. SSCP analysis was accurate and could be used to rapidly assess the proportions and dynamics of the various species during cheese ripening. Each cheese was clearly distinguished by its own microbial community dynamics.

© 2005 Elsevier GmbH. All rights reserved.

Keywords: Yeasts; Dynamics; SSCP; 18S rDNA genes; Salers cheese

Introduction

Bacteria and yeasts are an important determinant of the quality of cheeses made from raw milk because they enhance the richness and diversity of the taste of these products. Bacteria alone cannot explain the diversity of the sensorial properties of cheese [8,9]. A comprehensive study on yeast communities in cheeses could help to explain how the sensory qualities of cheeses develop. Several molecular methods are available for microbial

identification but only a few have been applied to yeasts isolated from cheeses. These methods include restriction fragment length polymorphism (RLFP) analysis of Intergenic Transcribed Spacers (ITS) [21], randomly amplified polymorphic DNA (RAPD) [1,22], and 26S rRNA gene sequencing with primers NL1 and NL4 [5]. These culture-dependent methods are now being replaced by direct molecular methods without culture. Indeed, Temperature Gradient Gel Electrophoresis (TGGE) has been used to profile wine yeasts [12] and Denaturing Gradient Gel Electrophoresis (PCR–DGGE) has been used to evaluate yeast biodiversity in raw milk [5]. By applying these methods, it is possible to evaluate the dynamics of yeast populations.

*Corresponding author. Tel.: +33 4 71 45 64 10;
fax: +33 4 71 45 64 13.

E-mail address: callon@clermont.inra.fr (C. Callon).

The objective of this study was to evaluate the diversity of yeasts and their dynamics during ripening in three different raw-milk RDO Salers cheese by combining a direct molecular approach relying on the 18S rDNA analysis by Single-Strand Conformation Polymorphism (SSCP) and phenotypic and molecular identification of isolates of the same cheeses. The SSCP methodology used is based on the direct analysis of 18S rRNA gene pool, after PCR amplification and separation in non-denaturing conditions. It provides a rapid fingerprint of a complex microbial community including dead and alive microbial cells. SSCP has been applied to study microbial communities in water [17], in compost of organic agricultural substrate [20], and in anaerobic digestors [16,29]. This technique has also been used in clinical microbiology to rapidly differentiate bacteria from blood culture [26]. SSCP analysis has never been applied to cheese yeast population but it has been used to investigate bacterial dynamics in the same Salers cheeses [4,9].

Materials and methods

Cheeses samples

Cheese samples were obtained from three farms (A, B, and C), in the Salers cheese production area and selected for different sensory features. Salers cheeses were exclusively produced with raw milk stored in a wooden container (the “gerle”). Milk was neither heated nor refrigerated after milking. No starter culture was added. Samples from three cheeses were collected 1 day after cheesemaking and after 8, 30, 90 and 150 days of ripening. Sub-samples were stored at -80°C until analyzed.

Total DNA isolation from cheese samples

One gram of cheese was mixed with 1 ml of 4 M guanidine thiocyanate–100 mM Tris (pH 7.5) and 125 μl of 10% *N*-lauroylsarcosine and blended with a reciprocating shaker (MM2000; Kurt Retsch) and frozen at -80°C . Phenol/chloroform extractions were used to extract total DNA from samples [9]. The DNA solutions were stored at -20°C .

Primer design

A set of specific primers was developed by comparing gene sequences encoding 18S rRNA (18S rDNA). These sequences, characteristic of most yeast species found in dairy products (26 different species), were aligned by using Multalin version 5.4.1 (multiple sequence alignment with hierarchical clustering) [6]. We designed two conserved primers flanking a variable region in the V4 portion of the 18S rDNA: forward primer Lev1 (5'-

AGTTGTTGCAGTTAAAAAG-3') and reverse primer Lev2 (5'-CCTATTAATCATTACGA-3'). Primer specificity was tested as part of the Ribosomal Database Project (RDP) [18] and Blast, and was experimentally tested with a series of 15 reference yeast strains and Salers isolates (Table 1), with *Geotrichum candidum* a positive control and several species of *Lactobacillus*, *Leuconostoc*, *Lactococcus* and *Pediococcus* as negative controls.

PCR–SSCP amplification and SSCP electrophoresis

The primers used to amplify the 270 bp of the 18S rRNA gene were labeled with 5'-hexachloro derivative of fluorescein (HEX: Lev1) and 5'-fluorescein phosphoramidite (NED: Lev2), purchased from Applied Biosystems (Courtaboeuf, France). PCR was performed in a total volume of 25 μl containing 1 \times Ez buffer (Applied Biosystems), 2.5 mM of Mn (OAc)₂ solution (Applied Biosystems), 0.5 μM of each primer, 0.5 mM dNTPs, 2 U of *rTth* polymerase (Applied Biosystems) and 10 ng of the extracted DNA. The thermocycler was programmed with an initial denaturing step at 94°C for 5 min, followed by 30 cycles of 94°C for 30 s, 47°C for 30 s and 72°C for 60 s, with a final extension at 72°C for 7 min. The amplification products were resolved on a 1% agarose gel and stained with ethidium bromide. PCR products were purified with a StrataPrepTM PCR purification Kit (Stratagene, Saint Quentin en Yvelines, France) following the manufacturer's instructions and then analyzed by SSCP [9].

Cheese 18S rDNA amplification and cloning

The 18S rDNA gene was amplified with unlabeled primers Lev1 and Lev2 from two different cheeses. Each reaction tube contained PfuUltra buffer 1 \times (Stratagene, Amsterdam, The Netherlands), 0.2 mM of each dNTP, 0.5 μM of each primer, 10 ng of purified DNA and 1.25 U of PfuUltra polymerase (Stratagene) adjusted to a total volume of 25 μl . After initial denaturation at 94°C for 5 min, 30 cycles were performed at 94°C for 30 s, 47°C for 30 s and 72°C for 60 s, followed by a final extension step of 72°C for 7 min. PCR products were ligated in the pCR4Blunt-TOPO vector and transformed into *Escherichia coli* TOPO10 OneShot cells as specified by the manufacturer (Invitrogen, Cergy Pontoise, France).

Identification of SSCP peaks

To assign SSCP pattern peaks to species, the 18S rRNA gene clone libraries were analyzed by SSCP. SSCP–PCR was performed on the clones and the resulting peaks were compared with the SSCP patterns of the cheese. Variable

Table 1. SSCP analysis of 18 S rRNA genes of 15 yeast strains and Salers isolates (three repetitions)

Species	Strains	A ^a	Species	Strains	B ^b
		Elution in scan ^c			Elution in scan
<i>C. catenulata</i>	CBS ^d 565	7246	<i>C. catenulata</i>	CBS 565	7378
<i>Y. lipolytica</i>	CLIB ^e 183	7327	<i>Y. lipolytica</i>	CLIB 183	7450
<i>C. rugosa</i>	CLIB 613	7780	<i>C. parapsilosis</i>	Salers isolate	7948
<i>D. hansenii</i> var <i>hansenii</i>	CLIB 197	7792	<i>D. hansenii</i> var <i>hansenii</i>	CLIB 197	7985
<i>C. intermedia</i>	CBS 572	7796	<i>C. intermedia</i>	CBS 572	7993
<i>C. krusei</i>	CLIB 308	7858	<i>C. krusei</i>	CLIB 308	8078
<i>P. membranifaciens</i>	CLIB 212	7859	<i>P. membranifaciens</i>	CLIB 212	8089
<i>P. fermentans</i>	CLIB 198	7883	<i>C. rugosa</i>	CLIB 613	8117
<i>C. parapsilosis</i>	Salers isolate	7949	<i>P. fermentans</i>	CLIB 198	8125
<i>C. silvae</i>	Salers isolate	8021	<i>C. zeylanoides</i>	CLIB 199	8276
<i>Kl. lactis</i>	CLIB 196	8060	<i>S. dairenensis</i>	CLIB 175	8336
<i>Kl. marxianus</i>	CLIB 282	8093	<i>T. delbrueckii</i>	CLIB 230	8339
<i>S. cerevisiae</i>	CLIB 227	8120	<i>Kl. lactis</i>	CLIB 196	8339
<i>T. delbrueckii</i>	CLIB 230	8129	<i>S. cerevisiae</i>	CLIB 227	8393
<i>S. dairenensis</i>	CLIB 175	8129	<i>S. unisporus</i>	CLIB 234	8402
<i>C. zeylanoides</i>	CLIB 199	8174	<i>Kl. marxianus</i>	CLIB 282	8411
<i>S. unisporus</i>	CLIB 234	8204	<i>C. silvae</i>	Salers isolate	8479

C. = Candida, S. = Saccharomyces, Kl. = Kluyveromyces, T. = Torulaspora, P. = Pichia.

^aElution on HEX-labeled forward strand.

^bElution on reverse strand NED.

^cUnit of GeneScan software on the SSCP pattern. Elution values represent the average of three analyses. Standard deviation values always < 5.

^dCentral Bureau voor Schimmelcultures, Neerland.

^eCollection de Levures d'Intérêt Biotechnologique. Grignon, France.

regions of the 18S rRNA gene from 15 representative clones were sequenced by using dye-terminator cycle sequencing ready reaction kits with *Taq* DNA polymerase and primers Lev1 and Lev2. Yeast reference strains and isolates from cheeses also were analyzed by SSCP to create a species pattern database.

Isolation and identification of isolates

Yeasts of the same three cheeses were counted from fresh sample plates. Yeast colonies were grown on an Oxytetracycline Glucose Agar base (OGA) (Biokar Diagnostics, Beauvais, France) supplemented with oxytetracycline (1 mg/ml). One hundred and two colonies were selected from plates at different times of ripening and purified in Yeast Extract Peptone Glucose (YPG) broth (yeast extract 10 g, bacto-peptone 20 g, glucose 40 g and demineralized water to 1000 ml, pH 5.2). On each plate, colonies were selected based on their aspect diversity. The total of picked colonies per plate was always equal to the root square of total number. All cultures were frozen (−80 °C) before identification.

Yeast isolates were identified by using the morphological, biochemical and physiological characters of Kurtzman and Fell [14] and assigned to a species with the dichotomous key of Barnett et al. [3]. Carbon compound assimilation was determined with the API ID

32C kit (Bio Merieux). All tests were performed in duplicate for non-identified isolates.

Molecular identification was performed by sequencing the 26S rRNA gene. Total DNA was extracted from cultures of each isolate according to the miniprep protocol of Hoffman and Winston [13]. The D1/D2 domain of the 26S rRNA gene was amplified by PCR, using primers NL1 and NL4 as described by Kurtzman and Robnett [15]. The sequencing reactions of the forward and reversed strands were carried out with a dye-terminator cycle sequencing ready reaction kit (Applied Biosystems) with AmpliTaq DNA polymerase FS and the NL1 and NL4 primers, respectively. The reaction products were sequenced with an automated DNA sequencer ABI PRISM 310 (Applied Biosystems) and analyzed with sequencing analysis software (Applied Biosystems). The partial 26S rDNA sequences obtained were compared to the sequences deposited in the GenBank DNA database, using the BLAST algorithm [2]. The percentage of similarity with the DNA sequences deposited was determined.

Results and discussion

Discrimination of yeast species by SSCP analysis

To create a species pattern database, the SSCP patterns of pure cultures (15 culture collection strains

belonging to different species frequently encountered in dairy products [11] were determined. In the same way, SSCP patterns of two strains identified below and originating from microbial analysis of the same Salers cheeses were determined too. 18S rDNA was amplified three times with the two primers Lev1 labeled with HEX and Lev2 labeled with NED. Each strain always produced the same pattern, thus confirming the reproducibility of PCR–SSCP analysis. Differences between the forward and reverse strand of the same strain is due to the fact that SSCP distinguishes DNA fragments according to their sequence on basis of single-strand conformation. Taking both strands into account, nearly all species were easily differentiated by SSCP analysis (Table 1). For example, the HEX-labeled strand from *Torulaspora delbrueckii* and *Saccharomyces dairenensis* had a similar pattern, but these two species could be distinguished by analyzing the NED-labeled strand. A similar result was obtained with *Candida krusei* and *Pichia membranifaciens*. *T. delbrueckii* and *Kluyveromyces lactis* had NED-labeled strands that were similar but could be differentiated by analyzing the HEX-labeled strand. Analysis of both strands generally was necessary to distinguish all 17 species.

Direct fingerprinting of yeast diversity in different Salers cheeses by SSCP–PCR

The reproducibility of the PCR–SSCP approach was evaluated by repeating the DNA analysis of a cheese yeast community two times. The two profiles of the same cheese aligned, with the same variations in each peak area (Fig. 1). The yeast community fingerprints from three different cheeses (A, B and C) were evaluated after 8 and 150 days of ripening (Fig. 2, HEX-labeled strand). The 8-day SSCP patterns differed in the number of peaks per sample. After 150 days of ripening, the fingerprints of the same cheese samples had different SSCP–PCR profiles. Some peaks appeared or disappeared in the course of ripening or differed with respect to their size.

Identification of peaks

To identify the sequences corresponding to peaks in the different patterns, a 260-clone library of 18S rRNA genes from the DNA extracted from cheese A at 150 days (Fig. 2 top right) was constructed and each clone was analyzed by SSCP. All of the clones corresponded to one of the four dominant peaks (a, b, d, e). For each peak, five co-eluting clones were sequenced and identified based on the similarity to 18S rDNA sequences deposited in Genbank. SSCP peak identification also relied on the elution of reference strains and yeast isolates (Table 2). Based on the sequence of the clone




Fig. 1. Reproducibility of PCR–SSCP analysis as assessed by repeating two times the DNA analysis of a cheese yeast community, forward primer labeled with HEX.

18S rRNA gene and co-elution with reference strains and isolates, peak a was *K. lactis*, peak b was *K. marxianus* and peak d was *C. zeylanoides*. Peak e co-eluted with clones identified as *D. hansenii* var. *hansenii* and *D. hansenii* var. *fabryii* with a same percentage of similarity with strains in Genbank database. However, peak e had a different pattern to that of the *D. hansenii* var. *hansenii* type strain. Peaks c and h co-eluted, respectively, with the reference strains for *S. cerevisiae* and *S. unisporus*. Peaks f and g co-eluted with isolates identified as *C. parapsilosis* and *C. silvae*. The other peaks present in patterns (Fig. 2) represent less than 1% of total area or in term of absolute area, less than the value considered as the background and were then considered as non-significant.

Dynamics of yeast populations in cheeses

To analyze the SSCP profiles of cheeses, the peak ratio (the proportion of one peak to the total area of all of the peaks) was calculated in each sample and used as an indicator of changes within the yeast populations. Considering the entire sample times (Table 3), the peaks corresponding to *K. lactis*, *K. marxianus*, *S. cerevisiae*, *D. hansenii* and *C. zeylanoides* were systematically present in the SSCP patterns of all three cheeses. Peaks corresponding to *C. parapsilosis*, *C. silvae* and *S.*




Fig. 2. SSCP patterns of PCR-amplified 18S rRNA gene fragments from yeast communities of three cheeses A, B and C at 8 and 150 d of ripening, forward primer labeled with HEX, Y-axis: fluorescence. X-axis: elution in scans (unit of Genescan software). The position and labeling of peaks discussed in Tables 2 and 3 and in text are indicated.

Table 2. Identification of peaks obtained from SSCP pattern of the yeast community (18S rDNA) by comparison with patterns of reference strains, clones, and isolates

Peaks (Fig. 2)	Assignment according to the elution of:		
	Reference and isolate SSCP patterns ^a	Corresponding clones showing more than 99% identity ^b	GenBank accession no
a	<i>K. lactis</i>	<i>K. lactis</i>	CSP508583
b	<i>K. marxianus</i>	<i>K. marxianus</i>	AF201299
c	<i>S. cerevisiae</i>		
d	<i>C. zeylanoides</i>	<i>C. zeylanoides</i>	AB013509.1
e		<i>D. hansenii</i> var. <i>Hansenii</i> <i>D. hansenii</i> var. <i>fabryii</i>	AB106349
f	<i>C. parapsilosis</i>		
g	<i>C. silvae</i>		
h	<i>S. unisporus</i>		

^aC., *Candida*, S., *Saccharomyces*, K., *Kluyveromyces*.

^bPercentage of identical nucleotides in the sequence retrieved from the clone and the sequences of the closest relatives found in GeneBank and RDP database.

Table 3. Evolution of the peak ratio of the SSCP patterns of the three cheeses (A, B, and C) at different times of ripening

Peaks	Species	Peak ratios (%) ^a											
		Cheese A				Cheese B				Cheese C			
		Stages of ripening				Stages of ripening				Stages of ripening			
		8 ^b	30	90	150	8	30	90	150	8	30	90	150
f	<i>C. parapsilosis</i>			4									
g	<i>C. silvae</i>	1.8							11				
a	<i>K. lactis</i>	23	22	13	30	28	43.4	33	5.0	11	60	66	48
b	<i>K. marxianus</i>	20	25	4.2	13	3.8	4.8	4.6		3.6	9.4	12	12
c	<i>S. cerevisiae</i>	16	2.8	5.6	3	4.7	9.9	18	0.5		11.5	15	13
d	<i>C. zeylanoides</i>	29	30	39	39	12	7.3	17	46	6.2	2.7	0.7	4
h	<i>S. unisporus</i>							14		38			
e	<i>D. hansenii</i>	11	20	37	12	53	35	9.7	47	30	17	6.9	23
		Total yeast counts (ufc/g) ^c											
		5.53	5.71	5.38	6.68	5.85	5.83	4.95	4.08	6.36	6.20	5.48	4.52

^aThe percent ratio was calculated as the area of one peak to the sum of the area of all peaks of the pattern $\times 100$.

^bStage of ripening expressed in days.

^cCounts on OGA medium + oxytetracyclin.

unisporus were only detected in some cheese at one time. The yeast community of cheese A had a high peak ratio for *C. zeylanoides*, which increased throughout the ripening period and had a peak for *C. parapsilosis*, which was detected only at 150 days. The yeast community of cheese B was dominated by *D. hansenii* after 8 days, but *C. zeylanoides* was dominant at 150 days. In this cheese, *S. unisporus* was not detected before 90 days. The high peak ratios for *C. silvae* and *S. unisporus* after 8 days and their lack of detection later in the ripening period characterized the yeast community of cheese C. During the ripening of this cheese *K. lactis* became dominant.

Dynamics of yeast populations as presented in Table 3 revealed changes in the relative abundance of DNA sequences in the yeast community. Cheese is a closed system and the number of species in the cheese core presumably does not increase throughout the ripening period. Thus, changes in peak ratios are not likely to be due to preferential PCR amplification of some sequences, but rather to changes in the balance of 18S rRNA genes of the species present. The changes of yeast population balance may result from many “events” such as development of one species or lysis of certain cells and degradation of DNA. The total count on OGA medium (Table 3) indicated that the yeast population level

decreased during ripening for two productions. This argues for a lysis of cells but does not provide information on the growth of each yeast species.

Comparison with Salers isolates identification

One hundred and two isolates were selected from the same three cheeses for identification based on phenotypic tests and 26S rRNA sequences. The majority of these partial 26S rRNA sequences were >98% similar to sequences in GenBank. Isolate identification by phenotypic testing and 26S rRNA gene sequencing was congruent for 92% of the isolates. Eight isolates assigned to *C. zeylanoides*, *C. parapsilosis*, and *K. lactis* based on their 26S rDNA sequences were difficult to identify by phenotypic testing because their carbon compound assimilation profiles were not consistent with those of any identified species.

This species inventory, based on isolate identification, was compared with that obtained by the cloning and SSCP methods (Table 4). Both approaches detected *K. lactis*, *K. marxianus*, *C. zeylanoides*, *S. cerevisiae* and *C. parapsilosis*. *K. lactis* and *K. marxianus* frequently are present in many types of cheese [7,10,23,25,27]. *C. zeylanoides* often is isolated from Portuguese ewe's cheese [19] and raw milk [5]. *S. cerevisiae* has been isolated from raw milk [5] and a number of cheeses including Mozzarella [7], Gouda [27], Cheddar [28], Camembert [23]. *C. parapsilosis* was found in Italian raw milk [5].

Table 4. Comparison of presence of different species in three cheeses A, B, and C (all times at once) detected using identification of 94 isolates or SSCP method

Species ^a	Cheese A		Cheese B		Cheese C	
	Isolates	SSCP	Isolates	SSCP	Isolates	SSCP
<i>Kl. lactis</i>	9	d ^b	12	d	9	d
<i>Kl. marxianus</i>	6	d	1	d	7	d
<i>S. cerevisiae</i>	2	d	3	d	7	d
<i>C. zeylanoides</i>	6	d	2	d	2	d
<i>D. hansenii</i>	3	d	3	d		d
<i>C. intermedia</i>	3	nd ^c	2	nd	3	nd
<i>C. silvae</i>		d			3	d
<i>P. guilliermondii</i>			1			
<i>C. parapsilosis</i>	5	d				
<i>C. tropicalis</i>					1	nd
<i>C. rugosa</i>			1	nd	1	nd
<i>S. unisporus</i>			1	d	1	nd
Total	34		26		34	

^aIdentification by 26S DNA sequencing in agreement with phenotypic tests.

^bDetected.

^cNon-detected.

The community composition determined by the SSCP method and by isolate identification differed in the detection of *D. hansenii*, *C. silvae* and *S. unisporus*. Neither *D. hansenii* nor *C. silvae* were detected by isolate identification but they were detected by SSCP. *D. hansenii* often is isolated from many types of cheeses. Several isolates were identified as *S. unisporus* whereas this species was seen in only on a few of the SSCP profiles. This species was found in Armada cheese [25]. We identified isolates of *C. intermedia*, *C. rugosa*, *C. tropicalis*, and *P. guilliermondii* from RDO Salers cheese by phenotypic identification that were not detected by SSCP. These species have been found sporadically in some dairy products: *P. guilliermondii* and *C. rugosa* were found in Italian raw milk [5]. *Y. lipolytica* previously was isolated from Feta cheese [10], Armada cheese [25], Gouda and Cheddar [27,28] but was not detected in Salers cheese by either method that we used.

The differences in community composition detected by each method reflect the strengths and limitations of each method. Community composition determined by identifying isolates depends on the number of isolates picked from the counting plates and the method used to select those isolates. If the number of isolates selected is too low, then the collection will be not representative of the population. Based on our results, this culture-dependent method is sensitive enough to detect species that are not detected by SSCP analysis. These results are consistent with those of Cocolin et al. [5], who found that more yeast species were detected in raw milk by a culture method than by DGGE. The difficulty in identifying less-common species by direct PCR-based molecular methods (SSCP, T(D)GGE) may be attributed to preferential amplification of dominant populations by PCR and/or to the bias of the SSCP detection system towards the most frequent sequences, i.e. the fluorescent signal of a less common sequence may not be easily distinguished from background noise. This can explain why *S. unisporus* was only detected after 90 days. At the beginning of ripening, it may be present but sub-dominant. But the SSCP method is more accurate and rapid to follow dominant population's dynamics.

This study shows the value of combining several methods (identification of isolates and direct molecular method such as SSCP) to describe cheese yeast communities. The use of multiple approaches to characterize populations has not commonly been used to analyze dairy products except by Cocolin et al. [5] who profiled yeast populations in raw milk by using both PCR–DGGE and isolate identification. To our knowledge, this is the first description of the dynamics of yeast communities in cheese by a culture-independent method.

The SSCP method quickly provides an image of the yeast population and permits the visual comparison of several samples. Thus PCR–SSCP is a powerful tool for tracking yeast community composition in fermented

dairy products. Each of the three cheeses had its own community composition profile. These results are consistent with those of Suzzi et al. [24] who also reported that yeast composition could change during the ripening period.

To complete this study, it would be necessary to determine the 18S rRNA SSCP profile, so as to assess the activity of yeasts during ripening. It will then be possible to integrate all these data with those obtained by Duthoit [8] on the bacterial community of this cheese, to better explain their sensorial properties.

Acknowledgements

We thank the Salers cheeses productions and the “Pôle Fromager AOC Massif Central” for their interest and Philip Rousseau-Cunningham for English proof-reading.

References

- [1] C. Andrighetto, E. Psomas, N. Tzanetakis, G. Suzzi, A. Lombardi, Randomly amplified polymorphic DNA (RAPD) PCR for the identification of yeasts isolated from dairy products, *Lett. Appl. Microbiol.* 30 (2000) 5–9.
- [2] S.F. Altschul, W. Gish, W. Miller, E.W. Myers, D.J. Lipman, Basic local alignment search tool, *J. Mol. Biol.* 215 (1990) 403–410.
- [3] J.A. Barnett, R.W. Payne, D. Yarrow, *Yeasts: Characteristics and identification*, second ed, Cambridge University Press, Cambridge, England, 1990.
- [4] C. Callon, L. Millet, M.C. Montel, Diversity of lactic acid bacteria isolated from AOC salers cheese, *J. Dairy Res.* 71 (2004) 231–244.
- [5] L. Cocolin, D. Aggio, M. Manzano, C. Cantoni, G. Comi, An application of PCR–DGGE analysis to profile the yeast populations in raw milk, *Int. Dairy J.* 12 (2002) 407–411.
- [6] F. Corpet, Multiple sequence alignment with hierarchical clustering, *Nucl. Acids Res.* 16 (1998) 10881–10890.
- [7] S. Coppola, G. Blaiotta, D. Ercolini, G. Moschetti, Molecular evaluation of microbial diversity occurring in different types of Mozzarella cheese, *J. Appl. Microbiol.* 90 (2001) 414–420.
- [8] F. Duthoit, C. Callon, L. Tessier, M.C. Montel, Relationships between sensorial characteristics and microbial dynamics in “Registered Designation of Origin” Salers cheese, *J. Appl. Microbiol.* 98 (2005) 1198–1208.
- [9] F. Duthoit, J.J. Godon, M.C. Montel, Bacterial community dynamics during production of Registered Designation of Origin Salers cheese as evaluated by 16S rRNA gene single strand conformation polymorphism analysis, *Appl. Environ. Microbiol.* 69 (2003) 3840–3848.
- [10] M.E. Fadda, S. Cosentino, M. Deplano, F. Palmas, Yeast populations in Sardinian feta cheese, *Int. J. Food Microbiol.* 69 (2001) 153–156.
- [11] G.H. Fleet, Yeasts in dairy products, *J. Appl. Bacteriol.* 68 (1990) 199–211.
- [12] S. Hernan-Gomez, J.C. Espinosa, J.F. Ubeda, Characterization of wine yeasts by temperature gradient gel electrophoresis (TGGE), *FEMS Microbiol. Lett.* 193 (2000) 45–50.
- [13] C.S. Hoffman, F. Winston, A ten minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*, *Gene* 57 (1987) 267–272.
- [14] C.P. Kurtzman, J.W. Fell, *The Yeasts: A Taxonomic Study*, fourth ed, Elsevier Science, Amsterdam, 1998.
- [15] P. Kurtzman, C.J. Robnett, Identification of clinically important ascomycetous yeasts based on nucleotide divergence in the 5′ end of the large-subunit (26S) ribosomal DNA gene, *J. Clin. Microbiol.* 35 (1997) 1216–1223.
- [16] M. Leclerc, C. Delbes, R. Moletta, J.J. Godon, Single strand conformation polymorphism of 16S rDNA Archae during start-up of an anaerobic digester, *FEMS Microbiol. Ecol.* 34 (2001).
- [17] H. Lee, Y.G. Zo, S.J. Kim, Non radioactive method to study genetic profiles of natural bacterial communities by PCR–single-strand-conformation polymorphism, *Appl. Environ. Microbiol.* 62 (1996) 3112–3120.
- [18] B.L. Maidak, J.R. Coles, C.T. Parker, G.M. Garrity, N. Larsen, B. Li, T.G. Lilburn, M.J. McCaughey, G.J. Olsen, R. Overbeek, S. Pramanik, T.M. Schmidt, J.M. Tiedje, C.R. Woese, A new version of the RDP (Ribosomal Database Project), *Nucl. Acids Res.* 27 (1999) 171–173.
- [19] S. Pereira-Dias, M.E. Potes, A. Marinho, M. Malfeito-Ferreira, V. Loureiro, Characterisation of yeast flora isolated from an artisanal Portuguese ewes’ cheese, *Int. J. Food Microbiol.* 60 (2000) 55–63.
- [20] S. Peters, S. Koschinsky, F. Schwieger, C.C. Tebbe, Succession of microbial communities during hot composting as detected by PCR–single-strand-conformation-polymorphism-based genetic profiles of small-subunit rRNA genes, *Appl. Environ. Microbiol.* 66 (2000) 930–936.
- [21] K.M. Petersen, P.L. Moller, L. Jespersen, DNA typing methods for differentiation of *Debaryomyces hansenii* strains and other yeasts related to surface ripened cheese, *Int. J. Food Microbiol.* 69 (2001) 11–24.
- [22] H. Prillinger, O. Molnar, F. Eliskases-Lechner, K. Lopandic, Phenotypic and genotypic identification of yeasts from cheese, *Antonie Van Leeuwenhoek* 75 (1999) 267–283.
- [23] R. Roostita, G.H. Fleet, The occurrence and growth of yeasts in Camembert and blue-veined cheeses, *Int. J. Food Microbiol.* 28 (1996) 393–404.
- [24] G. Suzzi, A. Lombardi, M.T. Lanorte, M. Caruso, C. Andrighetto, F. Gardini, Phenotypic and genotypic diversity of yeasts isolated from water-buffalo Mozzarella cheese, *J. Appl. Microbiol.* 88 (2000) 117–123.
- [25] M.E. Tornadijo, J.M. Fresno, R.M. Sarmiento, J. Carballo, Study of the yeasts during the ripening process of Armada cheeses from raw goat’s milk, *Lait* 78 (1998) 647–659.

- [26] C.Y. Turenne, E. Witwicki, D.J. Hoban, J.A. Karlowsky, A.M. Kabani, Rapid identification of bacteria from positive blood cultures by fluorescence-based PCR-single-strand conformation polymorphism analysis of the 16S rRNA gene, *J. Clin. Microbiol.* 38 (2000) 513–520.
- [27] J.J. Welthagen, B.C. Viljoen, Yeast profile in Gouda cheese during processing and ripening, *Int. J. Food Microbiol.* 41 (1998) 185–194.
- [28] M. Wojtatowicz, J. Chrzanowska, P. Juszczuk, A. Skiba, A. Gdula, Identification and biochemical characteristics of yeast microflora of Rokpol cheese, *Int. J. Food Microbiol.* 69 (2001) 135–140.
- [29] E. Zumstein, R. Moletta, J.J. Godon, Examination of two years of community dynamics in an anaerobic bioreactor using fluorescence polymerase chain reaction (PCR) single-strand conformation polymorphism analysis, *Environ. Microbiol.* 2 (2000) 69–78.