



AFLP-based genetic linkage maps of the blue mussel (*Mytilus edulis*)

Delphine Lallias, S. Lapegue, C. Hecquet, P. Boudry, A.R. Beaumont

► To cite this version:

Delphine Lallias, S. Lapegue, C. Hecquet, P. Boudry, A.R. Beaumont. AFLP-based genetic linkage maps of the blue mussel (*Mytilus edulis*). *Animal Genetics*, 2007, 38 (4), pp.340-349. 10.1111/j.1365-2052.2007.01611.x . hal-02660403

HAL Id: hal-02660403

<https://hal.inrae.fr/hal-02660403>

Submitted on 30 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AFLP-based genetic linkage maps of the blue mussel (*Mytilus edulis*)

D. Lallias^{*,†}, S. Lapègue^{*}, C. Hecquet^{*}, P. Boudry^{*} and A. R. Beaumont[†]

^{*}Ifremer, Laboratoire Génétique et Pathologie, Ronce-les-bains, 17390 La Tremblade, France. [†]School of Ocean Sciences, College of Natural Sciences, University of Wales, Bangor, Menai Bridge, Gwynedd LL59 5AB, UK

Summary

We report the construction of the first genetic linkage map in the blue mussel, *Mytilus edulis*. AFLP markers were used in 86 full-sib progeny from a controlled pair mating, applying a double pseudo-test cross strategy. Thirty-six primer pairs generated 2354 peaks, of which 791 (33.6%) were polymorphic in the mapping family. Among those, 341 segregated through the female parent, 296 through the male parent (type 1:1) and 154 through both parents (type 3:1). Chi-square goodness-of-fit tests revealed that 71% and 73% of type 1:1 and 3:1 markers respectively segregated according to Mendelian inheritance. Sex-specific linkage maps were built with MAPMAKER 3.0 software. The female framework map consisted of 121 markers ordered into 14 linkage groups, spanning 862.8 cM, with an average marker spacing of 8.0 cM. The male framework map consisted of 116 markers ordered into 14 linkage groups, spanning 825.2 cM, with an average marker spacing of 8.09 cM. Genome coverage was estimated to be 76.7% and 75.9% for the female and male framework maps respectively, rising to 85.8% (female) and 86.2% (male) when associated markers were included. Twelve probable homologous linkage group pairs were identified and a consensus map was built for nine of these homologous pairs based on multiple and parallel linkages of 3:1 markers, spanning 816 cM, with JOINMAP 4.0 software.

Keywords AFLP, blue mussel, genetic linkage map, *Mytilus edulis*.

Introduction

The blue mussel *Mytilus edulis* is a bivalve mollusc of major commercial importance with a worldwide production of around 1.5 million tons per year (FAO, Fisheries Department, 2002). Additionally, mussels are among the best studied species of the littoral and sublittoral communities and have been the focus of research into genetics, biochemistry, physiology and ecology (Gosling 1992). Despite their economical importance, most bivalves – including mussels – have not been domesticated like agricultural animals or crops and their production is mainly based on the collection of natural spat (i.e. juveniles). However, selective breeding programmes have been initiated in some bivalve species (e.g. Pacific oyster, <http://hmsc.oregonstate.edu/projects/mbp/>; green shell mussel, <http://www.cawthron.org.nz/aquaculture/selective-breeding.html>). In the blue

mussel, quantitative genetic studies of traits of economical importance for growth and length (Mallet *et al.* 1986; Stromgren & Nielsen 1989) suggest that significant improvement could be achieved by selective breeding. In this context, the development of genetic and genomic tools are likely to contribute to the development of selective breeding programmes, and, more generally, to improve knowledge about the genome of this species of aquacultural and ecological importance. Genetic linkage maps based on polymorphic markers such as AFLPs, RAPDs and microsatellites have been generated in several aquaculture species such as salmon (Moen *et al.* 2004), tilapia (Kocher *et al.* 1998) or shrimp (Li *et al.* 2006). In bivalves, genetic maps have been established in the Pacific oyster *Crassostrea gigas* (Hubert & Hedgecock 2004; Li & Guo 2004), the Eastern oyster *Crassostrea virginica* (Yu & Guo 2003) and the Zhikong scallop *Chlamys farreri* (Wang *et al.* 2004; Li *et al.* 2005; Wang *et al.* 2005). Such maps represent frameworks that enable the identification and localisation of quantitative trait locus (QTL) (e.g. Yu & Guo 2006) with the final aim of achieving genetic improvement through marker-assisted selection (Liu & Cordes 2004).

Only seven microsatellites have so far been published for the blue mussel (Presa *et al.* 2002) and, although more

Address for correspondence

A. R. Beaumont, School of Ocean Sciences, College of Natural Sciences, University of Wales, Bangor, Menai Bridge, Gwynedd LL59 5AB, UK.
E-mail: a.r.beaumont@bangor.ac.uk

Accepted for publication 26 March 2007

than 20 allozyme markers have been developed for the mussel, their relatively low polymorphism makes them unsuitable for extensive mapping (Beaumont 1994). Therefore, AFLP markers (Vos *et al.* 1995) were chosen as they require no preliminary knowledge of the genome, are highly reproducible (Jones *et al.* 1998) and can generate relatively quickly a high number of markers dispersed across the 14 pairs of chromosomes in the mussel genome.

Materials and methods

Mapping family

A full-sib mapping family was produced from two wild mussels collected from the Menai Strait, Wales, UK, as there has been no domestication of mussels and neither homozygous nor selected lines are available. The methods used for gamete release, fertilisation and larval development were essentially as described by Beaumont *et al.* (1988). Larvae were reared in 2l plastic beakers and fed 50 cells/ μ l of a 3:1 mixture *Pavlova lutheri* and *Rhinomonas reticulata*. Filtered (1 μ m) and UV-light treated water was changed three times a week when food was added. Ready-to-settle larvae were held on 80- μ m sieves in a downwelling system (Utting & Spencer 1991) to allow metamorphosis, and spat were fed a mixture of *P. lutheri*, *Chaetoceros calcitrans* and *Isochrysis galbana* clone T-Iso. Sieve mesh sizes were increased as mussel juveniles grew in size, and mussels were transferred to IFREMER, La Tremblade, France when 2 months old for further growing. Several full-sib families were produced, but only one was randomly chosen for study. DNA was extracted from 86 20-month-old F₁ mussels.

DNA extraction

DNA was extracted from gill tissue using a chloroform extraction followed by purification with the Wizard[®] DNA Clean-Up System (Promega), according to Wilding *et al.* (2001). Quality and concentration of DNA was assessed using a spectrophotometer and by running a small sample on a 2% agarose gel. High-quality extracted DNA was adjusted to a concentration of 100 μ g/ml.

AFLP analysis

AFLP analysis was performed by using a modified version of Vos *et al.* (1995), following Wilding *et al.*'s (2001) protocol, but digestion and ligation were achieved in the same mix and incubated 16 h at 16 °C. Electrophoresis and data collection were carried out on an ABI 3100-Avant (Applied Biosystems). Electrophoresis parameters were set at injection for 15 s at 15 kV, running for 25 min at 15 kV and 60 °C, with POP4 polymer. Repeatability of the technique was checked by comparing the band pattern of four replicates obtained independently (four different DNA

extractions: two gill, two muscle, four different AFLP amplifications (primer pair A1), performed on different days) on the same 20 samples of *M. edulis*. Results of this trial showed that consistent band patterns were obtained.

Thirty-six AFLP primer pairs were genotyped in the mapping family (Table S1). Two negative controls were included in each PCR reaction to detect any potential contamination. Data were analysed with GENEMAPPER[®] software version 3.7 and individuals were scored for the presence [A] or absence [a] of the amplified AFLP fragment.

Distortion of segregation ratios

Two kinds of segregating AFLP markers could be detected. Type 1:1 markers had one parent heterozygous for the band, the other parent homozygous for no band and the F₁ progeny were expected to segregate 1:1 (band:no band). Type 3:1 markers corresponded to AFLPs where both parents were heterozygous for the band and the progeny were expected to segregate 3:1 (band:no band). A chi-square goodness-of-fit test for the 1:1 or 3:1 segregation ratios was applied to each locus. All distorted markers ($P < 0.05$) were excluded from further linkage analysis.

Establishment of sex-specific framework linkage maps: MAPMAKER 3.0 software

MAPMAKER 3.0 software (Lander *et al.* 1987) was used to build sex-specific linkage maps, based on the markers segregating 1:1 in a double pseudo-test cross (F₂ backcross model, Aa = H, aa = A) (Grattapaglia & Sederoff 1994). Each dataset was duplicated and recoded to allow the detection of markers linked in repulsion phase ('r' added at the end of their names; e.g. marker A1f123r was the recoded marker of marker A1f123). Linkage groups were determined with the GROUP command of MAPMAKER, conducted at LOD score ≥ 4.0 and genetic distance ≤ 37.5 cM. Once linkage groups were determined, the markers were ordered by the COMPARE command for a limited number of markers ($n \leq 9$), or otherwise by the THREE POINT and ORDER commands. After the ordering of markers within each linkage group, the RIPPLE command allowed the testing of robustness of the map obtained. Markers that presented a conflict in map position (several map positions possible, with a small difference of LOD score) were placed as associated markers. The ERROR DETECTION command (Lincoln & Lander 1992) was on during all the analyses described above to detect genotyping errors. Map distances in centiMorgans were calculated using Kosambi's mapping function (Kosambi 1944) and linkage groups were drawn with MAPCHART software (Voorrips 2002).

Establishment of a consensus map: JOINMAP 4.0 software

Male and female maps based on 1:1 and 3:1 segregating AFLP markers and a consensus map were achieved using

JOINMAP 4.0 software (Van Ooijen 2006). Male and female datasets were treated independently, as a population type CP (composite). First, the two parental maps based on 1:1 and 3:1 markers were built. Then, homologous pairs of linkage groups were identified by multiple and parallel linkages of markers, i.e. several markers ordered in the same order in both parental maps. Only 3:1 markers that did not contradict the mapping order of framework markers, previously established with MAPMAKER, were retained for the establishment of a consensus map. Recombination rates were converted into genetic distances (in cM) using Kosambi's mapping function and linkage groups were drawn with MAPCHART software (Voorrips 2002).

Genome length and map coverage

Average marker spacing of the framework map was calculated by dividing the total length of the map by the number of intervals. In the same way, the average marker spacing for each linkage group was calculated by dividing the length of each linkage group by the number of intervals on that linkage group. The expected length of the genome was estimated using method 4 of Chakravarti *et al.* (1991), based on framework markers alone and then with framework and associated markers. Observed genome coverage estimates were determined by dividing the observed genome length (total length in cM of all the linkage groups) by the expected length of the genome. Two observed genome coverage estimates were computed, whether or not associated markers were taken into account.

Results

The 36 primer pairs, screened for 86 F₁ progeny and their two parents, generated a total of 2354 peaks, averaging 65 peaks per pair. The average number of segregating markers (among the two parents, including both types of markers) was 791, or 22 per primer pair, corresponding to 33.6% of polymorphic peaks. Among the 791 polymorphic markers in the mapping family, 341 were segregating through the female parent, 296 through the male parent and 154 through both parents. Chi-square analysis indicated that 243 (71.3%) and 210 (70.9%) of the markers segregated according to the expected 1:1 Mendelian ratio in the female and male respectively and that 112 (72.7%) markers segregated according to the expected 3:1 Mendelian ratio. Distorted markers ($P < 0.05$) were discarded from further linkage analysis.

Sex-specific linkage maps

The female framework map established with MAPMAKER 3.0, based on the 243 AFLP markers segregating through the female parent only, consisted of 121 markers (49.8%). Seven markers were not linked to the framework map

(2.9%). Additionally, 115 markers were linked to the framework map with a LOD score of 4.0 but not placed accurately and were therefore considered as 'associated markers'. Associated markers were located beside their closest framework marker (Table S2). Fourteen linkage groups were identified for the female map covering 862.8 cM (Fig. 1). The sizes of the linkage groups ranged from 9.5 to 101.5 cM. The number of framework markers per linkage group varied from 2 to 16 and the number of associated markers from 0 to 23. The average distance between 2 framework loci ranged from 4.09 (G10F) to 19.5 cM (G12F), with an average spacing of 8.06 cM. The maximum interval of the female map was 32.9 cM (G2F) (Table 1). Some clusters of AFLPs could be observed, containing from two to seven markers. A single linkage group could contain up to four clusters (e.g. G1F). The estimated genome length was 1125.3 cM. The observed coverage was therefore 76.7% for the female framework map. When associated markers were considered, the estimated genome length was 1006.0 cM and genome coverage became 85.8% for the female map.

The male framework map established with MAPMAKER 3.0 was based on the 210 AFLP markers segregating through the male parent only. The resulting map consisted of 116 framework markers (55.2%). Six markers were not linked to the framework map (2.9%). Additionally, 88 markers were placed as associated markers (Table S3). Fourteen linkage groups were set up for the male map covering 825.2 cM (Fig. 2). The sizes of the linkage groups ranged from 20.3 to 86.5 cM. The number of framework markers per linkage group varied from 3 to 20 and the number of associated markers from 1 to 14. The average distance between two framework loci ranged from 3.08 (G13M) to 15.4 cM (G6M), with an average spacing of 8.09 cM. The largest interval varied from 8.2 (G13M) to 37.6 cM (G1M) (Table 1). Some clusters of AFLPs could be observed, containing from two to seven markers. A single linkage group could contain up to six clusters (G8M). All framework markers were mapped with a LOD score of 4.0 except marker *A6f121r* of G5M, which was linked to this group with a LOD score of 2.79. For the male map, the estimated genome length was 1087.1 cM and observed coverage was 75.9%. Including associated markers, the estimated genome length was reduced to 957.6 cM with genome coverage of 86.2%.

Marker distribution

There was a random distribution among linkage groups of all markers generated by the three different EcoRI primers. A contingency chi-square test for the 14-groups \times 3-EcoRI primers was not significant for either female or male ($\chi^2 = 12.720$ or 24.604, 22 d.f., $P = 0.316$ or 0.941 respectively). The female and male framework maps (Figs 1 & 2) revealed a high number of clusters, 43 on the female and 36 on the male map. Uneven distribution of markers means that gaps remain to be filled because both maps show




Figure 1 AFLP linkage map of the blue mussel *Mytilus edulis*: female map obtained with MAPMAKER 3.0, including 121 framework markers for a total of 863 cM. AFLP markers are labelled with the primer pair name followed by the letter 'f' (for fragment) and a 3-digit fragment size in base pairs. Markers are indicated on the right and absolute positions on the left (in Kosambi cM). Numbers in brackets on the right of locus name correspond to number of associated markers (linked but unplaced). The letter 'r' at the end of some AFLPs refers to recoded markers.

intervals spanning more than 20 cM. The assumption of a random distribution of AFLP markers across the genome was tested using Spearman correlation coefficients and chi-square test for departure from a Poisson distribution. Spearman correlation coefficients (r_s) between genetic length and number of markers per group were 0.481 for the male ($P > 0.05$) and 0.635 for the female ($P < 0.05$). However, when a single outlier linkage group was removed from the male dataset, the correlation became significant ($P < 0.05$). Therefore, in spite of observed clusters, AFLP markers generally tended to be randomly distributed in the linkage maps.

Observed and expected distributions of AFLPs were compared over 20 cM intervals in female and male framework maps. A chi-square test for departure from a Poisson distribution was computed. The mean of the Poisson distribution was set up to the mean number of markers per 20 cM interval length: 2.61 for the female and 2.8 for the male maps. No significant departure from the Poisson distribution was observed for the female ($\chi^2 = 13.22$, 7 d.f., $P = 0.067$). However, this goodness-of-fit test was highly

significant for the male ($\chi^2 = 28.63$, 7 d.f., $P < 0.001$), mostly due to three intervals of 20 cM containing eight markers. This confirms that clustering of AFLPs was more important in the male framework map, with the presence of a few dense clusters of markers (containing up to seven markers).

Preliminary consensus map

Twelve probable homologous linkage groups were identified and for nine of them, a consensus map was established based on at least three markers of type 3:1 exhibiting multiple and parallel linkages (Fig. 3). Up to four 3:1 markers were used to build a consensus group. These consensus groups were named according to the names of the groups they derived from, e.g. consensus group G10F_G11M_comb issued from the joining of groups G10F of the female map and G11M of the male map. For three of the 12 probable homologous groups, a consensus map was difficult to construct (Fig. 4). For example, the homology of groups G7F and G14M was based on a single marker

Table 1 Length, number of markers (framework and associated), average spacing and largest interval of linkage groups of the female and male maps established with MAPMAKER 3.0.

Linkage group	Length (cM)	No. of framework markers	No. of associated markers	Average spacing (cM)	Largest interval (cM)
Female					
G1F	101.5	15	23	7.25	27.8
G2F	101.3	11	12	10.13	32.9
G3F	83.3	10	13	9.25	21.9
G4F	76.0	10	5	8.40	28.3
G5F	75.2	10	8	8.35	31.0
G6F	69.9	7	3	11.65	26.7
G7F	66.7	9	5	8.34	28.9
G8F	65.8	8	4	9.40	22.3
G9F	61.6	9	11	7.70	22.5
G10F	61.3	16	11	4.09	14.3
G11F	60.2	10	11	6.69	25.4
G12F	19.5	2	0	19.50	19.5
G13F	11.0	2	5	11.00	11.0
G14F	9.5	2	4	9.50	9.5
Total	862.8	121	115	8.06	32.9
Male					
G1M	86.5	13	9	7.21	37.6
G2M	84.9	10	10	9.43	22.1
G3M	79.5	7	14	13.25	30.8
G4M	77.8	8	3	11.11	23.7
G5M	74.9	10	5	8.32	35.2
G6M	61.6	5	10	15.40	23.9
G7M	61.2	6	5	12.24	24.1
G8M	60.2	20	7	3.17	16.9
G9M	57.6	7	5	9.60	12.9
G10M	57.0	8	5	8.14	25.2
G11M	53.0	5	5	13.25	20.4
G12M	26.4	5	1	6.60	16.0
G13M	24.3	9	7	3.08	8.2
G14M	20.3	3	2	10.15	11.5
Total	825.2	116	88	8.09	37.6

(B1f123*). The homology of groups G11F and G12M was based on the parallel and multiple linkage of three markers (B4f222*, E10f66* and E10f78*) but the alignment of these two groups according to these three markers indicated that G12M could be homologous to only the terminal part of G11F, making the establishment of a consensus map difficult and potentially unreliable. Finally, the male group G6M seemed to have two potential homologies in the female map: G13F and G7Fpartial, assessed by three and two markers of type 3:1 respectively. Despite the mapping of a few 3:1 markers, no clear homologues could be identified for the male groups G3M and G10M nor for the female groups G8F and G12F.

The observed genome lengths obtained for the female, the male and the consensus maps established with MAPMAKER and JOINMAP were very similar: 863 and 825 cM for the female and male maps (MAPMAKER), 871 and 799 cM for the

female and male maps (JOINMAP) and 816 cM for the consensus map.

Discussion

In our mapping family of *M. edulis*, segregation distortion (χ^2 , $P < 0.05$) averaged 29% for the type 1:1 markers and 27% for the type 3:1 markers. The observed numbers of distorted markers were 98 for the female 1:1 markers, 86 for the male 1:1 markers and 42 for the 3:1 markers and were higher than the expected numbers by chance only (17, 15 and 8 respectively at $\alpha = 5\%$).

High segregation distortion could be caused by technical artefacts in genotyping such as size homoplasy (i.e. AFLP fragments showing the same size but belonging to different loci). Incomplete enzyme digestion and/or inefficient PCR represent another technical artefact, leading towards null homozygous AFLP genotypes. In our study, around 60% of the distorted markers were deficient for homozygous null genotypes, ruling out technical artefacts as the main source of non-Mendelian segregation. Finally, the relatively high proportion of distorted markers (towards a homozygote deficiency) could be explained by linkage of markers with lethal or deleterious genes in recessive state that cause genotype-dependant mortalities. A high genetic load has previously been reported in bivalves (McGoldrick & Hedgecock 1997; Bierne *et al.* 1998; Launey & Hedgecock 2001) and is therefore to be the most likely hypothesis explaining our results.

To our knowledge, maps presented in our study represent the first genetic linkage maps established in the blue mussel *M. edulis*. Despite their preliminary nature, these male and female maps offer a good representation of the blue mussel genome. First, both maps contain 14 linkage groups, which correspond to the haploid number of chromosome of this species (Thiriot-Quievreux 1984). Secondly, total map length observed in this study is similar to the theoretical genetic length based on 1.0–1.3 crossing over per chromosome. The observed genetic length was 825.2 cM for the male map and 862.8 cM for the female map. Work on the Pacific and Eastern oysters ($2n = 20$) revealed an average number of chiasmata per chromosome of ~ 1.1 – 1.2 (X. Guo unpublished data cited in Li & Guo 2004). Based on these data, assuming a hypothetical range of 1.0–1.3 chiasmata per chromosome for *M. edulis*, the theoretical map length should range 700 to 910 cM (1.0 or $1.3 \times 50 \text{ cM} \times 14$ chromosomes). The observed total genetic length for both maps in our study falls into that range. Moreover, expected genome lengths estimated in this study (957–1006 cM) were 5–36% longer than the theoretical length based on cytological studies (700–910 cM). The discrepancy between both estimates (expected and theoretical genome lengths) is smaller than that identified in two separate studies on *C. gigas*: 42–99% (Hubert & Hedgecock 2004) and 32–79% (Li & Guo 2004).




Figure 2 AFLP linkage map of the blue mussel *Mytilus edulis*: male map obtained with MAPMAKER 3.0, including 116 framework markers for a total of 825 cM. AFLP markers are labelled with the primer pair name followed by the letter 'f' (for fragment) and a 3-digit fragment size in base pairs. Markers are indicated on the right and absolute positions on the left (in Kosambi cM). Numbers in brackets on the right of locus name correspond to number of associated markers (linked but unplaced). The letter 'r' at the end of some AFLPs refers to recoded markers.

The ratios of longest to shortest linkage groups are 10.7:1 and 4.3:1 in the female and male maps respectively. These ratios are greater than the cytological ratio (length of chromosome 1 to length of chromosome 14) observed in several karyological studies in *M. edulis*: 2:1 (Thiriot-Quiévreux 1984); 1.74–1.86:1 (Insua *et al.* 1994). This suggests that gaps remain to be filled and that more markers should be added for a better coverage of the genome. However, Hubert & Hedgecock (2004) reported slightly higher ratios in *C. gigas*, 7.6:1 in the female and 13.7:1 in the male, compared with the 2:1 cytological ratio (Thiriot-Quiévreux 1984). Therefore, the discrepancy observed between these two ratios is similar in *M. edulis* and *C. gigas*.

Genome coverage estimated for both maps is relatively good, at 76.7% and 75.9% for framework female and male maps respectively. Genome coverage increased to around 86% for both maps when associated markers (linked but not mapped) were taken into account. These estimates are similar to the one established in *C. gigas*: 70–79% (microsatellite markers; Hubert & Hedgecock 2004) or 81–92% (AFLPs; Li & Guo 2004); and in *C. virginica*: 70–84% (AFLPs; Yu & Guo 2003). Also, only 3% of markers were

unlinked to any other marker in both maps, another indicator that framework maps established in *M. edulis* cover a good proportion of the genome.

Our study reported a similarity of map lengths between sexes, in the range 800–870 cM, obtained with two different software (MAPMAKER 3.0 and JOINMAP 4.0). The linear relationship between linkage distance and recombination rate implies that recombination rates in males and females could be similar in *M. edulis*. Similar recombination rates were reported between males and females in *Penaeus japonicus* (Li *et al.* 2003). However, several studies reported large sex-specific differences in recombination rates, generally towards higher recombination rates in females in rainbow trout (Sakamoto *et al.* 2000), in *C. virginica* (Yu & Guo 2003) and in *P. monodon* (Wilson *et al.* 2002). Most of these studies showed congruence between sex-specific recombination rates and genetic distances, with a longer genetic distance reported in the sex exhibiting higher recombination rates. Nevertheless, caution must be taken in the inference of sex-specific recombination frequency from genetic distance because some studies reported sex-discrepancy between genetic map lengths due to the number of informative




Figure 3 Consensus map established in the blue mussel *Mytilus edulis*, in a mapping family including two parents and 86 F₁ progeny using JOINMAP 4.0 software. The consensus map is based on the finding of nine homologous pairs of linkage groups. Homologous markers are displayed in bold and underlined, ending with an asterisk (*). AFLP markers are labelled with the primer pair name followed by the letter 'f' (for fragment) and a 3-digit fragment size in base pairs. Markers are indicated on the right and absolute positions on the left (in Kosambi cM).




Figure 4 Probable homologies between female and male groups assessed by the mapping of 3:1 markers in *Mytilus edulis* using JOINMAP 4.0 software. Three pairs of likely homologous are represented for which no consensus map could be established. Pairs of homologous markers are displayed in bold and underlined, ending with an asterisk (*). AFLP markers are labelled with the primer pair name followed by the letter 'f' (for fragment) and a 3-digit fragment size in base pairs. Markers are indicated on the right and absolute positions on the left (in Kosambi cM).

markers (Agresti *et al.* 2000). However, in our study, the number of markers mapped in the female (121) and male (116) maps was similar so it is likely that recombination frequencies between sexes are similar in *M. edulis*. This could be confirmed by pairwise comparisons of average spacing between markers common to male and female maps, particularly after the addition of codominant markers.

Clustering of AFLPs is a common feature of AFLP-based genetic maps and has been observed in several species, e.g. in maize (Castiglioni *et al.* 1999), rainbow trout (Young *et al.* 1998), tilapia (Agresti *et al.* 2000) or channel catfish (Liu *et al.* 2003). Clustering of AFLPs could result from the non-random distribution of enzymatic restriction sites across the genome and therefore indirectly from the choice of enzymes of restriction used. In this study, EcoRI and MseI were used for digesting the DNA. EcoRI and MseI restriction sites are relatively AT-rich and so could reflect the variation in GC content among chromosomal regions (Yu & Guo 2003). Contrary to studies on oysters, clustering of AFLPs seems to occur more towards centromeric regions than telomeric regions in the blue mussel (G1M, G3M or G10M). As chromosomes of *M. edulis* are mostly metacentric or submetacentric (Thiriot-Quievreux 1984; Insua *et al.* 1994), this clustering could correspond to centromeric suppression of recombination, associated with heterochromatin (Tanksley *et al.* 1992).

The large intervals (>20 cM), observed in both maps, could be due to the medium-density of the maps obtained and it is expected that adding markers should reduce those

gaps. Alternatively, they could correspond to hot-spot regions of recombination in the genome.

The consensus map presented in this study, even though incomplete, shows the feasibility of an AFLP-based mapping strategy in an undomesticated marine species. However, to increase the accuracy of the consensus map, more 3:1 markers should be scored, to base the consensus map on more than three markers per group and markers should be chosen to be more evenly spaced throughout each linkage group. More importantly, adding codominant markers such as microsatellites, SNPs or ESTs (type I markers), serving as anchor loci between the two parental maps, will increase the accuracy of the consensus map as well as its portability in the context of QTL mapping. Efforts were made in our study to use the published microsatellite loci (Presa *et al.* 2002) but we could not achieve reliable results. The combination of dominant (AFLPs or RAPDs) and codominant (microsatellites) markers proved to be very useful for the construction of a consensus map in rainbow trout (Nichols *et al.* 2003), tilapia (Kocher *et al.* 1998), zebrafish (Johnson *et al.* 1996) and common carp (Sun & Liang 2004). To facilitate the mapping of type I markers, the DNA from the mapping family can be made available to the research community.

Future work could include the production of additional mapping families involving crosses between *M. edulis* and *M. galloprovincialis* that will be useful for mapping the numerous type I markers already developed in *M. galloprovincialis* (Venier *et al.* 2003, 2006). Such families could be used to

investigate QTL affecting production and life history traits that differ between the two taxa.

Acknowledgements

This work was funded by INTERREG IIIB (Atlantic Aquaculture Arc Group, AAAG). We particularly thank Dr Christophe Plomion and Dr Catherine Bodenes (INRA, France) for their useful advice on MAPMAKER and JOINMAP software. Thanks are due to Dr Richard Braithwaite and Serge Heurtebise for their help during the production of families of mussels.

References

- Agresti J.J., Seki S., Cnaani A., Poompuang S., Hallerman E.M., Umel N., Hulata G., Gall G.A.E. & May B. (2000) Breeding new strains of tilapia: development of an artificial center of origin and linkage map based on AFLP and microsatellite loci. *Aquaculture* **185**, 43–56.
- Beaumont A.R. (1994) Linkage studies in *Mytilus edulis*, the mussel. *Heredity* **72**, 557–62.
- Beaumont A.R., Beveridge C.M., Barnet E.A., Budd M.D. & Smyth-Chamosa M. (1988) Genetics studies of laboratory reared *Mytilus edulis*. I. Genotype specific selection in relation to salinity. *Heredity* **61**, 389–400.
- Bierne N., Launey S., Naciri-Graven Y. & Bonhomme F. (1998) Early effect of inbreeding as revealed by microsatellite analyses on *Ostrea edulis* larvae. *Genetics* **148**, 1893–906.
- Castiglioni P., Ajmone-Marsan P., Van Wijk R. & Motto M. (1999) AFLP markers in a molecular linkage map of maize: codominant scoring and linkage group distribution. *Theoretical and Applied Genetics* **99**, 425–31.
- Chakravarti A., Lasher L.K. & Reefer J.E. (1991) A maximum likelihood method for estimating genome length using genetic linkage data. *Genetics* **128**, 175–82.
- F.A.O. Fisheries Department (2002) *Aquaculture production: quantities*. Fishstart Plus Database, Rome.
- Gosling E.M. (1992) *The Mussel Mytilus: Ecology, Physiology, Genetics, and Culture*. Elsevier, Amsterdam.
- Grattapaglia D. & Sederoff R. (1994) Genetic linkage maps of *Eucalyptus grandis* and *Eucalyptus urophylla* using a pseudo-test-cross: mapping strategy and RAPD markers. *Genetics* **137**, 1121–37.
- Hubert S. & Hedgecock D. (2004) Linkage maps of microsatellite DNA markers for the Pacific oyster *Crassostrea gigas*. *Genetics* **168**, 351–62.
- Insua A., Labat J.P. & Thiriot-Quiévreux C. (1994) Comparative analysis of karyotypes and nucleolar organizer regions in different populations of *Mytilus trossulus*, *Mytilus edulis* and *Mytilus galloprovincialis*. *Journal of Molluscan Studies* **60**, 359–60.
- Johnson S.L., Gates M.A., Johnson M., Talbot W.S., Horne S., Baik K., Rude S., Wong J.R. & Postlethwait J.H. (1996) Centromere-linkage analysis and consolidation of the zebrafish genetic map. *Genetics* **142**, 1277–88.
- Jones C.J., Edwards K.J., Castaglione S., Winfield M.O., Sala F., Van Der Wiel C., Vosman B., Matthes M., Daly A. & Brettschneider R. (1998) Reproducibility testing of AFLPs by a network of European laboratories. In: *Molecular Tools for Screening Biodiversity* (Ed. by A. Karp, P.G. Isaac & D.S. Ingram), pp. 183–92. Chapman and Hall, London.
- Kocher T.D., Lee W.J., Sobolewska H., Penman D. & McAndrew B. (1998) A genetic linkage map of a cichlid fish, the tilapia (*Oreochromis niloticus*). *Genetics* **148**, 1225–32.
- Kosambi D.D. (1944) The estimation of map distances from recombination values. *Annals of Eugenics* **12**, 172–5.
- Lander E.S., Green P., Abrahamson J., Barlow A., Daly M.J., Lincoln S.E. & Newburg L. (1987) MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* **1**, 174–81.
- Launey S. & Hedgecock D. (2001) High genetic load in the Pacific oyster *Crassostrea gigas*. *Genetics* **159**, 255–65.
- Li L. & Guo X. (2004) AFLP-based genetic linkage maps of the Pacific oyster *Crassostrea gigas* Thunberg. *Marine Biotechnology* **6**, 26–36.
- Li Y., Byrne K., Miggiano E., Whan V., Moore S., Keys S., Crocos P., Preston N. & Lehnert S. (2003) Genetic mapping of the kuruma prawn *Penaeus japonicus* using AFLP markers. *Aquaculture* **219**, 143–56.
- Li L., Xiang J., Liu X., Zhang Y., Dong B. & Zhang X. (2005) Construction of AFLP-based genetic linkage map for Zhikong scallop, *Chlamys farreri* Jones et Preston and mapping of sex-linked markers. *Aquaculture* **245**, 63–73.
- Li Z., Li J., Wang Q., He Y. & Liu P. (2006) AFLP-based genetic linkage map of marine shrimp *Penaeus (Fenneropenaeus) chinensis*. *Aquaculture* **261**, 463–72.
- Lincoln S.E. & Lander E.S. (1992) Systematic detection of errors in genetic linkage data. *Genomics* **14**, 604–10.
- Liu Z.J. & Cordes J.F. (2004) DNA marker technologies and their applications in aquaculture genetics. *Aquaculture* **238**, 1–37.
- Liu Z., Karsi A., Li P., Cao D. & Dunham R. (2003) An AFLP-based genetic linkage map of channel catfish (*Ictalurus punctatus*) constructed by using an interspecific hybrid resource family. *Genetics* **165**, 687–94.
- Mallet A.L., Freeman K.R. & Dickie L.M. (1986) The genetics of production characters in the blue mussel *Mytilus edulis*. I. A preliminary analysis. *Aquaculture* **57**, 133–40.
- McGouldrick D.J. & Hedgecock D. (1997) Fixation, segregation and linkage of allozyme loci in inbred families of the Pacific oyster *Crassostrea gigas* (Thunberg): implications for the causes of inbreeding depression. *Genetics* **146**, 321–34.
- Moen T., Hoyheim B., Munck H. & Gomez-Raya L. (2004) A linkage map of Atlantic salmon (*Salmo salar*) reveals an uncommonly large difference in recombination rate between the sexes. *Animal Genetics* **35**, 81–92.
- Nichols K.M., Young W.P., Danzmann R.G. *et al.* (2003) A consolidated linkage map for rainbow trout (*Oncorhynchus mykiss*). *Animal Genetics* **34**, 102–15.
- Presa P., Pérez M. & Diz A.P. (2002) Polymorphic microsatellite markers for blue mussels (*Mytilus* spp.). *Conservation Genetics* **3**, 441–3.
- Sakamoto T., Danzmann R.G., Gharbi K. *et al.* (2000) A microsatellite linkage map of rainbow trout (*Oncorhynchus mykiss*) characterized by large sex-specific differences in recombination rates. *Genetics* **155**, 1331–45.

- Stromgren T. & Nielsen M.V. (1989) Heritability of growth in larvae and juveniles of *Mytilus edulis*. *Aquaculture* **80**, 1–6.
- Sun X. & Liang L. (2004) A genetic linkage map of common carp (*Cyprinus carpio* L.) and mapping of a locus associated with cold tolerance. *Aquaculture* **238**, 165–72.
- Tanksley S.D., Ganal M.W., Prince J.P. *et al.* (1992) High density molecular linkage maps of the tomato and potato genomes. *Genetics* **132**, 1141–60.
- Thiriot-Quiévreux C. (1984) Chromosome analysis of three species of *Mytilus* (Bivalvia: Mytilidae). *Marine Biology Letter* **5**, 265–73.
- Utting S.D. & Spencer B.E. (1991) *The Hatchery Culture of Bivalve Mollusc Larvae and Juveniles. Laboratory Leaflet Number 68*. Ministry of Agriculture, Fisheries and Food Directorate of Fisheries Research, Lowestoft.
- Van Ooijen J.W. (2006) *JoinMap 4, Software for the Calculation of Genetic Linkage Maps in Experimental Populations*. Wageningen, Netherlands, Kyazma, B.V.
- Venier P., Pallavicini A., De Nardi B. & Lanfranchi G. (2003) Towards a catalogue of genes transcribed in multiple tissues of *Mytilus galloprovincialis*. *Gene* **314**, 29–40.
- Venier P., De Pitta C., Pallavicini A., Marsano F., Varotto L., Romualdi C., Dondero F., Viarengo A. & Lanfranchi G. (2006) Development of mussel mRNA profiling: can gene expression trends reveal coastal water pollution? *Mutation Research* **602**, 121–34.
- Voorrips R.E. (2002) MapChart: software for the graphical presentation of linkage maps and QTLs. *The Journal of Heredity* **93**, 77–8.
- Vos P., Hogers R., Bleeker M. *et al.* (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research* **23**, 4407–14.
- Wang S., Bao Z., Pan J., Zhang L., Yao B., Zhan A., Bi K. & Zhang Q. (2004) AFLP linkage map of an intraspecific cross in *Chlamys farreri*. *Journal of Shellfish Research* **23**, 491–9.
- Wang L., Song L., Chang Y., Xu W., Ni D. & Guo X. (2005) A preliminary genetic map of Zhikong scallop (*Chlamys farreri* Jones et Preston 1904). *Aquaculture Research* **36**, 643–53.
- Wilding C.S., Butlin R.K. & Grahame J. (2001) Differential gene exchange between parapatric morphs of *Littorina saxatilis* detected using AFLP markers. *Journal of Evolutionary Biology* **14**, 611–9.
- Wilson K., Li Y., Whan V. *et al.* (2002) Genetic mapping of the black tiger shrimp *Penaeus monodon* with amplified fragment length polymorphism. *Aquaculture* **204**, 297–309.
- Young W.P., Wheeler P.A., Coryell V.H., Keim P. & Thorgaard G.H. (1998) A detailed linkage map of rainbow trout produced using doubled haploids. *Genetics* **148**, 839–50.
- Yu Z. & Guo X. (2003) Genetic linkage map of the eastern oyster *Crassostrea virginica* Gmelin. *Biological Bulletin* **204**, 327–38.
- Yu Z. & Guo X. (2006) Identification and mapping of disease-resistance QTLs in the eastern oyster, *Crassostrea virginica* Gmelin. *Aquaculture* **254**, 160–70.

Supplementary Material

The following supplementary material is available for this article online from <http://www.blackwell-synergy.com/doi/full/10.1111/j.1365-2052.2007.01611.x>

Table S1 Primer pairs used for scoring AFLPs, with their abbreviations.

Table S2 Associated markers of the female map (linked but not mapped).

Table S3 Associated markers of the male map (linked but not mapped).