



HAL
open science

MNF, an ankyrin repeat protein of myxoma virus, is part of a native cellular SCF complex during viral infection

Sophie Blanié, Jacqueline Gelfi, Stéphane Bertagnoli, Christelle Camus-Bouclainville

► To cite this version:

Sophie Blanié, Jacqueline Gelfi, Stéphane Bertagnoli, Christelle Camus-Bouclainville. MNF, an ankyrin repeat protein of myxoma virus, is part of a native cellular SCF complex during viral infection. *Virology Journal*, 2010, 7, pp.1-5. 10.1186/1743-422X-7-56 . hal-02660479

HAL Id: hal-02660479

<https://hal.inrae.fr/hal-02660479>

Submitted on 30 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

SHORT REPORT

Open Access

MNF, an ankyrin repeat protein of myxoma virus, is part of a native cellular SCF complex during viral infection

Sophie Błanié^{1,2}, Jacqueline Gelfi^{1,2}, Stéphane Bertagnoli^{1,2}, Christelle Camus-Bouclainville^{1,2*}

Abstract

Myxoma virus (MYXV), a member of the *Poxviridae* family, is the agent responsible for myxomatosis, a fatal disease in the European rabbit (*Oryctolagus cuniculus*). Like all poxviruses, MYXV is known for encoding multiple proteins that regulate cellular signaling pathways. Among them, four proteins share the same ANK/PRANC structure: M148R, M149R, MNF (Myxoma Nuclear factor) and M-T5, all of them described as virulence factors. This family of poxvirus proteins, recently identified, has drawn considerable attention for its potential role in modulating the host ubiquitin-proteasome system during viral infection. To date, many members of this novel protein family have been shown to interact with SCF components, *in vitro*. Here, we focus on MNF gene, which has been shown to express a nuclear protein presenting nine ANK repeats, one of which has been identified as a nuclear localization signal. In transfection, MNF has been shown to colocalise with the transcription factor NF- κ B in the nucleus of TNF α -stimulated cells. Functionally, MNF is a critical virulence factor since its deletion generates an almost apathogenic virus. In this study, to pursue the investigation of proteins interacting with MNF and of its mechanism of action, we engineered a recombinant MYXV expressing a GFP-linked MNF under the control of MNF native promoter. Infection of rabbits with MYXV-GFPMNF recombinant virus provided the evidence that the GFP fusion does not disturb the main function of MNF. Hence, cells were infected with MYXV-GFPMNF and immunoprecipitation of the GFPMNF fusion protein was performed to identify MNF's partners. For the first time, endogenous components of SCF (Cullin-1 and Skp1) were co-precipitated with an ANK myxoma virus protein, expressed in an infectious context, and without over-expression of any protein.

Findings

Myxoma virus (MYXV), a member of the *Poxviridae* family, is the agent responsible for myxomatosis, a fatal disease in the European rabbit (*Oryctolagus cuniculus*). Due to its host specificity, MYXV is a useful model to study *in vivo* the mechanisms by which the numerous virus-encoded virulence factors control host response to infection. Among the virulence factors, MYXV genome encodes four proteins which share the same structure: M148R, M149R, MNF and M-T5 [1-4]. They present Ankyrin repeats (ANK) and a C-terminal PRANC domain [5]. The Ankyrin repeat, a 33-residue sequence domain, is one of the most common protein-protein interaction motifs found in nature [6]. Extensively

identified in eukaryotes and bacteria, ANK motif is relatively rare among viruses, with the exception of poxviruses, especially chordopoxviruses [7]. Thus, poxviruses generally encode 4 or 5 ANK proteins, and more than 80% of these ANK proteins present an F-box-like domain at their C-terminus [5]. This domain is shorter than the typical cellular F-box and lacks helix 3 [8]. Furthermore, the poxviral F-box-like domain is located at the C-terminus of ANK protein whereas cellular F-boxes are typically located in the N-terminus of the protein. In addition, cellular F-boxes are associated with a second protein-protein interaction domain such as WD40 or Leucine Rich Repeat but no association with ANK motif has been reported yet. Due to these significant differences with eukaryotic F-boxes, the poxviral F-box domain has been defined as a new motif: the PRANC domain (Pox protein Repeats of Ankyrin C-terminal) [9].


* Correspondence: c.camus@envt.fr

¹INRA, UMR 1225 Interactions Hôtes - Agents Pathogènes F-31076 Toulouse, France

The F-box motif mediates protein-protein interactions and was first described as a recognition motif in the E3 ubiquitin ligase complex, known as SCF (Skp, Cullin, F-box) [10]. The SCF complex is a multi-protein complex that mediates the ubiquitination of substrates mainly destined for degradation by the 26S proteasome [11]. Targeted proteins are recognized by a variety of F-box proteins and are delivered to the E3 ligase complex via an adaptor protein, Skp1 [12]. Interaction between the adaptor protein and the F-box protein occurs via the F-box motif. The SCF complex plays a critical role in the selective degradation of regulatory proteins that mediate various cellular functions, such as signal transduction and cell cycle regulation [12]. Ten poxvirus ANK proteins have been shown to contain a functional F-box: myxoma virus protein M-T5 [13], orf virus proteins OV008, OV123, OV126, OV128 and OV129 [9], vaccinia virus protein MVA186R [14] and ectromelia virus proteins ECTV002, ECTV005 and ECTV154 [15]. Hence, among MYXV proteins, only M-T5 was identified as a cellular binding partner of Cullin-1 to date. Enhanced ubiquitination of p27/Kip1 and subsequent degradation via the proteasome pathway was observed through the interaction of M-T5 and Cullin-1. Consistent with this interaction, M-T5 was shown to promote cell cycle progression beyond the G0/G1 checkpoint during virus infection. In this study, we focused on MNF. MNF gene expresses a nuclear protein that presents nine ANK repeats, one of which exhibits significant sequence similarity with I κ B α nuclear localization signal. Moreover, in transfection, MNF colocalises with the transcription factor NF- κ B in the nucleus of TNF α -stimulated cells. Functionally, infection of rabbits with an MNF-deleted MYXV induced very mild clinical signs, with few discrete secondary myxomas, no respiratory infection, and no lethality. Histological analysis of the primary myxoma and parotid lymph node showed that deletion of MNF gene allowed a more rapid and quickly resolved inflammation [3]. To summarize, MNF was shown to inhibit the pro-inflammatory pathway, seemingly through an interaction with NF- κ B pathway.

To go further in the identification of the proteins interacting with MNF in a viral infectious context, different assays to produce antibodies directed against MNF were performed. All the rabbit sera obtained by immunization against part of or the whole MNF protein were specific to the MYXV infection but allowed the staining of cells infected with the MYXV Δ MNF too, indicating that they were not MNF-specific (data not shown). Thus, to overcome the difficulty to trace MNF by use of antibodies, a recombinant MYXV expressing a GFP-linked MNF, where GFP is in N-terminus of MNF, was engineered (MYXV-GFPMNF) (Figure 1A). The fusion protein GFPMNF was expressed under control of

MNF native promoter. The *Ecogpt* selection gene (*Escherichia coli* xanthine-guanine phosphoribosyl transferase), under the control of the P7.5 early poxviral promoter, was inserted between M149R and GFPMNF. The regulating region (R.R.) between M149R and MNF might contain sequences necessary to M149R expression but also contains MNF promoter which begins at the end of M149R-coding sequence. So as not to disturb the regulating region, and thus the expression of M149R, by the insertion of the selection gene, the R.R. was duplicated. Recombinant virus was selected upon gpt selection and purity was verified by PCR (data not shown). The correct expression of the fusion protein, its impact on MNF functionality, and thus on MYXV pathogenicity, were verified. The purified wild-type, MYXV-GFPMNF and MYXV Δ MNF [3] viruses were inoculated intradermally (5×10^3 Foci Forming Unit (FFU)) to the right ear of European rabbits (five animals per virus). Rabbits were monitored daily for clinical signs of myxomatosis. Rabbits infected with wild-type or MYXV-GFPMNF viruses developed a classic form of myxomatosis, totally different from the one induced by MYXV- Δ MNF virus which generated very mild clinical signs (Figure 1B). This experiment suggests that the main function of MNF is preserved even when MNF is fused to GFP. To identify the partners of MNF, immunoprecipitations of GFPMNF were performed. BGMK cells (Baby green monkey kidney) were infected with wild-type MYXV, MYXV-GFPMNF, MYXV-GFP or mock infected. Twenty-four hours post-infection (h.p.i.), cells were washed and lysed with hypotonic buffer. Lysates were cleared by centrifugation and the supernatants incubated with anti-GFP coupled to magnetic beads in order to precipitate GFPMNF fusion protein. Eluates were loaded on SDS/PAGE gels and analyzed by Coomassie blue staining. Three bands (Figure 2A a, b and 2A c) were clearly specific of MYXV-GFPMNF infection, and were thus excised, digested and analyzed by mass spectrometry. One of them (band c) was identified as the GFPMNF fusion protein and the two others as Cullin-1 (a and b). These data were confirmed by western-blot analysis of the lysates (Figure 2B left and middle panels). The presence of two GFPMNF bands suggests that during infection, MNF may be expressed under two forms, one being modified post-translation. Since the interaction between F-box protein and Cullin-1 usually occurs via the Skp1 protein, a western-blot analysis of the eluates was performed using anti-Skp1 antibodies (Figure 2B, middle panel). This experiment shows that Skp1 co-precipitates with MNF and suggests that Skp1 allows the interaction between MNF and Cullin-1. To identify a potential ubiquitinated substrate, an anti-ubiquitin western-blot was performed onto eluates (Figure 2B right panel). An ubiquitinated protein






Figure 2 Analysis of MNF partners by co-immunoprecipitation. Ten mm plates of BGMK cells (Baby green monkey kidney) were infected with MYXV-GFPMNF (1), wild-type MYXV (2) or MYXV-GFP (3), at m.o.i 3. The latter has been obtained by insertion of GFP gene, under control of the P7.5 poxviral promoter, in the thymidine kinase locus. 24 hours post-infection, cells were washed in PBS and lysed with hypotonic buffer (10 mM HEPES pH7.9, 150 mM NaCl, 600 mM KCl, 0.5% NP40 and proteases inhibitors). Lysates were cleared by centrifugation and the supernatants incubated with μ MACS anti-GFP MicroBeads. Washes and elution were performed according to manufacturer's instructions. Eluates were loaded on SDS/PAGE gels and analyzed by Simply Blue SafeStain (A) or by western blot (B). A. The 3 bands specific of MYXV-GFPMNF precipitation (a b and c) were analyzed by mass spectrometry. B. Western blot analysis of lysates with rabbit anti-GFP, rabbit anti-Cullin-1, rabbit anti-Skp1 or mouse anti-ubiquitin antibodies. Secondary antibodies were from anti-rabbit or anti-mouse WesternBreeze Chemiluminescent Kit, respectively. c and \rightarrow : GFPMNF, *: GFP, a and b: Cullin-1 bands, and \Rightarrow : Skp 1 band.

corresponding to band a (figure 2A), previously identified as Cullin-1 by mass spectrometry, was specifically co-precipitated with GFPMNF. This explains the two Cullin-1 bands identified by mass spectrometry, suggesting that the main part of MNF-co-precipitating Cullin-1 seems to be ubiquitinated. Cullin-1 auto-ubiquitination has already been reported [16] but its meaning remains to be clarified. However, in this experiment, no ubiquitinated substrate was co-precipitated with the SCF complex containing MNF. Since MNF is suspected to interact with NF κ B pathway and I κ B degradation occurs via its ubiquitination by SCF, variations in I κ B expression were measured in cells infected with MYXV or MYXV Δ MNF at various times post-infection. No differences were shown, suggesting that MNF does not interfere with degradation of I κ B (Data not shown).

For the first time, endogenous components of SCF (Cullin-1 and Skp1) were co-precipitated with an ANK myxoma virus protein, virally expressed under control of its own promoter, and without over-expression of any protein. Thus, our data demonstrate that MNF interacts with cellular SCF ubiquitin ligase complexes, in real

infectious context. Many viral proteins have been shown to interact with subunits of the SCF complex. In the case of ANK poxviral proteins, they seem to act as substrate adaptors, thus directing the ubiquitin ligase complex toward new targets or protecting cellular targets from ubiquitination. However, the substrates are mostly unknown. In some instances, viral proteins can also be targeted by SCF for their own destruction. The only identified substrates ubiquitinated by SCF containing an ANK/PRANC poxviral protein are p27 [13] and Akt [17], which are bound to the host SCF complex via Skp1 and MYXV M-T5 in infected cells. On the other hand, it has been shown that the 68k-ank protein contains another critical domain that may function independently from SCF ubiquitin ligase complex formation [18], suggesting that ANK/PRANC poxviral proteins may have multiple roles. In addition, poxviruses encode ANK proteins that do not present PRANC motif, such as Vaccinia virus K1L which mediates host-range function in RK-13 cells via ANK repeat. Moreover, K1L may interact with a cellular GTPase-activating protein [19] and inhibits host NF κ B activation by preventing I κ B α

degradation [20]. The identification of proteins bound by the N-terminal ANK domains and the outcome of these proteins could also greatly enhance our understanding of the multiple functions of the ANK/PRANC poxviral proteins during infection.

Acknowledgements

SBI was supported by funds from the French Ministry of Research and Technology. We thank Brigitte Peralta and Josyane Loupias for excellent technical assistance. We are grateful to B. Séverac for critical reading of the manuscript.

Author details

¹INRA, UMR 1225 Interactions Hôtes - Agents Pathogènes F-31076 Toulouse, France. ²Université de Toulouse; ENVT; UMR 1225 Interactions Hôtes - Agents Pathogènes; F-31076 Toulouse, France.

Authors' contributions

SBI carried out all the experiments and drafted the manuscript. CC, SBe and JG participated in the design of the study and revision of the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 9 December 2009

Accepted: 8 March 2010 Published: 8 March 2010

References

1. Cameron C, Hota-Mitchell S, Chen L, Barrett J, Cao JX, Macaulay C, Willer D, Evans D, McFadden G: **The complete DNA sequence of myxoma virus.** *Virology* 1999, **264**:298-318.
2. Blanié S, Mortier J, Delverdier M, Bertagnoli S, Camus-Bouclainville C: **M148R and M149R are two virulence factors for myxoma virus pathogenesis in the European rabbit.** *Vet Res* 2009, **40**:11.
3. Camus-Bouclainville C, Fiette L, Bouchiha S, Pignolet B, Counor D, Filipe C, Gelfi J, Messud-Petit F: **A virulence factor of myxoma virus colocalizes with NF-kappaB in the nucleus and interferes with inflammation.** *J Virol* 2004, **78**:2510-2516.
4. Mossman K, Lee SF, Barry M, Boshkov L, McFadden G: **Disruption of M-T5, a novel myxoma virus gene member of poxvirus host range superfamily, results in dramatic attenuation of myxomatosis in infected European rabbits.** *J Virol* 1996, **70**:4394-4410.
5. Mercer AA, Fleming SB, Ueda N: **F-box-like domains are present in most poxvirus ankyrin repeat proteins.** *Virus Genes* 2005, **31**:127-133.
6. Mosavi LK, Cammett TJ, Desrosiers DC, Peng ZY: **The ankyrin repeat as molecular architecture for protein recognition.** *Protein Sci* 2004, **13**:1435-1448.
7. Bork P: **Hundreds of ankyrin-like repeats in functionally diverse proteins: mobile modules that cross phyla horizontally?** *Proteins* 1993, **17**:363-374.
8. Sonnberg S, Fleming SB, Mercer AA: **A truncated two-alpha-helix F-box present in poxvirus ankyrin-repeat proteins is sufficient for binding the SCF1 ubiquitin ligase complex.** *J Gen Virol* 2009, **90**:1224-1228.
9. Sonnberg S, Seet BT, Pawson T, Fleming SB, Mercer AA: **Poxvirus ankyrin repeat proteins are a unique class of F-box proteins that associate with cellular SCF1 ubiquitin ligase complexes.** *Proc Natl Acad Sci USA* 2008, **105**:10955-10960.
10. Skowyra D, Craig KL, Tyers M, Elledge SJ, Harper JW: **F-box proteins are receptors that recruit phosphorylated substrates to the SCF ubiquitin-ligase complex.** *Cell* 1997, **91**:209-219.
11. Hershko A, Ciechanover A: **The ubiquitin system.** *Annu Rev Biochem* 1998, **67**:425-479.
12. Deshaies RJ: **SCF and Cullin/Ring H2-based ubiquitin ligases.** *Annu Rev Cell Dev Biol* 1999, **15**:435-467.
13. Johnston JB, Wang G, Barrett JW, Nazarian SH, Colwill K, Moran M, McFadden G: **Myxoma virus M-T5 protects infected cells from the stress of cell cycle arrest through its interaction with host cell cullin-1.** *J Virol* 2005, **79**:10750-10763.
14. Sperl KM, Schwantes A, Schnierle BS, Sutter G: **The highly conserved orthopoxvirus 68k ankyrin-like protein is part of a cellular SCF ubiquitin ligase complex.** *Virology* 2008, **374**:234-239.
15. van Buuren N, Couturier B, Xiong Y, Barry M: **Ectromelia virus encodes a novel family of F-box proteins that interact with the SCF complex.** *J Virol* 2008, **82**:9917-9927.
16. Maeda I, Ohta T, Koizumi H, Fukuda M: **In vitro ubiquitination of cyclin D1 by ROC1-CUL1 and ROC1-CUL3.** *FEBS Lett* 2001, **494**:181-185.
17. Werden SJ, Lanchbury J, Shattuck D, Neff C, Dufford M, McFadden G: **The myxoma virus m-t5 ankyrin repeat host range protein is a novel adaptor that coordinately links the cellular signaling pathways mediated by Akt and Skp1 in virus-infected cells.** *J Virol* 2009, **83**:12068-12083.
18. Sperl KM, Schwantes A, Staib C, Schnierle BS, Sutter G: **The orthopoxvirus 68-kilodalton ankyrin-like protein is essential for DNA replication and complete gene expression of modified vaccinia virus Ankara in nonpermissive human and murine cells.** *J Virol* 2009, **83**:6029-6038.
19. Bradley RR, Terajima M: **Vaccinia virus K1L protein mediates host-range function in RK-13 cells via ankyrin repeat and may interact with a cellular GTPase-activating protein.** *Virus Res* 2005, **114**:104-112.
20. Shisler JL, Jin XL: **The vaccinia virus K1L gene product inhibits host NF-kappaB activation by preventing IkkappaBalpha degradation.** *J Virol* 2004, **78**:3553-3560.

doi:10.1186/1743-422X-7-56

Cite this article as: Blanié *et al.*: MNF, an ankyrin repeat protein of myxoma virus, is part of a native cellular SCF complex during viral infection. *Virology Journal* 2010 **7**:56.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

