A molecular assay for the proactive detection of target site-based resistance to herbicides inhibiting acetolactate synthase in *Alopecurus myosuroides*.

Christophe C. Delye, Karelle Boucansaud

► To cite this version:

Christophe C. Delye, Karelle Boucansaud. A molecular assay for the proactive detection of target site-based resistance to herbicides inhibiting acetolactate synthase in *Alopecurus myosuroides*.. *Weed Research*, 2008, 48 (2), pp.97-101. 10.1111/j.1365-3180.2007.00615.x . hal-02660912

HAL Id: hal-02660912

<https://hal.inrae.fr/hal-02660912>

Submitted on 30 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A molecular assay for the proactive detection of target site-based resistance to herbicides inhibiting acetolactate synthase in *Alopecurus myosuroides*

C DÉLYE & K BOUCANSAUD

INRA, UMR1210 Biologie et Gestion des Adventices, Dijon, France

Received 26 July 2007

Revised version accepted 13 November 2007

Summary

Acetolactate synthase (ALS) inhibitors are the most resistance-prone herbicide group. Rapid resistance diagnosis is thus of importance for their optimal use. We formulate rules to use the derived cleaved amplified polymorphic sequence method to develop molecular tools detecting a change at a given codon, the nature of which is unknown. We applied them to *Alopecurus myosuroides* (black grass) to develop assays targeting ALS codons A122, P197, A205, W574 and S653 that are

crucial for herbicide sensitivity. These assays detected W574L or P197T, or both substitutions, in most plants analysed from a field where ALS inhibitors failed after 3 years of use. Similar assays can easily be set up for any species. Given the rapidity of selection for resistance to ALS inhibitors, these assays should be very useful in proactive herbicide resistance diagnosis.

Keywords: acetolactate synthase, ALS, AHAS, dCAPS, herbicide, PCR, proactive diagnosis, mutation, resistance, sulfonylurea, grass weed.

DÉLYE C & BOUCANSAUD K (2008). A molecular assay for the proactive detection of target site-based resistance to herbicides inhibiting acetolactate synthase in *Alopecurus myosuroides*. *Weed Research* **48**, 97–101.

Introduction

Herbicides targeting acetolactate synthase (ALS, EC 2.2.1.6) are among the most used herbicides worldwide (Tranel & Wright, 2002; Corbett & Tardif, 2006). They are also the most resistance-prone herbicide group to date, with resistant plants reported in at least 95 weed species (Heap, 2007). Resistance diagnosis is thus of importance for the optimal use of ALS inhibitors. ALS-based resistance is considered to play the major role in resistance to ALS inhibitors (Corbett & Tardif, 2006). Using *Arabidopsis thaliana* (L.) Heynh. ALS sequence as a reference for amino acid numbering (Tranel & Wright, 2002), most ALS-based resistance cases reported to date are due to substitutions at codons A122, P197, A205, W574 or S653 (Tranel *et al.*, 2007). Resistant ALS allele detection proved the

quickest and most accurate way of diagnosing resistance to ALS inhibitors (Corbett & Tardif, 2006). Molecular, DNA-based assays most often based upon the loss/gain of a restriction enzyme recognition site [polymerase chain reaction (PCR)–restriction fragment-length polymorphism (RFLP)], or upon direct specific amplification of the mutant allele (allele-specific PCR), have consequently been developed to detect resistant ALS alleles in various weed species where resistance arose (Corbett & Tardif, 2006). However, different possible substitutions have been reported at a given amino acid position, e.g. nine substitutions are known at codon 197 (Tranel *et al.*, 2007). The molecular tools hitherto developed were therefore set up on the basis of ALS sequences from resistant plants, after herbicide resistance confirmation (Corbett & Tardif, 2006). This takes time and is to be performed each time a new

primers about 40 nucleotide long to enable easy discrimination of undigested and digested (about 40 bp removed, see Fig. 1) amplicons by standard agarose gel electrophoresis; (3) no mismatch at the last dCAPS primer 3' nucleotide; (4) all nucleotides in the restriction enzyme recognition site exclusively located in the primer sequence and in the part of the targeted codon where any variation would cause amino-acid substitution; and (5) no other restriction enzyme recognition site in the amplicon. When several restriction enzymes fulfilled rules (4) and (5), the cheapest one was selected. Primers and restriction enzymes are given in Table 1. Positions of primers and restriction enzyme recognition sites in *A. myosuroides* ALS sequence are shown on Fig. 1.

Rule (4) was set up for two reasons. First, if nucleotides in the restriction enzyme recognition site were located elsewhere, silent changes or variation at positions not involved in herbicide resistance would lead to false detection of resistant ALS alleles. Second, the genetic code is degenerated. Thus, not all nucleotide substitutions cause amino acid variation. In *A. myosuroides*, P197 is encoded by CCC, and A122 and A205 are both encoded by GCC. A proline residue is encoded by CCN, an alanine by GCN (N is any nucleotide). Only changes at the two first nucleotides cause amino acid substitution at these codons. Recognition sites of the selected enzymes (*Bam*HI for P197, *Bgl*II for A122 and A205) thus encompass four nucleotides in dCAPS primers and the two first nucleotides in each targeted codon (Table 1, Fig. 1). Only codon TGG encodes a tryptophane residue. Any change at this codon causes amino acid substitution. *Bst*XI was selected, the recog-

nition site of which encompasses three nucleotides in primer W574F and the three nucleotides in codon 574 (Table 1, Fig. 1). In *A. myosuroides*, S653 is encoded by AGC. A serine residue is encoded by TCN or AGY. AGC-to-AGG and AGC-to-AGA changes cause an S-to-R substitution, while AGC-to-AGT is silent. Any other change at codon 653 causes amino acid substitution. *Bpu*10I was selected, the recognition site of which encompasses three nucleotides in dCAPS primer S653F and three in codon 653 (Table 1, Fig. 1). We deemed the possible false-positive diagnosis of the occurrence of mutant ALS caused by a silent AGC-to-AGT change to be a lesser flaw than not detecting the two other changes causing an S-to-R substitution.

DNA extraction and PCR mixes were as described (Délye *et al.*, 2002). Primers targeting A122 and W574 were used at 0.2 µM each. Primers targeting other codons were used at 0.4 µM each. Cycling programs consisted of 95°C for 5 min followed by 37 cycles at 95°C for 5 s, T_m (annealing temperature of the primer pair) (Table 1) for 10 s and 72°C for 30 s. Digestions were performed at 37°C for 3 h in PCR mixes added with 5 U enzyme (Fermentas, Vilnius, Lithuania), 0.7 µL of 10x enzyme buffer and 5 µL of water. dCAPS patterns were visualised by electrophoresis on 3% (wt/V) agarose gels run in 0.5x TBE buffer, except for A205 (3.5% wt/V agarose).

Testing and validating dCAPS

All assays were set up using 15 plants from an *A. myosuroides* accession never sprayed with ALS inhibitors. Digested and undigested amplicons were

Table 1 dCAPS primers and restriction enzymes

Target codons	Primers*	Sequence (5'–3')†	T _m (PCR)	Restriction enzyme, recognition site	Expected dCAPS patterns (fragment sizes in bp) WT‡	Mutant
A122	A122F	CCTACTCTCCCGCGCTACCTGCC	70°C	<i>Bgl</i> II, GCCN ₅ GGC	36, 208	244
	A122R§	TGGCCGGCGAGCGTGTGAGCGCCTGGTGGATCGCCATTGAG				
P197	P197F§	TTCTCGACTCCATCCCGATGGTGCCTATCACGGGACAGGAT	60°C	<i>Bam</i> HI, GGATCC	38, 200	238
	P197R	ATCTGCTGCTGGATGTCCTTGGG				
A205	A205F	TTCTCGACTCCATCCCGATGGTGGC	55°C	<i>Bgl</i> II, GCCN ₅ GGC	41, 69	110
	A205R§	GGTGATGGAGCGGGGACCTCTACAATGGGCGTGCCTGGAAG				
W574	W574F§	GGTGATGATACTGAACAATCAACATCTGGGAATGCCAGTGCCAG	60°C	<i>Bst</i> XI, CCAN ₆ TGG	39, 372	411
	WSR	ATACACCAGCATCATGCTGATCAGG				
S653	S653F§	ATCATCGTACCTACCAGGAGCAGTGCTGCCTATGATCCTA	65°C	<i>Bpu</i> 10I, CCTNAGC	40, 133	173
	WSR	ATACACCAGCATCATGCTGATCAGG				

*Primer name ending with F, forward primer; primer name ending with R, reverse primer.

†Nucleotide(s) in bold are modified from *A. myosuroides* ALS sequence in order to create a recognition site for the restriction enzyme used. Nucleotides belonging to the enzyme restriction site are underlined.

‡Wild-type.

§dCAPS primer.

Growing season	Crop	Herbicide(s) applied	% <i>A. myosuroides</i> control
2005–2006	Wheat	Mesosulfuron* + iodosulfuron*	0
		Imazamethabenz* + isoproturon†	30–40
		Mesosulfuron* + iodosulfuron*	5060
2004–2005	Wheat	Mesosulfuron* + iodosulfuron*	70
2003–2004	Wheat	Mesosulfuron* + iodosulfuron*	100
2002–2003	Wheat	Clodinafop‡	100
2001–2002	Wheat	Clodinafop‡	90–100
2000–2001	Wheat	Clodinafop‡	90

*Herbicide inhibiting ALS (HRAC group B).

†Herbicide inhibiting photosystem II (HRAC group C).

‡Herbicide inhibiting acetyl-CoA carboxylase (HRAC group A).

Table 2 Record of herbicide application on an *A. myosuroides* accession collected in 2006 in Sommepey (NE France)

electrophoresed side by side. After enzyme digestion, all plants yielded the expected wild-type patterns that were readily discriminated from undigested amplicons on agarose gels (not shown).

We then investigated a field where ALS-inhibiting herbicides gave very poor *A. myosuroides* control after only 3 years of use, with three applications during the last growing season (Table 2). Seeds were randomly sampled from about 150 surviving plants, allowed to germinate and 100 seedlings were analysed using the five dCAPS assays. No seedling contained mutations at codons A122, A205 and S653 (not shown). Six plants also did not contain mutations at codons P197 and W574. Ten plants were homozygous mutants at codon 197 and 77 plants at codon 574. The remaining seven plants were heterozygous mutants for both codons. dCAPS patterns obtained for codons 197 and 574 are illustrated in Fig. 2. dCAPS results were confirmed by

sequencing six seedlings in each of the four categories identified. No change at any of the five codons was observed in seedlings where no mutation was detected using dCAPS. Seedlings where a mutation at codon 574 was detected all contained a TGG-to-TTG change causing a W-to-L substitution, as described in 14 other species (Tranel *et al.*, 2007). Seedlings where a mutation at codon 197 was detected all contained a CCC-to-ACC change, causing a P-to-T substitution also described in five other species (Tranel *et al.*, 2007). Double-heterozygous mutant seedlings all contained both changes. In other species, W574L and P197T substitutions were both reported to confer resistance to sulfonylureas (like mesosulfuron and iodosulfuron), while only W574L conferred resistance to imidazolinones (like imazamethabenz) (Tranel *et al.*, 2007). Imazamethabenz use (Table 2) might be why L574 plants were predominant in the investigated field.

Fig. 2 dCAPS patterns of seven *A. myosuroides* seedlings analysed with dCAPS assays targeting codons 197 (top) and 574 (bottom). Lanes 1–4, homozygous mutants at codon 574; lane 5, homozygous mutant at codon 197; lane 6, double heterozygous mutant; lane 7, wild-type.

Conclusions

Similar to PCR-RFLP and allele-specific PCR, dCAPS enabled the discrimination of homozygous and heterozygous mutant plants. Allele-specific PCR requires known wild-type and mutant sequences to be developed. PCR-RFLP cannot be used for proactive resistance detection. First, a suitable restriction site including nucleotide position involved in resistance rarely exists in wild-type sequences. Second, restriction enzyme recognition sites contain at least four (most frequently six) nucleotides. There is therefore a significant risk for false detection of resistant alleles using PCR-RFLP, because of substitutions occurring outside nucleotide positions involved in resistance (see explanation to rule 4). dCAPS is therefore the only method so far detecting mutations of unknown nature at given codons. The assays described herein proved reliable for detecting mutations in *A. myosuroides* ALS. Following the five rules we set up, similar assays can easily be developed for any

other species. Given the surprisingly fast selection for resistance observed in the *A. myosuroides* accession investigated, dCAPS assays targeting ALS should prove immensely useful for quick and proactive diagnosis of ALS-based resistance.

Acknowledgements

We are grateful to B. Couloume (Bayer CropScience France) for financial support and for providing *A. myosuroides* seeds.

References

- CORBETT C-AL & TARDIF FJ (2006) Detection of resistance to acetolactate synthase inhibitors in weeds with emphasis on DNA-based techniques: a review. *Pest Management Science* **62**, 584–597.
- DÉLYE C, MATÉJICEK A & GASQUEZ J (2002) PCR-based detection of resistance to acetyl- CoA carboxylase-inhibiting herbicides in black-grass (*Alopecurus myosuroides* Huds) and ryegrass (*Lolium rigidum* Gaud). *Pest Management Science* **58**, 474–478.
- HEAP I (2007) *International Survey of Herbicide-resistant Weeds* (available at <http://www.weedscience.org>; last accessed 11 June 2007).
- KAUNDUN SS & WINDASS JD (2006) Derived cleaved amplified polymorphic sequence, a simple method to detect a key point mutation conferring acetyl CoA carboxylase inhibitor herbicide resistance in grass weeds. *Weed Research* **46**, 34–39.
- NEFF MM, NEFF JD, CHORY J & PEPPER AE (1998) dCAPS, a simple technique for the genetic analysis of single nucleotide polymorphisms: experimental applications in *Arabidopsis thaliana* genetics. *The Plant Journal* **14**, 387–392.
- NEFF MM, TURK E & KALISHMAN M (2002) Web-based primer design for single nucleotide polymorphism analysis. *Trends in Genetics* **18**, 613–615.
- TRANEL PJ & WRIGHT TR (2002) Resistance of weeds to ALS-inhibiting herbicides: what have we learned? *Weed Science* **50**, 700–712.
- TRANEL PJ, WRIGHT TR & HEAP IN (2007) *ALS Mutations from Resistant Weeds* (available at <http://www.weed-science.org>; last accessed 24th July 2007).