



HAL
open science

A method for assessing the vectorial competence of field collected culicoides spp. for bluetongue virus

C. Ninio, D. Augot, H. Ferté, Emmanuel Breard, Thomas Lilin, Stéphan Zientara, P.S. Mellor, Barbara Dufour, J. Depaquit

► **To cite this version:**

C. Ninio, D. Augot, H. Ferté, Emmanuel Breard, Thomas Lilin, et al.. A method for assessing the vectorial competence of field collected culicoides spp. for bluetongue virus. *Parasite*, 2010, 17 (2), pp.172-172. 10.1051/parasite/2010172172 . hal-02662703

HAL Id: hal-02662703

<https://hal.inrae.fr/hal-02662703>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

LETTRE À LA RÉDACTION

A METHOD FOR ASSESSING THE VECTORIAL COMPETENCE OF FIELD COLLECTED *CULICOIDES* SPP. FOR BLUETONGUE VIRUS

NINIO C.*, AUGOT D.*, FERTÉ H.*, BREARD E.**,
LILIN T.***, ZIENTARA S.**, MELLOR P.S.****,
DUFOUR B.***** & DEPAQUIT J.*

Culicoides (Diptera: Ceratopogonidae) transmit Bluetongue virus (BTV). The identity of the competent species in the northern Europe is still uncertain. To help resolve this issue, two sheep were experimentally infected with BTV8. Viremia was confirmed by quantitative RT-PCR: Ct = 32 on day 4 post infection (p.i.) and 26 on day 5 p.i. On day 4 p.i. the sheep were placed individually into 4 m³ netted cages under P3. 192 field collected females captured at two cattle holdings in northern France were released into one cage, and 139 *Culicoides*, that emerged in the laboratory from breeding site material, such as manure, mud around trough and pounds, collected from both holdings, into the second. After 24 hours surviving *Culicoides* were collected and the blood-fed females were transferred into individual boxes. The blood engorged females were placed in an incubator (21 °C, 80 % humidity), supplied with a damp filter paper for egg laying and fed with sucrose solution.

Four field collected *C. obsoletus* s. l., had taken a blood meal but none of the *Culicoides* that emerged in the laboratory was blood-fed. A blood-fed female laid eggs six days after blood feeding and was assayed for virus eight days post feeding. The other blood-fed midges were killed before egg laying. All surviving *Culicoides* were dissected for identification, thorax and abdomen being stored individually at - 80 °C, then were crushed in 300 µl MEM. 100 µL was used for qRT-PCR, and 100 µL was inoculated intravenously into 10 days old embryonated chicken eggs. Three *C. sonorensis*, a major vector of BTV in the USA, were used as positive controls. They come from a colony kept in the Institute of Animal Health in Pirbright (UK) and were intrathoracically infected with BTV8. The female which laid eggs was weakly positive for BTV antigen using the qRT-PCR assay (Ct = 36.9) but live virus was not isolated via the chicken egg inoculation assay. All other experimental *Culicoides* were negative in both assays, but all

C. sonorensis were positive with Ct values ranging from 29 to 30 for individual midges, and between 18 and 19 for the qRT-PCR when run on the infected chicken eggs (which died at 2 and 3 days post inoculation).

The low feeding rate of the midges used in this study can be explained by the stress of captivity and an important percentage of *Culicoides festivipennis*, not known to feed on sheep. Based on a limited sampling, these results are not representative. Nevertheless, this method could be useful for future vectorial competence studies, though a major improvement in blood-feeding rates will be required to achieve meaningful results and detailed behavioural studies on *Culicoides* spp. are required to achieve this.

ACKNOWLEDGEMENTS

We would like to thank Simon Carpenter and Eva Veronesi, Institute for Animal Health, Vector-borne Disease Programme, who kindly provided us positive controls.

Reçu le 1^{er} mars 2010
Accepté le 14 avril 2010

Species or species group	Field collected midges' batch				Emerged midges' batch			
	Dead during the experiment		Killed after experiment		Dead during the experiment		Killed after experiment	
	♀	♂	♀	♂	♀	♂	♀	♂
<i>C. festivipennis</i>	–	–	1	–	17	55	22	12
<i>C. circumscriptus obsoletus</i> complex ¹	11	–	1	–	–	–	–	–
<i>C. dewulfi</i>	4	2	3**	–	–	–	–	–
<i>pulicaris</i> complex ²	8	2	1	–	7	–	1	–
<i>C. punctatus</i>	58	2	1	–	–	–	3	2
<i>C. achrayi</i>	5	1	–	–	–	1	2	–
<i>C. brunnicans</i>	3	1	–	–	–	–	–	–
<i>C. vexans</i>	13	–	–	–	–	–	–	–
<i>C. nubeculosus</i>	21	8	15	–	–	2	2	–
<i>C. kibunensis</i>	–	–	–	–	1	4	3	1
<i>C. stigma</i>	–	–	–	–	–	–	1	–
<i>C. unknown</i> ³	7	–	1	–	–	–	2	1
Sub-total	167	20	25	–	25	62	36	16
Total	212			139				

* Two females belonging to this species group had a blood meal at 4 days p.i. One of them laid eggs and was slightly positive for BTV antigen by qRT-PCR. ** Two females belonging to this species group had a blood meal at 4 days p.i.

¹ The *Obsoletus* complex includes *C. obsoletus* and *C. scoticus* females (males have all been identified as *C. obsoletus*). ² The *Pulicaris* complex includes *C. pulicaris* and *C. lupicaris*. ³ Damaged individual not suitable for morphological identification.

Table – Sampling *Culicoides* species collected inside the net cages after having spent 24 hours with a viraemic sheep.

* Usc Vecpar-AFSSA, JE 2533, Faculté de Pharmacie, Université de Reims Champagne-Ardenne, 51, rue Cognacq Jay, 51096 Reims, France.

** AFSSA-LERPAZ, Unité de virologie, Maisons-Alfort, France.

*** Centre de Recherches Biomédicales, École Nationale Vétérinaire, Maisons-Alfort, France.

**** Institute for Animal Health, Vector-borne Disease Programme, Ash Road, Pirbright, Woking, Surrey, GU24 0NF, UK.

***** Unité de recherche Epimai, École Nationale Vétérinaire, Maisons-Alfort, France.

Correspondence: Jérôme Depaquit.

E-mail: jerome.depaquit@univ-reims.fr