



**HAL**  
open science

## Behavioural, endocrine and immune responses to repeated social stress in pregnant gilts

David Couret, Wilfred Otten, B. Puppe, Armelle Prunier, Elodie Merlot

► **To cite this version:**

David Couret, Wilfred Otten, B. Puppe, Armelle Prunier, Elodie Merlot. Behavioural, endocrine and immune responses to repeated social stress in pregnant gilts. *Animal*, 2009, 3 (1), pp.118-127. 10.1017/S1751731108003236 . hal-02663422

**HAL Id: hal-02663422**

**<https://hal.inrae.fr/hal-02663422v1>**

Submitted on 31 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Behavioural, endocrine and immune responses to repeated social stress in pregnant gilts

D. Couret<sup>1,2†</sup>, W. Otten<sup>3</sup>, B. Puppe<sup>3</sup>, A. Prunier<sup>1,2</sup> and E. Merlot<sup>1,2</sup>

<sup>1</sup>INRA, UMR1079, Systèmes d'Élevage Nutrition Animale et Humaine, F-35000 Rennes, France; <sup>2</sup>Agrocampus Rennes, UMR1079, SENAH, F-35000 Rennes, France; <sup>3</sup>Behavioural Physiology Research Unit, Research Institute for the Biology of Farm Animals, Wilhelm-Stahl-Allee 2, 18196 Dummerstorf, Germany

(Received 14 March 2008; Accepted 14 August 2008; First published online 23 September 2008)

*Pregnant sows are exposed to various stressors in intensive pig husbandry that may have negative consequences on their health, reproductive performances and welfare. Social stress is one of these challenges, because gestating sows have to be housed in groups according to EU guidelines (2001/88/CE). The purpose of this study was to determine the consequences of repeated social stress in pregnant female pigs on their behavioural, endocrine and immunological responses and on pregnancy outcome. Pregnant gilts were submitted to a repeated social stress procedure induced by housing unfamiliar gilts in pairs changed twice a week between days 77 and 105 of gestation (S group, n = 18). Control gilts were housed in stable pairs during the same period (C group, n = 18). Agonistic behaviour was observed during the first 3 h after each grouping. Skin lesions were numbered 2 h after each grouping. Salivary cortisol was measured before and repeatedly during the 4 weeks of grouping. Gilts were immunized against keyhole limpet haemocyanin (KLH) on days 81 and 95 of gestation. Immunoglobulins G against KLH, proliferative responses to concanavalin A, lipopolysaccharide, pokeweed mitogen and KLH and peripheral blood leukocyte numbers were evaluated 1 week before the first grouping and 3 days after the last one. Agonistic interactions and skin lesions were observed in S gilts at each grouping, although there was a decline between the first and the last grouping (P < 0.05). The repeated social stress induced a sustained endocrine response as shown by elevated salivary cortisol levels from 1 to 48 h after grouping in S gilts compared to C gilts. The cellular as well as the humoral immunity and the leukocyte numbers were not influenced by social stress. Gestation length tended to be shorter in S gilts (P = 0.09), but litter size, piglet weight or mortality at birth were not affected. Variability of the response of S gilts to groupings was partly explained by their average success value determined according to the outcome (defeat or win) of all the groupings. In conclusion, our study demonstrates that the application of repeated social stress to pregnant gilts during the last third of their gestation repeatedly activates their hypothalamo–pituitary–adrenal axis but does not impair their immune function and pregnancy outcome.*

**Keywords:** cortisol, immunity, pig, social stress, pregnancy

### Introduction

In intensive pig husbandry, pregnant sows are exposed to various stressors such as diet and space restriction, which may have negative consequences on their welfare, health (Salak-Johnson and McGlone, 2007) and reproductive performances (Von Borell *et al.*, 2007). One potent stressor that pigs are exposed to is social stress, i.e. stress caused by composition or modification of their social group (de Groot *et al.*, 2001; Otten *et al.*, 2002; Coutellier *et al.*, 2007). In farms, the grouping of unfamiliar animals is usually accomplished to form cohorts of homogeneous age or body weight, but actual EU guidelines (2001/88/CE) also

impose the housing of gestating sows in groups for welfare reasons. The formation of a new social group leads to fights during the first days to establish a social hierarchy (Puppe *et al.*, 2008). The physical and psychological stress associated with the fights activates stress systems such as the hypothalamo–pituitary–adrenal (HPA) axis and the sympathetic system, leading to increased plasma levels of cortisol and catecholamines (Otten *et al.*, 2002), which in turn modulate animal physiology in order to cope with the new situation. In pigs, social stress or high cortisol concentrations can decrease weight gain (McGlone *et al.*, 1993; Coutellier *et al.*, 2007), lymphocyte proliferation (Kanitz *et al.*, 2004), natural killer cell activity (McGlone *et al.*, 1993; Salak-Johnson *et al.*, 1996), neutrophilic chemotaxis (Salak-Johnson *et al.*, 1996) and antiviral immunity (de Groot *et al.*, 2001).

† E-mail: david.couret@rennes.inra.fr

Most of the studies were performed on young animals and little is known about the influence of stress on the immune capacity of pregnant sows. Considering that pregnancy is a particular physiological state associated with immunological changes (Wegmann *et al.*, 1993) and a decreased sensitivity of the HPA axis to stress (Neumann *et al.*, 1998), one can wonder whether chronic stress has similar immune effects in pregnant gilts as in younger non-pregnant females or males.

Evaluation of the endocrine and immune consequences of stress in gestating sows can be of particular importance because stress during pregnancy in mammals can affect not only maternal welfare and health but also embryonic and foetal development. Although Soede *et al.* (2007) failed to observe any difference in fertilization rate and embryonic development in sows, data in rodents showed that stress during early gestation can lead to abortions (Wiebold *et al.*, 1986). In species with multiple fetuses like pigs, maternal stress is suspected to reduce the number of young born alive, suggesting an increased foetal mortality (von Borell *et al.*, 2007). In addition, maternal stress during gestation can have detrimental effects on growth (Fowden *et al.*, 2006), neuroendocrine (Weinstock, 1997) and immune (Merlot *et al.*, 2008) functions in the offspring after birth. The deleterious effects of stress on pregnancy and foetal development seem to be related to maternal immune changes (Arck, 2001; Joachim *et al.*, 2003; Coussons-Read *et al.*, 2007) and to maternal neuroendocrine activity, especially to glucocorticoid secretions (Barbazanges *et al.*, 1996).

The purpose of this study was to determine whether exposing pregnant gilts to repeated social stress was able to induce a prolonged state of stress as reported for non-pregnant growing pigs (Coutellier *et al.*, 2007) and therefore to determine the consequences of this stressor on their behaviour, endocrine and immune responses. The second aim was to investigate the effects of social stress on their reproductive performances when gestation is maintained despite stress. Because the risk of abortion is higher at the beginning of gestation, the social stress procedure was performed during the last third of pregnancy, and the litter characteristics were observed at birth.

## Material and methods

### *Animals and treatments*

A total of 36 nulliparous Large White × Landrace gilts from 11 litters were used for the experiment in three independent replicates of 12 females. Gilts were inseminated with fresh Pietrain semen after oestrus synchronisation with Regumate<sup>®</sup> (20 mg of 17  $\alpha$ -allyl-trenbolone; Roussel-Uclaf, Romainville, France). During gestation, gilts had free access to water. They were fed once a day at 0830 h with a standard pig diet for pregnant sows, which covered 145% of their metabolic requirements (1.1 kg per 100 kg live weight of a diet containing 13.5% crude protein, 2280 kcal/kg of metabolizable energy and 0.48% lysine). Animals were blocked in individual feeders for 30 min during the meal in order to prevent competition for food. Within each

replicate, gilts were equally allocated to one of the two treatment groups (Stress group, S; Control group, C) at  $77 \pm 1$  days of gestation (mean  $\pm$  s.d.). They were  $396 \pm 9$  days of age and weighed  $215 \pm 10$  kg at the beginning of the experiment. Assignment of gilts to treatment groups was realized in order to balance litter of origin and live weight between S and C groups. At least 4 months before allocation to the treatment groups, all gilts were housed in stable groups of four in a single room on a concrete floor with straw bedding (11.5 m<sup>2</sup>). On the first day of the stress procedure, two gilts per pen were moved to new pens (11.5 m<sup>2</sup>) located in another room for allocation to the S group. The other two gilts, allocated to the C group, remained undisturbed in the initial pens. Gilts from the S group were exposed to a repeated social stress routine. Over a period of 4 weeks, between days 77 and 101 of gestation, pairs of S gilts were changed twice a week (on Mondays and Thursdays at 1400 h). In order to form pairs of unfamiliar animals at each grouping, six additional non-experimental gilts with similar age and weight as S gilts were used. Pairs of C gilts were left undisturbed during the experimental period. On day 105 of gestation, all gilts were moved to individual farrowing crates (5 m<sup>2</sup>) with straw bedding and stayed there with their litter until weaning of piglets at 4 weeks of age. Parturition was not induced.

On days 70 and 105 of gestation, gilts were weighed and back fat thickness was measured ultrasonically (back at 65 mm from the midline). All gilts were immunized intra-muscularly in the neck with a solution of 2.5 mg DNP-keyhole limpet haemocyanin (KLH) (2,4-dinitrophenyl conjugated to keyhole limpet haemocyanin; Merck, Nottingham, UK) diluted in 1 ml sterilized water and 1 ml Montanide ISA 206 adjuvant (SEPPIC; Paris, France) on days 81 and 95 of gestation.

No animal was severely injured, required veterinary treatment or died during the experiment. Piglets were weighed at birth and the average weight of piglets was calculated. Stillborn piglets and mummified piglets were numbered.

### *Blood and saliva collection*

Blood samples were collected from the anterior vena cava of gilts restrained by snaring on days 70 and 105 of gestation. The sampling procedure took less than 2 min. EDTA-blood samples were collected for blood formula analyses, heparinized-blood samples for the proliferation assays and blood samples without anti-coagulant for immunoglobulins G (IgG) measurement. Blood was kept at 4°C until use.

Salivary samples were collected at 0900 h and 1500 h 1 week before the first grouping (day 72 of gestation). For the 1st, 4th and 8th groupings, samples were collected before (at 1330 h), after 1 h (at 1500 h) and after 19 h (at 0900 h) of grouping. Additional samples were collected 2 days after the 3rd and 7th groupings (i.e. the day before the 4th and 8th groupings) at 0700, 1100, 1500 and 1900 h in order to analyse diurnal variations of cortisol. Salivary samples were collected by allowing gilts to chew on cotton buds until they

were moistened. Cotton buds were rapidly centrifuged at  $3000 \times g$  for 15 min at 4°C. Saliva samples were stored at -20°C until cortisol analysis.

#### Measurements

**Agonistic behaviour and skin lesions.** In the S group, behaviour was video recorded for the first 3 h of each grouping. Agonistic behaviour of the dyads was analysed from the videotapes by continuous event sampling to determine the number and duration of agonistic interactions, the initiating sow and the outcome (win, defeat or undecided). An agonistic interaction was defined as an overt fight or a displacement event with physical contact for more than 3 s. If an agonistic sequence lasted longer than 8 s, the next fight was classified as a new interaction. Agonistic interactions comprised 'bitings', 'head-to-head knocks', 'head-to-body knocks' or 'parallel/inverse pressing' (Puppe, 1998; Tuchscherer *et al.*, 1998). A sow won the agonistic interaction when the other (defeated) sow first stopped fighting, turned away from an attack or tried to flee. The agonistic interaction was defined as undecided when there was no clear win or defeat. In addition, the initiating sow for each agonistic interaction was registered. For each S gilt a success value (SV) was evaluated for each grouping day according to the formula: (number of wins - number of defeats)/(number of wins + number of defeats) ranging from -1 to +1. An aggression value (AV) was calculated according to the formula: (number of agonistic interactions initiated by the gilt)/(total number of agonistic interactions) and ranged from 0 (not aggressive) to +1 (aggressive).

Skin lesions on the entire body (legs, sides, back and ears) were numbered 2 h after the beginning of each grouping in both S and C groups by an observing person in the pen. Only fresh lesions of more than 2 cm length were numbered.

**Anti-KLH IgG.** Blood samples without anti-coagulant were left to clot for 3 h at room temperature and were centrifuged at  $3000 \times g$  for 15 min at 4°C. Sera samples were stored at -20°C until analysis for anti-KLH IgG by ELISA as described previously (Schrama *et al.*, 1997). Measurements were performed in duplicates. Briefly, serial diluted sera were applied to antigen-coated wells (KLH, 0.1 µg/150 µl per well) of a 96-well round-bottomed plate (Greiner, Courtaboeuf, France). After 2 h of incubation at room temperature and subsequent washing with phosphate-buffered saline (Tris 0.05% Tween20), a one-step conjugation was performed. Conjugation consisted of incubation for 1 h with a 1:8000 diluted peroxidase-conjugated rabbit anti-swine IgG Fc (RaSw-IgG (Fc)/PO; Nordic, Tilburg, The Netherlands). After washing, tetramethylbenzidine substrate (TMB; Sigma-Aldrich, Saint Quentin Fallavier, France) was added. After 10 min, the reaction was stopped with 2.5 N sulphuric acid. Absorbance was read with a microplate reader at a test wavelength of 450 nm. The titer of immunoglobulins against KLH was calculated as the dilution that showed 50% extinction of Emax, where Emax

represents the saturation optical density of a positive standard serum. In the absence of immunization, there was no detectable level of IgG anti-KLH in sera of gilts. Inter- and intra-assay CV were 9.9% and 2.2%, respectively. Gilts from the first replicate were not immunized.

**Lymphocyte proliferation.** Peripheral blood mononuclear cells (PBMC) were isolated from heparinized blood samples by density gradient centrifugation. Briefly, 3 ml of blood diluted in an equal volume of Hank's balanced salts (HBSS) culture medium was layered on 3 ml histopaque 1077. After centrifugation (30 min,  $400 \times g$ , 4°C), PBMC were collected from the interface and washed twice in HBSS medium. Erythrocytes were lysed in 1 ml of sterilized water. Lysis reaction was stopped after 15 s by adding 8 ml of HBSS medium. PBMC were spun down ( $100 \times g$ , 10 min) and re-suspended in plain growth medium (RPMI 1640 supplemented with 10% FBS, 1% penicillin/streptomycin and 1% L-glutamine). Viability of cells was determined by trypan blue dye exclusion and cell concentration was determined using a Malassez counting chamber. Cell viability was always higher than 90%. Cell concentration was adjusted to  $5 \times 10^6$  cells/ml in growth medium. T-cell-specific mitogen concanavalin A (ConA; 1.5 µg/ml), B-cell-specific mitogen lipopolysaccharid (LPS; 6.25 µg/ml), T-cell-dependent B-cell mitogen pokeweed (PWM; 1.5 µg/ml) mitogen and KLH (10 µg/ml) were used for the proliferation assays. All media and mitogens were obtained from Sigma-Aldrich, except KLH, which was purchased from Merck Chemicals (Nottingham, UK). Cells ( $0.5 \times 10^6$  cells/well) were pipetted into 96-well flat-bottomed cell culture plates (Nunc, VWR International SAS, Fontenay sous bois, France) and 50 µl of the appropriate mitogen or growth medium was added in triplicates. After 68 h of incubation in a 5% CO<sub>2</sub>-humidified incubator at 37°C, cultures were pulsed with 10 µl of MTT labelling reagent (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide, final concentration 0.5 µg/ml) for 4 h at 37°C. A 10% sodium dodecyl sulphate buffer in 0.01 M HCl was used to solubilize the dark-blue crystals overnight at 37°C. The proliferation was proportional to the mitochondrial activity of viable cells evaluated by the reduction of tetrazolium salt into a blue formazan product. The optical density (OD) was measured by a micro-plate reader at a test wavelength of 550–600 nm and a reference wavelength of 630 nm. Results were expressed as proliferative index (PI) calculated according to the formula:  $PI = (OD(570/630 \text{ nm}) \text{ of stimulated cells}) / (OD(570/630 \text{ nm}) \text{ of non-stimulated cells})$ . Lymphocyte proliferative response to PWM was not assessed on day 70 of gestation in gilts from the first replicate and response to KLH was not assessed in gilts from the first replicate on days 70 and 105 of gestation.

**Blood formula.** Blood collected in EDTA tubes was used for blood formula determination. The total numbers of lymphocytes and granulocytes were measured with a haematology automatic cell counter calibrated for pigs (MS-9<sup>®</sup>, Melet Schloesing Laboratories, Osny, France).

**Cortisol.** Cortisol was measured in salivary samples with a luminescence immunoassay kit (LIA, IBL, Hamburg, Germany). The measurements were performed without replicates. The inter- and intra-assay coefficients of variation were 7.8% and 6.1% at 2.1 ng/ml, respectively. The assay sensitivity was 0.15 ng/ml.

#### *Sub-classification of gilts*

In order to explain the variability in the results obtained within the S group, we tried to determine whether particular behavioural or cortisol response patterns could be identified and used as main factors of variation in the statistical analyses. First, the S group was divided into three homogenous groups of similar size (High H, Medium M and Low L,  $n = 6/\text{group}$ ) according to the average success value (AvSV) calculated on the eight groupings. Thereafter, it was determined whether behavioural, cortisol, immune and reproductive results differed between these groups by ANOVA (see section Statistical analysis). The same procedure was applied to classifications based on average aggression value (AvAV), acute cortisol response (mean of cortisol levels at 1 and 19 h after the 1st, 4th and 8th groupings), diurnal cortisol (mean of cortisol levels at 0700, 1100, 1500 and 1900 h 2 days after the 3rd and 7th groupings) or total cortisol (mean of all acute and diurnal levels). Differences between sub-groups were more marked for the classifications based on success and aggressive values. Both classifications led to nearly similar sub-groups. Therefore, the AvSV sub-classification was chosen.

#### *Statistical analyses*

Statistical analyses were performed with Statistical Analysis Systems Institute Software (SAS, 1999) on 36 gilts except for gestation length and litter data where one gilt of the S group was excluded because of a leg injury, which occurred during transfer to the farrowing crate. A square root transformation was used for skin lesions and behavioural data; a logarithmic transformation was used for cortisol concentrations and proliferative indexes to fit a normal distribution. Data measured only once per individual were analysed by ANOVA using the MIXED procedure of SAS, including replicate as a random effect in the model and treatment as a main effect. Litter size was included in the statistical model as a covariate for gestation length, number of stillborn piglets, mummified piglets, piglet weight and mean litter weight. For data measured on the same animals at different times, replicate was included as a main effect in the model and the repeated statement of SAS was used with individual females as experimental units. The best covariance structure was chosen for each variable, using the structure minimizing the absolute value of the Akaike and Bayesian information criteria (AIC and BIC). The complete model included the main effects of treatment, time, replicate, and the treatment  $\times$  time interaction. The time effect refers either to the number of grouping (skin lesions, behavioural data, cortisol at 1 and 19 h after grouping), to the hour within a day (diurnal variations of cortisol) or to the stage of gestation (blood formula, proliferative indexes).

Statistical analyses for behavioural data and skin lesions were performed only within the S group because there were no behavioural observations in the C group and because the distribution of skin lesions in the C group was not normal since the lesions were observed in very small numbers and in a sporadic way. The different sub-classifications for S gilts were tested by including the gilt status (H, L, M or C) instead of the treatment as the main effect in the model of analysis. Correlation analyses were performed using the CORR procedure of SAS. When a significant effect was found, comparison between groups was made with the Bonferroni *t*-test. For all comparisons,  $P < 0.05$  was considered significant and  $0.05 < P < 0.1$  as a tendency. Data are presented as mean  $\pm$  s.e. before square root and logarithmic transformation.

## Results

#### *Agonistic behaviour and skin lesions*


Within the C group, agonistic behaviour was not recorded and number of skin lesions was very low all over the experiment (Figure 1a). Within the S group, numerous agonistic interactions and skin lesions were observed during all groupings (Figure 1b–d). However, the number ( $F_{7,109} = 4.90$ ;  $P < 0.001$ ) and duration ( $F_{7,109} = 2.24$ ;  $P < 0.05$ ) of agonistic interactions and the number of skin lesions ( $F_{7,105} = 3.72$ ;  $P < 0.01$ ) varied significantly over time. Generally, the number of skin lesions and agonistic behaviour was very high at the first grouping and decreased over the successive groupings. Compared to the first grouping, the number of agonistic interactions was lower from the 3rd to the 8th grouping ( $P < 0.05$ ), the duration of agonistic interactions was shorter after the 7th grouping ( $P < 0.05$ ) and the number of skin lesions was lower after the 6th ( $P = 0.07$ ) and 8th ( $P < 0.01$ ) groupings.

Success and aggressive values were tightly correlated at each grouping ( $r = 0.75$  to  $0.99$  depending on the grouping,  $P < 0.05$ ,  $n = 18/\text{group}$ ), indicating that winning sows were also the most aggressive ones.


#### *Cortisol*

Before the beginning of the experimental procedure, salivary cortisol levels at 0900 and 1500 h were similar in both groups (data not shown). Cortisol responses were measured at the 1st, 4th and 8th groupings. The grouping day  $\times$  treatment interaction was significant at 1 h ( $F_{2,68} = 3.45$ ;  $P < 0.05$ ) but not at 30 min before and 19 h after grouping (Figure 2). One hour after grouping, cortisol was higher in S than in C gilts at the 1st and 4th groupings ( $P < 0.001$ ). Nineteen hours after grouping, cortisol levels were higher in the S group regardless of the number of grouping ( $F_{1,32} = 19.23$ ;  $P < 0.001$ ).


For the investigation of diurnal rhythm, four samples were taken between 0700 and 1900 h, 2 days after the 3rd and 7th grouping. The sampling hour  $\times$  treatment interaction was not significant, which indicates that S and C groups followed the same pattern of diurnal variations. However, on both sampling days, salivary cortisol was higher in



**Figure 1** Number of skin lesions in C (a) and S (b) gilts 2 h after grouping. In the S group, number (c) and duration (d) of agonistic interactions (AI) were evaluated during the first 3 h of grouping. Values are mean  $\pm$  s.e. Different letters (a, b, c) indicate significant differences at  $P < 0.05$ .



**Figure 2** Salivary cortisol measured 1 h (a) and 19 h (b) after grouping. For cortisol measured 1 h after the beginning of grouping, a significant interaction between time (number of grouping) and treatment (T) was found, and comparisons between treatments within each time were realized using the Bonferroni *t*-test. Values are mean  $\pm$  s.e. \* $P < 0.05$ , \*\*\* $P < 0.0001$ .




**Figure 3** Salivary cortisol measured 2 days after the 3rd and the 7th groupings. Significant effects of treatment (T) and time but not time  $\times$  treatment interaction were found. Values are mean  $\pm$  s.e.

the S group than in the C group ( $F_{1,30} = 13.73$ ;  $P < 0.05$ , Figure 3). Compared to 0700 h, cortisol was lower at 1100, 1500 and 1900 h after the 3rd and 7th grouping ( $P < 0.05$ ). After the 3rd grouping, cortisol was also lower at 1100 and 1900 h than at 1500 h ( $P < 0.05$ ).


*Immune parameters*

The treatment  $\times$  stage of gestation interaction and the treatment effect were not significant for the numbers of lymphocytes and granulocytes (Figure 4). Towards the end of gestation, the number of lymphocytes decreased

( $F_{1,23} = 7.07$ ;  $P < 0.05$ ) whereas the number of granulocytes increased ( $F_{1,23} = 17.78$ ;  $P < 0.05$ ). Both groups of gilts had detectable levels of IgG against KLH on day 105 of gestation ( $3.7 \pm 0.7$  arbitrary units) but there was no difference between groups ( $F_{1,20} = 1.56$ ;  $P = 0.2$ ). The treatment  $\times$  stage of gestation interaction and the treatment effect were not significant for the proliferative index regardless of mitogen. Between days 70 and 105 of gestation, the proliferative index tended to increase for



**Figure 4** Numbers of blood lymphocytes and granulocytes before and after the 4-week stress procedure (on days 70 and 105 of gestation, DG70 and DG105). Values are mean  $\pm$  s.e. \* $P < 0.05$ , \*\*\* $P < 0.0001$ .



**Figure 5** Lymphocyte proliferative responses to lipopolysaccharide (LPS,  $6.25 \mu\text{g/ml}$ ), pokeweed mitogen (PWM,  $1.5 \mu\text{g/ml}$ ) or keyhole haemocyanin (KLH,  $10 \mu\text{g/ml}$ ) measured before and after the 4-week stress procedure (on days 70 and 105 of gestation). Values are mean  $\pm$  s.e. † $P < 0.1$ , \* $P < 0.05$ , \*\*\* $P < 0.0001$ .

ConA ( $F_{1,30} = 4.55$ ;  $P = 0.08$ ) and significantly increased for PWM ( $F_{1,19} = 8.56$ ;  $P < 0.01$ ), but decreased for LPS ( $F_{1,29} = 27.66$ ;  $P < 0.001$ ) (Figure 5).

#### Reproductive performances

Weight and back fat gains during the experimental period were similar in both groups (Table 1). None of the sows aborted. Litter size, number of stillborn and mummified piglets as well as litter weight and the average piglet weight were also similar in both groups. However, there was a tendency for a shorter length of gestation ( $F_{1,30} = 2.93$ ;  $P = 0.09$ ) in the S group.

#### Sub-classification of S gilts according to their success values

The mean AvSV in the different sub-groups of S gilts were as follows: H:  $0.44 \pm 0.12$ ; M:  $-0.32 \pm 0.13$ ; L:  $-0.86 \pm 0.06$ . The effects of AvSV classification on other behavioural variables and salivary cortisol are presented in Table 2. There was a significant effect of AvSV classification on AVs, showing that H gilts were the most aggressive ones ( $F_{2,13} = 23.34$ ;  $P < 0.001$ ). The duration of agonistic interactions was significantly lower in L gilts compared to M and

**Table 1** Effect of repeated social stress on weight gain between days 70 and 105 of gestation and reproductive performances of pregnant gilts

	Treatment groups		P values	
	Stress	Control	Treatment	Litter size
Weight gain (kg)	$22.9 \pm 2.3$	$23.6 \pm 0.9$	$>0.1$	
Back fat gain (cm)	$0.7 \pm 0.4$	$0.3 \pm 0.3$	0.1	
Gestation length (days)	$115.1 \pm 0.3$	$115.7 \pm 0.2$	0.09	$>0.1$
Litter size	$15.1 \pm 0.5$	$14.2 \pm 0.6$	$>0.1$	
Stillborn	$0.7 \pm 0.2$	$0.9 \pm 0.3$	$>0.1$	$>0.1$
Mummified piglets	$0.4 \pm 0.2$	$0.4 \pm 0.2$	$>0.1$	$>0.1$
Litter weight (kg)	$20.9 \pm 0.7$	$21.2 \pm 0.8$	$>0.1$	$<0.001$
Mean piglet weight (kg)	$1.40 \pm 0.04$	$1.50 \pm 0.03$	$>0.1$	$<0.001$

Litter size was included in the statistical model as a covariate for gestation length, number of stillborn piglets, mummified piglets, litter weight and mean piglet weight. Values are mean  $\pm$  s.e.

**Table 2** Effect of average success value on behavioural parameters (aggressive value, number of skin injuries, number and duration of agonistic interactions by grouping) and salivary cortisol (before, 1 and 19 h after the 1st, 4th and 8th groupings and between 0700 and 1900 h 2 days after the 3rd and 7th groupings)

	Control	H	M	L	P-value
Agonistic behaviour					
Aggressive value	–	0.70 ± 0.04 <sup>a</sup>	0.37 ± 0.07 <sup>b</sup>	0.12 ± 0.04 <sup>c</sup>	<0.001
Number of skin injuries	–	47 ± 6	39 ± 5	24 ± 5	>0.1
Number of AI	–	6.0 ± 1.1	4.5 ± 0.6	5.8 ± 0.6	>0.1
Duration of AI (min)	–	5.1 ± 0.4 <sup>a</sup>	4.8 ± 0.7 <sup>a</sup>	1.1 ± 0.2 <sup>b</sup>	<0.001
Salivary cortisol at grouping					
Just before grouping	1.8 ± 0.2 <sup>a</sup>	3.9 ± 0.9 <sup>b</sup>	2.2 ± 0.3 <sup>a</sup>	2.2 ± 0.3 <sup>a</sup>	0.05
One hour after grouping	1.7 ± 0.2 <sup>a</sup>	7.1 ± 2.4 <sup>b</sup>	3.4 ± 0.8 <sup>b</sup>	3.7 ± 0.5 <sup>b</sup>	<0.001
19 h after grouping	1.9 ± 0.2 <sup>a,t</sup>	3.3 ± 0.7 <sup>a,b,t</sup>	3.7 ± 0.3 <sup>b</sup>	2.5 ± 0.3 <sup>b</sup>	<0.01
Diurnal rhythm of salivary cortisol					
0700 h	2.8 ± 0.3 <sup>a</sup>	10.8 ± 2.5 <sup>b,t</sup>	4.5 ± 1.0 <sup>a,b,t</sup>	2.7 ± 0.6 <sup>a</sup>	<0.001
1100 h	1.5 ± 0.2 <sup>a</sup>	4.4 ± 0.9 <sup>b</sup>	2.9 ± 0.6 <sup>a,b</sup>	1.9 ± 0.3 <sup>a,b</sup>	<0.05
1500 h	2.4 ± 0.5	6.2 ± 2.6	2.3 ± 0.4	2.6 ± 0.8	>0.1
1900 h	1.7 ± 0.2 <sup>a</sup>	3.6 ± 1.0 <sup>b</sup>	2.0 ± 0.5 <sup>a,b</sup>	2.3 ± 0.8 <sup>a,b</sup>	<0.05

Values are mean ± s.e. Within a row, different superscript letters indicate significant differences at  $P < 0.05$  and values with superscript t tend to differ each other ( $0.05 < P < 0.1$ ).

H=high; M=medium; L=low; AI=agonistic interactions; AvSV=average success value.

$n = 6/\text{group}$ .

H gilts ( $P < 0.01$ ) but AvSV did not influence the numbers of skin lesions and agonistic interactions.

For cortisol levels at 30 min before grouping, AvSV ( $F_{3,30} = 2.92$ ;  $P = 0.05$ ) and the interaction between the grouping number × AvSV ( $F_{6,64} = 2.22$ ;  $P = 0.05$ ) were significant. This latter effect was due to the fact that H gilts displayed higher cortisol levels than C and L gilts on the 4th grouping (H:  $6.8 \pm 3.2$  ng/ml, M:  $1.9 \pm 0.5$  ng/ml, L:  $1.4 \pm 0.3$  ng/ml, C:  $1.3 \pm 0.2$  ng/ml;  $P < 0.05$ ). Cortisol levels were also significantly higher in all sub-groups of S gilts than in C gilts at 1 and 19 h after grouping except in H gilts at 19 h where only a tendency was observed ( $P = 0.09$ ). For diurnal cortisol variations, AvSV effect was significant at 0700 h ( $F_{3,30} = 9.48$ ;  $P < 0.001$ ), 1100 h ( $F_{3,30} = 5.40$ ;  $P < 0.05$ ) and 1900 h ( $F_{3,30} = 2.97$ ;  $P < 0.05$ ), whereas the interaction day of sampling × AvSV was not. H gilts had higher cortisol levels than C gilts at 0700, 1100 and 1900 h ( $P < 0.05$ ), and than L gilts ( $P < 0.01$ ) only at 0700 h. M and L gilts never differed from C gilts at any of these time points.

Immune measures, weight of the sows and reproductive data were not affected by AvSV except for gestation length ( $F_{1,30} = 2.93$ ;  $P < 0.05$ ). L gilts had shorter gestation ( $114.5 \pm 0.3$  days) than C gilts ( $115.7 \pm 0.2$  days,  $P < 0.05$ ).

## Discussion

To our knowledge, the present study is the first one to report the influence of repeated social stress applied to pregnant gilts during the last third of their gestation on acute and delayed cortisol response, immune function of sows and pregnancy outcome. Our data demonstrate a cortisol response to successive groupings in all gilts without marked negative effects on the health and pregnancy outcome.

In the S group, grouping and relocation of gilts with an unfamiliar and unrelated partner led to vigorous fights during the first hours following the grouping. This agonistic behaviour after grouping is common in animals that form social groups and is part of the process leading to the establishment of a social hierarchy in the newly formed group (Ewbank and Meese, 1971; Meese and Ewbank, 1972). Moreover, our data show that grouping with an unfamiliar conspecifics induced a stimulation of the HPA axis in pregnant gilts because high levels of salivary cortisol were observed 1 and 19 h after grouping during the whole experimental period. An increased cortisol response due to social stress was already reported in growing pigs submitted to repeated groupings (Olsson *et al.*, 1999; Coutellier *et al.*, 2007) and in pregnant sows submitted to one or two groupings (Tsuma *et al.*, 1996; Jarvis *et al.*, 2006). In contrast, Soede *et al.* (2006) failed to demonstrate a clear increase in salivary cortisol in gilts submitted to repeated groupings around insemination. The increase in cortisol shortly after social grouping can be explained by the physical activity and the injuries associated with the fights and by the psychological stress associated with the appraisal of the fighting ability of the unfamiliar partner. Our data show that high levels of cortisol were still observed 2 days (diurnal rhythm assessment) and 3 days (measurement 30 min before the 4th and 8th grouping) after mixing in gilts presenting a high AvSV. Long-term effects of grouping were also observed by Coutellier *et al.* (2007) in growing pigs submitted to repeated groupings and by Tsuma *et al.* (1996) in pregnant gilts after a single mixing. However, in our study, salivary cortisol measured 2 and 3 days after grouping was similar in C gilts, medium- and low-success S gilts. The H gilts were more aggressive, and it can be hypothesized that additional fights occurred after the period



of observation, inducing new stress reactions with adrenal axis stimulation. Such variability associated with the social rank of the pigs was also described by Otten *et al.* (1997) in growing pigs of high or low social rank reintroduced in a new group after 10 days of social isolation and by Mendl *et al.* (1992) in pregnant gilts submitted to a single grouping during mid-gestation. In agreement with our results, Otten *et al.* (1997) found that aggressive high-ranking animals showed higher cortisol release in response to a grouping with unfamiliar conspecifics than low-ranking pigs. In contrast, Mendl *et al.* (1992) reported higher basal cortisol levels in gilts of intermediate rank in comparison to high- and no-success gilts 5 weeks after grouping. This discrepancy can be explained by the fact that Otten *et al.* (1997) and the present study investigated the cortisol response relatively shortly after grouping, whereas Mendl *et al.* (1992) investigated it 5 weeks later. Shortly after grouping, the social hierarchy is still being established, whereas several weeks after grouping, social interactions between individuals were more predictable and the level of aggression was lower.

We observed that the number and length of agonistic interactions as well as cortisol levels short- (1 h) and long-term (19 h and 2 days) after grouping declined as the number of groupings increased. Such a decrease in agonistic behaviour has already been reported in growing pigs submitted to repeated groupings (Giersing and Andersson, 1998; Coutellier *et al.*, 2007). Contrarily, Olsson *et al.* (1999) in growing pigs and Soede *et al.* (2006) in gilts did not show a reduction of the endocrine response to repeated grouping. However, in both studies, grouping was only performed once a week over a period of 5 weeks and this frequency might have been insufficient to induce habituation. Moreover, in the study from Soede *et al.* (2006), grouping had only minor effects on behaviour and cortisol release. The observed decline of offensive acts in our study could reflect the adoption of a new coping strategy to a changing and aversive environment (Schouten and Wiegant, 1997; Bolhuis *et al.*, 2004). Gilts may have developed a more efficient strategy to establish a social hierarchy with less agonistic interactions (Puppe *et al.*, 2008) and hence to limit the energetic cost and the number of injuries. In the same way, the decline of cortisol response to grouping towards the end of the experiment can be attributed to the decrease in the intensity of the stress perceived by the gilts and/or to a decrease in the adrenal axis sensitivity. The first hypothesis is supported by the decrease in the number and duration of agonistic behaviours and in the number of injuries. ACTH or corticotropin-releasing hormone challenges would be needed to confirm the second hypothesis. Finally, despite a decreased response at the end of the stress procedure, acute endocrine and behavioural responses to grouping were still observed during the last grouping and cortisol levels remained higher than in the control group between the 7th and the last grouping.

Acute and chronic social stress are known to change the distribution of immune cells and to alter the immune status

in pigs (Tuchscherer *et al.*, 1998; de Groot *et al.*, 2001; Niekamp *et al.*, 2007) and in other mammal species (for review see Bohus *et al.*, 1991; Bartolomucci, 2007). Typically, the lymphocytes/granulocytes ratio and the lymphocyte abilities to proliferate in the presence of mitogens decrease after acute or chronic stress. It is also assumed that stress inhibits the cellular adaptive immunity (Th1) in favour of Th2 humoral immunity (Elenkov, 2004). In the present study, no alteration in immune cell distribution and no shift from cellular to humoral immunity were found in stressed gilts. An absence of social stress effects on humoral immunity was also reported in 9-month-old pregnant gilts (Mendl *et al.*, 1992). The lacking effect of cortisol elevation on the immune function of the pregnant gilts in our study may be explained by the particular physiological state of the adult pregnant female. First, immune effects of social stress were observed in non-pregnant pigs; however, it is possible that pregnancy, which is associated with strong immune modifications (Wegmann *et al.*, 1993), abrogated the stress-induced immune shift. For instance, the decrease of lymphocytes/granulocytes ratio towards the end of gestation could explain why we and others (Stefansky *et al.*, 2005) could not find any additional effects of stress on this parameter. Moreover, females seem to be less sensitive than castrated males to social stress-induced immune suppression. For example, de Groot *et al.* (2001) found a suppressed humoral immunity in mixed barrows, whereas no differences were found in mixed gilts. Finally, the lack of effect of cortisol elevation on the immune function of the pregnant gilts can be explained by the age-related decreased sensitivity of porcine leukocytes to corticosteroids (Yang and Schultz, 1986).

Stress can negatively influence reproductive processes in pigs such as implantation and foetal growth and may lead to abortion (for review see von Borell *et al.*, 2007). In our experiment, the stress model had no negative impact on pregnancy maintenance or intrauterine mortality since the size of the litter at birth was similar in S and C groups. A non-significant decrease of 0.6 day in the gestation length was observed in S gilts, which was due to a shorter gestation in low-AvSV individuals, but this had no obvious consequences on piglet weight or mortality at birth. Cortisol elevation during pregnancy is suspected to generate prenatal stress and alter the birth weight and viability of the progeny (Hausmann *et al.*, 2000; Tuchscherer *et al.*, 2002). A high variability is observed for piglet birth weight among prenatal stress studies, probably due to differences in the methodologies used. Effects varied from higher birth weight in the case of maternal ACTH administration (Kanitz *et al.*, 2006) to lack of effect in case of maternal restraint and social stress (Tuchscherer *et al.*, 2002; Jarvis *et al.*, 2006) and to lower birth weight in the case of maternal cortisol treatment (Kranendonk *et al.*, 2006). A lower birth weight of the litter was observed in pregnant sows that displayed higher cortisol levels in relation to their social rank (Mendl *et al.*, 1992). Therefore, additional work is needed to clarify the relationships between the adrenal activity of the pregnant mother and foetal growth in pigs.

## Conclusion

The present study demonstrates that repeated social grouping during the last third of pregnancy in gilts is able to induce behavioural and cortisol responses as described in growing non-pregnant pigs. Among stressed gilts, the outcome of agonistic interactions seems to influence the cortisol release with sows of high success having elevated cortisol levels for a longer time after grouping than other gilts. Assuming that cortisol is a reliable marker of stress, this result suggests that repeated groupings induce a state of stress, especially for pregnant sows that are dominant and more aggressive. However, the social stress paradigm used did not affect either the immune functions of the pregnant gilts nor foetal viability or growth. Considering the effects of maternal-mediated stress during pregnancy on the physiology of the progeny in many species, further work should focus on the physiological, immunological and behavioural consequences of maternal social stress on the offspring in pigs.

## Acknowledgements

The authors thank the staff of the SENAH Research and Experimental pig unit, Mounier Anne-Marie, Lefebvre Michel and Massard Michel for their technical assistance and the AFSSA unit of Ploufragan, Kuntz Gaëlle, Jamin Agnès, for their assistance for the haematological analyses. This study received financial support from the INRA PHASE department.

## References

Arck PC 2001. Stress and pregnancy loss: role of immune mediators, hormones and neurotransmitters. *American Journal of Reproductive Immunology* 46, 117–123.

Barbazanges A, Piazza PV, Le Moal M and Maccari S 1996. Maternal glucocorticoid secretion mediates long-term effects of prenatal stress. *Journal of Neuroscience* 16, 3943–3949.

Bartolomucci A 2007. Social stress, immune functions and diseases in rodents. *Frontiers in Neuroendocrinology* 28, 28–49.

Bohus B, Koolhaas JM, De Ruiter AJH and Heijnen CJ 1991. Stress and differential alterations in immune system functions: conclusions from social stress studies in animals. *Netherlands Journal of Medicine* 39, 306–315.

Bolhuis JE, Schouten WG, De Leeuw JA, Schrama JW and Wiegant VM 2004. Individual coping characteristics, rearing conditions and behavioural flexibility in pigs. *Behavioural Brain Research* 152, 351–360.

Coussons-Read ME, Okun ML and Nettles CD 2007. Psychosocial stress increases inflammatory markers and alters cytokine production across pregnancy. *Brain, Behaviour, and Immunity* 21, 343–350.

Coutellier L, Arnould C, Boissy A, Orgeur P, Prunier A, Veissier I and Meunier-Salaün MC 2007. Pig's responses to repeated social regrouping and relocation during the growing-finishing period. *Applied Animal Behaviour Science* 105, 102–115.

De Groot J, Ruis MAW, Scholten JW, Koolhaas JM and Boersma WJA 2001. Long-term effects of social stress on antiviral immunity in pigs. *Physiology and Behavior* 73, 145–158.

Elenkov IJ 2004. Glucocorticoids and the Th1/Th2 balance. *Annals of the New York Academy of Sciences* 1024, 138–146.

Ewbank R and Meese GB 1971. Aggressive behaviour in groups of domesticated pigs on removal and return of individuals. *Animal Production* 13, 685–693.

Fowden AL, Dino AG and Forhead AJ 2006. Intrauterine programming of physiological systems: causes and consequences. *Physiology* 21, 29–37.

Giersing M and Andersson A 1998. How does former acquaintance affect aggressive behaviour in repeatedly mixed male and female pigs? *Applied Animal Behaviour Science* 59, 297–306.

Hausmann MF, Carroll JA, Weesner GD, Daniels MJ, Matteri RL and Lay DC 2000. Administration of ACTH to restrained, pregnant sows alters their pigs' hypothalamic-pituitary-adrenal (HPA) axis. *Journal of Animal Science* 78, 2399–2411.

Jarvis S, Moinard C, Robson SK, Baxter E, Ormandy E, Douglas AJ, Seckl JR, Russell JA and Lawrence AB 2006. Programming the offspring of the pig by prenatal social stress: neuroendocrine activity and behaviour. *Hormones and Behavior* 49, 68–80.

Joachim R, Zencussen AC, Polgar B, Douglas AJ, Fest S, Knackstedt M, Klapp BF and Arck PC 2003. The progesterone derivative dydrogesterone abrogates murine stress-triggered abortion by inducing a Th2 biased local immune. *Steroids* 68, 931–940.

Kanitz E, Tuchscherer M, Puppe B, Tuchscherer A and Stabenow B 2004. Consequences of repeated early isolation in domestic piglets (*Sus scrofa*) on their behavioural, neuroendocrine, and immunological responses. *Brain, Behaviour, and Immunity* 18, 35–45.

Kanitz E, Otten W and Tuchscherer M 2006. Changes in endocrine and neurochemical profiles in neonatal pigs prenatally exposed to increased maternal cortisol. *Journal of Endocrinology* 191, 207–220.

Kranendonk G, Hopster H, Fillerup M, Ekkel ED, Mulder EJJ, Wiegant VM and Taverne MAM 2006. Lower birth weight and attenuated adrenocortical response to ACTH in offspring from sows that orally received cortisol during gestation. *Domestic Animal Endocrinology* 30, 218–238.

McGlone JJ, Salak JL, Lumpkin EA, Nicholson RI, Gibson M and Norman RL 1993. Shipping stress and social status effects on pig performance, plasma cortisol, natural killer cell activity, and leukocyte numbers. *Journal of Animal Science* 71, 888–896.

Meese GB and Ewbank R 1972. A note on instability of the dominance hierarchy and variations in level of aggression within groups of fattening pigs. *Animal Production* 14, 359–362.

Mendl M, Zanella AJ and Broom DM 1992. Physiological and reproductive correlates of behavioural strategies in female domestic pigs. *Animal Behaviour* 44, 1107–1121.

Merlot E, Couret D and Otten W 2008. Prenatal stress, fetal imprinting and immunity. *Brain, Behavior, and Immunity* 22, 42–51.

Neumann ID, Johnstone HA, Hatzinger M, Liebsch G, Shipston M, Russell JA, Landgraf R and Douglas AJ 1998. Attenuated neuroendocrine responses to emotional and physical stressors in pregnant rats involve adenohipophysial changes. *Journal of Physiology* 508, 289–300.

Niekamp SR, Sutherland MA, Dahl GE and Salak-Johnson JL 2007. Immune responses of piglets to weaning stress: impact of photoperiod. *Journal of Animal Science* 85, 93–100.

Olsson IAS, De Jonge FH, Schuurman T and Helmond FA 1999. Poor rearing conditions and social stress in pigs: repeated social challenge and the effect on behavioural and physiological responses to stressors. *Behavioural Processes* 46, 201–215.

Otten W, Puppe B, Stabenow B, Kanitz E, Schön PC, Brüßow KP and Nürnberg G 1997. Agonistic interactions and physiological reactions of top- and bottom-ranking pigs confronted with a familiar and an unfamiliar group: preliminary results. *Applied Animal Behaviour Science* 55, 79–90.

Otten W, Puppe B, Kanitz E, Schön PC and Stabenow B 2002. Physiological and behavioral effects of different success during social confrontation in pigs with prior dominance experience. *Physiology and Behaviour* 75, 127–133.

Puppe B 1998. Effects of familiarity and relatedness on agonistic pair relationships in newly mixed domestic pigs. *Applied Animal Behaviour Science* 58, 233–239.

Puppe B, Langbein J, Bauer J and Hoy S 2008. A comparative view on social hierarchy formation at different stages of pig production using sociometric measures. *Livestock Science* 113, 155–162.

Salak-Johnson JL and McGlone JJ 2007. Making sense of apparently conflicting data: stress and immunity in swine and cattle. *Journal of Animal Science* 85, 81–88.

Salak-Johnson JL, McGlone JJ and Norman RL 1996. In vivo glucocorticoid effects on porcine natural killer cell activity and circulating leukocytes. *Journal of Animal Science* 74, 584–592.

- Schouten WG and Wiegant VM 1997. Individual responses to acute and chronic stress in pigs. *Acta Physiologica Scandinavica Supplementum* 640, 88–91.
- Schrama JW, Schouten JM, Swinkels JWGM, Gentry JL, De Vries Reilingh G and Parmentier HK 1997. Effect of hemoglobin status on humoral immune response of weanling pigs differing in coping styles. *Journal of Animal Science* 75, 2588–2596.
- Soede NM, Van Sleuwen MJW, Molenaar R, Rietveld FW, Schouten WPG, Hazeleger W and Kemp B 2006. Influence of repeated regrouping on reproduction in gilts. *Animal Reproduction Science* 96, 133–145.
- Soede NM, Roelofs JB, Verheijen RJ, Schouten WP, Hazeleger W and Kemp B 2007. Effect of repeated stress treatments during the follicular phase and early pregnancy on reproductive performance of gilts. *Reproduction in Domestic Animal* 42, 135–142.
- Statistical Analysis Systems Institute 1999. *Statistical analysis systems*, released 8.01. Statistical Analysis Systems Institute Inc., Cary, NC, USA.
- Stefansky V, Raabe C and Schulte M 2005. Pregnancy and social stress in female rats: influences on blood leukocytes and corticosterone concentrations. *Journal of Neuroimmunology* 162, 81–88.
- Tsuma VT, Einarsson S, Madej A, Kindahl H, Lundeheim N and Rojkkittikhun T 1996. Endocrine changes during group housing of primiparous sows in early pregnancy. *Acta Veterinaria Scandinavica* 37, 481–489.
- Tuchscherer M, Puppe B, Tuchscherer A and Kanitz E 1998. Effects of social status after grouping on immune, metabolic and endocrine responses in pigs. *Physiology and Behavior* 64, 353–360.
- Tuchscherer M, Kanitz E, Otten W and Tuchscherer A 2002. Effects of prenatal stress on cellular and humoral immune responses in neonatal pigs. *Veterinary Immunology and Immunopathology* 86, 195–203.
- Von Borell E, Dobson H and Prunier A 2007. Stress, behaviour and reproductive performance in female cattle and pigs. *Hormones and Behavior* 52, 130–138.
- Wegmann TG, Lin H, Guilbert L and Mosmann TR 1993. Bidirectional cytokine interactions in the maternal–fetal relationship: is successful pregnancy a Th2 phenomenon? *Immunology Today* 14, 353–356.
- Weinstock M 1997. Does prenatal stress impair coping and regulation of hypothalamo–pituitary–adrenal axis? *Neuroscience and Biobehavioral Reviews* 21, 1–10.
- Wiebold JL, Stanfield PH, Becker WC and Hillers JK 1986. The effect of restraint stress in early pregnancy in mice. *Journal of Reproduction and Fertility* 78, 185–192.
- Yang WC and Schultz RD 1986. Effect of corticosteroids on porcine leukocytes: age-related effects of corticosteroid inhibition on porcine lymphocyte responses to mitogens. *Veterinary Immunology and Immunopathology* 13, 19–29.