



HAL
open science

Nutritional values of sugarcane products in local Caribbean growing pigs

Xavier Xande, Carole Regnier, Harry Archimède, Bruno Bocage, Jean Noblet,
David Renaudeau

► **To cite this version:**

Xavier Xande, Carole Regnier, Harry Archimède, Bruno Bocage, Jean Noblet, et al.. Nutritional values of sugarcane products in local Caribbean growing pigs. *Animal*, 2010, 4 (5), pp.745-754. 10.1017/S175173110999173X . hal-02665280

HAL Id: hal-02665280

<https://hal.inrae.fr/hal-02665280v1>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nutritional values of sugarcane products in local Caribbean growing pigs

X. Xandé¹, C. Régnier¹, H. Archimède¹, B. Bocage², J. Noblet³ and D. Renaudeau^{1†}

¹INRA, Unité de Recherches Zootechniques, UR143, 97170 Petit-Bourg, Guadeloupe, French West Indies, France; ²INRA, Plateforme Tropicale de l'Expérimentation sur l'Animal, UE503, 97170 Petit-Bourg, Guadeloupe, French West Indies, France; ³INRA, UMR 1079 Systèmes d'Élevage, Nutrition Animale et Humaine, 35590 St-Gilles, France

(Received 25 August 2008; Accepted 10 December 2009; First published online 13 January 2010)

A total of 24 castrated male Creole were used for digestibility studies on three sugarcane (SC) products: final molasses, SC juice and ground SC. Digestibility of macronutrients were determined in three consecutive experimental trials between 35 and 60 kg BW. The first trial measured the effect of a constant rate of incorporation of those SC products in a control diet (23% on a dry matter (DM) basis) on digestibility of energy and nutrients. The second and the third trials were designed to determine the effects of increasing rates of inclusion of SC juice and ground SC in the control diet on the digestibility of nutrients and energy. The DM content of molasses, SC juice and ground SC were 74.3%, 19.3% and 25.8%, respectively. Free sugar contents were 73.8%, 81.6% and 51.5% of DM for molasses, SC juice and ground SC, respectively. In contrast with molasses and SC juice, NDF content of ground SC was very high (40.3% DM). Energy digestibility coefficients (DC) were 99.0% and 83.6% for SC juice and molasses, respectively. For ground SC, amounts and composition of intakes differed from feed allowance as a consequence of chewing-then-spitting out most of the fibrous contents of ground SC. Expressed as a percentage of energy intake and energy allowance, DC of energy for ground SC were 68.6% and 31.9%, respectively. On an intake basis, digestible energy (DE) and metabolisable energy (ME) values were higher for SC juice (17.3 and 17.2 MJ/kg DM, respectively) than for molasses (13.1 and 12.6 MJ/kg DM, respectively) or ground SC (12.3 and 11.8 MJ/kg DM, respectively). On an allowance basis, the corresponding values for ground SC were only 5.7 and 5.6 MJ/kg DM, respectively. The gradual inclusion rate of SC juice in the diet up to 66% resulted in a linear increase of the DC of the diet organic matter and energy ($P < 0.001$) by 0.10% per 1% in SC juice DM. On the other hand, the partial substitution of the control diet by ground SC up to 74% resulted in a linear ($P < 0.05$) reduction of digestibility of nutrients and energy. Each 1% increase of ground SC incorporation rate resulted in a 0.13% and 0.61% linear decrease of DC of the energy, expressed as a percentage of energy intake or energy allowance, respectively. In conclusion, our study provides updated energy values of SC products usable as energy sources for feeding pigs.

Keywords: Creole pig, sugarcane products, digestibility, energy value

Implications

The aim of this study was to increase the current knowledge on nutritional values of sugarcane (SC) products and by-products. Our results provide original data especially for digestibility values of nutrients and energy values of SC juice and ground SC. In particular for ground SC, our results suggest that the digestive utilisation of nutrients and energy was closely dependant on changes in the individual feeding behaviour. From this point of view, additional studies are warranted to test if the genotype (local v. exotic breeds) could affect the digestive utilisation of ground SC. Data obtained in this study should also allow practical feed

formulation of mixed diets for pig production from chemical characteristics of these SC products. In particular, these data should be essential to define a dynamic model to analyse the bio-technical and socio-economic interactions in mixed farming systems based on the use of SC as an energy source in the diet. However, SC products, particularly juice and ground SC that contains juice, are easily fermentable stuffs. Therefore, care must be taken to avoid fermentation of these materials, as it would alter their nutritional value.

Introduction

Cereal grains which are the basis of livestock production in the more-developed countries face a competing demand to

† E-mail: David.Renaudeau@antilles.inra.fr

feed the increasing human population (Preston, 1995). This scenario will give opportunities to alternative feeds that can replace cereals in livestock diets. Sugarcane (SC) is a perennial tropical crop with a high-energy yield per unit area (80 tons dry matter (DM)/ha; Preston and Leng, 1989). In several tropical countries (Cuba, Dominican Republic, Costa Rica, Guadeloupe, Vietnam), many studies and development of feeding strategies for pig production have been based on the use of this local resource and/or its by-products. Most studies were focused on the growth performance obtained when pigs were fed SC-based diets, especially as juice or molasses but also as ground cane stalks (Preston *et al.*, 1968; González *et al.*, 2006; Xandé *et al.*, 2009). The determination of digestible and metabolisable energy (ME) values of these SC products is a first step to their evaluation as feedstuffs for pig feeding. However, there is little information on nutritional values of SC products available that can allow practical feed formulation of mixed diets. Regarding those products, digestive utilisation of SC molasses in pigs is the most described in the literature (Ly and Preston, 1969; Christon and Le Dividich, 1978; Bayley *et al.*, 1983). In contrast, little is published about the digestive utilisation of nutrients and energy of SC juice and ground SC stalks.

In SC production area, small scale farmers traditionally use whole or chopped SC as a feeding source for pigs (Bravo *et al.*, 1996). The Caribbean native pigs, that is, Creole (CR) pigs are well-known for their rusticity and according to their low nutrient requirements (Renaudeau *et al.*, 2006), they are likely to have comparative advantages over 'exotic' high-performance genotypes, when the feeding system is based on unconventional feedstuffs such as SC products and by-products.

The objectives of our study were: (1) to measure the *in vivo* total tract digestibility of dietary energy or nutrients of SC final molasses, SC juice and ground SC stalks in growing CR pigs, (2) to measure *in vivo* the effect of increased inclusion rates of SC juice or ground SC stalks in the diet, on total tract digestibility values in growing CR pigs.

Material and methods

Care and use of animal were performed according to the certificate of authorisation to experiments on living animals, number A-971-18-01 (issued by the French ministry of agriculture to the head of the experiment unit).

Experimental design and animal management

Nutritional values of SC products (final molasses, juice and ground cane stalks) were studied on a total of 24 castrated CR pigs at the experimental facilities of the Institut National de la Recherche Agronomique in Guadeloupe (latitude 16°N, longitude 61°W). The experiment was conducted in two replication of 12 pigs each. Within each replication, three blocks of four littermates were chosen where each littermate was randomly assigned to one dietary treatment. All in all, 6 replicates per dietary treatments were obtained. At about 30 kg BW, pigs were moved to individual digestibility cages in an open front experimental room. Climatic parameters were not controlled and followed those of the outdoor conditions. Nutritional values of SC products (final molasses, SC juice and ground SC) were studied in three consecutive experimental trials. The first trial was designed to measure the effect of a constant rate (23% on a DM basis) of incorporation of SC products (SC molasses, SC juice and ground SC) in a control soybean meal-corn diet (19.8% DM of CP and 16.6 MJ DE/kg DM; DE, digestible energy) on feed digestibility. The second and the third trials were designed to determine the effects of increasing rates of inclusion of SC juice and ground SC on digestibility of nutrients and energy. Within each trial, pigs were adapted to the diet for 10 days before total collection of faeces and urine for 8 days. During the adaptation and the collection periods, the amount of feed offered daily was 35 g DM/kg per BW. The basal diet was a commercially available pig-grower feed in pellet form (Table 1). The molasses were also produced commercially (Gardel SA, Moule, Guadeloupe, French West Indies). Twice a week, 1 ton of fresh SC was delivered to our experimental facilities, cleaned and

Table 1 Chemical composition of the control diet and the SC products¹

Item	Control	SC molasses	SC juice	Ground SC allowed	Ground SC intake ²
DM (%)	88.4	74.3	19.3	25.8	43.7
Chemical composition (% DM)					
Ash	5.9	15.1	1.6	1.5	1.0
CP	19.8	4.7	1.4	1.2	0.5
Crude fat	4.0	–	–	–	–
NDF	11.2	–	–	40.3	21.2
ADF	3.6	–	–	26.6	14.2
ADL	–	–	–	4.4	0.0
Starch	51.8	–	–	–	–
Free sugars	8.4	73.8	81.6	51.5	78.0
Gross energy (MJ/kg DM)	18.7	15.7	17.5	17.9	17.6

SC = sugarcane; DM = dry matter.

¹Mean chemical composition measured on a total of 18, 6, 24 and 24 samples for the soybean meal-corn control diet, SC molasses, SC juice and ground SC, respectively.

²Average values calculated from results obtained during trials 1 and 3 according to the difference between measured values of ground SC allowed and refusals.

milled for juice production or ground in small size particles (less than 1 cm). The SC juice and the ground SC were stored at 4°C to prevent any fermentation reactions. The average chemical composition of the SC products is presented in Table 1. For each experimental trial, one littermate was fed with the control diet and the other three littermates were fed with an experimental diet in which a constant (trial 1) or a gradual amount (trials 2 and 3) of the control diet was replaced by a SC product. The experimental diets were prepared by mixing molasses, SC juice or ground SC with the basal diet (8, 8 and 13 g DM/day per kg BW; trial 1), various amounts of SC juice with basal diet (8, 16 and 24 g DM/day per kg BW; trial 2), and various amounts of ground SC with basal diet (13, 26 and 39 g DM/day per kg BW; trial 3) just before feeding. For ground SC, the following protocol was applied for the distribution of the diet: first, the control diet was distributed and once it was completely consumed, the ground SC was distributed. According to Xandé *et al.* (2009), with a ground SC diet pigs show a specific 'sucking-spitting' feeding behaviour and the DM extraction rate, that is, DM intake/DM offered, of ground SC averaged 62%; this ratio was taken into account in order to allow a constant DM intake of SC products (i.e. 8 g DM/day per kg BW; trial 1) or a similar variation of DM intake from ground SC in trial 3 than from SC juice in trial 2 (8, 16 and 24 g DM/day per kg BW). Between each experimental trial, each littermate was randomly assigned to one dietary treatment. All animals had free access to water from one individual low-pressure nipple drinker.

Measurements

All the pigs were weighed at the beginning and at the end of the adaptation period, and at the end of the collection period. Every morning, any feed refusal and spillage were collected. Samples of allowed feed and refusals were dried at 103°C for 24 h to measure the DM content. One sample of the control diet was also taken daily and pooled at the end of the sampling period for DM determination and further chemical analyses. Samples of molasses, SC juice and ground SC were daily and individually collected, stored at -20°C and pooled at the end of the collection period. One representative sample of SC products was dried at 65°C for 96 h for further chemical analyses. During the 8-day collection period, faeces and urine were daily and individually collected and stored at 4°C. Urine was collected into a flask containing sulphuric acid (0.1 Nitrogen (N); 10% v/v) in order to avoid ammonia losses during collection and storage. Faeces and urine were separately weighed, homogenised and sub-sampled at the end of the collection period for analysis (digestibility sample). One faeces sample was heat dried (48 h at 103°C) for DM determination and the second was freeze-dried for further chemical analyses.

Chemical analyses

The diet samples were analysed for DM, ash, organic matter (OM), crude protein (CP, N × 6.25), ether extract (EE) and starch, according to the AOAC methods (AOAC, 1990), for

free sugars according to Tollier and Robin (1979) and for gross energy (GE) content using an adiabatic bomb calorimeter (IKA, C5000, Staufen, Germany). Dry faecal samples were analysed for DM, ash, EE and CP and fresh samples of urine for N. Cell wall components (NDF, ADF and ADL) in diet samples were determined according to the Van Soest and Wine method (1967).

Calculations and statistical analyses

From the results of the digestibility trial, apparent digestibility coefficients (DC) of DM, OM, CP, ash, EE, free sugars and energy were calculated for each pig. In the case of ground SC, it was expressed relative to allowance or intake quantities. DE content was estimated as the difference between measured GE and estimated energy loss in faeces (Noblet, personal communication).¹ ME content of diets was calculated as the difference between DE and energy loss in urine estimated from N loss (g/day) according to the equation of Le Goff and Noblet (2001).² N retained was obtained as the difference between N intake and N losses in faeces and urine.

In a first approach, we statistically controlled that measurements performed in trials 2 and 3 were not affected by the dietary treatment offered in trials 1 and 2, respectively. For that purpose, we used the MIXED procedure of SAS (1997) with an unstructured variance co-variance matrix. According to Wald Z test (SAS Institute, 1997), the covariance estimates were not significantly different from zero. We concluded there was no correlation between two successive measurements carried out on the same animal, suggesting that the adaptation period allowed suppressing the remnant effects. Then, in a second approach, we analysed the results obtained for each trial independently using an ANOVA (Proc GLM, SAS Institute, 1997) including the effects of diet, replicate, block within replicate and the interaction between diet and replicate. Comparison of means was performed according to the PDIFF option using Tukey test for contrasts (SAS Institute, 1997). In trial 1, the digestive utilisation of SC products was calculated according to the difference method (Noblet and Shi, 1994). Assuming that energy and nutrient digestibility of the control diet was constant in the experimental diets, we calculated DC of energy and dietary chemical fractions of SC products from the rate of incorporation in feed and the DC of control and mixed (control + SC products) diets. For trials 2 and 3, linear or quadratic regressions between DC measured on complete diet and rate of incorporation of SC juice or ground SC were calculated. According to that chemical composition of allowed and intake feed for dietary treatments including ground SC, DC of ground SC were calculated on a intake basis (method 1) and allowance basis (method 2).

¹ Energy loss in faeces (MJ/kg DM) = 18.73 - 1.92 × A + 2.23 × EE + 4.07 × N, with A, EE and N for ash, ether extract and nitrogen contents in faeces (g/kg DM), $R^2 = 0.93$, r.s.d. = 0.35.

² Energy loss in (kJ/day) = 345 + 31.1 × N loss in urine (g/day), $R^2 = 0.94$, r.s.d. = 110. This equation was calculated from a 641 sets of data.

Results

General aspects and chemical composition of the dietary ingredients

One pig fed diet 4 (control + ground SC) was removed during the first replicate of trial 1 after 5 days of sampling period because of leg problems, which had consequences on its daily consumption. For trial 2, this pig was replaced by a littermate pig with a similar initial BW. The results obtained with another pig fed diet 2 (control + molasses) during the second replicate of trial 1 were removed from the ANOVA because of 2 days of diarrhoea observed after the 5th day of sampling period. This pig was successfully treated and then used in the following two trials. For both replicates, sampling period in trial 3 lasted only 7 days instead of 8 days for practical reasons (day off in the first replicate), and weather reasons (tropical hurricane in the second replicate).

As shown in Table 1, ground SC refusal composition was obviously different from allowed ground SC composition, with a chewed aspect (trial 1 and 3). With ground SC, pigs showed a peculiar feeding behaviour: they chew the cane stalks to extract the juice and then spit out most of the fibrous residue. In connection with a ratio intake/allowance for ground SC DM of 50.8%, the composition of DM intake from ground SC was different from the DM composition of ground SC offered; NDF and ADF contents decreased from 40.3% to 21.2% and from 26.6% to 14.2%, respectively, while free sugar contents increased from 51.5% to 78.0%. In addition, the amount of ground SC refusals linearly increased when ground SC allowance increased from 0% to 74% of total feed offered (trial 3; Table 6). Then, expressed as a percentage of ground SC allowance, DM extraction rate from ground SC (DM intake/DM allowed) was maintained constant and averaged 36%. In SC products, measured GE was numerically lower in SC molasses than in SC juice and ground SC allowed and intake (15.7 v. 17.7 MJ/kg DM, respectively).

Effect of SC products on total tract digestibility of energy and nutrients (trial 1)

According to the experimental design, there was no statistical difference for average BW and total DM intake ($P > 0.05$) between diets (33.6 kg BW and 33.1 g DM/day per kg BW, respectively) during the first trial (Table 2). In this trial, the faeces DM was not affected by the inclusion of SC products in the control diet ($P > 0.05$). The incorporation of molasses did not significantly affect the total tract digestibility of DM, OM and energy when compared with the control diet. However, DC of CP was lower (82.7% v. 86.4%; $P < 0.01$) and DC of EE and sugars were higher than in the control diet (88.3% v. 82.1%, $P < 0.05$; and 97.6% v. 96.1%, $P < 0.001$, respectively). In comparison with the control treatment, the inclusion of SC juice in the diet increased DC of DM, OM, free sugars and energy ($P < 0.001$). For the experimental diet in which a fraction of the control diet was replaced by ground SC, DC of nutrients and energy were calculated on nutrients or energy intake basis; when compared to the control diet, the DC of DM,

OM and energy were significantly reduced (83.6% v. 87.5%, 85.3% v. 89.6% and 84.6% v. 88.7%, respectively; $P < 0.05$). The ME/DE ratio was not affected ($P > 0.05$) by inclusion of molasses or ground SC in the control diet; it averaged 96.0%. This ratio was significantly higher ($P < 0.01$) with SC juice treatment (i.e. 96.5%). According to their low N content, daily N intake in diets with SC products was lower than with the control diet (28.0 v. 38.7 g/day; $P < 0.01$). Whatever the dietary treatment, < 50% of N intake was retained.

Estimated DC of nutrients and energy of SC products using the difference method are presented in Table 3. The DC of OM and energy of molasses averaged 84.4% and 83.6%, respectively. The corresponding values for SC juice were equal to 99.3% and 99.0%, respectively. The DC calculated for ground SC offered or consumed differed significantly with lower values for ground SC offered. On average, DE values of the control diet, molasses, SC juice and ground SC intake and offered were 16.55, 13.11, 17.34, and 12.31 and 5.72 MJ/kg DM, respectively.

Effect of gradual increase of SC juice on total tract digestibility of energy and nutrients (trial 2)

The effects of a gradual increase in the rate of incorporation of SC juice (trial 2) on total tract digestibility of nutrients and energy are presented in Table 4. There was no statistical difference in average BW ($P > 0.05$) between diets. SC juice incorporation in diet increased from 0 to 66 g/100 DM between diets 1 and 4, respectively. The response of SC juice incorporation rate on DC was linear and the quadratic effect was not significant ($P > 0.05$). Despite an important numerical increase (33.6% to 42.1% between diets 1 and 4), faeces DM was not affected by the level of SC juice. The gradual decrease of N retained from diet 1 to 4 was related to the dilution of the N source from control diet with the progressive inclusion of SC juice. Except for CP and EE, the gradual inclusion rate of SC juice from diet 1 to diet 4 increased the DC of all components ($P < 0.001$). The DC of energy increased from 88.9% to 95.4%. In contrast, the DC of EE significantly decreased ($P < 0.01$) from 88.9% to 80.5% between diets 1 and 4. The ME/DE ratio increased from 95.9% to 97.6%, respectively, between diets 1 and 4.

According to the regression analysis of the DC of the nutrients or energy with the rate of SC juice inclusion, each 1% increase of SC juice inclusion rate resulted in a linear increase of 0.10% of the DC of OM and energy (Table 5). In contrast, the digestibility of CP was not affected by SC juice levels in diet. From these relationships, DC of nutrient and energy of SC juice can be calculated; they averaged 99.8%, 85.4% and 99.1% for OM, CP and energy, respectively. These values are similar to the values obtained in trial 1, (i.e. 99.0% for energy). Each 1% increase of SC juice inclusion rate resulted in a linear increase of 0.007 and 0.011 MJ/kg DM of DE and ME, respectively; average DE and ME values of SC juice obtained by regressions were then 17.32 and 17.08 MJ/kg DM, respectively.

Table 2 Effect of feeding diets containing SC products on digestibility of nutrients, ME and N retention in CR pigs (trial 1; least square means)¹

Item	Control	Control + SC molasses	Control + SC juice	Control + ground SC	r.s.d.	Statistical analyses ²
Number of pigs	6	5	6	5	–	–
Average BW (kg)	35.5	32.2	34.6	32.1	4.3	B**
DM offered (g/day per kg BW)						
Control	34.3	26.1	26.5	26.3	–	–
Sugar cane products	0	7.7	7.7	10.7	–	–
Total	34.3	33.9	34.3	37.0	–	–
Incorporation of SC products (%) ³	0	22.9	22.5	29.0	–	–
DM refused (g/day per kg BW) ⁴						
Sugar cane products	0	0.4	1.1	5.5	1.4	D***, R*, B*
Total DM intake (g/day per kg BW)	34.3	33.5	33.1	31.5	1.4	B*
Faeces DM (%)	28.5 ^{ab}	34.8 ^a	33.1 ^a	25.6 ^b	4.3	D*, D × R*
Digestibility coefficient (%) ⁵						
DM	87.5 ^a	87.0 ^a	90.6 ^b	83.6 ^c	0.4	D***, R**, D × R***
Organic matter	89.6 ^a	88.6 ^a	92.2 ^b	85.3 ^c	1.5	D***, R**
CP	86.4 ^a	82.7 ^b	86.1 ^a	84.1 ^a	2.0	D*, R***
Ether extract	82.1 ^a	88.3 ^b	83.8 ^a	84.7 ^a	1.7	D**, R***, B*, D × R**
Free sugars	96.1 ^a	97.6 ^b	99.1 ^c	97.8 ^b	0.2	D***, B*, D × R**
Energy	88.7 ^{ab}	87.7 ^{ac}	91.3 ^b	84.6 ^c	1.6	D***, R**
DE (MJ/kg DM)	16.55 ^a	15.77 ^b	16.82 ^a	15.63 ^b	0.28	D***, R**
ME (MJ/kg DM)	15.87 ^a	15.13 ^b	16.23 ^a	14.99 ^b	0.30	D***, R**
ME/DE (%)	95.9 ^a	96.0 ^a	96.5 ^b	95.9 ^a	0.3	D**, B**
N balance (g/day)						
N intake	38.7 ^a	28.4 ^b	28.8 ^b	26.9 ^b	4.6	D**, B*
N losses						
In faeces	5.3 ^a	4.9 ^a	4.0 ^b	4.2 ^a	0.5	D**, R***, B**, D × R*
In urine	15.4 ^a	10.6 ^b	10.2 ^b	9.3 ^b	2.1	D**, B*
N retention	18.1 ^a	12.7 ^b	14.2 ^{ab}	14.5 ^{ab}	2.7	D*, B*

SC = sugarcane; r.s.d. = residual standard deviation; ME = metabolisable energy; CR = Creole; DM = dry matter; DE = digestible energy; D = diet; R = replicate; B = block within replicate; D × R = diet × replicate; NS = not significant.

¹Diets were prepared by mixing molasses, SC juice or ground SC with the control diet.

²Levels of significance: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Within a line, least square means with different superscripts differed ($P < 0.05$) according to the dietary treatment.

³Expressed as percentage of total DM allowed.

⁴No control diet was refused whatever the dietary treatment.

⁵Digestibility coefficients of nutrients = digested nutrients or energy expressed as a percentage of nutrient or energy intake.

Table 3 Estimated apparent digestibility coefficients of SC products according to the difference method

Item	Control	SC molasses ¹	SC juice ¹	Ground SC offered ¹	Ground SC intake ²
Digestibility coefficient (%) ³					
Organic matter	89.6	84.4	99.3	30.4	68.6
CP	86.4	68.1	82.9	70.3	73.7
Energy	88.7	83.6	99.0	31.9	68.6
DE (MJ/kg DM)	16.55	13.11	17.34	5.72	12.31
ME (MJ/kg DM)	15.87	12.56	17.22	5.56	11.82

SC = sugarcane; DE = digestible energy; ME = metabolisable energy.

¹Values calculated according to the difference method: digestibility coefficients (DC) of energy and organic matter and CP of SC products were calculated from the rate of incorporation in feed and the DC of control and mixed (control + SC products) diets and expressed as a percentage of nutrient or energy offered.

²Values calculated according to the difference method and expressed as a percentage of nutrient or energy intake.

³DC of free sugars was found 100% whatever the SC product considered.

Effect of gradual increase of ground SC on total tract digestibility of energy and nutrients (trial 3)

The effects of a gradual increasing rate of incorporation of ground SC (trial 3) on nutrient intake and extraction rate are presented in Table 6. Average BW was not significantly

affected ($P > 0.05$) by ground SC incorporation level. The total DM intake was significantly reduced between diets 1 and 4 in connection with the increase of ground SC refusals. Expressed in g/100 g ground SC intake, the nutrient extraction rate was not influenced by the increase of

Table 4 Effect of feeding diets including various levels of SC juice on digestibility of nutrients, ME and N retention in CR pigs (trial 2; least square means)¹

Item	SC juice inclusion (%) ²				r.s.d.	Statistical analyses ³
	0	21	43	66		
Number of pigs	6	6	6	6	—	—
Average BW (kg)	47.7	44.5	41.7	39.7	6.2	B*
DM offered (g/day per kg BW)						
Control	35.1	26.9	18.8	10.9	—	—
SC juice	0	7.3	14.4	21.5	—	—
Total	35.1	34.2	33.2	32.4	—	—
Total DM intake (g/day per kg BW) ¹	34.9 ^a	34.1 ^b	33.2 ^c	32.4 ^d	0.4	D***, R***, D×R***
Faeces DM (%)	33.6	36.9	41.7	42.1	5.2	NS
Digestibility coefficient (%) ¹						
DM	87.4 ^a	90.4 ^b	93.1 ^c	95.4 ^d	0.8	D***, R*
Organic matter	89.5 ^a	92.2 ^b	94.4 ^c	96.1 ^d	0.8	D***, R*
CP	88.1	87.6	88.1	85.5	2.0	NS
Ether extract	88.9 ^a	88.5 ^a	85.9 ^{ab}	80.5 ^b	3.1	D**, R***
Free sugars	96.7 ^a	98.9 ^b	99.5 ^c	99.7 ^d	0.1	D***, R***, D×R***
Energy	88.9 ^a	91.4 ^b	93.6 ^c	95.4 ^d	0.9	D***
DE (MJ/kg DM)	16.58 ^a	16.82 ^{ab}	16.98 ^b	17.05 ^b	0.15	D**
ME (MJ/kg DM)	15.91 ^a	16.27 ^b	16.53 ^c	16.63 ^d	0.18	D**
ME/DE (%)	95.9 ^a	96.7 ^b	97.3 ^c	97.6 ^c	0.3	D***, R**, B*
N balance (g/day)						
N intake	52.6 ^a	38.8 ^b	26.2 ^c	15.4 ^d	5.5	D***, B*
N losses						
In faeces	6.2 ^a	4.8 ^b	3.1 ^c	2.2 ^c	0.6	D***, R**, B*
In urine	24.6 ^a	15.8 ^b	8.8 ^c	5.4 ^c	3.3	D***
N retention	21.7 ^a	18.2 ^{ab}	14.4 ^b	7.8 ^c	3.2	D***

SC = sugarcane; r.s.d. = residual standard deviation; ME = metabolisable energy; CR = Creole; r.s.d. = residual standard deviation; DM = dry matter; DE = digestible energy; D = diet; R = replicate; B = block within replicate; D × R = diet × replicate; NS = not significant.

¹Diets were prepared by mixing molasses, SC juice or ground SC with the control diet.

²Diet 1 was 35 g DM/day per kg BW of control diet and diets 2, 3 and 4 were a mix of 8, 16 and 24 g DM/day per kg BW of SC juice with control diet, respectively.

³Levels of significance: **P* < 0.05, ***P* < 0.01, ****P* < 0.001. Within a line, least square means with different superscripts differed (*P* < 0.05) according to the dietary treatment.

Table 5 Relationship between incorporation rates of SC juice in the diet and digestibility coefficients of energy and nutrients (trial 2)

Y	Parameters ¹				r.s.d.	Adj. R ²	Digestive or energy value ²
	a	s.e. of a	b	s.e. of b			
Digestibility coefficient (%)							
Organic matter	89.8	0.2	0.10	0.01	0.7	0.96	99.8
CP	88.4	0.7	−0.03	0.02	1.9	0.13	85.4
Energy	89.1	0.3	0.10	0.01	0.8	0.92	99.1
Energy values (MJ/kg DM)							
DE	16.62	0.04	0.007	0.001	0.13	0.65	17.32
ME	15.96	0.07	0.011	0.001	0.19	0.71	17.08

SC = sugarcane; Y = digestibility coefficient values or energy value; a = digestibility coefficient or energy value of control diet; b = the linear variation of digestibility coefficient or energy value with the increase of SC juice; r.s.d. = residual standard deviation; DM = dry matter; DE = digestible energy; ME = metabolisable energy.

¹Nutritional and energy values calculated using the regression analysis with the following model: Y = a + b × (SC juice incorporation rate), with Y, a and b. Incorporation rate = SC juice intake/total intake (g DM/g DM).

²Calculated from the regression equation for a 100% incorporation rate of SC juice in the diet.

ground SC incorporation rate. As a consequence, the amount of daily nutrient or energy intake from ground SC linearly increased with the rate of ground SC incorporation.

As the chemical composition of ground SC refusals differed from that of ground SC allowance with an increase of

fibre content and a decrease of soluble free sugars, the DC was calculated on ground SC allowance basis or on actual ground SC intake. The increase of ground SC inclusion in the diet from 0% to 74% resulted in a linear decrease (*P* < 0.05) of faecal DM from 33.8% to 17.4% and in

Table 6 Effect of feeding diets including various levels of ground SC on intake and extraction rate of nutrients in CR pigs (trial 3; least square means)¹

Item	Ground SC inclusion (%) ²				r.s.d.	Statistical analyses ³
	0	28	52	74		
Number of pigs	6	6	6	6	–	–
Average BW (kg)	58.9	49.7	49.2	50.9	7.5	B*
DM offered (g/day per kg BW)						
Control	37.1	27.4	19.1	10.7	–	–
Ground SC	0	10.7	21.1	30.6	–	–
Total	37.1	38.1	40.3	41.2	–	–
Ground SC refused (g DM/day per kg BW)	0	6.9	13.7	19.4	2.3	D***
Total DM intake (g/day per kg BW) ¹	37.1 ^a	30.2 ^b	26.4 ^{bc}	21.8 ^c	3.8	D**
Nutrient extraction rate from ground SC (%)						
DM	–	36.0	35.4	36.6	11.2	
Organic matter	–	36.3	35.6	36.7	11.2	
Free sugars	–	61.8	58.2	59.0	16.4	
NDF	–	19.5	19.4	18.7	8.7	
Gross energy	–	35.3	34.7	35.8	11.1	

SC = sugarcane; CR = Creole; r.s.d. = residual standard deviation; DM = dry matter.

¹Diets were prepared by mixing molasses, SC juice or ground SC with the control diet.

²Diet 1 was 35 g DM/day per kg BW of control diet and diets 2, 3 and 4 were a mix of 13, 26 and 39 g DM/day per kg BW of ground SC with control diet, respectively.

³Levels of significance: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Within a line, least square means with different superscripts differed ($P < 0.05$) according to the dietary treatment.

a linear reduction ($P < 0.05$) of DM, OM and energy digestibility coefficients (Table 7). In contrast, with a progressive inclusion of ground SC, the free sugars DC increased when expressed on an intake basis (96.4% v. 98.7%; $P < 0.001$). The regression analysis of the nutrients DC with incorporation rate of ground SC in the diet is presented in Table 8. Whatever the expression of the DC, a linear decrease of the DC was shown for OM, CP and energy. Each 1% increase of ground SC incorporation rate resulted in a 0.12% and 0.13% linear decrease of the DC of OM and energy, respectively, when expressed as a percentage of energy intake (method 1). The average DC of energy, DE and ME values were lower when method 2 was used instead of method 1 (28.2% v. 77.6%, 4.90 v. 13.58 MJ/kg DM and 4.74 v. 13.15 MJ/kg DM, respectively).

Discussion

Nutritional and energy values of SC juice

The chemical composition of SC juice determined in this study is similar to the values published in Brazilian tables (Rostagno, 2005). Total soluble sugars represented the main part of the identified OM (i.e. 83%). In these sugars, sucrose is the predominant carbohydrate (about 90%) and fructose and glucose represent less than 10% (Meade and Chen, 1977). The non-sugar OM in SC juice (i.e. 17.0%) is represented by chlorophyll, tannins, wax and fibre (Meade and Chen, 1977). In our experiment, these non-sugars OM content in SC juice was about 16.8 g/100 g DM.

Our results show that OM and energy digestibility of SC juice is close to 100% according to its high content of water soluble sugars, which are known to be entirely digestible

(Noblet *et al.*, 2003). Logically, total tract digestibility of energy of the diet is linearly enhanced as more as the inclusion rate of SC juice in the diet is increased. This result also suggests that SC juice does not interact with the control corn-soybean meal diet. However, our DE and ME values for SC juice (17.3 and 17.1 MJ/kg DM) are higher than those calculated in the Brazilian tables (15.9 and 15.2 MJ/kg DM; Rostagno, 2005) despite a GE value similar to ours (17.1 v. 17.5 MJ/kg DM). In fact, in Brazilian tables, DE and ME values were not measured but estimated from the chemical composition of SC juice. From that, it can be suggested that equations used to estimate energy values of SC juice in those tables underestimate the actual values. Our results also show a higher DC of OM for the SC juice than in the Brazilian tables (99.5% v. 93.4%; Rostagno, 2005).

Nutritional and energy values of final SC molasses

The chemical composition of molasses is closely dependant on the technological process applied for the extraction of raw sugar. Bayley *et al.* (1983) showed that the range for ash and soluble sugars contents in Cuban commercial conditions was from 5.7 to 16.3 g/100 DM and from 46 to 76 g/100 DM, respectively. In our study, the SC molasses contained 15.1 g/100 g DM of ash and 71.0 g/100 g DM of soluble sugars (sucrose, fructose and glucose). During the crystallisation processes, non-sugars OM were concentrated in molasses and represented more than 13% of the total OM. The DC of OM and energy are lower in molasses than in SC juice (–15%). This result can be related to the high ash content that increases endogenous energy losses (Noblet and Shi, 1994) and the low digestibility of the non-sugars OM in molasses. Comparing the digestive utilisation

Table 7 Effect of feeding diets including various levels of ground SC on digestibility of nutrients, ME and N retention in CR pigs (trial 3; least square means)¹

Item	Ground SC inclusion (%) ²				r.s.d.	Statistical analyses ³
	0	28	52	74		
Number of pigs	6	6	6	6	—	—
Faeces DM (%)	33.8 ^a	27.7 ^{ab}	26.0 ^b	17.4 ^c	4.2	D***
Digestibility coefficient (%) ⁴						
DM	88.3 ^a	87.0 ^{ab}	86.6 ^{ab}	83.1 ^b	2.4	D*
Organic matter	90.5 ^a	88.6 ^a	88.0 ^{ab}	84.5 ^b	2.3	D**
CP	89.9 ^a	88.5 ^a	87.9 ^a	82.4 ^b	2.6	D**
Free sugars	96.4 ^a	97.9 ^b	98.7 ^c	98.7 ^c	0.5	D***, R***, D × R**
Energy	90.1 ^a	88.1 ^a	87.4 ^{ab}	83.5 ^b	2.4	D**
Digestibility coefficient (%) ⁵						
DM	88.3 ^a	68.8 ^b	56.8 ^c	44.0 ^d	6.8	D***
Organic matter	90.5 ^a	69.7 ^b	57.2 ^c	44.4 ^d	7.0	D***
CP	89.9 ^a	82.3 ^b	81.6 ^b	71.3 ^c	5.9	D*
Free sugars	96.4 ^a	70.3 ^b	62.5 ^b	60.5 ^b	12.8	D*
Energy	90.1 ^a	70.0 ^b	57.6 ^c	44.2 ^d	6.8	D***
DE (MJ/kg DM) ³	16.81 ^a	16.32 ^a	16.05 ^a	15.13 ^b	0.45	D***
DE (MJ/kg DM) ⁴	16.81 ^a	12.90 ^b	10.53 ^c	8.01 ^d	1.27	D***
ME (MJ/kg DM) ³	16.21 ^a	15.72 ^{ab}	15.39 ^{bc}	14.71 ^c	0.46	D***
ME (MJ/kg DM) ⁴	16.21 ^a	12.43 ^b	10.15 ^c	7.74 ^d	1.21	D***
ME/DE (%)	96.5	96.4	96.4	96.7	0.4	D × R*

SC = sugarcane; r.s.d. = residual standard deviation; ME = metabolisable energy; CR = Creole; DM = dry matter; DE = digestible energy; D = diet; R = replicate; D × R = diet × replicate.

¹Diets were prepared by mixing molasses, SC juice or ground SC with the control diet.

²Diet 1 was 35 g DM/day per kg BW of control diet and diets 2, 3 and 4 were a mix of 13, 26 and 39 g DM/day per kg BW of ground SC with control diet, respectively.

³Levels of significance: **P* < 0.05, ***P* < 0.01, ****P* < 0.001. Within a line, least square means with different superscripts differed (*P* < 0.05) according to the dietary treatment.

⁴Digestibility coefficient values expressed as percentage of nutrient or energy intake.

⁵Digestibility coefficient values expressed as percentage of nutrient or energy offered.

Table 8 Relationship between incorporation rates of allowed ground SC in the diet and digestibility coefficients of energy and nutrients

Y	Parameters				r.s.d.	Adj. R ²	Digestibility or energy value ³
	a	s.e. of a	b	s.e. of b			
Digestibility coefficient (%) ¹							
Organic matter	90.6	0.6	−0.12	0.02	2.0	0.59	79.0
CP	90.4	0.8	−0.14	0.03	2.4	0.59	76.1
Energy	90.1	0.6	−0.13	0.02	2.0	0.61	77.6
Digestibility coefficient (%) ²							
Organic matter	89.2	1.5	−0.61	0.03	4.2	0.95	27.9
CP	90.0	1.2	−0.23	0.03	3.5	0.77	67.3
Energy	89.0	1.4	−0.61	0.03	3.9	0.95	28.2
Energy value (MJ/kg DM)							
DE ¹	16.81	0.12	−0.032	0.004	0.38	0.75	13.58
DE ²	16.58	0.25	−0.117	0.005	0.72	0.96	4.90
ME ¹	16.20	0.13	−0.031	0.004	0.40	0.71	13.15
ME ²	15.99	0.25	−0.113	0.005	0.71	0.95	4.74

SC = sugarcane; Y = digestibility coefficient values of nutrients or energy expressed as a percentage of nutrient or energy intake; a = digestibility coefficient of control diet; b = the linear variation of digestibility coefficient with the increase of ground SC incorporation rate; DM = dry matter; DE = digestible energy; ME = metabolisable energy.

¹Parameters estimated from the following model: Y = a + b × (ground SC incorporation rate), with Y, a and b. Incorporation rate = ground SC intake/total intake (g DM/g DM).

²Parameters estimated from the following model: Y = a + b × (ground SC incorporation rate), with Y, a and b. Incorporation rate = ground SC allowance/total allowance (g DM/g DM).

³Calculated from the regression equation for a 100% incorporation rate of ground SC in the diet.

of SC molasses with a mixture of soluble sugars representative of sugars composition of molasses, Bayley *et al.* (1983) showed a reduced DC of energy with SC molasses. In fact, Figueroa *et al.* (1990) showed that the non-identified OM in SC molasses is poorly digested, which considerably reduces the dietary DE density of this material. In this study, the DC for N was higher than the value reported in the INRA tables (68.1% *v.* 40.0%; Sauvant *et al.*, 2002). However, these values should be used with caution according to the low accuracy of the N digestibility due to both low N content in molasses and possible effects of molasses products (minerals) on digestion and transit (Brooks, 1967; Christon and Le Dividich, 1978). The mechanism by which protein digestibility was depressed by molasses is not fully understood. Christon and Le Dividich (1978) reported that the depressive effect of molasses on N retention in rat is mainly related to an excretion of endogenous N in faeces. They concluded that this high endogenous N losses could be explained an abrasive effect of ash and minerals like potassium (K).

The energy DC (83.6%) is similar to the value reported in the French tables (85%; Sauvant *et al.*, 2002). The DE values of molasses (13.1 MJ/kg DM) are in accordance with Bayley *et al.* (1983) study (13.6 MJ/kg DM) and with French (Sauvant *et al.*, 2002; 12.7 MJ/kg DM) and Brazilian (Rostagno, 2005; 13.5 MJ/kg DM) feeding tables.

Nutritional and energy values of ground SC stalks

The DM of faeces was reduced when inclusion of ground SC increased. Similarly, Henry and Etienne (1969) reported lower faeces DM when increasing levels of pure cellulose were included in growing pig's diet. According to these latter authors, the increase of faecal DM is directly related to the high water holding capacity of fibre.

To our knowledge, no study has been published on the digestive utilisation of energy and nutrients of SC ground stalks in growing pigs. Our results show that DC and energy values of ground SC vary with the method used for their calculations (intake *v.* allowance basis). In fact, only one part of ground SC allowed was effectively consumed because the pigs chew the cane stalks to extract juice and spit out most of the fibrous residue (Mederos *et al.*, 2004). This peculiar feeding behaviour could be considered as an adaptation to increase the energy extraction rate in ground SC. In fact, with this feeding behaviour, pigs avoid very bulky intakes and subsequent low energy intakes. As a result, the chemical composition of the actual ground SC intake widely differed from that of ground SC allowed: it contained a lower fibre and higher free sugar contents. From that, two expressions of nutrients and energy DC were needed to characterise the nutritional value of ground SC (intake *v.* allowance basis). In our work, the chemical composition of ground SC intake was maintained constant at a level of about 60% whatever its incorporation rate. This suggests that the feeding behaviour remains comparable whatever the level of proposed ground SC. In contrast, Bravo *et al.* (1996) showed that sugar extraction rate decreased from 85.4% to 68.8% when ground SC allow-

ance increased from 17 to 51 g DM/day per kg BW. In this latter study, SC particle size averaged 3 cm. According to Mederos *et al.* (2004), the juice extraction rate was more efficient in 3 cm sized ground SC particles than in milled SC. Thus, it can be assumed that the small size of ground SC particles used in the present experiment could have limited the ability of the pigs to extract juice from ground SC. From a practical point of view, this result suggests that grinding conditions and/or particle size could influence the sugar extraction rate and then the energy value of ground SC. However, other criteria such as the SC variety (hard *v.* smooth stalk), the pigs BW and/or breed could also affect the sugar extraction rate. According to Bravo *et al.* (1996), the efficiency of sugar extraction by the pigs appeared to be higher than when the juice was extracted mechanically with a 3 roll-mill (Vencedora Magtron, Joaçaba, SC 89600-000, Brazil). All in all, in a small scale farming system context, we have to take into account this point for the global efficiency of the production system.

From our observations, the coefficient of variation for the extraction rate of nutrients and energy from ground SC within each dietary treatment was high (> 30%) suggesting important between-animal variability. In addition, energy digestibility was negatively correlated with NDF extraction rate ($r = -0.42$; $P = 0.08$) suggesting that a large between-pig variability exists in the ability to suck and spit ground SC and in the ability to ingest dietary fibre from the ground SC. In consequence, the digestive utilisation of nutrients and energy from ground SC was closely dependant on the inter-individual feeding behaviour variations.

In comparison with the other SC products, the energy DC of ground SC was lower mainly because of its high dietary fibre content. The reduced digestibility of high fibre diets is explained by low digestibility of the fibre fraction and possible negative interactions with the other nutrients (Le Goff *et al.*, 2002). From the latter authors, each extra 1% increase in NDF content decreases by one percentage point the DC of energy in growing pigs. From that, it can be suggested that the reduced energy DC related to the inclusion of ground SC in the experimental diet could be related to its high consecutive fibre content. In addition, the negative effect of dietary fibre on energy DC was emphasised by the high ADF content of ground SC.

In our experimental conditions, the actual DM and energy intake from ground SC represented less than 40% and 35% of DM and energy allowance, respectively. First, it can be suggested that time dedicated for the sugar extraction chewing would limit the ability of the pigs to increase their feeding level (Xandé *et al.*, 2009). Second, in connection to their high water holding capacity, dietary fibre increases gut fill and satiety. From that, it can be also suggested that the low DM and energy intake from ground SC was related to the high dietary fibre content. According to the net energy (NE) system, the metabolic utilisation of high fibre diets is accompanied by an increase of heat expenditure and a decrease of retained energy (Noblet *et al.*, 1994). From that, it can be expected that the NE/ME ratio would be

much lower for ground SC than for juice in connection with a high-energy expenditure connected to chewing activity. Then, the low growth performance obtained in our previous study with pigs fed ground SC (i.e. 200 g/day; Xandé *et al.*, 2009) is mainly related to the fact that a small part of energy allowed in ground SC (about 25%) was available for growth purposes.

Acknowledgements

This work was supported by the Guadeloupe Region and the European funds (FSE FEOGA). The authors gratefully acknowledge C. Anais, K. Benony, M. Bructer, E. M. Giorgi, K. A. Racon and F. Silou for their technical assistance, Y. Jaguelin, M. Baucal, S. Calif, T. Etienne and G. Saminadin for the laboratory analysis.

References

- Association of Official Analytical Chemists 1990. Official methods of analysis, 15th edition. AOAC, Arlington, VA, USA.
- Bayley HS, Figueroa V, Ly J, Maylin A and Perez A 1983. Utilisation of sugarcane final molasses by the pig: energy metabolism. *Canadian Journal of Animal Science* 63, 455–462.
- Bravo M, Lasso M, Esnaola MA and Preston TR 1996. Preliminary studies on the use of chopped sugarcane stalks as the basal diet for fattening pigs. *Livestock Research for Rural Development* 8. Retrieved from <http://www.cipav.org.co/lrrd/lrrd8/3/bravo83.htm>
- Brooks CC 1967. Effect of sex, fat, fibre, molasses and thyroprotein on digestibility of nutrients and performance of growing swine. *Journal of Animal Science* 26, 495–499.
- Christon R and Le Dividich J 1978. Utilisation de la mélasse de canne à sucre dans l'alimentation du porc: essai d'interprétation des acquisitions récentes. *Annales de Zootechnie* 27, 267–288.
- Figueroa V, Maylin A, Ly J, Pérez A, Carrillo O and Bayley HS 1990. Ileal and total digestibility studies in pigs fed molasses type A or starch diets supplemented with torula yeast or soybean meal. *Livestock Production Science* 25, 151–161.
- González D, González C, Ojeda A, Machado W and Ly J 2006. Growth performance of pigs feed sugarcane juice (*Saccharum officinarum*) and mulberry leaf meal (*Morus alba*). *Archivos Latinoamericanos de Producción Animal* 14, 42–48.
- Henry Y and Etienne M 1969. Effets nutritionnels de l'incorporation de cellulose purifiée dans le régime du porc en croissance-finition. 1- Influence sur l'utilisation digestive des nutriments. *Annales de Zootechnie* 18, 337–357.
- Le Goff G and Noblet J 2001. Comparative total tract digestibility of dietary energy and nutrients in growing pigs and adult sows. *Journal of Animal Science* 79, 2418–2427.
- Le Goff G, van Milgen J and Noblet J 2002. Influence of dietary fibre on digestive utilisation and rate of passage in growing pigs, finishing pigs and adult sows. *Animal Science* 74, 503–515.
- Ly MVJ and Preston TR 1969. Digestible and metabolisable energy values for pigs of diets based on high-test molasses or final molasses and sugar. *Journal of Animal Science* 29, 578–580.
- Meade GP and Chen JCP 1977. Cane sugar handbook; a manual for cane sugar manufacturers and their chemists, 10th edition. Wiley-Interscience, New York, NY, USA.
- Mederos CM, Figueroa V, Garcia A, Aleman E, Martinez RM and Ly J 2004. Growth performance of pigs fed hand-chopped sugarcane stalks. *Livestock Research for Rural Development* 16. Retrieved from <http://www.cipav.org.co/lrrd/lrrd16/3/med16014.htm>
- Noblet J and Shi XS 1994. Effect of body weight on digestive utilisation of energy and nutrients of ingredients and diets in pigs. *Livestock Production Science* 72, 323–338.
- Noblet J, Bontems V and Tran G 2003. Estimation de la valeur énergétique des aliments chez le porc. *INRA Productions Animales* 16, 197–210.
- Noblet J, Shi XS, Fortune H, Dubois S, Lechevestrier Y, Corniaux C, Sauvart D and Henry Y 1994. Teneur en énergie nette des aliments chez le porc. Mesure, prédiction et validation aux différents stades de sa vie. *Journées de la Recherche Porcine en France* 26, 235–250.
- Preston TR 1995. Research, extension and training for sustainable farming systems in the tropics. *Livestock Research for Rural Development* 7. Retrieved from <http://www.cipav.org.co/lrrd/lrrd7/2/1.htm>
- Preston TR and Leng RA 1989. The greenhouse effect and its implications for world agriculture. The need for environmentally friendly development. *Livestock research for rural development*. 1. Retrieved from <http://www.cipav.org.co/lrrd/lrrd1/1/preston.htm>
- Preston TR, MacLeod NA, Lassota L, Willis MB and Velazquez M 1968. Sugar cane products as energy sources for pigs. *Nature* 219, 727–730.
- Renaudeau D, Bocage B and Noblet J 2006. Influence of energy intake on protein and lipid deposition in Creole and Large White growing pigs in a humid tropical climate. *Animal Science* 82, 937–945.
- Rostagno HS 2005. Brazilian tables for poultry and swine: composition of feedstuffs and nutritional requirements, 2th edition. Universidade Federal de Viçosa. Departamento de zootecnia, Viçosa, Minas Gerais, Brazil, 186p.
- SAS Institute Inc. 1997. SAS/STAT user's guide: version 7, 4th edition. SAS Institute Inc., Cary, NC, USA.
- Sauvart D, Perez JM and Tran G 2002. Tables de composition et de valeur nutritive des matières premières destinées aux animaux d'élevage. AFZ-INRA, pp. 1–301.
- Tollier MTh and Robin JP 1979. Adaptation de la méthode à l'orcino sulfurique au dosage automatique des glucides neutres totaux: conditions d'application aux extraits d'origine végétale. *Annales de Technologie Agricole* 28, 1–15.
- Van Soest PJ and Wine RH 1967. Use of detergents in the analysis of fibrous feeds. IV. Determination of plant cell-wall constituents. *Journal of the AOAC* 50, 50–55.
- Xandé X, Despois E, Giorgi M, Gouridine JL, Archimède H and Renaudeau D 2009. Influence of sugarcane diets and a high fibre commercial diet on growth and carcass performance in local Caribbean pigs. *Asian-Australasian Journal of Animal Sciences* 22, 90–98.